

# 9-9.6-17

EE24BTECH11064 - Harshil Rathan

## Question:

Find the equation of a curve passing through the point (0, 2) given that the sum of the coordinates of any point on the curve exceeds the magnitude of the slope of the tangent to the curve at that point by 5.

## Solution:

Let us solve the given equation theoretically and then verify the solution computationally According to the question,

Sum of the coordinates of any point say  $(x, y)$  on the curve = Magnitude of the slope of the tangent to the curve + 5

$$x + y = \frac{dy}{dx} + 5 \quad (0.1)$$

$$\frac{dy}{dx} - y = x - 5 \quad (0.2)$$

Comparing 0.2 with  $\frac{dy}{dx} + Py = Q$

$$P = -1 \text{ and } Q = x - 5 \quad (0.3)$$

I.F is 0.4

$$e^{\int P dx} = e^{-x} \quad (0.4)$$

Solution is 0.6

$$y (I.F) = \int Q (I.F) dx + c \quad (0.5)$$

$$y \cdot e^{-x} = \int (x - 5) e^{-x} dx + c \quad (0.6)$$

Apply product rule :

$$\int A \cdot B dx = A \int B dx - \int \frac{d}{dx}(A) \left( \int B dx \right) dx \quad (0.7)$$

$$ye^{-x} = -(x - 5)e^{-x} + \int e^{-x} dx + c \quad (0.8)$$

$$ye^{-x} = -(x - 5)e^{-x} + \frac{e^{-x}}{-1} + c \quad (0.9)$$

$$\frac{y}{(e^x)} = -\frac{(x-5)}{(e^x)} - \frac{1}{e^x} + c \quad (0.10)$$

On Simplifying

$$y = -(x-5) - 1 + ce^x \quad (0.11)$$

$$x + y = 4 + ce^x \quad (0.12)$$

To find c, put x=0 and y=2 in 0.12

$$c = -2 \quad (0.13)$$

The curve is

$$x + y = 4 - 2e^x \quad (0.14)$$

Now lets verify the solution computationally from the definition of  $\frac{dy}{dx}$

$$y_{n+1} = y_n + \frac{dy}{dx} \cdot h \quad (0.15)$$

From the differential equation given,

$$\frac{dy}{dx} = x + y - 5 \quad (0.16)$$

Substituting 0.16 in 0.15

$$y_{n+1} = y_n + (x_n + y_n - 5) \cdot h \quad (0.17)$$

From the figure it is clearly verified that the theoretical solution matches with the computational solution

