

# 2021-February Session-02-24-2021-shift-1

ee24btech11064 - Harshil Rathan

1) Let

$$P = \begin{pmatrix} 3 & -1 & -2 \\ 2 & 0 & \alpha \\ 3 & -5 & 0 \end{pmatrix}$$

where  $\alpha \in R$ . Suppose  $Q = [q_{ij}]$  is a matrix satisfying  $PQ = kI_3$  for some non-zero  $k \in R$ . If  $q_{23} = \frac{-k}{8}$  and  $|Q| = \frac{k^2}{2}$ , then  $\alpha^2 + k^2$  is equal to

2) Let  $B_i (i = 1, 2, 3)$  be three independent events in a sample space. The probability that only  $B_1$  occur is  $\alpha$ , only  $B_2$  occurs is  $\beta$  and only  $B_3$  occurs is  $\gamma$ . Let  $p$  be the probability that none of the events  $B_i$  occurs and these 4 probabilities satisfy the equations  $(\alpha - 2\beta)p = \alpha\beta$  and  $(\beta - 3\gamma)p = 2\beta\gamma$ . All the probabilities are assumed to lie in the interval  $(0, 1)$ . Then  $\left(\frac{P(B_1)}{P(B_3)}\right)$  is equal to

3) The minimum value of  $\alpha$  for which the equation  $\frac{4}{\sin x} + \frac{1}{1 - \sin x} = \alpha$  has at least one solution in  $\left(0, \frac{\pi}{2}\right)$  is

4) If one of the diameters of the circle  $x^2 + y^2 - 2x - 6y + 6 = 0$  is a chord of another circle 'C' whose centre is at  $(2, 1)$ , then its radius is

5)  $\lim_{n \rightarrow \infty} \tan\left(\sum_{r=1}^n \tan^{-1}\left(\frac{1}{1+r+r^2}\right)\right)$  is equal to

6) If

$$\int_{-a}^a (|x| + |x - 2|) dx = 22$$

and  $[x]$  denotes the greatest integer  $\leq x$ , then  $\int_a^{-a} (x + [x]) dx$  is equal to

7) Let three vectors  $\mathbf{a}, \mathbf{b}$  and  $\mathbf{c}$  be such that  $\mathbf{c}$  is coplanar with  $\mathbf{a}$  and  $\mathbf{b}$ ,  $\mathbf{a} \cdot \mathbf{c} = 7$  and  $\mathbf{b}$  is perpendicular to  $\mathbf{c}$ , where  $\mathbf{a} = -\hat{i} + \hat{j} + \hat{k}$  and  $\mathbf{b} = 2\hat{i} + \hat{k}$ , then the value of  $2|\mathbf{a} + \mathbf{b} + \mathbf{c}|^2$  is

8) Let  $A = \{n \in N : n \text{ is a 3-digit number}\}$ ,  $B = \{9k + 2 : k \in N\}$  and  $C = \{9k + I : k \in N\}$  for some  $I$  ( $0 < I < 9$ ). If the sum of all the elements of the set  $A \cap (B \cup C)$  is  $274 \times 400$ , then  $I$  is equal to

9) If the least and the largest real values of  $\alpha$ , for which the equation  $z + \alpha|z - 1| + 2i = 0$  ( $z \in \mathbb{C}$  and  $i = \sqrt{-1}$ ) has a solution, are  $p$  and  $q$  respectively; then  $4(p^2 + q^2)$  is equal to

10) Let  $M$  be any  $3 \times 3$  matrix with entries from the set  $\{0, 1, 2\}$ . The maximum number of such matrices, for which the sum of diagonal elements of  $M^T M$  is seven, is