Lecture Notes on Numerical Weather Prediction

Harshit Prashant Dhanwalkar (SC21B164)1*

Abstract

Notes of Lectures and addional information from books.

¹MTech, Earth System Sciences (ESS), 1st year, Department of Physics, Indian Institute Of Spacescience and Technology (IIST) *email: harshitpd1729@gamil.com

Contents

1 Lecture 1 06/01/2025

3

1. Lecture 1 06/01/2025

Numerical Weathering Problem (NWP) was first proposed by Bjerkives around 1900. It is mathematical initial value problem (IVP).

Initial value Problem (IVP) \rightarrow simple pendulum.

$$\ddot{\theta} + \omega^2 \theta = 0 \tag{1}$$

$$\frac{d^2\theta}{dt^2} + \omega^2\theta = 0\tag{2}$$

$$\theta(t) = A\cos(\omega t) + B\sin(\omega t) \tag{3}$$

Eq.(1) and (2) are second order linear ordinary differential equation, whose solution Eq.(3) has 2 constants of integration A and B. Here θ and t are tge dependent and independent variable since Eq.(1) and (2) have only one independent variable.

Values of *A* and *B* will depend on initial condition.

Since ODE is second order, 2 initial condition are needed at initial time, say t = 0. Which are:

$$\frac{\theta(t=0)=1}{\theta(t=0)} = 0$$
(4)

Eq.(2) and initial conditions Eq.(4) are together called **Mathematical IVP**. For any physical system the following two requirements are needed:

- 1. The equation (ODE or PDE) that governs the evolution of the above system.
- 2. The initial state of the system.

7 independent variables (\mathbf{u} , \mathbf{v} , \mathbf{w} , \mathbf{T} , ρ , \mathbf{p} , \mathbf{q}).

Surface area of Earth = $4\pi R^2 = 4\pi (6.37 \times 10^{12}) \approx 5.1 \times 10^{14} \text{ m}^2$