

Lecture Notes on Numerical Weather Prediction

Harshit Prashant Dhanwalkar (SC21B164)^{1*}

Abstract

Notes of Lectures and additional information from books.

¹ MTech, Earth System Sciences (ESS), 1st year, Department of Physics, Indian Institute Of Space Science and Technology (IIST)

*email: harshitpd1729@gamil.com

Contents	
1	Lecture 1 06/01/2025 3
1.1	Introduction to Numerical Weather Prediction . . 3
2	Lecture 2 07/01/2025 4
3	Lecture 3 08/01/2025 5
4	Lecture 4 15/01/2025 6
4.1	Forward Difference 6
4.2	Backward Difference 6
4.3	Current Difference 6
5	Lecture 5 20/01/2025 8
5.1	Space Difference 8
5.2	Time Difference 8
5.3	Explicit Form Of Second Order PDE 8
5.4	Implicit Form Of Second Order PDE 8
6	Lecture 6 21/01/2025 10
6.1	1-D Linear Advection Equation 10
6.2	Triangular Property 10
7	Lecture 7 22/01/2025 11
8	Lecture 08/01/2025 12
8.1	Heat Equation 12
8.2	1-D Wave Equation 12
8.3	Laplace Equation 13
9	Lecture 9 28/01/2025 14
9.1	Wave Equation 14
9.2	Heat Transfer Equation 14

List of Figures	
1	Figure showing the grid 4
2	Path line or trajectory of fluid element 10
3	Triangular property distribution 10

1. Lecture 1 06/01/2025

1.1 Introduction to Numerical Weather Prediction

Numerical Weathering Problem (NWP) was first proposed by Bjerkvies around 1900. It is mathematical initial value problem (IVP).

Initial value Problem (IVP) → simple pendulum.

$$\ddot{\theta} + \omega^2 \theta = 0 \quad (1.1)$$

$$\frac{d^2 \theta}{dt^2} + \omega^2 \theta = 0 \quad (1.2)$$

$$\theta(t) = A \cos(\omega t) + B \sin(\omega t) \quad (1.3)$$

Eq.(1.1) and Eq.(1.2) are second order linear ordinary differential equation, whose solution .Eq.(1.3) has 2 constants of integration A and B . Here θ and t are the dependent and independent variable since Eq.(1.1) and Eq.(1.2) have only one independent variable.

Values of A and B will depend on initial condition.

Since ODE is second order, 2 initial condition are needed at initial time, say $t = 0$. Which are:

$$\left. \begin{aligned} \theta(t=0) &= 1 \\ \frac{\theta(t=0)}{dt} &= 0 \end{aligned} \right\} \quad (1.4)$$

Eq.(1.2) and initial conditions Eq.(1.4) are together called **Mathematical IVP**. For any physical system the following two requirements are needed:

1. The equation (ODE or PDE) that governs the evolution of the above system.
2. The initial state of the system.

7 independent variables ($\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{T}, \rho, \mathbf{p}, \mathbf{q}$).

Surface area of Earth = $4\pi R^2 = 4\pi(6.37 \times 10^{12}) \approx 5.1 \times 10^{14} \text{ m}^2$

2. Lecture 2 07/01/2025

7 independent variables (u, v, w, T, ρ, p, q) therefore we need 7 Governing equations (system of 7 coupled non-linear partial differential equations):

1. Conservation of mass (continuity equation).
2. Conservation of momentum in rotating frame of reference (3 scalar equations, one each corresponding to scalar component of velocity).
3. Conservation of energy (Thermodynamic energy equation).
4. Conservation of moisture (moisture continuity equation).
5. Equation of state (Ideal gas equation).

Euler description of fluid motion is more convenient because of dependence on time and above 7 equations.

Total advective and convective time of lagrangian is given by:

$$\underbrace{\frac{DT}{Dt}}_{\text{Lagrangian Derivative}} = \underbrace{\frac{\partial T}{\partial t}}_{\text{Local derivative}} + \underbrace{\vec{V} \cdot \nabla T}_{\text{Advective Term}} \quad (2.1)$$

$$\frac{DT}{Dt} = \frac{\partial T}{\partial t} + u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} + w \frac{\partial T}{\partial z} \quad (2.2)$$

Using first law of Thermodynamics, Rate of heat is given by:

$$\begin{aligned} d\dot{q} &= d\dot{u} + d\dot{w} \\ \frac{DU}{Dt} &= \frac{Dq}{Dt} - \frac{Dw}{Dt} \\ C_v \frac{DT}{Dt} &= \frac{Dq}{Dt} - p \frac{D\alpha}{Dt} \end{aligned}$$

where $\frac{Dq}{Dt}$ is rate at which heating of air parcel due to non-adiabatic process, this change can happen via radiation, convection, conduction, latent heat while phase change.

$$\frac{DU}{Dt} = \vec{F}_{\text{net}} + \vec{F}_{\text{coriolis}} \quad (2.3)$$

This above Eq.(2.3) is convective derivative equation involving non-linear terms (i.e. $u \frac{\partial T}{\partial x}, v \frac{\partial T}{\partial y}, w \frac{\partial T}{\partial z}$).

Continuity equation:

$$\frac{1}{\rho} \frac{D\rho}{Dt} + \nabla \cdot \vec{V} = 0 \quad (2.4)$$

Consider grid on globe as shown in figure(1).

Let grid of following resolutions:

- $1^\circ \times 1^\circ \rightarrow 3 \times 10^6$ grid cells \therefore no. of variables $\rightarrow 7 \times 3 \times 10^6$.
- $5^\circ \times 5^\circ \rightarrow 1.3 \times 10^5$ grid cells \therefore no. of variables $\rightarrow 7 \times 1.3 \times 10^5$.
- $20^\circ \times 20^\circ \rightarrow 9 \times 10^3$ grid cells \therefore no. of variables $\rightarrow 7 \times 9 \times 10^3$.

- $25^\circ \times 25^\circ \rightarrow 6 \times 10^3$ grid cells \therefore no. of variables $\rightarrow 7 \times 6 \times 10^3$.

These are even larger than entire country, which means that we can't hope to find the change of variables with these grids. This we don't have a way to determine initial condition, if we try to use interpolation, it will cause errors which will grow with time since atmosphere is chaotic and dynamic system.



Figure 1. Figure showing the grid

3. Lecture 3 08/01/2025

$$u = \bar{u} + u' \quad (3.1)$$

Here, u is the velocity field, which is decomposed into a mean component \bar{u} and a fluctuating component u' .

Navier-Stokes Equation The general Navier-Stokes equation is given by:

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} - f v = \frac{1}{\rho} \frac{\partial \bar{P}}{\partial x} + \gamma \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right) \quad (3.2)$$

Reynolds-Averaged Navier-Stokes (RANS) Equation Applying Reynolds decomposition ($u = \bar{u} + u'$) and averaging leads to the RANS equation:

$$\underbrace{\frac{\partial \bar{u}}{\partial t}}_{\text{Local acceleration}} + \underbrace{\bar{u} \frac{\partial \bar{u}}{\partial x} + \bar{v} \frac{\partial \bar{u}}{\partial y} + \bar{w} \frac{\partial \bar{u}}{\partial z}}_{\text{Advection}} - \underbrace{f \bar{v}}_{\text{Coriolis force}} = \underbrace{\frac{1}{\rho} \frac{\partial \bar{P}}{\partial x}}_{\text{Pressure gradient force}} + \underbrace{\gamma \left(\frac{\partial^2 \bar{u}}{\partial x^2} + \frac{\partial^2 \bar{u}}{\partial y^2} + \frac{\partial^2 \bar{u}}{\partial z^2} \right)}_{\text{Viscous dissipation}} + \underbrace{\frac{1}{\rho} \left(\frac{\partial (-\rho \bar{u}'u')}{\partial x} + \frac{\partial (-\rho \bar{u}'v')}{\partial y} + \frac{\partial (-\rho \bar{u}'w')}{\partial z} \right)}_{\text{Reynolds stress tensor}} \quad (3.3)$$

The Reynolds stress tensor represents the transport of momentum due to turbulent fluctuations.

Nonlinear Term Expansion Expanding the nonlinear term $u \frac{\partial u}{\partial x}$ using Reynolds decomposition:

$$\begin{aligned} u \frac{\partial u}{\partial x} &= (\bar{u} + u') \frac{\partial (\bar{u} + u')}{\partial x} \\ &= \bar{u} \frac{\partial \bar{u}}{\partial x} + u' \frac{\partial \bar{u}}{\partial x} + \bar{u} \frac{\partial u'}{\partial x} + u' \frac{\partial u'}{\partial x} \end{aligned}$$

Applying Reynolds averaging rules:

$$\begin{aligned} \bar{u} &= \overline{\bar{u} + u'} \\ \bar{u} &= \bar{\bar{u}} + \bar{u'} \\ \bar{u} &= \bar{u} + \bar{u'} \Rightarrow \bar{u'} = 0. \end{aligned}$$

Thus, the fluctuating component u' averages out to zero over time, leaving only the mean component \bar{u} in the averaged equations.

We have,

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0 \quad (3.4)$$

Substituting u , v and w in above Eq.(3.4), we get:

$$\begin{aligned} \frac{\partial (\bar{u} + u')}{\partial x} + \frac{\partial (\bar{v} + v')}{\partial y} + \frac{\partial (\bar{w} + w')}{\partial z} &= 0 \\ \frac{\partial \bar{u}}{\partial x} + \frac{\partial \bar{v}}{\partial y} + \frac{\partial \bar{w}}{\partial z} &= 0 \end{aligned} \quad (3.5)$$

The term $\frac{\partial (\overline{u'u'})}{\partial t}$ represents the rate of change of kinetic energy per unit mass due to turbulent fluctuations. It can be expressed as:

$$\frac{\partial (\overline{u'u'})}{\partial t} = \frac{\partial (\rho \overline{u'u'})}{\partial y} + \dots$$

This term involves higher-order correlations between velocity fluctuations, which complicates the equation system.

The closure problem arises in the Reynolds-Averaged Navier-Stokes (RANS) equations because the number of dependent variables (unknowns) exceeds the number of equations available. For instance: $-\bar{u}$ is an unknown. $-\bar{u'v'}$ (a Reynolds stress term) introduces additional unknowns.

To resolve this, closure models are used, which provide approximations for higher-order terms based on known variables.

For example, consider the term $\bar{u'w'}$. Using a simple closure model:

$$\bar{u'w'} = -k \frac{\partial \bar{u}}{\partial x},$$

where k is a proportionality constant (often related to eddy viscosity). Here, \bar{u} is already an unknown, so no additional variables are introduced, avoiding further complexity.

This is an example of first-order closure where higher-order terms are approximated using first-order variables.

Types of Closure Models

1. First-Order Closure:

- Simplifies higher-order terms using known variables and gradients (e.g., eddy viscosity models).
- Example: $\bar{u'w'} = -k \frac{\partial \bar{u}}{\partial x}$.
- Advantage: Computationally efficient but may lack accuracy in complex flows.

2. One-Point Closure:

- Approximates turbulence at a single point using local flow properties.
- Example: Mixing length models, where turbulent viscosity is proportional to local shear.

3. Second-Order Closure:

- Directly models second-order correlations like $\bar{u'u'}$ and $\bar{u'v'}$ by solving additional transport equations.
- Provides higher accuracy but increases computational cost.
- Example: Reynolds stress models (RSM), where additional equations are solved for Reynolds stresses.

4. Lecture 4 15/01/2025

$$\frac{d^2\theta}{dt^2} + \omega^2\theta = 0 \quad (4.1)$$

Initial conditions:

$$\theta(t=t_0) = \theta_0$$

$$\dot{\theta}(t=t_0) = \omega_0$$

We can rewrite this equation into two 1st-order ODEs,

Let $\frac{\theta}{dt} = p$, then

$$\frac{dp}{dt} + \omega^2\theta = 0 \quad (i)$$

$$\frac{d\theta}{dt} = p \quad (ii)$$

Initial conditions become:

$$\theta(t=t_0) = \theta_0$$

$$p(t=t_0) = \omega_0$$

4.1 Forward Difference

Invoking Taylor series expansion for $f(t+t_0)$:

$$\left. \begin{aligned} f(t+t_0) &= f(t_0) + (t+t_0) \frac{\partial f}{\partial t} \Big|_{t=t_0} \\ &\quad + \frac{(t+t_0)^2}{2!} \frac{\partial^2 f}{\partial t^2} \Big|_{t=t_0} + \dots \\ f(t+\Delta t) &= f(t_0) + \Delta t \frac{\partial f}{\partial t} \Big|_{t=t_0} \\ &\quad + \frac{(\Delta t)^2}{2!} \frac{\partial^2 f}{\partial t^2} \Big|_{t=t_0} + \dots \end{aligned} \right\} \quad (4.2)$$

we will get;

$$\left. \begin{aligned} p(t+t_0) &\approx p(t_0) + \Delta t \frac{\partial p}{\partial t} \Big|_{t=t_0} \\ p(t+t_0) &\approx p(t_0) + (t-t_0) \frac{\partial p}{\partial t} \Big|_{t=t_0} \\ \frac{\partial p}{\partial t} \Big|_{t=t_0} &\approx \frac{p(t+t_0) - p(t_0)}{(t-t_0)} \\ \frac{\partial p}{\partial t} \Big|_{t=t_0} &\approx \frac{p(t_1) - p(t_0)}{\Delta t} \end{aligned} \right\} \quad (4.3)$$

Eq.(4.3) is called **Forward difference**.

Order of Forward difference is $O(\Delta t)$.

Similarly Forward difference for θ ,

$$\left. \begin{aligned} \frac{\partial \theta}{\partial t} \Big|_{t=t_0} &\approx \frac{\theta(t+t_0) - \theta(t_0)}{(t-t_0)} \\ \frac{\partial \theta}{\partial t} \Big|_{t=t_0} &\approx \frac{\theta(t_1) - \theta(t_0)}{\Delta t} \end{aligned} \right\} \quad (4.4)$$

From Eq.(i) and Eq.(4.3):

$$\left. \begin{aligned} \frac{p(t+t_0) - p(t_0)}{(t-t_0)} + \omega^2\theta(t_0) &= 0 \\ p(t+t_0) &= p(t_0) - (t-t_0)\omega^2\theta(t_0) \\ p(t_1) &= p(t_0) - \Delta t\omega^2\theta(t_0) \end{aligned} \right\} \quad (4.5)$$

From Eq.(ii) and Eq.(4.4):

$$\left. \begin{aligned} \frac{\theta(t+t_0) - \theta(t_0)}{(t-t_0)} &= p(t_0) \\ \theta(t+t_0) &= \theta(t_0) - (t-t_0)p(t_0) \\ \theta(t_1) &= \theta(t_0) - \Delta t p(t_0) \end{aligned} \right\} \quad (4.6)$$

4.2 Backward Difference

Invoking Taylor series expansion for $f(t-t_0)$:

$$\left. \begin{aligned} f(t-t_0) &= f(t_0) - (t-t_0) \frac{\partial f}{\partial t} \Big|_{t=t_0} \\ &\quad + \frac{(t-t_0)^2}{2!} \frac{\partial^2 f}{\partial t^2} \Big|_{t=t_0} + \dots \\ f(t-\Delta t) &= f(t_0) - \Delta t \frac{\partial f}{\partial t} \Big|_{t=t_0} \\ &\quad + \frac{(\Delta t)^2}{2!} \frac{\partial^2 f}{\partial t^2} \Big|_{t=t_0} + \dots \end{aligned} \right\} \quad (4.7)$$

we will get;

$$\left. \begin{aligned} p(t-t_0) &\approx p(t_0) - \Delta t \frac{\partial p}{\partial t} \Big|_{t=t_0} \\ p(t-t_0) &\approx p(t_0) - (t-t_0) \frac{\partial p}{\partial t} \Big|_{t=t_0} \\ \frac{\partial p}{\partial t} \Big|_{t=t_0} &\approx \frac{p(t_0) - p(t-t_0)}{(t-t_0)} \\ \frac{\partial p}{\partial t} \Big|_{t=t_0} &\approx \frac{p(t_0) - p(t_{-1})}{\Delta t} \end{aligned} \right\} \quad (4.8)$$

Eq.(4.8) is called **Backward difference**.

Order of Backward difference is $O(\Delta t)$.

Similarly Backward difference for θ ,

$$\left. \begin{aligned} \frac{\partial \theta}{\partial t} \Big|_{t=t_0} &\approx \frac{\theta(t_0) - \theta(t-t_0)}{(t-t_0)} \\ \frac{\partial \theta}{\partial t} \Big|_{t=t_0} &\approx \frac{\theta(t_0) - \theta(t_{-1})}{\Delta t} \end{aligned} \right\} \quad (4.9)$$

From Eq.(i) and Eq.(4.8):

$$\left. \begin{aligned} \frac{p(t+t_0) - p(t_0)}{(t-t_0)} + \omega^2\theta(t_0) &= 0 \\ p(t+t_0) &= p(t_0) - (t-t_0)\omega^2\theta(t_0) \\ p(t_1) &= p(t_0) - \Delta t\omega^2\theta(t_0) \end{aligned} \right\} \quad (4.10)$$

From Eq.(ii) and Eq.(4.9):

$$\left. \begin{aligned} \frac{\theta(t+t_0) - \theta(t_0)}{(t-t_0)} &= p(t_0) \\ \theta(t+t_0) &= \theta(t_0) - (t-t_0)p(t_0) \\ \theta(t_1) &= \theta(t_0) - \Delta t p(t_0) \end{aligned} \right\} \quad (4.11)$$

4.3 Current Difference

The current difference method is a combination of the forward and backward difference methods, where we approximate the

values at the current time using both the forward and backward information.

Using forward difference:

$$\left. \frac{d\theta}{dt} \right|_{t=t_0} = p(t_0) = \text{known} = p(t_0) - \Delta t \omega^2 \theta(t_0) \quad (4.12)$$

Using backward difference:

$$\left. \frac{d\theta}{dt} \right|_{t=t_0} = \frac{\theta(t_1) - \theta(t - \Delta t)}{\Delta t} \quad (4.13)$$

From Eq.(4.12) and Eq.(4.13), we can combine both the equations to get:

$$\begin{aligned} \frac{\theta(t_1) - \theta(t - \Delta t)}{\Delta t} &= p(t_0) - \Delta t \omega^2 \theta(t_0) \\ \theta(t_1) &= \theta(t - \Delta t) + \Delta t [p(t_0) - \Delta t \omega^2 \theta(t_0)] \end{aligned} \quad (4.14)$$

In Eq.(4.14) all the terms on the RHS are known, allowing us to compute the value of θ at time (t_1) .

Thus, we obtain a method to compute the new value of θ based on the current and past time steps.

5. Lecture 5 20/01/2025

$$\left. \begin{aligned} f(x+\Delta x) &= f(x) + \Delta x \frac{\partial f}{\partial x} \Big|_{x=x_0} \\ &\quad + \frac{(\Delta x)^2}{2!} \frac{\partial^2 f}{\partial x^2} \Big|_{x=x_0} \\ &\quad + \frac{(\Delta x)^3}{3!} \frac{\partial^3 f}{\partial x^3} \Big|_{x=x_0} + \dots \end{aligned} \right\} \quad (5.1)$$

$$\left. \begin{aligned} f(x-\Delta x) &= f(x) - \Delta x \frac{\partial f}{\partial x} \Big|_{x=x_0} \\ &\quad + \frac{(\Delta x)^2}{2!} \frac{\partial^2 f}{\partial x^2} \Big|_{x=x_0} \\ &\quad - \frac{(\Delta x)^3}{3!} \frac{\partial^3 f}{\partial x^3} \Big|_{x=x_0} + \dots \end{aligned} \right\} \quad (5.2)$$

Using Eq.(5.1),

$$\frac{df}{dx} \Big|_{x=x_0} \simeq \left[\frac{f(x+\Delta x) - f(x)}{\Delta x} \right] + O(\Delta x) \quad (5.3)$$

Similarly, using Eq.(5.2), one obtain,

$$\frac{df}{dx} \Big|_{x=x_0} \simeq \left[\frac{f(x) - f(x-\Delta x)}{\Delta x} \right] + O(\Delta x) \quad (5.4)$$

Subtracting Eq.(5.2) from Eq.(5.1), we get,

$$\frac{df}{dx} \simeq \left[\frac{f(x+\Delta x) + f(x-\Delta x)}{2\Delta x} \right] + O(\Delta x^2) \quad (5.5)$$

Adding Eq.(5.1) and Eq.(5.2), we get,

$$\frac{d^2 f}{dx^2} \simeq \left[\frac{f(x+\Delta x) - 2f(x) + f(x-\Delta x)}{(\Delta x)^2} \right] + O(\Delta x^2) \quad (5.6)$$

$$\left. \begin{aligned} f(x+\Delta x, t) &= f(x, t) + \Delta x \frac{\partial f(x, t)}{\partial x} \Big|_{x=x_0} \\ &\quad + \frac{(\Delta x)^2}{2!} \frac{\partial^2 f(x, t)}{\partial x^2} \Big|_{x=x_0} \\ &\quad + \frac{(\Delta x)^3}{3!} \frac{\partial^3 f(x, t)}{\partial x^3} \Big|_{x=x_0} + \dots \end{aligned} \right\} \quad (5.7)$$

$$\left. \begin{aligned} f(x-\Delta x, t) &= f(x, t) - \Delta x \frac{\partial f(x, t)}{\partial x} \Big|_{x=x_0} \\ &\quad + \frac{(\Delta x)^2}{2!} \frac{\partial^2 f(x, t)}{\partial x^2} \Big|_{x=x_0} \\ &\quad - \frac{(\Delta x)^3}{3!} \frac{\partial^3 f(x, t)}{\partial x^3} \Big|_{x=x_0} + \dots \end{aligned} \right\} \quad (5.8)$$

5.1 Space Difference

From Eq.(5.7) and Eq.(5.8) respectively by difference equation w.r.t 1-direction, say x, we obtain **Space difference**:

$$\frac{df}{dx} \Big|_{x=x_0} \simeq \left[\frac{f(x+\Delta x, t) - f(x, t)}{\Delta x} \right] + O(\Delta x) \quad (5.9)$$

$$\frac{df}{dx} \Big|_{x=x_0} \simeq \left[\frac{f(x+\Delta x, t) - f(x, t)}{\Delta x} \right] + O(\Delta x) \quad (5.10)$$

Subtracting Eq.(5.8) from Eq.(5.7), we get,

$$\frac{df}{dx} \simeq \left[\frac{f(x+\Delta x, t) + f(x-\Delta x, t)}{2\Delta x} \right] + O(\Delta x^2) \quad (5.11)$$

Adding Eq.(5.7) and Eq.(5.8), we get,

$$\frac{d^2 f}{dx^2} \simeq \left[\frac{f(x+\Delta x, t) - 2f(x, t) + f(x-\Delta x, t)}{(\Delta x)^2} \right] + O(\Delta x^2) \quad (5.12)$$

5.2 Time Difference

Similarly, From Eq.(5.7) and Eq.(5.8) respectively, difference equation w.r.t time (t), we obtain **Time difference**:

$$\frac{df}{dt} \Big|_{t=t_0} \simeq \left[\frac{f(x, t+\Delta t) - f(x, t)}{\Delta t} \right] + O(\Delta t) \quad (5.13)$$

$$\frac{df}{dt} \Big|_{t=t_0} \simeq \left[\frac{f(x, t+\Delta t) - f(x, t)}{\Delta t} \right] + O(\Delta t) \quad (5.14)$$

$$\frac{df}{dt} \simeq \left[\frac{f(x, t+\Delta t) + f(x, t-\Delta t)}{2\Delta t} \right] + O(\Delta t^2) \quad (15)$$

$$\frac{d^2 f}{dt^2} \simeq \left[\frac{f(x, t+\Delta t) - 2f(x, t) + f(x, t-\Delta t)}{(\Delta t)^2} \right] + O(\Delta t^2) \quad (5.16)$$

5.3 Explicit Form Of Second Order PDE

$$A \frac{\partial^2 f}{\partial x^2} + B \frac{\partial^2 f}{\partial x \partial y} + C \frac{\partial^2 f}{\partial y^2} + D \frac{\partial f}{\partial x} + E \frac{\partial f}{\partial y} + F f = G \quad (5.17)$$

Cases:

1. If A, B, C, D, E, F and G are either constant or function of x and y, Eq.(5.17) is called **Linear PDE**.
2. If A, B and C are function of x, y and f, Eq.(5.17) is called **Quasi-linear PDE**.
3. If A, B and C are function of x and y only, Eq.(5.17) is called **Semi-linear PDE**.

Example: Momentum equation, which is,

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} - f v = -\frac{1}{\rho} \frac{\partial P}{\partial x}$$

is a quasi-linear PDE.

5.4 Implicit Form Of Second Order PDE

$$G\left(f, \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial^2 f}{\partial x^2}, \frac{\partial^2 f}{\partial y^2}, \frac{\partial^2 f}{\partial x \partial y}\right) = 0 \quad (5.18)$$

Eq.(5.18) is Implicit form of PDE. Example: Continuity equation in 1-D, which is,

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial x}(\rho u) = 0$$

1-D linear advection equation:

$$\frac{\partial \rho}{\partial t} + u \frac{\partial \rho}{\partial x} = 0$$

in Euler description.

$$\frac{D\rho}{Dt} = 0$$

in Lagrangian description, represents change in density (ρ) following the motion.

6. Lecture 6 21/01/2025

6.1 1-D Linear Advection Equation

$$\frac{\partial f}{\partial t} + u \frac{\partial f}{\partial x} = 0 \quad (6.1)$$

where

f is fluid property

u is x-component of fluid velocity




Figure 2. Path line or trajectory of fluid element

$$\frac{dx}{dt} = u = \text{constant} \quad (6.2)$$

Eq.(6.2) is called **Characteristic equation**.

Integrating Eq.(6.2) w.r.t time, we get:

$$x = x_0 + \int_{t_0}^t u dt \quad (6.3)$$

Eq.(6.3) is called equation of pathline as shown in Fig2.

Substituting, $u = \frac{dx}{dt}$ in Eq.(6.1), we obtain:

$$\frac{\partial f}{\partial t} + \frac{dx}{dt} \frac{\partial f}{\partial x} = 0 \quad (6.4)$$

$$\frac{Df}{Dt} = 0 \quad (6.4)$$

Eq.(6.4) implies the property f is conserved following the motion of fluid element.

6.2 Triangular Property

1. **Hyperbolic:** 1st order partial equation → 1 family of characteristic equation in real domain.
2. **Hyperbolic:** 2nd order partial equation → 2 distinct and real set of characteristic equations.
3. **Parabolic:** 2nd order partial equation → 2 equal and real set of characteristic equations.
4. **Elliptical:** 2nd order partial equation → 2 distinct and complex set of characteristic equations.




Figure 3. Triangular property distribution

$$\frac{\partial f}{\partial t} + u \frac{\partial f}{\partial x} = 0 \quad \begin{cases} u = \text{constant} & \Rightarrow \text{linear PDE} , \\ u = u(x, t) & \Rightarrow \text{linear PDE} , \\ u = u(x, t, f) & \Rightarrow \text{quasi-linear PDE} \end{cases}$$

$$a(x, t, f) \frac{\partial f}{\partial t} + b(x, t, f) \frac{\partial f}{\partial x} = 0 \quad (6.5)$$

Above Eq.(6.5) is quasi-linear PDE, having characteristic equation in real domain $\Rightarrow \frac{dx}{dt} = \frac{b}{a}$

$$a(x, t, f) \frac{\partial f}{\partial t} + b(x, t, f) \frac{\partial f}{\partial x} = c(x, t, f) \quad (6.6)$$

Above Eq.(6.6) is semi-linear PDE, having characteristic equation in real domain.

7. Lecture 7 22/01/2025

General quasi-linear 2nd order PDE:

$$A \frac{\partial^2 f}{\partial x^2} + B \frac{\partial^2 f}{\partial x \partial y} + C \frac{\partial^2 f}{\partial y^2} + D \frac{\partial f}{\partial x} + E \frac{\partial f}{\partial y} + Ff = G \quad (7.1)$$

i.e. A, B, C can be function of $x, y, f, \frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$.

To find sign of discriminant $B^2 - 4AC$ tells the classification of equation:

$$B^2 - 4AC \begin{cases} > 0, & \text{Hyperbolic PDE} \\ = 0, & \text{Parabolic PDE} \\ < 0, & \text{Elliptical PDE} \end{cases}$$

This is analogous to general equation of conic-section curves. Which is general by Eq.(7.2)

$$Ay^2 + Bxy + Cx^2 + Dx + Ey + F = 0 \quad (7.2)$$

$$B^2 - 4AC \begin{cases} > 0, & \text{Hyperbola} \\ = 0, & \text{Parabola} \\ < 0, & \text{Ellipse} \end{cases}$$

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy \quad (7.3)$$

$$d\left(\frac{\partial f}{\partial x}\right) = \frac{\partial^2 f}{\partial x^2} dx + \frac{\partial^2 f}{\partial x \partial y} dy \quad (7.4)$$

$$d\left(\frac{\partial f}{\partial y}\right) = \frac{\partial^2 f}{\partial x \partial y} dx + \frac{\partial^2 f}{\partial y^2} dy \quad (7.5)$$

Unknown in Eq.(7.3), Eq.(7.4) and Eq.(7.5) are 2nd order derivatives.

From Eq.(7.1),

$$A \frac{\partial^2 f}{\partial x^2} + B \frac{\partial^2 f}{\partial x \partial y} + C \frac{\partial^2 f}{\partial y^2} = -D \frac{\partial f}{\partial x} - E \frac{\partial f}{\partial y} - Ff + G \quad (7.6)$$

From Eq.(7.4),

$$\frac{\partial^2 f}{\partial x^2} dx + \frac{\partial^2 f}{\partial x \partial y} dy = d\left(\frac{\partial f}{\partial x}\right) \quad (7.7)$$

From Eq.(7.5),

$$\frac{\partial^2 f}{\partial x \partial y} dx + \frac{\partial^2 f}{\partial y^2} dy = d\left(\frac{\partial f}{\partial y}\right) \quad (7.8)$$

Matrix representation of Eq.(7.6), Eq.(7.7) and Eq.(7.8):

$$\underbrace{\begin{bmatrix} A & B & C \\ dx & dy & 0 \\ 0 & dx & dy \end{bmatrix}}_{\det(\mathbf{A})=0} \begin{bmatrix} \frac{\partial^2 f}{\partial x^2} \\ \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y^2} \end{bmatrix} = \begin{bmatrix} -D \frac{\partial f}{\partial x} - E \frac{\partial f}{\partial y} - Ff + G \\ d\left(\frac{\partial f}{\partial x}\right) \\ d\left(\frac{\partial f}{\partial y}\right) \end{bmatrix}.$$

Determinant of (\mathbf{A}) should be equal to zero, i.e.,

$$\begin{aligned} \det(\mathbf{A}) &= 0 \\ A(dy)^2 - B(dxdy) + C(dx)^2 &= 0 \\ A\left(\frac{dy}{dx}\right)^2 - B\left(\frac{dy}{dx}\right) + C &= 0 \end{aligned} \quad (7.9)$$

Solution of above Eq.(7.9) are 2 characteristic equations:

$$\frac{dy}{dx} = \frac{B \pm \sqrt{B^2 - 4AC}}{2A}$$

if $B^2 - 4AC > 0 \Rightarrow 2$ distinct Real roots

Family of char. curve $\in \mathbb{R}$

Hyperbolic PDE

if $B^2 - 4AC = 0 \Rightarrow$ Real and equal roots

1 Family of char. curve $\in \mathbb{R}$


Parabolic PDE

if $B^2 - 4AC < 0 \Rightarrow$ Imaginary roots

Family of char. curves $\in \mathbb{C}$

Elliptic PDE

Examples:



8. Lecture 08/01/2025

8.1 Heat Equation

$$k \frac{\partial^2 T}{\partial x^2} - \frac{\partial T}{\partial t} = 0 \quad (8.1)$$

Eq.(8.1) can be decomposed into pair of PDEs:

$$\frac{\partial^2 T}{\partial x \partial t} dx + \frac{\partial^2 T}{\partial t^2} dt = d\left(\frac{\partial T}{\partial t}\right) \quad (8.2)$$

$$\frac{\partial^2 T}{\partial x^2} dx + \frac{\partial^2 T}{\partial x \partial t} dt = d\left(\frac{\partial T}{\partial x}\right) \quad (8.3)$$

Rewriting Eq.(8.1), Eq.(8.2) and Eq.(8.3) into matrix form:

$$\underbrace{\begin{bmatrix} k & 0 & 0 \\ 0 & dx & dt \\ dx & dt & 0 \end{bmatrix}}_A \begin{bmatrix} \frac{\partial^2 T}{\partial x^2} \\ \frac{\partial^2 T}{\partial x \partial t} \\ \frac{\partial^2 T}{\partial t^2} \end{bmatrix} = \begin{bmatrix} d\left(\frac{\partial T}{\partial t}\right) \\ d\left(\frac{\partial T}{\partial x}\right) \end{bmatrix}$$

$$\det(A) = 0$$

$$k(0 - (dt)^2) - 0(0 - dxdt) + 0(0 - (dx)^2) = 0$$

$$-k(dt)^2 = 0$$

$$\therefore dt = 0$$

$$t = \text{const.}$$

Therefore the equation represents Parabolic PDE (2ⁿ order quasi-linear PDE).

8.2 1-D Wave Equation

$$\frac{\partial^2 f}{\partial t^2} = c^2 \frac{\partial^2 f}{\partial x^2} \quad (8.4)$$

$$\underbrace{\left(\frac{\partial^2}{\partial t^2} - c^2 \frac{\partial^2}{\partial x^2}\right)}_{\text{Operator acting on } f} f = 0$$

$$\left[\left(\frac{\partial}{\partial t} - c \frac{\partial}{\partial x}\right)\left(\frac{\partial}{\partial t} + c \frac{\partial}{\partial x}\right)\right] f = 0 \quad (8.5)$$

Above Eq.(8.5) is pair two 1-D advection equations with constant speed c in **negative** x-direction and **positive** x-direction respectively.



Discriminant of Eq.(8.4), for $A = 1$, $B=0$, $C=-1$:

$$B^2 - 4AC$$

$$0^2 - 4c^2(-1)$$

$$4c^2 > 0$$

Therefore Eq.(8.4) represents Hyperbolic PDE.



$$\frac{\partial^2 f}{\partial x \partial t} dx + \frac{\partial^2 f}{\partial t^2} dt = d\left(\frac{\partial f}{\partial t}\right) \quad (8.5)$$

$$\frac{\partial^2 f}{\partial x^2} dx + \frac{\partial^2 f}{\partial x \partial t} dt = d\left(\frac{\partial f}{\partial x}\right) \quad (8.6)$$

Rewriting Eq.(8.4), Eq.(8.5) and Eq.(8.6) into matrix form:

$$\underbrace{\begin{bmatrix} c^2 & 0 & -1 \\ 0 & dx & dt \\ dx & dt & 0 \end{bmatrix}}_A \begin{bmatrix} \frac{\partial^2 f}{\partial x^2} \\ \frac{\partial^2 f}{\partial x \partial t} \\ \frac{\partial^2 f}{\partial t^2} \end{bmatrix} = \begin{bmatrix} d\left(\frac{\partial f}{\partial t}\right) \\ d\left(\frac{\partial f}{\partial x}\right) \end{bmatrix}$$

$$\det(A) = 0$$

$$c^2(0 - (dt)^2) - 0(0 - dxdt) + (-1)(0 - (dx)^2) = 0$$

$$-c^2(dt)^2 + (dx)^2 = 0$$

$$-c^2\left(\frac{dt}{dx}\right)^2 + 1 = 0$$

$$c^2\left(\frac{dt}{dx}\right)^2 - 1 = 0$$

Slope $\frac{dt}{dx}$ is equal to:

$$\frac{dt}{dx} = \frac{-0 \pm \sqrt{0^2 - 4 \cdot c^2 \cdot (-1)}}{2 \cdot c^2}$$

$$\frac{dt}{dx} = \pm \frac{1}{2c}$$

8.3 Laplace Equation

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0 \quad (8.7)$$

$$\frac{\partial^2 f}{\partial x \partial y} dx + \frac{\partial^2 f}{\partial y^2} dy = d \left(\frac{\partial f}{\partial y} \right) \quad (8.8)$$

$$\frac{\partial^2 f}{\partial x^2} dx + \frac{\partial^2 f}{\partial x \partial y} dy = d \left(\frac{\partial f}{\partial x} \right) \quad (8.9)$$

Discriminant of Eq.(8.7), for A = 1, B=0, C=1:

$$B^2 - 4AC$$

$$0^2 - 4(1)(1)$$

$$-4 < 0$$

Therefore Eq.(8.7) represents Elliptical PDE.

Rewriting Eq.(8.7), Eq.(8.8) and Eq.(0) into matrix form:

$$\underbrace{\begin{bmatrix} 1 & 0 & 1 \\ 0 & dx & dy \\ dx & dy & 0 \end{bmatrix}}_A \begin{bmatrix} \frac{\partial^2 f}{\partial x^2} \\ \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y^2} \end{bmatrix} = \begin{bmatrix} 0 \\ d \left(\frac{\partial f}{\partial y} \right) \\ d \left(\frac{\partial f}{\partial x} \right) \end{bmatrix}$$

$$\det(A) = 0$$

$$1(0 - (dy)^2) - 0(0 - dx dy) + 1(0 - (dx)^2) = 0$$

$$-(dy)^2 - (dx)^2 = 0$$

$$\left(\frac{dy}{dx} \right)^2 + 1 = 0$$

$$\frac{dy}{dx} = \pm i$$

9. Lecture 9 28/01/2025

- Domain of dependence at a given point in the solution domain would corresponding to the region where solution will impact solution will be impacted by the solution at the above given point.
- Range of independence at a given point in the solution domain would corresponding to the region where solution will impact solution will be impacted by the solution at the above given point.

9.1 Wave Equation

Hyperbolic PDE:


$$\frac{\partial^2 f}{\partial t^2} = c^2 \frac{\partial^2 f}{\partial x^2} \quad (9.1)$$

Parabolic PDE:

$$\frac{\partial T}{\partial t} = c^2 \frac{\partial T}{\partial x} \quad (9.2)$$

Ellipse PDE:

$$\frac{\partial^2 f}{\partial t^2} + \frac{\partial^2 f}{\partial x^2} = 0 \quad (9.3)$$



$$a \frac{\partial f}{\partial x} + b \frac{\partial f}{\partial x} + c \frac{\partial g}{\partial t} + d \frac{\partial g}{\partial x} = e \quad (9.4)$$

$$A \frac{\partial f}{\partial x} + B \frac{\partial f}{\partial x} + C \frac{\partial g}{\partial t} + D \frac{\partial g}{\partial x} = E \quad (9.5)$$

Using chain rule;

$$\frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dx = df \quad (9.6)$$

$$\frac{\partial g}{\partial t} dt + \frac{\partial g}{\partial x} dx = dg \quad (9.7)$$

Rewriting Eq.(9.4), Eq.(9.5), Eq.(9.6) and Eq.(9.7) into matrix form:

$$\underbrace{\begin{bmatrix} a & b & c & d \\ A & B & C & D \\ dt & dx & 0 & 0 \\ 0 & 0 & dt & dx \end{bmatrix}}_A \begin{bmatrix} \frac{\partial f}{\partial t} \\ \frac{\partial f}{\partial x} \\ \frac{\partial g}{\partial t} \\ \frac{\partial g}{\partial x} \end{bmatrix} = \begin{bmatrix} e \\ E \\ df \\ dg \end{bmatrix}$$

$$\det(A) = 0$$

$$(dx)^2 \underbrace{(aC - Ac)}_{\bar{A}} - (dx)(dt) \underbrace{(aD - Ad + bC - Bc)}_{\bar{B}} + (dt)^2 \underbrace{(bD - Bd)}_{\bar{C}} = 0$$

Discriminant of Eq.(8.7), for $\bar{A} = (aC - Ac)$, $\bar{B} = (aD - Ad + bC - Bc)$, $\bar{C} = (bD - Bd)$:

$$\bar{B}^2 - 4\bar{A}\bar{C}$$

$$(aD - Ad + bC - Bc)^2 - 4(aC - Ac)(bD - Bd)$$

$$\frac{dx}{dt} = \frac{\bar{B} \pm \sqrt{\bar{B}^2 - 4\bar{A}\bar{C}}}{2\bar{A}} \quad (9.8)$$


Depending upon sign of $\bar{B}^2 - 4\bar{A}\bar{C}$ the following classification can be made:

- Negative \rightarrow Elliptic
- Zero \rightarrow Parabolic
- positive \rightarrow Hyperbolic

9.2 Heat Transfer Equation

$$\frac{\partial T}{\partial t} = k \frac{\partial^2 T}{\partial x^2} \quad (9.9)$$

where, k is coefficient of thermal expansion.



Applying FTCS (Forward Time and Central in Space) scheme,

$$\begin{aligned} t_n &= n\Delta t & , n &= 0, 1, 2, \dots \\ x_m &= x_0 + m\Delta x & , m &= 0, 1, 2, \dots, N \end{aligned}$$

Backward difference:

$$\left(\frac{\partial T}{\partial t} \right)_{x,t} = \frac{T_{m,n+1} - T_{m,n}}{\Delta t} \quad (9.10)$$

Central difference:

$$\left(\frac{\partial^2 T}{\partial t^2} \right)_{x,t} = \frac{T_{m+1,n} - 2 \cdot T_{m,n} + T_{m-1,n}}{\Delta t} \quad (9.11)$$

Substituting Eq.(9.10) and Eq.(9.11) into Eq.(9.9), we obtain,

$$\left(\frac{T_{m,n+1}-T_{m,n}}{\Delta t}\right)=k\left(\frac{T_{m+1,n}-2\cdot T_{m,n}+T_{m-1,n}}{\Delta t}\right)$$

$$T_{m,n+1}=T_{m,n}+\underbrace{\frac{k\Delta t}{(\Delta x)^2}}_{\lambda}(T_{m+1,n}-2T_{m,n}+T_{m-1,n})$$

$$T_{m,n+1}=(1-2\lambda)T_{m,n}+\lambda T_{m-1,n}+\lambda T_{m+1,n}$$

