
Started on Monday, 27 October 2025, 4:10 PM

State Finished

Completed on Monday, 27 October 2025, 4:39 PM

Time taken 29 mins 23 secs

Marks 1.00/1.00

Grade **10.00** out of 10.00 (**100%**)

Question 1 | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1 #include<stdio.h>
2 int main(){
3     int n,f=0;
4     scanf("%d",&n);
5     int a[n];
6     for(int i=0;i<n;i++)
7         scanf("%d",&a[i]);
8     int k;
9     scanf("%d",&k);
10    int i=0,j=1;
11    while(i<n && j<n){
12        if(i!=j){
13            if(a[j]-a[i]==k){
14                f=1;
15                i++;
16                j++;
17            }
18            else if(a[j]-a[i]<k)
19                j++;
20            else
21                i++;
22        }
23        else
24            j++;
25    }
26    printf("%d",f);
27 }
28
29

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.