



सत्यमेव जयते
Government of India

Ministry of Health and Family Welfare

NATIONAL FAMILY HEALTH SURVEY - 6

2023-24

COMPENDIUM OF FACT SHEETS KEY INDICATORS OF STATE AND DISTRICTS UTTAR PRADESH



(स्थापना/ Established in 1956)
बेहतर भविष्य के लिए क्षमता निर्माण
Capacity Building for a Better Future

International Institute for Population Sciences
(Deemed University)

NATIONAL FAMILY HEALTH SURVEY (NFHS-6)

2023-24

FACT SHEETS

AUGUST 2026

Suggested citation: International Institute for Population Sciences (IIPS) 2026.
National Family Health Survey (NFHS-6), 2023-24: State and District Fact Sheets, Uttar Pradesh. Mumbai:
IIPS.

CONTRIBUTORS

Laxmi Kant Dwivedi

Preeti Dhillon

Somnath Jana

Nirbhay Singh

Kaviyarasan M.

Vaidehi Yelamanchili

Ashapurna Goswami

For additional information about the 2023-24 National Family Health Survey (NFHS-6), please contact:

International Institute for Population Sciences, Govandi Station Road, B.S. Devashi Marg,
Deonar, Mumbai- 400 088

Telephone: 022 - 42372549

Email: nfhs6pmu@gmail.com, nfhsdata@iipsindia.ac.in

For related information, visit <https://www.iipsindia.ac.in> and <https://www.nfhsiips.in>

TABLE OF CONTENTS

S. No.	Content	Page No.
STATE FACT SHEET		
	UTTAR PRADESH	3
DISTRICT FACT SHEET		
1.	Agra	9
2.	Aligarh	13
3.	Ambedkar Nagar	17
4.	Amethi	21
5.	Amroha	25
6.	Auraiya	29
7.	Ayodhya	33
8.	Azamgarh	37
9.	Baghpat	41
10.	Bahraich	45
11.	Ballia	49
12.	Balrampur	53
13.	Banda	57
14.	Barabanki	61
15.	Bareilly	65
16.	Basti	69
17.	Bhadohi	73
18.	Bijnor	77
19.	Budaun	81
20.	Bulandshahr	85
21.	Chandauli	89
22.	Chitrakoot	93
23.	Deoria	97
24.	Etah	101
25.	Etawah	105
26.	Farrukhabad	109
27.	Fatehpur	113
28.	Firozabad	117
29.	Gautam Buddha Nagar	121
30.	Ghaziabad	125
31.	Ghazipur	129
32.	Gonda	133
33.	Gorakhpur	137
34.	Hamirpur	141
35.	Hapur	145
36.	Hardoi	149
37.	Hathras	153
38.	Jalaun	157

S. No.	Content	Page No.
DISTRICT FACT SHEET		
39.	Jaunpur	161
40.	Jhansi	165
41.	Kannauj	169
42.	Kanpur Dehat	173
43.	Kanpur Nagar	177
44.	Kasganj	181
45.	Kaushambi	185
46.	Kheri	189
47.	Kushinagar	193
48.	Lalitpur	197
49.	Lucknow	201
50.	Maharajganj	205
51.	Mahoba	209
52.	Mainpuri	213
53.	Mathura	217
54.	Mau	221
55.	Meerut	225
56.	Mirzapur	229
57.	Moradabad	233
58.	Muzaffarnagar	237
59.	Pilibhit	241
60.	Pratapgarh	245
61.	Prayagraj	249
62.	Rae Bareli	253
63.	Rampur	257
64.	Saharanpur	261
65.	Sambhal	265
66.	Sant Kabir Nagar	269
67.	Shahjahanpur	273
68.	Shamli	277
69.	Shrawasti	281
70.	Siddharth Nagar	285
71.	Sitapur	289
72.	Sonbhadra	293
73.	Sultanpur	297
74.	Unnao	301
75.	Varanasi	305
Appendix: State and Districtwise Sample Size		309



NATIONAL FAMILY HEALTH SURVEY (NFHS-6), 2023-24

FACT SHEET **State**

Introduction

The National Family Health Survey 2023-24 (NFHS-6), the sixth in the NFHS series, provides information on population, health, and nutrition for India and each State/Union Territory (UT) except Manipur. Like NFHS-4 and NFHS-5, NFHS-6 also provides District-level estimates for many important indicators. NFHS-6 is not designed to provide rural/urban estimates at the District level.

As in the earlier rounds, the Ministry of Health and Family Welfare (MoHFW), Government of India, designated the International Institute for Population Sciences (IIPS), Mumbai, as the nodal agency to conduct NFHS-6. The NFHS-6 architecture has been designed and implemented indigenously, and it was conducted without any technical or financial support from external organizations.

The main objective of each successive round of the NFHS has been to provide high quality data on health and family welfare and emerging issues in this area. NFHS-6 data will be useful in setting benchmarks and examining the progress the health sector has made over time. Besides providing evidence for the effectiveness of ongoing programmes, the data from NFHS-6 help in identifying the need for new programmes with an area specific focus and identifying groups that are most in need of essential services.

Four Survey Schedules - Household, Woman's, Man's, and Biomarker - were canvassed in local languages using Computer Assisted Personal Interviewing (CAPI). In the Household Schedule, information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as socio-economic characteristics of the household; health insurance coverage; land ownership, etc. The Woman's Schedule covered a wide variety of topics, including the woman's characteristics, marriage, fertility, contraception, children's immunizations and healthcare, nutrition, reproductive health, sexual behaviour, women's empowerment, and domestic violence. The Man's Schedule covered the man's characteristics, marriage, health issues, nutrition, sexual behaviour and attitudes towards gender roles. The Biomarker Schedule covered measurements of length/height and weight for children under age 5 years; measurements of height and weight for women age 15-49 years and men age 15-54 years; and blood pressure and random blood glucose levels for women and men age 15 years and over. In addition, women and men were requested to provide a few additional drops of blood from a finger prick for laboratory testing for HIV, Hepatitis-B and Hepatitis-C. Additionally, Dried Blood Spot (DBS) was collected from children age 4-5 years for Hepatitis-B testing.

Like the previous NFHS surveys, a two-stage sample design was used to collect data from both urban and rural areas. However, unlike the previous rounds, the Urban Frame Survey (UFS) blocks (2012-17) were selected in first stage in urban areas instead of Census Enumeration Blocks (CEB). This strategy improved the identification of boundaries of the Primary Sampling Units (PSUs). Several standard protocols and strategies were adopted to minimize non-sampling errors and maintain data quality such as multi-level monitoring and supervision of fieldwork; use of Computer Assisted Personal Interviewing (CAPI) for collecting data; centralized training for uniformity in survey implementation; standardized tools and procedures; Clinical, Anthropometric, and Biochemical (CAB) protocols followed as per international standards; secondary editing of data in the field; feedback on data quality to field teams based on real-time data; etc.

This compendium of fact sheets provides information on key indicators and trends for Uttar Pradesh and its districts. NFHS-6 fieldwork for Uttar Pradesh was conducted from 12 January, 2024 to 25 December, 2024 by Research & Development Initiative Private Limited, Academy of Management Studies and Network for Engineering and Economics Research and Management Pvt. Ltd. The results presented in the NFHS-6 Fact Sheets are provisional. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)			NFHS-5 (2019-21)
	Urban	Rural	Total	Total
Population and Household Profile				
1. Population below age 5 years (%)	7.7	9.9	9.4	9.5
2. Population below age 15 years (%)	25.4	31.3	29.9	31.0
3. Population age 60 years and above (%)	9.7	10.4	10.3	10.0
4. Population living in households with electricity (%)	98.8	94.9	95.8	91.0
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	99.2	99.3	99.2
6. Households using iodized salt (%)	92.7	84.6	86.5	92.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.7	37.0	37.2	15.9
8. Households with any usual member having a bank account/post office account (%)	97.1	98.8	98.4	96.6
9. Female population age 6 years and above who ever attended school (%)	78.4	67.6	70.1	67.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.8	19.8	20.1	12.2
11. Children age 2-4 years who attended pre-school (%)	33.9	29.6	30.4	15.1
Characteristics of Adults (age 15-49 years)				
12. Women with 10 or more years of schooling (%)	55.2	38.6	42.5	39.3
13. Men with 10 or more years of schooling (%)	53.5	47.4	49.1	48.6
14. Women who have ever used the internet (%)	79.0	67.3	70.0	30.6
15. Men who have ever used the internet (%)	85.9	79.1	81.0	59.1
Marriage and Fertility				
16. Women age 20-24 years married before age 18 years (%)	8.7	15.1	13.7	15.8
17. Men age 25-29 years married before age 21 years (%)	11.1	22.4	19.5	23.0
18. Total fertility rate (children per woman)	1.8	2.3	2.2	2.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.1	3.8	3.5	2.9
Current Use of Family Planning Methods (currently married women age 15-49 years)				
20. Any method ² (%)	66.1	61.3	62.4	62.4
21. Any modern method ² (%)	37.9	31.9	33.2	44.5
22. Any traditional method (%)	28.2	29.4	29.1	17.9
23. Female sterilization (%)	12.3	18.3	17.0	16.9
24. Male sterilization (%)	0.2	0.1	0.1	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)				
25. Total unmet need ³ (%)	7.3	11.8	10.8	12.8
26. Unmet need for spacing ³ (%)	3.2	5.3	4.8	4.7
27. Unmet need for limiting ³ (%)	4.1	6.5	6.0	8.1
Maternal and Child Health				
Antenatal Care (for last birth in the 5 years before the survey)				
28. Mothers who had an antenatal check-up in the first trimester (%)	76.2	69.1	70.6	62.5
29. Mothers who had any antenatal care visits (%)	96.8	96.9	96.9	94.4
30. Mothers who had at least 4 antenatal care visits (%)	58.6	50.0	51.8	42.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.7	92.8	93.0	92.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	40.7	26.7	29.6	22.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	26.7	15.3	17.7	9.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	89.3	94.1	93.1	95.7
Delivery Care (for births in the 5 years before the survey)				
35. Institutional births (%)	86.5	85.8	85.9	83.4
36. Institutional births in public facility (%)	36.6	55.3	51.8	57.7
37. Births attended by skilled health personnel ⁵ (%)	89.4	87.2	87.6	84.8
38. Births delivered by caesarean section (%)	33.6	16.2	19.5	13.7
39. Births in a private health facility that were delivered by caesarean section (%)	53.9	44.7	47.3	39.4
40. Births in a public health facility that were delivered by caesarean section (%)	18.4	4.6	6.5	6.2

Note: Major indicators are highlighted in grey.

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)			NFHS-5 (2019-21)
	Urban	Rural	Total	Total
Postnatal Care (for last birth in the 5 years before the survey)				
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery ⁺ (%)	81.4	75.4	76.7	72.0
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.3	3.2	3.3	2.4
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/ other health personnel within 2 days of delivery ⁺ (%)	82.8	77.6	78.7	70.2
Child Vaccinations and Vitamin A Supplementation				
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	72.7	83.5	81.4	69.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	82.6	90.0	88.6	78.4
46. Children age 12-23 months who have received any vaccine (%)	94.8	96.2	95.9	95.1
47. Children age 12-23 months who have received BCG (%)	92.8	94.7	94.3	93.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	76.2	86.6	84.5	74.3
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.4	89.0	88.1	81.3
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	87.1	91.2	90.4	83.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	67.1	74.0	72.6	58.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	56.7	57.4	57.3	50.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	79.8	87.4	85.9	49.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	61.3	66.5	65.4	73.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	88.1	96.5	94.8	94.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	6.0	1.2	2.2	2.2
Treatment of Childhood Diseases (children under age 5 years)				
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.9	13.2	13.0	5.6
58. Prevalence of severe diarrhoea in the 2 weeks preceding the survey (%)	0.9	1.2	1.1	0.5
59. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.1	2.7	2.6	3.5
60. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	75.3	62.4	64.6	63.0
Child Feeding Practices and Nutritional Status of Children				
61. Children under age 3 years breastfed within one hour of birth ⁹ (%)	35.2	45.0	43.1	23.9
62. Children under age 6 months exclusively breastfed ¹⁰ (%)	35.9	34.4	34.6	59.7
63. Children under age 6 months currently breastfeeding ¹⁰ (%)	88.2	93.3	92.5	93.9
64. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water only/other milk ¹⁰ (%)	85.6	82.4	82.9	82.0
65. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	58.8	59.8	59.6	31.0
66. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.7	18.2	18.1	5.8
67. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	12.5	13.5	13.2	6.7
68. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	16.4	17.3	17.1	5.9
69. Children under 5 years who are stunted (height-for-age) ¹² (%)	28.8	32.1	31.5	39.7
70. Children under 5 years who are wasted (weight-for-height) ¹² (%)	18.7	19.4	19.2	17.3
71. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.5	4.7	4.8	7.3
72. Children under 5 years who are underweight (weight-for-age) ¹² (%)	32.0	35.0	34.5	32.1
73. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	0.5	0.5	3.1

⁺LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)		NFHS-5 (2019-21)	
	Urban	Rural	Total	Total
Nutritional Status of Adults (age 15-49 years)				
74. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	16.5	23.9	22.2	19.0
75. Men whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) (%)	19.0	25.1	23.5	18.0
76. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁹ (%)	36.7	23.7	26.7	21.4
77. Men who are overweight or obese (BMI ≥25.0 kg/m ²) (%)	29.4	19.1	21.8	18.5
Blood Sugar Level among Adults (age 15 years and above)				
Women				
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.8	7.4	7.5	4.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.0	7.0	7.6	4.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.7	14.7	15.5	10.0
Men				
81. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.1	9.5	9.4	5.8
82. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.2	9.0	9.5	5.0
83. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.3	18.9	19.5	11.6
Hypertension among Adults (age 15 years and above)				
Women				
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.0	5.6	5.7	11.5
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	3.6	3.6	4.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.6	11.4	12.1	18.3
Men				
87. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.6	7.9	8.1	15.2
88. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	2.3	2.4	5.2
89. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.3	13.4	14.1	21.7
Women's Empowerment (women age 15-49 years)				
90. Currently married women who usually participate in three household decisions ¹⁷ (%)	90.1	84.7	85.9	87.6
91. Women who worked in the last 12 months and were paid in cash (%)	23.8	20.2	21.1	15.5
92. Women having a bank or savings account that they themselves use (%)	81.0	84.3	83.5	75.4
93. Women having a mobile phone that they themselves use (%)	67.1	53.1	56.4	46.5
94. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁸ (%)	81.7	65.9	69.2	72.9
Gender Based Violence (age 18-49 years)				
95. Ever-married women age 18-49 years who have ever experienced spousal violence ¹⁹ (%)	20.7	30.7	28.5	34.9
96. Ever-married women age 18-49 years who have experienced physical violence during any pregnancy (%)	3.1	3.9	3.7	3.7
97. Young women age 18-29 years who experienced sexual violence by age 18 (%)	0.3	0.6	0.5	0.7
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)				
98. Women age 15 years and above who use any kind of tobacco (%)	6.7	10.5	9.6	8.5
99. Men age 15 years and above who use any kind of tobacco (%)	34.7	48.9	45.2	44.0
100. Women age 15 years and above who consume alcohol (%)	0.3	0.3	0.3	0.3
101. Men age 15 years and above who consume alcohol (%)	15.2	19.9	18.7	14.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Decisions about health care for herself, making major household purchases, and visits to her family or relatives.

¹⁸Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

¹⁹Spousal violence is defined as physical and/or sexual violence.



NATIONAL FAMILY HEALTH SURVEY (NFHS-6), 2023-24

FACT SHEETS **Districts**

Agra, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.9	9.8
2. Population below age 15 years (%)	29.9	30.5
3. Population age 60 years and above (%)	9.7	8.4
4. Population living in households with electricity (%)	97.7	99.1
5. Population living in households with an improved drinking-water source ¹ (%)	98.5	99.8
6. Households using iodized salt (%)	90.3	93.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	25.4	9.4
8. Households with any usual member having a bank account/post office account (%)	97.5	95.9
9. Female population age 6 years and above who ever attended school (%)	68.3	69.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.2	15.6
11. Children age 2-4 years who attended pre-school (%)	27.6	35.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	40.3	36.1
13. Men with 10 or more years of schooling (%)	43.8	44.0
14. Women who have ever used the internet (%)	63.4	32.9
15. Men who have ever used the internet (%)	81.2	59.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	15.4	17.9
17. Men age 25-29 years married before age 21 years (%)	(18.4)	*
18. Three or more births in the 5 years preceding the survey (%)	1.7	3.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.1	3.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	68.2	67.7
21. Any modern method ² (%)	37.3	47.1
22. Any traditional method (%)	30.9	20.6
23. Female sterilization (%)	22.2	20.0
24. Male sterilization (%)	0.2	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.5	9.0
26. Unmet need for spacing ³ (%)	3.5	4.0
27. Unmet need for limiting ³ (%)	6.0	5.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	75.2	62.1
29. Mothers who had any antenatal care visits (%)	98.2	89.8
30. Mothers who had at least 4 antenatal care visits (%)	50.0	42.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.7	91.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	29.8	20.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	20.2	11.3
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	90.2	85.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	93.1	89.1
36. Institutional births in public facility (%)	51.2	43.0
37. Births attended by skilled health personnel ⁵ (%)	94.7	90.1
38. Births delivered by caesarean section (%)	23.2	18.8
39. Births in a private health facility that were delivered by caesarean section (%)	48.1	31.6
40. Births in a public health facility that were delivered by caesarean section (%)	6.0	9.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.
() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Agra, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.8	77.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	(1.9)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.6	75.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	80.8	74.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	88.7	78.2
46. Children age 12-23 months who have received any vaccine (%)	100.0	100.0
47. Children age 12-23 months who have received BCG (%)	97.9	99.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	83.0	75.8
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	98.9	92.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	94.4	96.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	64.8	67.9
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	62.4	47.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	91.2	45.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	64.8	71.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.4	87.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.6	5.7
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	6.2	6.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.4	4.5
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	89.7	80.7
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	39.6	19.5
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	48.0	43.1
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	91.6	96.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	86.2	77.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	13.0	10.7
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	(10.1)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	13.4	10.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	33.8	35.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	11.7	16.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.9	4.6
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	29.3	26.2
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.8	3.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Agra, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	18.3	15.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	27.9	25.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.7	3.7
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.6	3.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.5	7.2
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.1	4.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.8	3.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.9	8.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.1	10.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.3	5.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.3	16.8
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	11.3	16.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	6.1	4.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.9	20.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.9	75.5
88. Women having a mobile phone that they themselves use (%)	58.9	39.9
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	79.5	76.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.2	8.9
91. Men age 15 years and above who use any kind of tobacco (%)	43.4	46.1
92. Women age 15 years and above who consume alcohol (%)	0.2	0.6
93. Men age 15 years and above who consume alcohol (%)	20.3	20.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Aligarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.7	10.1
2. Population below age 15 years (%)	30.9	32.1
3. Population age 60 years and above (%)	10.0	9.9
4. Population living in households with electricity (%)	95.5	95.3
5. Population living in households with an improved drinking-water source ¹ (%)	99.5	100.0
6. Households using iodized salt (%)	91.0	96.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.1	17.6
8. Households with any usual member having a bank account/post office account (%)	97.0	97.4
9. Female population age 6 years and above who ever attended school (%)	71.3	68.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.9	11.6
11. Children age 2-4 years who attended pre-school (%)	28.6	12.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	41.6	37.5
13. Men with 10 or more years of schooling (%)	52.4	51.4
14. Women who have ever used the internet (%)	60.5	35.9
15. Men who have ever used the internet (%)	78.2	60.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	18.5	15.8
17. Men age 25-29 years married before age 21 years (%)	(31.3)	(18.2)
18. Three or more births in the 5 years preceding the survey (%)	4.4	3.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	6.7	4.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	70.0	68.2
21. Any modern method ² (%)	37.1	39.7
22. Any traditional method (%)	32.9	28.5
23. Female sterilization (%)	13.5	12.7
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	7.7	6.4
26. Unmet need for spacing ³ (%)	3.9	1.6
27. Unmet need for limiting ³ (%)	3.8	4.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.6	69.5
29. Mothers who had any antenatal care visits (%)	96.8	98.5
30. Mothers who had at least 4 antenatal care visits (%)	39.5	44.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	89.9	93.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	23.5	25.2
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	15.4	12.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	88.6	98.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	84.6	82.6
36. Institutional births in public facility (%)	56.0	49.2
37. Births attended by skilled health personnel ⁵ (%)	86.7	85.5
38. Births delivered by caesarean section (%)	16.5	18.7
39. Births in a private health facility that were delivered by caesarean section (%)	47.3	45.6
40. Births in a public health facility that were delivered by caesarean section (%)	5.2	7.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.
() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Aligarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	76.0	82.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	2.7	1.4
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.6	80.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	77.9	70.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	88.9	83.9
46. Children age 12-23 months who have received any vaccine (%)	99.0	93.3
47. Children age 12-23 months who have received BCG (%)	98.0	93.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	78.8	72.4
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	99.0	78.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	99.0	81.8
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	65.0	69.9
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	57.0	50.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	81.8	69.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	75.1	69.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.9	89.7
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.1	3.7
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.6	5.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	5.1	3.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	70.7	(67.6)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	53.9	43.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	58.8	(57.0)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	93.2	(91.9)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	85.0	(86.0)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(58.3)	(46.5)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.6	9.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(18.2)	(12.3)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.7	9.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	34.6	35.0
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	10.1	10.9
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.8	4.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	23.9	26.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.0	2.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Aligarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.8	18.6
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.1	29.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	4.7	7.4
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.3	5.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	10.5	13.1
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.7	7.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.0	6.1
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.3	14.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.9	11.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.1	4.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.5	18.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.2	16.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.4	4.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.7	22.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	79.6	79.1
88. Women having a mobile phone that they themselves use (%)	53.9	42.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	73.8	79.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.9	5.1
91. Men age 15 years and above who use any kind of tobacco (%)	38.4	37.0
92. Women age 15 years and above who consume alcohol (%)	0.3	0.1
93. Men age 15 years and above who consume alcohol (%)	19.8	16.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Ambedkar Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.7	8.0
2. Population below age 15 years (%)	27.5	29.2
3. Population age 60 years and above (%)	11.4	11.6
4. Population living in households with electricity (%)	95.4	94.2
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	93.0	98.3
7. Households with any usual member covered under a health insurance/financing scheme (%)	43.4	23.0
8. Households with any usual member having a bank account/post office account (%)	99.0	98.1
9. Female population age 6 years and above who ever attended school (%)	75.5	71.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.9	13.1
11. Children age 2-4 years who attended pre-school (%)	35.9	9.6
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	57.0	53.0
13. Men with 10 or more years of schooling (%)	57.3	48.8
14. Women who have ever used the internet (%)	73.5	27.7
15. Men who have ever used the internet (%)	77.6	61.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	6.4	5.5
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	1.2	2.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.5	0.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	58.3	48.9
21. Any modern method ² (%)	22.5	20.7
22. Any traditional method (%)	35.8	28.2
23. Female sterilization (%)	10.8	12.7
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.7	22.9
26. Unmet need for spacing ³ (%)	5.4	7.0
27. Unmet need for limiting ³ (%)	4.3	15.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	72.6	61.1
29. Mothers who had any antenatal care visits (%)	97.1	96.6
30. Mothers who had at least 4 antenatal care visits (%)	43.4	44.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.9	94.6
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	35.7	25.2
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	25.7	11.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.5	97.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	92.8	92.2
36. Institutional births in public facility (%)	45.8	49.8
37. Births attended by skilled health personnel ⁵ (%)	93.5	87.9
38. Births delivered by caesarean section (%)	26.8	16.0
39. Births in a private health facility that were delivered by caesarean section (%)	48.9	29.0
40. Births in a public health facility that were delivered by caesarean section (%)	8.4	7.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Ambedkar Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.1	76.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.9	74.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	87.0	75.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	89.1	(91.3)
46. Children age 12-23 months who have received any vaccine (%)	96.7	90.0
47. Children age 12-23 months who have received BCG (%)	95.3	90.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	96.7	76.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	88.4	84.9
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.1	83.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	71.3	62.7
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	68.8	38.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	88.4	55.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	77.7	80.0
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.4	91.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.6	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	20.0	3.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.5	1.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	61.7	(62.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	31.9	20.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(12.1)	(68.0)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.4)	(95.7)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(55.4)	(91.1)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	40.1	2.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	42.2	5.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	23.1	31.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	20.2	17.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.7	3.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	35.0	29.2
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.9	1.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Ambedkar Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	26.8	26.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.1	20.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.5	7.2
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.8	6.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.6	14.6
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.0	8.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.2	9.3
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	22.1	18.4
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.0	9.0
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.2	5.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.5	17.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.5	13.6
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.6	6.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.6	22.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	91.6	81.0
88. Women having a mobile phone that they themselves use (%)	61.2	47.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	71.0	77.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	12.0	12.8
91. Men age 15 years and above who use any kind of tobacco (%)	49.4	52.9
92. Women age 15 years and above who consume alcohol (%)	0.1	0.2
93. Men age 15 years and above who consume alcohol (%)	21.0	16.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Amethi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.1	10.4
2. Population below age 15 years (%)	30.6	32.1
3. Population age 60 years and above (%)	12.0	11.7
4. Population living in households with electricity (%)	95.9	92.3
5. Population living in households with an improved drinking-water source ¹ (%)	98.6	98.7
6. Households using iodized salt (%)	82.0	97.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	44.4	20.9
8. Households with any usual member having a bank account/post office account (%)	99.3	97.9
9. Female population age 6 years and above who ever attended school (%)	67.1	66.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	25.4	16.4
11. Children age 2-4 years who attended pre-school (%)	31.3	7.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	39.8	34.6
13. Men with 10 or more years of schooling (%)	35.2	58.4
14. Women who have ever used the internet (%)	63.6	35.2
15. Men who have ever used the internet (%)	65.0	74.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	9.2	14.9
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	2.8	3.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	0.9	1.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	55.1	48.4
21. Any modern method ² (%)	24.7	30.5
22. Any traditional method (%)	30.4	17.9
23. Female sterilization (%)	10.9	14.4
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	15.0	19.9
26. Unmet need for spacing ³ (%)	7.0	7.0
27. Unmet need for limiting ³ (%)	8.0	12.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	70.5	72.6
29. Mothers who had any antenatal care visits (%)	98.0	96.0
30. Mothers who had at least 4 antenatal care visits (%)	47.8	31.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.5	93.5
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.0	15.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.9	10.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.5	96.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	92.9	90.8
36. Institutional births in public facility (%)	71.1	71.3
37. Births attended by skilled health personnel ⁵ (%)	93.1	91.4
38. Births delivered by caesarean section (%)	10.7	11.1
39. Births in a private health facility that were delivered by caesarean section (%)	33.7	40.3
40. Births in a public health facility that were delivered by caesarean section (%)	4.6	4.5

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Amethi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	86.1	76.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	(12.7)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	87.3	73.7
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	91.3	75.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	93.9	86.8
46. Children age 12-23 months who have received any vaccine (%)	99.1	96.8
47. Children age 12-23 months who have received BCG (%)	97.2	96.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	93.2	78.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	97.2	82.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	97.2	85.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	77.7	70.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	77.7	58.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	95.0	65.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	73.8	80.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	95.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.6	4.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.2	3.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	70.6	57.3
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	55.6	21.6
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(72.0)	72.6
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(95.1)	96.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(93.1)	81.8
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(42.9)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	16.0	5.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.5	5.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.4	35.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	7.3	19.9
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.3	5.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	23.6	33.1
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.8	1.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Amethi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	21.5	21.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	24.0	23.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.9	5.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.1	6.5
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.6	12.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.8	5.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.9	5.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.2	12.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.6	12.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.9	5.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.6	19.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.5	14.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	6.6	5.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.9	21.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.5	70.4
88. Women having a mobile phone that they themselves use (%)	56.0	51.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	66.0	71.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	9.1	11.5
91. Men age 15 years and above who use any kind of tobacco (%)	52.5	49.3
92. Women age 15 years and above who consume alcohol (%)	0.4	0.4
93. Men age 15 years and above who consume alcohol (%)	13.3	9.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Amroha, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.3	11.0
2. Population below age 15 years (%)	30.3	32.5
3. Population age 60 years and above (%)	8.7	6.6
4. Population living in households with electricity (%)	96.9	96.4
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.7
6. Households using iodized salt (%)	78.4	90.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	32.2	8.3
8. Households with any usual member having a bank account/post office account (%)	98.7	96.8
9. Female population age 6 years and above who ever attended school (%)	65.7	63.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.2	8.9
11. Children age 2-4 years who attended pre-school (%)	35.0	17.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	41.4	35.4
13. Men with 10 or more years of schooling (%)	45.3	44.8
14. Women who have ever used the internet (%)	67.2	25.6
15. Men who have ever used the internet (%)	79.0	65.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.7	11.6
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.8	4.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.2	4.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	69.9	65.7
21. Any modern method ² (%)	39.5	45.3
22. Any traditional method (%)	30.4	20.4
23. Female sterilization (%)	10.9	13.0
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.5	6.8
26. Unmet need for spacing ³ (%)	3.9	2.4
27. Unmet need for limiting ³ (%)	4.6	4.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	71.2	70.4
29. Mothers who had any antenatal care visits (%)	97.4	95.8
30. Mothers who had at least 4 antenatal care visits (%)	49.4	43.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.0	87.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	31.4	23.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.0	4.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	90.2	95.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	91.4	81.1
36. Institutional births in public facility (%)	27.2	31.3
37. Births attended by skilled health personnel ⁵ (%)	92.3	86.7
38. Births delivered by caesarean section (%)	37.3	20.3
39. Births in a private health facility that were delivered by caesarean section (%)	53.7	36.4
40. Births in a public health facility that were delivered by caesarean section (%)	10.4	6.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Amroha, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.4	77.7
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(6.2)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	87.6	79.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	94.4	72.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	95.3	91.1
46. Children age 12-23 months who have received any vaccine (%)	98.9	96.0
47. Children age 12-23 months who have received BCG (%)	98.9	95.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	94.4	78.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	98.9	86.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	98.9	84.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	79.4	67.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	46.9	36.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	94.2	61.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	69.6	76.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	75.3	92.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.5	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	18.4	5.5
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.3	0.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	80.2	(79.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	27.3	30.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(6.3)	62.0
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(83.4)	91.6
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(85.2)	86.6
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.4	17.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(8.2)	(13.2)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	13.0	16.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	26.0	42.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	24.7	22.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.4	12.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	38.9	35.4
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	4.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Amroha, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.8	16.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.5	19.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.3	2.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.1	3.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.7	6.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.7	6.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.8	3.9
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.9	11.1
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.2	13.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.4	4.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.4	19.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.2	18.5
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	5.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.0	24.6
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.3	79.7
88. Women having a mobile phone that they themselves use (%)	50.0	42.2
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	68.0	74.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.1	4.4
91. Men age 15 years and above who use any kind of tobacco (%)	35.8	38.9
92. Women age 15 years and above who consume alcohol (%)	0.4	0.4
93. Men age 15 years and above who consume alcohol (%)	17.1	14.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Auraiya, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.4	9.4
2. Population below age 15 years (%)	29.5	29.3
3. Population age 60 years and above (%)	12.7	10.1
4. Population living in households with electricity (%)	96.0	94.0
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.5
6. Households using iodized salt (%)	43.4	93.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.3	18.4
8. Households with any usual member having a bank account/post office account (%)	98.4	96.7
9. Female population age 6 years and above who ever attended school (%)	77.4	75.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.2	11.3
11. Children age 2-4 years who attended pre-school (%)	32.8	10.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	47.9	46.2
13. Men with 10 or more years of schooling (%)	53.6	52.2
14. Women who have ever used the internet (%)	72.5	27.3
15. Men who have ever used the internet (%)	65.2	60.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	21.1	18.1
17. Men age 25-29 years married before age 21 years (%)	(17.4)	*
18. Three or more births in the 5 years preceding the survey (%)	3.7	2.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.8	4.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	66.5	51.4
21. Any modern method ² (%)	31.3	32.3
22. Any traditional method (%)	35.2	19.1
23. Female sterilization (%)	12.0	13.3
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.2	18.7
26. Unmet need for spacing ³ (%)	4.0	6.2
27. Unmet need for limiting ³ (%)	6.2	12.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	73.0	72.2
29. Mothers who had any antenatal care visits (%)	98.0	96.9
30. Mothers who had at least 4 antenatal care visits (%)	44.3	50.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	96.5	91.3
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.4	23.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.7	10.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.5	98.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	84.5	90.1
36. Institutional births in public facility (%)	59.9	72.8
37. Births attended by skilled health personnel ⁵ (%)	87.0	90.9
38. Births delivered by caesarean section (%)	13.6	8.0
39. Births in a private health facility that were delivered by caesarean section (%)	51.0	36.1
40. Births in a public health facility that were delivered by caesarean section (%)	1.8	2.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Auraiya, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.6	77.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.3)	(5.5)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.4	73.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	84.5	82.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(90.9)	84.3
46. Children age 12-23 months who have received any vaccine (%)	100.0	97.8
47. Children age 12-23 months who have received BCG (%)	96.5	95.6
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	86.2	84.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	94.2	89.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	96.4	89.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	74.7	47.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	82.4	70.3
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	94.2	74.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.9	84.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	99.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.9
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	14.4	6.5
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9	3.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	74.4	(68.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	60.1	19.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(11.7)	(69.4)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	(94.7)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(87.9)	(71.5)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(7.9)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.3	5.2
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	7.7	6.1
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	36.2	39.7
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	24.6	19.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.4	9.2
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	40.9	32.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	7.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Auraiya, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.4	18.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	23.9	19.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.8	3.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.0	3.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.8	6.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.6	3.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.0	2.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.0	7.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.2	14.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.3	5.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.2	20.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.0	16.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	5.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.5	22.1
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	79.7	67.4
88. Women having a mobile phone that they themselves use (%)	55.6	42.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	61.5	77.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	6.2	9.3
91. Men age 15 years and above who use any kind of tobacco (%)	49.7	46.8
92. Women age 15 years and above who consume alcohol (%)	0.2	0.3
93. Men age 15 years and above who consume alcohol (%)	19.7	11.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Ayodhya, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.7	9.7
2. Population below age 15 years (%)	27.5	29.1
3. Population age 60 years and above (%)	11.0	11.1
4. Population living in households with electricity (%)	95.4	90.3
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	91.1	94.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	40.5	19.8
8. Households with any usual member having a bank account/post office account (%)	98.6	98.8
9. Female population age 6 years and above who ever attended school (%)	72.5	71.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	23.1	12.3
11. Children age 2-4 years who attended pre-school (%)	36.5	7.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	49.1	49.0
13. Men with 10 or more years of schooling (%)	43.6	55.7
14. Women who have ever used the internet (%)	72.5	29.1
15. Men who have ever used the internet (%)	79.1	57.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	6.1	7.6
17. Men age 25-29 years married before age 21 years (%)	(16.3)	*
18. Three or more births in the 5 years preceding the survey (%)	1.5	1.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.7	2.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	59.8	46.0
21. Any modern method ² (%)	23.0	21.0
22. Any traditional method (%)	36.8	25.0
23. Female sterilization (%)	9.8	11.3
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.2	25.9
26. Unmet need for spacing ³ (%)	5.1	12.2
27. Unmet need for limiting ³ (%)	4.1	13.7
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	74.7	48.4
29. Mothers who had any antenatal care visits (%)	96.9	95.0
30. Mothers who had at least 4 antenatal care visits (%)	37.5	33.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	91.6	94.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.8	20.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	18.2	9.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	89.9	94.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	86.5	89.1
36. Institutional births in public facility (%)	56.4	63.3
37. Births attended by skilled health personnel ⁵ (%)	87.1	91.6
38. Births delivered by caesarean section (%)	21.4	21.2
39. Births in a private health facility that were delivered by caesarean section (%)	55.3	59.3
40. Births in a public health facility that were delivered by caesarean section (%)	8.5	9.3

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

1Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

2Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

3Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Ayodhya, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.8	75.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.9	71.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	79.9	68.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	86.1	82.9
46. Children age 12-23 months who have received any vaccine (%)	94.8	91.1
47. Children age 12-23 months who have received BCG (%)	90.0	88.6
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	94.8	71.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.8	86.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.1	82.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(75.7)	43.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	62.5	30.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	79.9	62.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	75.6	65.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.1	97.3
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.0	1.4
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	17.6	2.5
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9	0.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	57.5	(88.6)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	42.4	12.5
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(11.7)	64.2
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	90.1
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(69.4)	81.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(71.4)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	32.4	3.5
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	34.2	2.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	29.0	30.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	25.9	12.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.4	2.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	40.0	27.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	2.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Ayodhya, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	26.4	20.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.0	24.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.8	7.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.7	8.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.7	17.0
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.8	10.6
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.9	9.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	22.4	20.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.6	8.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	4.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.2	14.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.3	10.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.1	4.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.2	17.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	88.4	83.7
88. Women having a mobile phone that they themselves use (%)	57.1	52.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	71.0	71.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	9.4	8.7
91. Men age 15 years and above who use any kind of tobacco (%)	48.3	41.2
92. Women age 15 years and above who consume alcohol (%)	0.3	0.2
93. Men age 15 years and above who consume alcohol (%)	19.0	10.6

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Azamgarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.7	8.9
2. Population below age 15 years (%)	28.0	29.3
3. Population age 60 years and above (%)	11.6	12.2
4. Population living in households with electricity (%)	98.8	96.5
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	100.0
6. Households using iodized salt (%)	96.2	99.3
7. Households with any usual member covered under a health insurance/financing scheme (%)	38.3	9.0
8. Households with any usual member having a bank account/post office account (%)	98.8	97.4
9. Female population age 6 years and above who ever attended school (%)	75.2	71.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.4	17.1
11. Children age 2-4 years who attended pre-school (%)	30.0	11.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	55.9	52.6
13. Men with 10 or more years of schooling (%)	67.9	65.5
14. Women who have ever used the internet (%)	79.0	31.9
15. Men who have ever used the internet (%)	92.0	70.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	4.7	12.0
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	0.7	5.2
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	0.7	0.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.6	52.7
21. Any modern method ² (%)	28.8	31.4
22. Any traditional method (%)	32.8	21.3
23. Female sterilization (%)	21.4	23.3
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.8	20.7
26. Unmet need for spacing ³ (%)	5.5	8.1
27. Unmet need for limiting ³ (%)	5.3	12.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	71.4	53.7
29. Mothers who had any antenatal care visits (%)	96.6	96.9
30. Mothers who had at least 4 antenatal care visits (%)	36.8	38.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.0	94.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	33.5	19.4
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	23.7	9.9
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	92.0	96.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	96.3	93.2
36. Institutional births in public facility (%)	39.6	44.7
37. Births attended by skilled health personnel ⁵ (%)	96.5	89.8
38. Births delivered by caesarean section (%)	25.8	16.8
39. Births in a private health facility that were delivered by caesarean section (%)	38.9	27.4
40. Births in a public health facility that were delivered by caesarean section (%)	9.3	7.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.
() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Azamgarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.0	72.9
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	84.8	72.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	70.0	64.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	82.4	80.3
46. Children age 12-23 months who have received any vaccine (%)	94.3	95.7
47. Children age 12-23 months who have received BCG (%)	91.4	92.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	72.7	69.8
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	76.7	82.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	85.1	80.0
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	64.9	50.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	43.8	28.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	74.3	43.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	70.8	67.8
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	88.7	86.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	11.3	8.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.7	2.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.1	1.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	73.8	(77.5)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	31.7	18.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(17.4)	61.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(94.6)	100.0
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(67.1)	91.1
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(67.1)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	24.3	0.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	28.4	0.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	20.3	33.4
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	13.7	14.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	2.0	4.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	23.7	28.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.3	0.5

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Azamgarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.5	22.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	27.8	21.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.7	6.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.7	6.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.7	14.2
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.8	6.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.1	10.1
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.9	17.1
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.7	11.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	5.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.0	19.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.9	18.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.6	8.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.2	28.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.0	83.3
88. Women having a mobile phone that they themselves use (%)	65.2	48.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	71.9	79.3
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	8.6	9.4
91. Men age 15 years and above who use any kind of tobacco (%)	45.0	44.1
92. Women age 15 years and above who consume alcohol (%)	0.1	0.3
93. Men age 15 years and above who consume alcohol (%)	24.0	14.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Baghpat, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.6	8.9
2. Population below age 15 years (%)	27.4	29.8
3. Population age 60 years and above (%)	13.2	10.4
4. Population living in households with electricity (%)	98.8	98.5
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.9
6. Households using iodized salt (%)	97.7	96.6
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.7	12.5
8. Households with any usual member having a bank account/post office account (%)	99.3	97.1
9. Female population age 6 years and above who ever attended school (%)	72.5	71.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	23.3	11.2
11. Children age 2-4 years who attended pre-school (%)	46.1	15.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	50.2	44.0
13. Men with 10 or more years of schooling (%)	54.4	60.2
14. Women who have ever used the internet (%)	83.2	46.4
15. Men who have ever used the internet (%)	88.7	66.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.6	8.1
17. Men age 25-29 years married before age 21 years (%)	(25.1)	(18.8)
18. Three or more births in the 5 years preceding the survey (%)	4.4	5.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.0	2.7
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	67.7	71.7
21. Any modern method ² (%)	39.5	45.2
22. Any traditional method (%)	28.2	26.5
23. Female sterilization (%)	13.0	15.6
24. Male sterilization (%)	0.4	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	7.9	5.2
26. Unmet need for spacing ³ (%)	2.5	2.2
27. Unmet need for limiting ³ (%)	5.4	3.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	76.0	79.4
29. Mothers who had any antenatal care visits (%)	99.0	95.1
30. Mothers who had at least 4 antenatal care visits (%)	63.8	42.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.8	93.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	41.0	33.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	23.1	8.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.6	96.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	85.8	80.9
36. Institutional births in public facility (%)	31.1	44.0
37. Births attended by skilled health personnel ⁵ (%)	88.9	82.3
38. Births delivered by caesarean section (%)	22.1	12.6
39. Births in a private health facility that were delivered by caesarean section (%)	37.2	23.7
40. Births in a public health facility that were delivered by caesarean section (%)	5.5	8.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Baghpat, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.9	79.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(11.2)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.6	82.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	88.1	83.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	94.1	93.0
46. Children age 12-23 months who have received any vaccine (%)	97.3	94.1
47. Children age 12-23 months who have received BCG (%)	97.3	92.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	88.1	86.4
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	97.3	91.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	97.3	87.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	82.1	62.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	73.9	81.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	92.1	81.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	70.3	73.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	88.5	92.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.5	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	12.1	4.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.5	0.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	74.4	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	30.8	18.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(18.8)	58.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(93.7)	92.0
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(89.7)	87.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.0	10.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(6.8)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.7	9.1
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	25.3	25.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.4	10.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.2	3.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	32.9	26.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.4	1.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Baghpat, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.2	14.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	32.9	34.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.2	5.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.6	5.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.0	11.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	12.4	6.0
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.1	6.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	22.9	12.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.5	13.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.7	6.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.1	21.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.8	18.0
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.4	6.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.8	25.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	79.0	80.0
88. Women having a mobile phone that they themselves use (%)	56.5	46.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	79.7	81.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.7	4.7
91. Men age 15 years and above who use any kind of tobacco (%)	31.4	31.8
92. Women age 15 years and above who consume alcohol (%)	0.4	0.3
93. Men age 15 years and above who consume alcohol (%)	11.5	12.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Bahraich, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	12.9	13.1
2. Population below age 15 years (%)	39.2	41.0
3. Population age 60 years and above (%)	8.2	8.5
4. Population living in households with electricity (%)	90.6	77.8
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	100.0
6. Households using iodized salt (%)	74.5	85.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	29.8	10.1
8. Households with any usual member having a bank account/post office account (%)	98.9	94.5
9. Female population age 6 years and above who ever attended school (%)	58.0	48.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.0	12.6
11. Children age 2-4 years who attended pre-school (%)	13.9	11.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	21.1	14.4
13. Men with 10 or more years of schooling (%)	27.4	22.1
14. Women who have ever used the internet (%)	65.7	8.6
15. Men who have ever used the internet (%)	79.0	32.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	31.5	37.5
17. Men age 25-29 years married before age 21 years (%)	(31.4)	*
18. Three or more births in the 5 years preceding the survey (%)	4.3	4.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	9.5	8.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	46.1	38.4
21. Any modern method ² (%)	20.4	33.4
22. Any traditional method (%)	25.7	5.0
23. Female sterilization (%)	7.6	5.8
24. Male sterilization (%)	0.0	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	19.5	27.6
26. Unmet need for spacing ³ (%)	7.7	11.1
27. Unmet need for limiting ³ (%)	11.8	16.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	62.7	40.2
29. Mothers who had any antenatal care visits (%)	96.9	81.9
30. Mothers who had at least 4 antenatal care visits (%)	70.3	34.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.4	86.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	23.6	22.4
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	12.4	14.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.3	85.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	79.9	67.7
36. Institutional births in public facility (%)	63.4	58.6
37. Births attended by skilled health personnel ⁵ (%)	82.7	70.7
38. Births delivered by caesarean section (%)	8.8	5.2
39. Births in a private health facility that were delivered by caesarean section (%)	33.6	24.2
40. Births in a public health facility that were delivered by caesarean section (%)	5.2	5.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Bahraich, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	67.1	52.4
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.1	4.7
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	71.7	54.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	83.1	51.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.3	62.2
46. Children age 12-23 months who have received any vaccine (%)	91.9	92.9
47. Children age 12-23 months who have received BCG (%)	88.9	89.4
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	91.1	65.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.5	64.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	86.8	70.3
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	82.8	39.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	54.7	44.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	81.4	35.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	76.3	69.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	96.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	21.9	9.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.1	6.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	51.0	60.8
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	47.8	17.0
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	8.2	70.5
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	95.5	93.8
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	61.5	82.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(70.8)	(32.3)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	24.0	9.5
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(22.1)	(11.5)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	23.8	9.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	41.7	52.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.2	14.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.2	7.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	43.2	38.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	4.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Bahraich, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	27.2	29.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	19.8	12.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.2	2.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.0	3.1
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.6	6.1
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.5	4.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.2	2.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.0	8.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.2	13.4
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.9	7.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.1	24.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.6	18.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.9	8.5
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.9	28.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	79.3	61.9
88. Women having a mobile phone that they themselves use (%)	54.0	39.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	50.4	48.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	25.9	26.6
91. Men age 15 years and above who use any kind of tobacco (%)	61.4	59.0
92. Women age 15 years and above who consume alcohol (%)	0.3	0.6
93. Men age 15 years and above who consume alcohol (%)	14.2	12.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Ballia, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.7	7.9
2. Population below age 15 years (%)	29.0	29.0
3. Population age 60 years and above (%)	11.3	11.1
4. Population living in households with electricity (%)	96.7	89.1
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	99.8
6. Households using iodized salt (%)	94.9	96.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	46.9	10.6
8. Households with any usual member having a bank account/post office account (%)	98.2	97.3
9. Female population age 6 years and above who ever attended school (%)	75.3	68.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.1	8.9
11. Children age 2-4 years who attended pre-school (%)	34.4	15.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	55.6	50.1
13. Men with 10 or more years of schooling (%)	63.5	60.7
14. Women who have ever used the internet (%)	64.5	29.3
15. Men who have ever used the internet (%)	78.7	53.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	9.7	9.9
17. Men age 25-29 years married before age 21 years (%)	(23.2)	*
18. Three or more births in the 5 years preceding the survey (%)	3.5	3.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	0.4	1.7
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.5	55.2
21. Any modern method ² (%)	33.5	41.9
22. Any traditional method (%)	28.0	13.3
23. Female sterilization (%)	22.1	15.5
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.7	20.2
26. Unmet need for spacing ³ (%)	4.3	5.7
27. Unmet need for limiting ³ (%)	7.4	14.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	65.6	64.0
29. Mothers who had any antenatal care visits (%)	90.5	96.5
30. Mothers who had at least 4 antenatal care visits (%)	31.8	30.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.9	90.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.5	15.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.8	2.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.7	95.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.0	90.0
36. Institutional births in public facility (%)	58.0	73.9
37. Births attended by skilled health personnel ⁵ (%)	89.2	93.1
38. Births delivered by caesarean section (%)	18.0	7.4
39. Births in a private health facility that were delivered by caesarean section (%)	52.6	32.4
40. Births in a public health facility that were delivered by caesarean section (%)	2.0	2.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Ballia, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.1	76.6
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(17.6)	(8.7)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.2	72.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	80.2	62.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	91.2	65.7
46. Children age 12-23 months who have received any vaccine (%)	96.4	96.8
47. Children age 12-23 months who have received BCG (%)	96.4	96.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	82.6	67.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.9	69.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.3	80.3
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	68.2	49.7
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	58.3	37.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	87.7	31.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	73.8	75.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.2	94.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.8	2.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	7.5	10.3
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	7.2	7.1
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	54.8	55.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	45.6	8.0
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(71.5)	(69.8)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(93.5)	(91.3)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(82.2)	(77.2)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	21.9	1.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.7	0.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	25.9	43.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	11.0	21.9
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.6	10.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	21.2	42.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.8	2.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Ballia, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.9	16.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	30.0	16.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.4	5.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.9	6.1
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.7	12.2
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.9	6.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.4	6.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.6	14.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.6	10.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.1	3.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.0	17.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	16.3	15.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	8.2	3.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	26.9	20.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	88.3	79.0
88. Women having a mobile phone that they themselves use (%)	63.4	52.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	66.7	71.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.0	2.2
91. Men age 15 years and above who use any kind of tobacco (%)	39.3	35.6
92. Women age 15 years and above who consume alcohol (%)	0.5	0.5
93. Men age 15 years and above who consume alcohol (%)	20.4	9.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Balrampur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.8	11.9
2. Population below age 15 years (%)	37.2	38.9
3. Population age 60 years and above (%)	8.2	10.6
4. Population living in households with electricity (%)	89.8	78.4
5. Population living in households with an improved drinking-water source ¹ (%)	99.6	99.8
6. Households using iodized salt (%)	87.0	96.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	30.4	13.0
8. Households with any usual member having a bank account/post office account (%)	99.4	96.1
9. Female population age 6 years and above who ever attended school (%)	55.6	53.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	23.2	11.4
11. Children age 2-4 years who attended pre-school (%)	13.5	10.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	22.0	16.8
13. Men with 10 or more years of schooling (%)	33.8	34.9
14. Women who have ever used the internet (%)	73.3	19.7
15. Men who have ever used the internet (%)	85.2	44.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	21.4	35.0
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	2.9	3.2
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.7	4.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	37.4	49.0
21. Any modern method ² (%)	12.0	39.4
22. Any traditional method (%)	25.4	9.6
23. Female sterilization (%)	3.7	6.6
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	23.9	22.3
26. Unmet need for spacing ³ (%)	9.8	7.8
27. Unmet need for limiting ³ (%)	14.1	14.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	61.4	55.2
29. Mothers who had any antenatal care visits (%)	96.0	93.4
30. Mothers who had at least 4 antenatal care visits (%)	48.1	41.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	91.3	91.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	18.6	23.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	12.3	12.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.1	95.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	70.7	69.7
36. Institutional births in public facility (%)	47.7	58.0
37. Births attended by skilled health personnel ⁵ (%)	72.1	67.4
38. Births delivered by caesarean section (%)	10.8	5.3
39. Births in a private health facility that were delivered by caesarean section (%)	31.2	32.4
40. Births in a public health facility that were delivered by caesarean section (%)	7.6	2.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Balrampur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	63.6	62.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.7	2.3
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	62.2	61.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	77.2	57.4
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	88.3	67.9
46. Children age 12-23 months who have received any vaccine (%)	90.4	88.7
47. Children age 12-23 months who have received BCG (%)	87.5	86.4
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	80.0	64.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	82.2	67.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	86.0	74.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	65.5	49.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	48.1	42.5
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	79.4	34.0
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	50.3	76.2
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	97.3	88.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.8	0.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	17.4	5.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9	5.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	53.5	64.3
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	41.6	14.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	12.8	70.6
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	96.1	98.8
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	68.1	80.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(68.5)	(41.3)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	24.3	7.4
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(15.6)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	23.0	6.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	36.5	41.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	26.0	24.9
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.0	14.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	43.2	37.2
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	3.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Balrampur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	28.4	22.2
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	23.3	14.8
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.2	3.3
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.9	4.0
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.3	8.2
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	14.4	4.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	12.4	3.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	27.7	8.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.2	13.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.1	7.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.7	23.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.2	16.6
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.3	7.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.0	25.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	75.4	72.4
88. Women having a mobile phone that they themselves use (%)	60.6	56.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	53.5	53.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	24.8	23.6
91. Men age 15 years and above who use any kind of tobacco (%)	60.8	59.8
92. Women age 15 years and above who consume alcohol (%)	0.1	0.4
93. Men age 15 years and above who consume alcohol (%)	11.4	10.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Banda, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.8	9.1
2. Population below age 15 years (%)	31.6	30.6
3. Population age 60 years and above (%)	11.4	10.4
4. Population living in households with electricity (%)	97.7	89.2
5. Population living in households with an improved drinking-water source ¹ (%)	99.2	98.2
6. Households using iodized salt (%)	65.9	90.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.3	13.3
8. Households with any usual member having a bank account/post office account (%)	98.2	94.5
9. Female population age 6 years and above who ever attended school (%)	64.8	61.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.5	8.2
11. Children age 2-4 years who attended pre-school (%)	22.0	8.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	31.4	27.1
13. Men with 10 or more years of schooling (%)	35.3	33.0
14. Women who have ever used the internet (%)	60.9	15.4
15. Men who have ever used the internet (%)	78.6	55.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	17.9	19.0
17. Men age 25-29 years married before age 21 years (%)	*	(32.1)
18. Three or more births in the 5 years preceding the survey (%)	2.6	5.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	7.1	2.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	64.3	62.4
21. Any modern method ² (%)	31.1	45.6
22. Any traditional method (%)	33.2	16.8
23. Female sterilization (%)	18.0	16.7
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	14.1	15.3
26. Unmet need for spacing ³ (%)	6.5	5.1
27. Unmet need for limiting ³ (%)	7.6	10.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	70.1	66.9
29. Mothers who had any antenatal care visits (%)	97.0	94.1
30. Mothers who had at least 4 antenatal care visits (%)	50.5	39.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.0	90.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.6	18.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.5	2.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.0	95.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.0	89.8
36. Institutional births in public facility (%)	80.7	85.9
37. Births attended by skilled health personnel ⁵ (%)	91.6	92.5
38. Births delivered by caesarean section (%)	6.7	4.2
39. Births in a private health facility that were delivered by caesarean section (%)	(63.0)	*
40. Births in a public health facility that were delivered by caesarean section (%)	1.8	2.3

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Banda, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	80.5	73.7
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(9.4)	(3.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.8	65.1
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	90.5	62.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	86.6	64.9
46. Children age 12-23 months who have received any vaccine (%)	97.0	96.3
47. Children age 12-23 months who have received BCG (%)	97.0	95.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	97.0	65.6
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.5	73.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.8	80.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	68.9	43.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	85.5	58.5
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	89.4	39.3
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	54.8	81.2
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	97.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	1.2
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.6	2.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	5.3	3.5
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	75.7	(45.7)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	67.4	16.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	39.1	46.7
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	98.2	96.9
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	96.1	81.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(58.4)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.7	5.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.7	9.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	37.7	51.0
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	16.8	25.7
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.5	13.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	39.7	49.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	6.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Banda, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	26.0	25.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	17.9	13.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.4	6.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.1	3.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.9	11.5
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.5	6.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.4	3.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.6	13.1
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.9	12.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.8	4.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.5	20.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.1	12.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	4.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.1	18.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	83.6	69.8
88. Women having a mobile phone that they themselves use (%)	52.4	29.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	57.4	52.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	30.0	19.7
91. Men age 15 years and above who use any kind of tobacco (%)	69.6	63.3
92. Women age 15 years and above who consume alcohol (%)	0.5	0.5
93. Men age 15 years and above who consume alcohol (%)	19.3	12.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Barabanki, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.8	10.3
2. Population below age 15 years (%)	31.9	32.6
3. Population age 60 years and above (%)	8.4	9.7
4. Population living in households with electricity (%)	84.3	74.8
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	98.8
6. Households using iodized salt (%)	80.3	78.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	40.1	14.6
8. Households with any usual member having a bank account/post office account (%)	98.4	92.4
9. Female population age 6 years and above who ever attended school (%)	69.2	65.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.0	12.3
11. Children age 2-4 years who attended pre-school (%)	37.9	21.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	34.3	29.5
13. Men with 10 or more years of schooling (%)	35.2	37.4
14. Women who have ever used the internet (%)	58.9	21.0
15. Men who have ever used the internet (%)	78.3	32.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	19.3	20.4
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	4.0	3.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.9	3.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	57.4	45.4
21. Any modern method ² (%)	33.4	38.3
22. Any traditional method (%)	24.0	7.1
23. Female sterilization (%)	14.1	9.8
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.5	21.4
26. Unmet need for spacing ³ (%)	4.2	7.8
27. Unmet need for limiting ³ (%)	6.3	13.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.3	43.3
29. Mothers who had any antenatal care visits (%)	96.3	68.5
30. Mothers who had at least 4 antenatal care visits (%)	53.8	22.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.0	83.7
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	27.5	12.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.8	5.3
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.7	91.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.9	76.0
36. Institutional births in public facility (%)	65.2	63.1
37. Births attended by skilled health personnel ⁵ (%)	92.1	73.3
38. Births delivered by caesarean section (%)	16.9	9.0
39. Births in a private health facility that were delivered by caesarean section (%)	46.3	29.3
40. Births in a public health facility that were delivered by caesarean section (%)	7.7	8.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Barabanki, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.3	41.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(17.4)	7.2
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.2	42.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	91.3	64.4
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.2	67.2
46. Children age 12-23 months who have received any vaccine (%)	97.2	96.2
47. Children age 12-23 months who have received BCG (%)	96.3	94.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	97.2	74.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	91.3	78.3
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.4	78.2
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	83.7	64.4
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	68.5	47.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.4	30.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	61.2	64.8
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.3	93.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.7	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.0	5.2
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	6.3	4.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	66.0	44.9
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	52.7	30.2
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(63.9)	60.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.7)	96.8
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(93.3)	73.7
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(62.4)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.8	7.2
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(8.9)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.8	6.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	34.3	41.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	8.5	18.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.6	7.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	25.5	31.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	3.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Barabanki, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.7	26.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.6	17.1
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.4	3.2
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.7	4.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.4	9.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.1	5.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.1	5.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.8	12.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.0	11.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.2	6.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.6	19.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	12.2	14.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.9	6.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.9	22.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.3	62.7
88. Women having a mobile phone that they themselves use (%)	46.3	48.0
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	63.0	58.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	11.4	11.2
91. Men age 15 years and above who use any kind of tobacco (%)	55.0	50.2
92. Women age 15 years and above who consume alcohol (%)	0.2	0.2
93. Men age 15 years and above who consume alcohol (%)	15.6	11.3

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Bareilly, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.4	8.8
2. Population below age 15 years (%)	29.2	31.3
3. Population age 60 years and above (%)	8.4	7.1
4. Population living in households with electricity (%)	95.2	92.5
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	94.6	93.5
7. Households with any usual member covered under a health insurance/financing scheme (%)	39.8	18.7
8. Households with any usual member having a bank account/post office account (%)	98.0	96.1
9. Female population age 6 years and above who ever attended school (%)	63.2	55.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.1	11.0
11. Children age 2-4 years who attended pre-school (%)	25.3	11.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	30.0	22.1
13. Men with 10 or more years of schooling (%)	38.9	34.9
14. Women who have ever used the internet (%)	67.3	31.5
15. Men who have ever used the internet (%)	79.5	59.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	10.8	11.4
17. Men age 25-29 years married before age 21 years (%)	(28.3)	*
18. Three or more births in the 5 years preceding the survey (%)	3.8	3.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.6	5.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	62.0	68.8
21. Any modern method ² (%)	29.5	43.5
22. Any traditional method (%)	32.5	25.3
23. Female sterilization (%)	10.2	16.7
24. Male sterilization (%)	0.0	0.4
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	7.6	5.9
26. Unmet need for spacing ³ (%)	3.7	1.8
27. Unmet need for limiting ³ (%)	3.9	4.1
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	79.0	72.2
29. Mothers who had any antenatal care visits (%)	96.3	98.2
30. Mothers who had at least 4 antenatal care visits (%)	52.8	43.1
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.6	96.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.3	18.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	18.2	6.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	90.6	94.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	78.0	75.1
36. Institutional births in public facility (%)	34.4	40.3
37. Births attended by skilled health personnel ⁵ (%)	80.1	77.8
38. Births delivered by caesarean section (%)	23.5	15.5
39. Births in a private health facility that were delivered by caesarean section (%)	49.8	37.9
40. Births in a public health facility that were delivered by caesarean section (%)	5.3	5.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Bareilly, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	70.9	75.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	72.6	75.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	72.5	70.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.0	(91.0)
46. Children age 12-23 months who have received any vaccine (%)	97.0	95.4
47. Children age 12-23 months who have received BCG (%)	93.4	95.4
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	73.6	73.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	96.0	85.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	97.0	88.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	81.2	68.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	70.2	52.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	89.7	57.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	64.2	80.2
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	93.4	91.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	6.6	5.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.6	4.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.8	2.5
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	73.9	(60.6)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	41.9	42.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(10.0)	(58.1)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(79.0)	(95.4)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(96.7)	(95.2)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.7	11.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(9.6)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	6.7	10.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.4	45.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	26.0	15.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.2	5.6
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	36.7	35.2
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	1.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Bareilly, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.5	18.2
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.3	25.0
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.1	3.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.7	5.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.1	9.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.9	5.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.4	5.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.1	11.8
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.9	10.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.7	4.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.8	16.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.9	12.0
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.3	4.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.7	17.6
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.7	77.5
88. Women having a mobile phone that they themselves use (%)	48.2	36.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	64.9	71.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.2	5.5
91. Men age 15 years and above who use any kind of tobacco (%)	36.8	39.3
92. Women age 15 years and above who consume alcohol (%)	0.4	0.1
93. Men age 15 years and above who consume alcohol (%)	15.9	17.8

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Basti, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.7	10.2
2. Population below age 15 years (%)	30.4	31.7
3. Population age 60 years and above (%)	10.9	11.6
4. Population living in households with electricity (%)	98.9	94.6
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	92.9	97.5
7. Households with any usual member covered under a health insurance/financing scheme (%)	44.0	15.8
8. Households with any usual member having a bank account/post office account (%)	98.6	97.8
9. Female population age 6 years and above who ever attended school (%)	70.3	64.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.8	14.4
11. Children age 2-4 years who attended pre-school (%)	28.5	19.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	45.7	40.9
13. Men with 10 or more years of schooling (%)	52.4	47.2
14. Women who have ever used the internet (%)	78.1	36.1
15. Men who have ever used the internet (%)	88.3	67.4
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	6.5	15.9
17. Men age 25-29 years married before age 21 years (%)	(13.7)	*
18. Three or more births in the 5 years preceding the survey (%)	2.0	2.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.5	1.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	59.7	72.0
21. Any modern method ² (%)	22.1	53.8
22. Any traditional method (%)	37.6	18.2
23. Female sterilization (%)	11.4	12.2
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.0	11.1
26. Unmet need for spacing ³ (%)	4.9	4.0
27. Unmet need for limiting ³ (%)	5.1	7.1
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	71.3	45.0
29. Mothers who had any antenatal care visits (%)	95.3	99.2
30. Mothers who had at least 4 antenatal care visits (%)	40.9	31.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.0	96.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.4	15.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	20.7	7.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	88.8	99.0
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.3	93.2
36. Institutional births in public facility (%)	58.7	79.2
37. Births attended by skilled health personnel ⁵ (%)	90.3	92.0
38. Births delivered by caesarean section (%)	28.0	11.6
39. Births in a private health facility that were delivered by caesarean section (%)	63.7	49.5
40. Births in a public health facility that were delivered by caesarean section (%)	14.5	5.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Basti, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.6	83.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.7)	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.9	81.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	81.7	73.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	95.8	77.4
46. Children age 12-23 months who have received any vaccine (%)	97.9	95.7
47. Children age 12-23 months who have received BCG (%)	96.4	93.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	85.4	84.6
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	91.3	76.3
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	94.1	86.0
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	76.2	52.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	53.2	31.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	92.4	45.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	66.4	78.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	96.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	2.2
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	16.5	10.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.4	7.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	46.4	53.7
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	46.3	12.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(15.6)	72.1
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.5)	100.0
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(79.8)	86.1
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(20.9)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	36.5	6.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	36.0	5.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	24.6	35.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	26.6	24.2
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.5	15.2
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	36.6	39.2
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	2.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Basti, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	23.0	20.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.3	16.8
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.4	4.3
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.6	4.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.5	10.0
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.3	5.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	14.2	5.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	26.3	11.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.0	11.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	1.8	6.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.2	19.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.9	16.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	7.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.4	25.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.8	80.4
88. Women having a mobile phone that they themselves use (%)	60.8	54.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	76.7	73.3
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	9.2	10.2
91. Men age 15 years and above who use any kind of tobacco (%)	50.3	51.3
92. Women age 15 years and above who consume alcohol (%)	0.2	0.2
93. Men age 15 years and above who consume alcohol (%)	17.2	14.8

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Bhadohi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.8	11.0
2. Population below age 15 years (%)	30.5	33.7
3. Population age 60 years and above (%)	10.6	9.7
4. Population living in households with electricity (%)	97.5	96.5
5. Population living in households with an improved drinking-water source ¹ (%)	95.4	96.7
6. Households using iodized salt (%)	94.7	98.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.8	18.8
8. Households with any usual member having a bank account/post office account (%)	99.7	97.6
9. Female population age 6 years and above who ever attended school (%)	70.2	67.3
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.2	14.4
11. Children age 2-4 years who attended pre-school (%)	29.5	4.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	47.6	46.5
13. Men with 10 or more years of schooling (%)	56.9	55.0
14. Women who have ever used the internet (%)	73.5	23.9
15. Men who have ever used the internet (%)	84.0	69.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	12.8	19.6
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	4.2	3.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.3	1.7
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.0	53.8
21. Any modern method ² (%)	39.0	40.2
22. Any traditional method (%)	22.0	13.6
23. Female sterilization (%)	31.7	30.9
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	13.5	22.4
26. Unmet need for spacing ³ (%)	6.3	10.1
27. Unmet need for limiting ³ (%)	7.2	12.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.5	53.6
29. Mothers who had any antenatal care visits (%)	95.1	93.0
30. Mothers who had at least 4 antenatal care visits (%)	48.4	25.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.5	89.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	26.0	12.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	15.1	5.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.8	92.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.2	90.3
36. Institutional births in public facility (%)	42.0	47.8
37. Births attended by skilled health personnel ⁵ (%)	92.2	92.3
38. Births delivered by caesarean section (%)	18.2	18.9
39. Births in a private health facility that were delivered by caesarean section (%)	35.7	37.8
40. Births in a public health facility that were delivered by caesarean section (%)	2.3	5.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Bhadohi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	68.5	69.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	70.2	67.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	69.2	62.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.7	72.1
46. Children age 12-23 months who have received any vaccine (%)	87.9	93.6
47. Children age 12-23 months who have received BCG (%)	86.7	87.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	72.4	69.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	74.0	78.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	77.8	80.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	60.8	39.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	16.9	28.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	74.1	59.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	59.0	63.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	96.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	17.5	4.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.4	1.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	55.2	(77.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	34.6	7.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(32.4)	66.6
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.9)	93.5
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(85.4)	84.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(56.4)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.5	3.5
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	(0.0)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.6	2.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	27.1	42.7
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	19.7	9.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.5	2.4
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	33.7	26.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.6	2.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Bhadohi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	19.3	15.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	27.6	26.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.9	4.3
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.7	4.0
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.9	8.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.7	5.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.7	7.3
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	22.9	13.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.2	9.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	4.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.6	14.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.9	15.9
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.4	5.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.9	23.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.6	81.0
88. Women having a mobile phone that they themselves use (%)	60.8	52.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	73.9	74.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	11.1	7.4
91. Men age 15 years and above who use any kind of tobacco (%)	43.7	36.5
92. Women age 15 years and above who consume alcohol (%)	0.1	0.1
93. Men age 15 years and above who consume alcohol (%)	14.2	7.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Bijnor, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.3	9.0
2. Population below age 15 years (%)	30.4	30.7
3. Population age 60 years and above (%)	10.2	8.2
4. Population living in households with electricity (%)	96.2	96.1
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.7
6. Households using iodized salt (%)	89.0	95.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.0	18.8
8. Households with any usual member having a bank account/post office account (%)	97.9	96.5
9. Female population age 6 years and above who ever attended school (%)	70.2	69.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.3	12.2
11. Children age 2-4 years who attended pre-school (%)	43.2	17.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	40.8	39.7
13. Men with 10 or more years of schooling (%)	51.6	42.7
14. Women who have ever used the internet (%)	72.2	41.3
15. Men who have ever used the internet (%)	76.4	66.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.4	5.2
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	2.9	3.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.6	1.2
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	60.9	68.1
21. Any modern method ² (%)	36.8	47.3
22. Any traditional method (%)	24.1	20.8
23. Female sterilization (%)	12.4	11.3
24. Male sterilization (%)	0.3	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.5	5.0
26. Unmet need for spacing ³ (%)	4.7	1.7
27. Unmet need for limiting ³ (%)	5.8	3.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	71.8	68.7
29. Mothers who had any antenatal care visits (%)	99.7	97.6
30. Mothers who had at least 4 antenatal care visits (%)	62.2	48.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.1	93.8
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.8	21.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	13.9	6.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.4	96.0
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	85.8	83.7
36. Institutional births in public facility (%)	28.5	30.6
37. Births attended by skilled health personnel ⁵ (%)	89.6	89.4
38. Births delivered by caesarean section (%)	29.8	22.4
39. Births in a private health facility that were delivered by caesarean section (%)	48.8	40.1
40. Births in a public health facility that were delivered by caesarean section (%)	6.6	3.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Bijnor, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.8	82.4
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(6.9)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.2	81.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	85.1	90.6
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	89.5	96.7
46. Children age 12-23 months who have received any vaccine (%)	96.6	96.7
47. Children age 12-23 months who have received BCG (%)	94.6	96.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	87.0	90.6
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	86.8	92.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	87.5	92.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	75.3	86.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	46.3	67.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	87.7	87.0
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	69.0	78.2
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	86.0	89.6
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.4	1.2
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.9	3.2
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.7	1.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	60.2	(74.8)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	30.2	31.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(36.9)	(57.9)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(79.7)	(93.2)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(97.7)	(80.1)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(47.7)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	13.2	6.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(6.4)	(6.6)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.6	6.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	23.4	36.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.7	9.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.3	2.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	32.3	21.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.1	2.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Bijnor, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.3	17.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.5	28.0
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.9	4.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.3	6.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.9	11.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.7	6.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.3	6.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.6	13.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.5	13.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	4.5
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.4	19.7
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	16.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	5.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.3	24.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.7	85.5
88. Women having a mobile phone that they themselves use (%)	55.5	40.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	82.5	77.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.3	3.6
91. Men age 15 years and above who use any kind of tobacco (%)	37.3	36.5
92. Women age 15 years and above who consume alcohol (%)	0.3	0.2
93. Men age 15 years and above who consume alcohol (%)	21.8	15.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Budaun, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.7	11.3
2. Population below age 15 years (%)	34.7	35.3
3. Population age 60 years and above (%)	6.8	8.6
4. Population living in households with electricity (%)	89.8	80.7
5. Population living in households with an improved drinking-water source ¹ (%)	98.9	99.8
6. Households using iodized salt (%)	85.5	91.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.1	13.6
8. Households with any usual member having a bank account/post office account (%)	97.5	95.5
9. Female population age 6 years and above who ever attended school (%)	59.1	54.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	15.1	14.3
11. Children age 2-4 years who attended pre-school (%)	18.1	20.6
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	25.2	21.6
13. Men with 10 or more years of schooling (%)	40.2	35.2
14. Women who have ever used the internet (%)	59.0	16.7
15. Men who have ever used the internet (%)	74.4	39.2
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	20.9	22.9
17. Men age 25-29 years married before age 21 years (%)	(32.9)	*
18. Three or more births in the 5 years preceding the survey (%)	4.7	4.1
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.5	7.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	65.3	76.9
21. Any modern method ² (%)	25.2	42.0
22. Any traditional method (%)	40.1	34.9
23. Female sterilization (%)	6.4	7.3
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.2	4.3
26. Unmet need for spacing ³ (%)	3.7	1.5
27. Unmet need for limiting ³ (%)	4.5	2.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	61.8	58.5
29. Mothers who had any antenatal care visits (%)	92.0	93.7
30. Mothers who had at least 4 antenatal care visits (%)	39.6	40.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	88.2	90.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	24.4	15.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	10.8	1.9
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	92.5	94.4
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	74.7	72.3
36. Institutional births in public facility (%)	52.2	57.4
37. Births attended by skilled health personnel ⁵ (%)	78.5	73.1
38. Births delivered by caesarean section (%)	10.1	5.7
39. Births in a private health facility that were delivered by caesarean section (%)	32.5	20.9
40. Births in a public health facility that were delivered by caesarean section (%)	5.3	4.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Budaun, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	66.0	62.4
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.9	2.1
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	69.0	59.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	84.5	69.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	89.9	74.5
46. Children age 12-23 months who have received any vaccine (%)	94.6	94.8
47. Children age 12-23 months who have received BCG (%)	93.8	93.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	85.3	73.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	88.3	82.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	89.9	86.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	68.1	61.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	38.9	55.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.5	35.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	64.9	71.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	70.8	97.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.0	1.1
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	12.1	10.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.9	4.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	74.5	50.0
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	44.7	33.6
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	33.0	52.4
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	82.2	87.1
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	81.0	80.5
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(55.5)	(21.3)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.3	8.4
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(3.4)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.7	8.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	38.6	51.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	16.3	18.2
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.4	6.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	33.8	43.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	3.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Budaun, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	23.2	19.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.8	15.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.5	5.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	4.0	3.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	10.6	9.0
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.5	6.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.8	3.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.4	10.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.6	8.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.5	3.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.4	13.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.6	10.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.7	3.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.9	13.8
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.4	67.9
88. Women having a mobile phone that they themselves use (%)	45.2	29.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	53.2	58.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.1	7.1
91. Men age 15 years and above who use any kind of tobacco (%)	42.3	41.9
92. Women age 15 years and above who consume alcohol (%)	0.4	0.5
93. Men age 15 years and above who consume alcohol (%)	17.4	14.0

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Bulandshahr, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.8	10.0
2. Population below age 15 years (%)	29.3	31.3
3. Population age 60 years and above (%)	10.3	10.9
4. Population living in households with electricity (%)	98.0	96.3
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	100.0
6. Households using iodized salt (%)	87.3	95.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.2	20.5
8. Households with any usual member having a bank account/post office account (%)	98.0	98.3
9. Female population age 6 years and above who ever attended school (%)	67.2	67.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.6	11.9
11. Children age 2-4 years who attended pre-school (%)	32.8	15.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	40.0	37.8
13. Men with 10 or more years of schooling (%)	55.5	59.5
14. Women who have ever used the internet (%)	62.2	44.9
15. Men who have ever used the internet (%)	73.7	63.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	10.2	12.8
17. Men age 25-29 years married before age 21 years (%)	(18.4)	*
18. Three or more births in the 5 years preceding the survey (%)	5.5	4.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.2	5.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	66.6	72.9
21. Any modern method ² (%)	30.6	40.4
22. Any traditional method (%)	36.0	32.5
23. Female sterilization (%)	12.5	13.7
24. Male sterilization (%)	0.0	0.3
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.2	5.1
26. Unmet need for spacing ³ (%)	3.5	1.8
27. Unmet need for limiting ³ (%)	4.7	3.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.6	78.3
29. Mothers who had any antenatal care visits (%)	97.0	98.9
30. Mothers who had at least 4 antenatal care visits (%)	45.1	47.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.9	94.8
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	36.2	30.4
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	18.0	7.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.1	97.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.2	83.5
36. Institutional births in public facility (%)	44.1	49.1
37. Births attended by skilled health personnel ⁵ (%)	90.8	86.7
38. Births delivered by caesarean section (%)	18.7	17.0
39. Births in a private health facility that were delivered by caesarean section (%)	38.9	40.0
40. Births in a public health facility that were delivered by caesarean section (%)	2.5	6.5

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Bulandshahr, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.8	82.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	1.8
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.9	82.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	89.5	67.3
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	89.0	84.7
46. Children age 12-23 months who have received any vaccine (%)	96.3	93.9
47. Children age 12-23 months who have received BCG (%)	93.6	92.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	90.9	70.9
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	96.3	80.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	96.3	77.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	67.4	68.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	53.7	79.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	92.2	65.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	71.1	69.6
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.7	92.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.3	0.5
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	16.8	5.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9	1.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	95.2	(82.2)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	41.9	21.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(22.8)	(49.7)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(77.0)	(92.7)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(86.5)	(88.6)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(46.4)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	14.7	9.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(0.0)	(5.1)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.5	8.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	29.2	37.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.7	13.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.0	2.6
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	35.7	26.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.3	2.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Bulandshahr, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.5	15.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	30.8	32.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.0	6.7
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.5	5.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.7	12.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.7	6.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.3	7.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.4	14.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.0	13.0
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	6.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.1	21.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.7	18.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.1	6.5
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.2	26.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.0	77.2
88. Women having a mobile phone that they themselves use (%)	51.6	49.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	70.5	79.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.4	4.1
91. Men age 15 years and above who use any kind of tobacco (%)	35.4	34.4
92. Women age 15 years and above who consume alcohol (%)	0.1	0.2
93. Men age 15 years and above who consume alcohol (%)	18.9	17.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Chandauli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.4	9.7
2. Population below age 15 years (%)	30.1	30.5
3. Population age 60 years and above (%)	11.7	11.3
4. Population living in households with electricity (%)	97.0	91.9
5. Population living in households with an improved drinking-water source ¹ (%)	98.8	97.5
6. Households using iodized salt (%)	95.2	98.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	46.4	28.1
8. Households with any usual member having a bank account/post office account (%)	99.0	97.9
9. Female population age 6 years and above who ever attended school (%)	72.2	68.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.6	12.8
11. Children age 2-4 years who attended pre-school (%)	29.8	10.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	51.1	47.2
13. Men with 10 or more years of schooling (%)	62.5	58.5
14. Women who have ever used the internet (%)	70.4	27.6
15. Men who have ever used the internet (%)	88.3	58.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	16.6	17.2
17. Men age 25-29 years married before age 21 years (%)	(15.3)	(26.7)
18. Three or more births in the 5 years preceding the survey (%)	2.8	3.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.5	2.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	65.2	60.4
21. Any modern method ² (%)	45.0	48.7
22. Any traditional method (%)	20.2	11.7
23. Female sterilization (%)	37.9	34.9
24. Male sterilization (%)	0.4	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.8	15.4
26. Unmet need for spacing ³ (%)	5.3	5.5
27. Unmet need for limiting ³ (%)	5.5	9.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	68.8	69.3
29. Mothers who had any antenatal care visits (%)	95.2	94.8
30. Mothers who had at least 4 antenatal care visits (%)	49.5	32.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	88.5	92.3
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.2	26.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	17.7	15.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.1	96.5
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	86.0	86.2
36. Institutional births in public facility (%)	50.6	53.4
37. Births attended by skilled health personnel ⁵ (%)	86.2	85.1
38. Births delivered by caesarean section (%)	24.8	18.5
39. Births in a private health facility that were delivered by caesarean section (%)	59.1	48.0
40. Births in a public health facility that were delivered by caesarean section (%)	7.7	5.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Chandauli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	66.8	65.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.7)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	70.8	63.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	81.2	70.6
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	91.5	82.6
46. Children age 12-23 months who have received any vaccine (%)	92.4	91.2
47. Children age 12-23 months who have received BCG (%)	92.4	90.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	81.2	72.6
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	85.2	82.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	87.6	82.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	63.3	51.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	36.9	22.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	82.5	64.0
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	58.4	75.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	100.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	14.0	4.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.4	1.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	63.3	(53.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	36.0	28.0
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(8.6)	65.7
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(96.2)	97.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(76.9)	82.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	25.4	3.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(25.0)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	25.3	3.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	27.9	39.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.9	17.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.5	7.4
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	33.4	29.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	0.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Chandauli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.6	21.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	25.1	23.1
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.1	5.2
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.5	4.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.7	10.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.7	6.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.5	4.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.4	11.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.0	11.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.1	5.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.6	18.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.7	15.6
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.3	6.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.9	23.6
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.8	84.7
88. Women having a mobile phone that they themselves use (%)	59.1	44.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	69.1	80.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	11.4	7.1
91. Men age 15 years and above who use any kind of tobacco (%)	47.2	44.8
92. Women age 15 years and above who consume alcohol (%)	0.3	0.4
93. Men age 15 years and above who consume alcohol (%)	24.6	16.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Chitrakoot, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.5	8.8
2. Population below age 15 years (%)	30.5	31.3
3. Population age 60 years and above (%)	11.7	10.1
4. Population living in households with electricity (%)	98.2	92.0
5. Population living in households with an improved drinking-water source ¹ (%)	95.2	97.0
6. Households using iodized salt (%)	80.2	75.8
7. Households with any usual member covered under a health insurance/financing scheme (%)	39.6	17.4
8. Households with any usual member having a bank account/post office account (%)	97.9	94.0
9. Female population age 6 years and above who ever attended school (%)	65.7	63.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	16.4	7.6
11. Children age 2-4 years who attended pre-school (%)	21.7	6.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	34.8	30.2
13. Men with 10 or more years of schooling (%)	46.1	39.7
14. Women who have ever used the internet (%)	62.8	20.6
15. Men who have ever used the internet (%)	67.0	45.2
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	20.7	21.8
17. Men age 25-29 years married before age 21 years (%)	*	(27.2)
18. Three or more births in the 5 years preceding the survey (%)	4.6	2.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.7	3.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	71.4	57.1
21. Any modern method ² (%)	45.9	49.0
22. Any traditional method (%)	25.5	8.1
23. Female sterilization (%)	32.4	32.8
24. Male sterilization (%)	0.3	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.1	12.7
26. Unmet need for spacing ³ (%)	5.1	4.7
27. Unmet need for limiting ³ (%)	5.0	8.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	73.3	53.7
29. Mothers who had any antenatal care visits (%)	97.7	83.6
30. Mothers who had at least 4 antenatal care visits (%)	36.2	30.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.6	79.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.8	13.4
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.6	7.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.7	93.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	84.1	81.1
36. Institutional births in public facility (%)	72.9	75.0
37. Births attended by skilled health personnel ⁵ (%)	84.9	78.4
38. Births delivered by caesarean section (%)	7.9	5.3
39. Births in a private health facility that were delivered by caesarean section (%)	(55.6)	(51.2)
40. Births in a public health facility that were delivered by caesarean section (%)	2.2	3.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Chitrakoot, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.3	50.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(10.1)	5.4
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.0	49.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	89.4	63.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.1	68.9
46. Children age 12-23 months who have received any vaccine (%)	100.0	86.2
47. Children age 12-23 months who have received BCG (%)	98.9	84.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	100.0	68.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.5	76.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	93.4	80.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	91.0	43.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	75.5	50.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.5	54.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	74.6	80.6
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	92.3
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	4.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.6	3.2
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.0	1.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	71.1	(43.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	54.5	26.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	60.2	41.6
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	97.5	84.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	81.9	66.7
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(51.6)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	14.4	8.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	14.0	8.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	36.9	47.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.2	24.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.9	12.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	36.8	41.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	6.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Chitrakoot, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	25.6	24.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	16.2	14.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.2	1.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.2	1.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.7	5.1
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.7	3.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.0	2.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.8	9.1
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.2	8.4
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.2	3.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	6.6	15.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.3	10.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.4	3.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.3	16.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.9	73.8
88. Women having a mobile phone that they themselves use (%)	51.4	38.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	61.4	55.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	16.2	12.7
91. Men age 15 years and above who use any kind of tobacco (%)	65.0	59.6
92. Women age 15 years and above who consume alcohol (%)	0.3	0.5
93. Men age 15 years and above who consume alcohol (%)	17.4	13.6

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Deoria, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.7	8.6
2. Population below age 15 years (%)	30.9	30.3
3. Population age 60 years and above (%)	11.2	10.7
4. Population living in households with electricity (%)	97.0	95.3
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.8
6. Households using iodized salt (%)	90.4	97.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	35.6	10.5
8. Households with any usual member having a bank account/post office account (%)	98.7	95.9
9. Female population age 6 years and above who ever attended school (%)	73.4	69.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.5	8.5
11. Children age 2-4 years who attended pre-school (%)	37.7	9.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	51.0	50.4
13. Men with 10 or more years of schooling (%)	57.2	57.3
14. Women who have ever used the internet (%)	68.1	48.4
15. Men who have ever used the internet (%)	82.1	71.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.2	13.7
17. Men age 25-29 years married before age 21 years (%)	(17.5)	*
18. Three or more births in the 5 years preceding the survey (%)	3.2	1.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.9	2.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	56.4	56.2
21. Any modern method ² (%)	28.8	45.9
22. Any traditional method (%)	27.6	10.3
23. Female sterilization (%)	19.0	18.0
24. Male sterilization (%)	0.0	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	12.6	20.4
26. Unmet need for spacing ³ (%)	4.7	6.3
27. Unmet need for limiting ³ (%)	7.9	14.1
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	57.4	55.9
29. Mothers who had any antenatal care visits (%)	91.7	97.6
30. Mothers who had at least 4 antenatal care visits (%)	31.5	42.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	88.8	93.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	19.3	13.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	13.8	7.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.0	96.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	91.8	93.4
36. Institutional births in public facility (%)	58.2	76.4
37. Births attended by skilled health personnel ⁵ (%)	94.2	91.6
38. Births delivered by caesarean section (%)	23.2	12.7
39. Births in a private health facility that were delivered by caesarean section (%)	50.1	51.6
40. Births in a public health facility that were delivered by caesarean section (%)	11.0	5.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Deoria, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.7	77.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.7	73.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	83.3	64.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	91.3	73.2
46. Children age 12-23 months who have received any vaccine (%)	92.7	98.7
47. Children age 12-23 months who have received BCG (%)	92.7	92.6
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	84.7	73.9
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.9	79.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	87.7	77.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	71.6	51.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	62.2	37.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.2	61.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	66.2	68.0
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.0	98.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.9	5.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.8	1.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	47.0	(89.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	47.4	17.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(68.7)	(54.8)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(94.0)	(97.8)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.1)	(73.0)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	14.1	3.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	13.2	2.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	26.0	36.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	9.5	26.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.9	10.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	20.8	39.4
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.2	3.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Deoria, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	18.1	16.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	33.4	19.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.2	4.2
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.4	4.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.6	9.6
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.6	6.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.7	5.3
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.2	12.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	12.4	10.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	6.0	5.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	22.5	17.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	20.0	13.0
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	10.2	5.8
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	32.8	21.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.0	79.6
88. Women having a mobile phone that they themselves use (%)	62.7	59.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	76.6	76.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.7	2.3
91. Men age 15 years and above who use any kind of tobacco (%)	42.8	36.9
92. Women age 15 years and above who consume alcohol (%)	0.5	0.3
93. Men age 15 years and above who consume alcohol (%)	20.0	12.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Etah, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.0	10.0
2. Population below age 15 years (%)	31.2	33.2
3. Population age 60 years and above (%)	10.2	9.8
4. Population living in households with electricity (%)	93.8	88.0
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.8
6. Households using iodized salt (%)	79.3	74.8
7. Households with any usual member covered under a health insurance/financing scheme (%)	34.1	12.8
8. Households with any usual member having a bank account/post office account (%)	97.8	93.7
9. Female population age 6 years and above who ever attended school (%)	71.6	68.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.7	13.4
11. Children age 2-4 years who attended pre-school (%)	32.7	32.4
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	42.3	38.8
13. Men with 10 or more years of schooling (%)	60.3	44.8
14. Women who have ever used the internet (%)	54.9	18.7
15. Men who have ever used the internet (%)	79.7	34.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	16.6	20.1
17. Men age 25-29 years married before age 21 years (%)	(34.2)	(20.6)
18. Three or more births in the 5 years preceding the survey (%)	3.3	4.1
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.9	4.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	65.9	73.1
21. Any modern method ² (%)	28.4	39.7
22. Any traditional method (%)	37.5	33.4
23. Female sterilization (%)	7.4	7.8
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.1	7.0
26. Unmet need for spacing ³ (%)	5.0	2.0
27. Unmet need for limiting ³ (%)	6.1	5.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.1	54.1
29. Mothers who had any antenatal care visits (%)	97.1	90.6
30. Mothers who had at least 4 antenatal care visits (%)	42.5	33.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.8	92.3
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	23.6	20.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	13.3	4.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.1	96.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	81.4	76.7
36. Institutional births in public facility (%)	46.8	50.5
37. Births attended by skilled health personnel ⁵ (%)	83.8	80.8
38. Births delivered by caesarean section (%)	11.3	7.5
39. Births in a private health facility that were delivered by caesarean section (%)	30.3	23.3
40. Births in a public health facility that were delivered by caesarean section (%)	1.7	2.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Etah, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	66.5	60.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	1.6	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	68.5	61.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	81.3	55.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	89.4	71.9
46. Children age 12-23 months who have received any vaccine (%)	94.3	98.1
47. Children age 12-23 months who have received BCG (%)	92.8	98.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	84.1	60.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	87.8	75.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.1	88.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	67.4	45.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	43.1	53.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	84.5	19.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	75.4	69.6
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.4	90.2
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.6	1.1
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.7	9.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.7	9.1
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	67.1	46.5
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	49.2	29.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	50.6	56.5
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	92.2	94.9
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	90.7	91.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(31.5)	(26.1)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	7.0	4.2
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(20.3)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.5	5.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	34.9	48.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	13.4	15.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.9	8.4
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	28.6	30.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.1	3.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Etah, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	21.6	18.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.0	20.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.0	3.7
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.2	2.1
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	11.5	6.2
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.8	4.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.2	2.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.3	7.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.4	11.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	3.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.6	16.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.2	14.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.4	4.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.0	19.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	78.9	63.6
88. Women having a mobile phone that they themselves use (%)	50.4	40.2
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	68.2	64.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.4	5.0
91. Men age 15 years and above who use any kind of tobacco (%)	44.1	46.0
92. Women age 15 years and above who consume alcohol (%)	0.4	0.6
93. Men age 15 years and above who consume alcohol (%)	19.7	15.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Etawah, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.0	9.7
2. Population below age 15 years (%)	29.1	29.4
3. Population age 60 years and above (%)	13.1	11.4
4. Population living in households with electricity (%)	98.3	95.7
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.5
6. Households using iodized salt (%)	82.2	77.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.7	17.9
8. Households with any usual member having a bank account/post office account (%)	97.9	97.5
9. Female population age 6 years and above who ever attended school (%)	76.0	74.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.2	11.8
11. Children age 2-4 years who attended pre-school (%)	40.0	38.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	49.9	45.7
13. Men with 10 or more years of schooling (%)	58.6	58.0
14. Women who have ever used the internet (%)	72.7	37.9
15. Men who have ever used the internet (%)	82.3	66.2
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	18.9	16.3
17. Men age 25-29 years married before age 21 years (%)	(23.5)	(19.2)
18. Three or more births in the 5 years preceding the survey (%)	1.7	2.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.5	2.2
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	67.8	71.0
21. Any modern method ² (%)	30.4	48.9
22. Any traditional method (%)	37.4	22.1
23. Female sterilization (%)	13.5	14.4
24. Male sterilization (%)	0.2	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.1	8.3
26. Unmet need for spacing ³ (%)	3.4	2.4
27. Unmet need for limiting ³ (%)	6.7	5.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	72.8	67.9
29. Mothers who had any antenatal care visits (%)	98.6	98.1
30. Mothers who had at least 4 antenatal care visits (%)	41.0	43.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.1	93.6
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.8	28.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	20.6	13.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.1	94.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.2	85.2
36. Institutional births in public facility (%)	61.9	70.1
37. Births attended by skilled health personnel ⁵ (%)	90.3	88.0
38. Births delivered by caesarean section (%)	21.1	11.2
39. Births in a private health facility that were delivered by caesarean section (%)	67.4	45.1
40. Births in a public health facility that were delivered by caesarean section (%)	4.4	6.3

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Etawah, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	74.9	79.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.6	79.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	80.6	77.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.2	87.2
46. Children age 12-23 months who have received any vaccine (%)	97.3	100.0
47. Children age 12-23 months who have received BCG (%)	96.3	100.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	81.5	82.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	83.5	88.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	88.9	92.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	78.9	67.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	83.1	64.0
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	83.5	32.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	70.2	73.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.5	97.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.5	2.6
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	5.2	5.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.3	4.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	43.7	68.8
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	48.5	30.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(45.4)	57.6
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(99.0)	94.6
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.6)	81.0
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	6.6	7.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.6	6.2
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	30.9	38.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	12.4	13.9
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.7	6.2
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	27.9	24.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	2.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Etawah, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.9	16.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.2	23.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.5	4.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.5	3.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.4	8.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.4	5.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.7	4.1
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.3	10.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.2	11.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.3	3.9
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.5	16.7
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.6	15.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.4	3.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.4	19.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.5	76.7
88. Women having a mobile phone that they themselves use (%)	63.0	52.3
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	73.7	76.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.4	5.2
91. Men age 15 years and above who use any kind of tobacco (%)	41.0	45.5
92. Women age 15 years and above who consume alcohol (%)	0.2	0.1
93. Men age 15 years and above who consume alcohol (%)	17.8	16.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Farrukhabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.2	10.5
2. Population below age 15 years (%)	31.2	32.7
3. Population age 60 years and above (%)	10.2	9.1
4. Population living in households with electricity (%)	95.7	91.8
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.7
6. Households using iodized salt (%)	84.9	77.5
7. Households with any usual member covered under a health insurance/financing scheme (%)	38.1	13.5
8. Households with any usual member having a bank account/post office account (%)	98.0	96.8
9. Female population age 6 years and above who ever attended school (%)	69.2	67.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.8	11.3
11. Children age 2-4 years who attended pre-school (%)	34.9	23.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	34.7	33.3
13. Men with 10 or more years of schooling (%)	47.0	40.5
14. Women who have ever used the internet (%)	58.5	23.3
15. Men who have ever used the internet (%)	78.4	48.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	13.7	18.8
17. Men age 25-29 years married before age 21 years (%)	(27.5)	*
18. Three or more births in the 5 years preceding the survey (%)	3.9	4.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.9	4.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	57.9	67.2
21. Any modern method ² (%)	22.5	40.0
22. Any traditional method (%)	35.4	27.2
23. Female sterilization (%)	4.7	8.3
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.1	7.7
26. Unmet need for spacing ³ (%)	4.3	3.5
27. Unmet need for limiting ³ (%)	6.8	4.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.7	71.4
29. Mothers who had any antenatal care visits (%)	95.3	97.6
30. Mothers who had at least 4 antenatal care visits (%)	49.6	41.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.3	94.8
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	25.8	24.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.8	10.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.2	91.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	70.2	69.3
36. Institutional births in public facility (%)	40.6	48.8
37. Births attended by skilled health personnel ⁵ (%)	74.2	74.7
38. Births delivered by caesarean section (%)	13.9	10.0
39. Births in a private health facility that were delivered by caesarean section (%)	43.7	38.8
40. Births in a public health facility that were delivered by caesarean section (%)	2.4	4.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Farrukhabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	68.5	72.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.6	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	68.2	72.0
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	72.4	68.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	79.5	83.7
46. Children age 12-23 months who have received any vaccine (%)	95.8	98.2
47. Children age 12-23 months who have received BCG (%)	94.7	95.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	76.1	72.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	79.5	82.7
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	84.4	85.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	63.4	51.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	48.1	51.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	79.5	34.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.5	71.0
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	98.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	14.5	4.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.3	4.1
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	68.3	67.5
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	35.9	32.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	26.9	58.6
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	95.0	98.3
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	95.3	85.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(51.0)	(48.5)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.2	16.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(7.9)	(7.9)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.6	14.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	34.8	47.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	14.5	14.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.2	6.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	31.5	31.1
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.6	4.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Farrukhabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.2	18.3
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.4	15.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.0	3.3
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.2	2.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	10.8	6.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.4	3.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.4	2.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.3	7.1
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.8	9.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.8	3.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.3	13.8
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.1	12.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	1.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.9	14.8
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	79.7	74.9
88. Women having a mobile phone that they themselves use (%)	50.1	40.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	61.3	61.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.3	6.4
91. Men age 15 years and above who use any kind of tobacco (%)	45.9	49.0
92. Women age 15 years and above who consume alcohol (%)	0.0	0.3
93. Men age 15 years and above who consume alcohol (%)	18.0	19.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Fatehpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.5	9.5
2. Population below age 15 years (%)	30.1	29.0
3. Population age 60 years and above (%)	11.5	12.4
4. Population living in households with electricity (%)	92.1	76.8
5. Population living in households with an improved drinking-water source ¹ (%)	99.0	98.8
6. Households using iodized salt (%)	52.9	76.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	35.4	16.8
8. Households with any usual member having a bank account/post office account (%)	98.7	96.1
9. Female population age 6 years and above who ever attended school (%)	68.6	63.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	23.1	14.2
11. Children age 2-4 years who attended pre-school (%)	29.7	21.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	38.1	37.3
13. Men with 10 or more years of schooling (%)	55.6	46.0
14. Women who have ever used the internet (%)	56.2	24.1
15. Men who have ever used the internet (%)	74.8	54.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	12.6	10.4
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	2.2	3.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.0	3.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	65.3	66.2
21. Any modern method ² (%)	31.4	58.0
22. Any traditional method (%)	33.9	8.2
23. Female sterilization (%)	13.0	8.4
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.1	9.7
26. Unmet need for spacing ³ (%)	5.3	2.7
27. Unmet need for limiting ³ (%)	5.8	7.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	61.7	56.3
29. Mothers who had any antenatal care visits (%)	90.5	91.1
30. Mothers who had at least 4 antenatal care visits (%)	41.8	38.1
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	89.7	90.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	29.8	17.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	15.9	6.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.7	96.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	83.9	81.4
36. Institutional births in public facility (%)	63.7	64.6
37. Births attended by skilled health personnel ⁵ (%)	86.6	85.6
38. Births delivered by caesarean section (%)	10.9	7.0
39. Births in a private health facility that were delivered by caesarean section (%)	41.9	26.7
40. Births in a public health facility that were delivered by caesarean section (%)	3.8	3.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Fatehpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	69.0	66.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	3.6
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	73.4	65.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	92.5	55.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	92.9	63.5
46. Children age 12-23 months who have received any vaccine (%)	98.7	93.9
47. Children age 12-23 months who have received BCG (%)	98.7	90.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	98.7	61.8
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	92.5	76.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	96.0	71.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	70.3	60.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	65.2	47.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	92.4	20.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	76.1	76.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	95.4	91.3
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	4.6	6.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	4.6	2.9
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.3	1.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	61.6	(43.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	63.2	13.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(47.3)	48.0
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	87.7
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(98.0)	64.3
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(42.1)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.7	3.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.8	3.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	40.6	51.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	20.0	17.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.3	9.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	41.7	38.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	3.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Fatehpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	28.8	24.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	20.6	15.8
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.2	2.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.9	2.5
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.2	5.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.6	4.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.9	4.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	22.8	8.8
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.4	10.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.5	4.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.1	17.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.6	13.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.5	3.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.3	18.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.3	67.4
88. Women having a mobile phone that they themselves use (%)	55.6	49.2
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	61.9	70.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	24.1	20.8
91. Men age 15 years and above who use any kind of tobacco (%)	62.5	60.8
92. Women age 15 years and above who consume alcohol (%)	0.4	0.8
93. Men age 15 years and above who consume alcohol (%)	23.9	18.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Firozabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.1	10.2
2. Population below age 15 years (%)	28.8	31.9
3. Population age 60 years and above (%)	10.1	10.0
4. Population living in households with electricity (%)	96.9	96.3
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	99.4
6. Households using iodized salt (%)	90.5	88.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.0	13.4
8. Households with any usual member having a bank account/post office account (%)	97.9	95.2
9. Female population age 6 years and above who ever attended school (%)	74.4	72.5
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.5	14.6
11. Children age 2-4 years who attended pre-school (%)	49.2	33.4
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	47.5	42.4
13. Men with 10 or more years of schooling (%)	51.5	51.4
14. Women who have ever used the internet (%)	67.8	26.8
15. Men who have ever used the internet (%)	82.8	48.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	15.5	24.8
17. Men age 25-29 years married before age 21 years (%)	*	(25.7)
18. Three or more births in the 5 years preceding the survey (%)	2.1	4.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.4	3.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	69.1	66.9
21. Any modern method ² (%)	31.9	34.6
22. Any traditional method (%)	37.2	32.3
23. Female sterilization (%)	15.7	15.0
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	6.8	9.2
26. Unmet need for spacing ³ (%)	2.9	3.2
27. Unmet need for limiting ³ (%)	3.9	6.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.1	58.3
29. Mothers who had any antenatal care visits (%)	95.2	92.1
30. Mothers who had at least 4 antenatal care visits (%)	51.0	39.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.1	84.7
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	20.1	14.9
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	12.9	4.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.1	92.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	84.7	80.1
36. Institutional births in public facility (%)	50.4	56.4
37. Births attended by skilled health personnel ⁵ (%)	89.5	83.3
38. Births delivered by caesarean section (%)	16.7	10.7
39. Births in a private health facility that were delivered by caesarean section (%)	43.6	38.5
40. Births in a public health facility that were delivered by caesarean section (%)	3.4	2.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Firozabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	72.8	73.7
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	1.5
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	74.9	72.0
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	76.1	67.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	78.7	72.9
46. Children age 12-23 months who have received any vaccine (%)	98.1	97.6
47. Children age 12-23 months who have received BCG (%)	98.1	94.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	76.1	72.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	98.1	79.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	98.1	79.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	74.0	48.7
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	55.7	44.5
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	81.5	19.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	72.1	52.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.2	91.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.8	2.6
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	12.2	7.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.5	3.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	57.8	72.5
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	29.5	28.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(42.5)	57.2
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.4)	98.0
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(97.1)	85.6
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(38.7)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	4.5	8.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(3.4)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	4.3	6.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	38.8	46.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	9.4	9.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.7	3.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	27.8	25.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.3	4.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Firozabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.9	18.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.5	24.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.0	4.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	4.3	2.5
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	9.9	7.5
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.8	5.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.5	3.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.4	9.4
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.2	11.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	5.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.2	18.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.7	15.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	4.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.3	20.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.3	72.7
88. Women having a mobile phone that they themselves use (%)	51.3	42.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	72.9	70.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.9	5.2
91. Men age 15 years and above who use any kind of tobacco (%)	40.9	45.6
92. Women age 15 years and above who consume alcohol (%)	0.1	0.3
93. Men age 15 years and above who consume alcohol (%)	15.8	17.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Gautam Buddha Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.3	8.3
2. Population below age 15 years (%)	28.4	28.5
3. Population age 60 years and above (%)	4.7	9.0
4. Population living in households with electricity (%)	99.8	99.4
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.9
6. Households using iodized salt (%)	90.5	97.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.3	23.8
8. Households with any usual member having a bank account/post office account (%)	95.6	96.1
9. Female population age 6 years and above who ever attended school (%)	82.6	78.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	11.3	9.7
11. Children age 2-4 years who attended pre-school (%)	24.1	16.6
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	52.8	51.9
13. Men with 10 or more years of schooling (%)	59.4	58.3
14. Women who have ever used the internet (%)	79.9	63.4
15. Men who have ever used the internet (%)	78.4	76.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	18.0	13.5
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	1.7	2.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.3	2.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	80.2	76.3
21. Any modern method ² (%)	47.6	55.9
22. Any traditional method (%)	32.6	20.4
23. Female sterilization (%)	16.6	24.2
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	4.1	5.3
26. Unmet need for spacing ³ (%)	1.7	2.4
27. Unmet need for limiting ³ (%)	2.4	2.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.7	75.3
29. Mothers who had any antenatal care visits (%)	97.9	98.5
30. Mothers who had at least 4 antenatal care visits (%)	61.6	52.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.8	96.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	42.0	34.1
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	26.7	19.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.0	97.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	83.4	86.6
36. Institutional births in public facility (%)	43.4	32.4
37. Births attended by skilled health personnel ⁵ (%)	88.6	90.1
38. Births delivered by caesarean section (%)	17.6	19.8
39. Births in a private health facility that were delivered by caesarean section (%)	40.0	31.1
40. Births in a public health facility that were delivered by caesarean section (%)	3.5	9.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Gautam Buddha Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.3	85.4
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.6)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.4	84.0
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	72.0	68.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	76.9	(90.4)
46. Children age 12-23 months who have received any vaccine (%)	94.4	94.6
47. Children age 12-23 months who have received BCG (%)	91.6	94.6
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	74.0	69.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	94.4	86.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	94.4	87.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	62.2	(67.2)
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	56.7	70.1
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	78.4	74.0
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	55.3	72.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	82.9	83.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	8.1	6.5
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	7.2	4.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.7	5.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	94.2	(83.4)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	48.8	33.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(32.7)	*
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(91.9)	*
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(85.7)	*
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.9	7.4
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.5	6.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	40.1	25.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.1	12.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.3	3.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	46.8	21.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	0.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Gautam Buddha Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	16.0	15.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	30.2	34.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.9	6.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.2	7.0
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.3	14.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.5	6.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.9	9.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.6	17.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	2.5	9.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	4.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.0	16.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.2	14.0
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.8	5.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.0	20.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.6	74.3
88. Women having a mobile phone that they themselves use (%)	69.4	61.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	91.3	89.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	2.3	3.0
91. Men age 15 years and above who use any kind of tobacco (%)	38.7	29.7
92. Women age 15 years and above who consume alcohol (%)	0.2	0.2
93. Men age 15 years and above who consume alcohol (%)	21.8	17.3

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Ghaziabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.5	7.2
2. Population below age 15 years (%)	25.5	24.9
3. Population age 60 years and above (%)	9.4	9.1
4. Population living in households with electricity (%)	99.5	99.9
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.1
6. Households using iodized salt (%)	94.0	96.3
7. Households with any usual member covered under a health insurance/financing scheme (%)	34.5	20.0
8. Households with any usual member having a bank account/post office account (%)	97.7	97.8
9. Female population age 6 years and above who ever attended school (%)	80.8	78.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	16.1	11.1
11. Children age 2-4 years who attended pre-school (%)	42.4	16.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	56.8	55.2
13. Men with 10 or more years of schooling (%)	59.8	62.2
14. Women who have ever used the internet (%)	80.5	64.2
15. Men who have ever used the internet (%)	88.3	75.4
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	7.4	8.0
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	1.2	3.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.1	0.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	73.5	72.7
21. Any modern method ² (%)	43.7	51.1
22. Any traditional method (%)	29.8	21.6
23. Female sterilization (%)	12.7	15.9
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	3.3	5.3
26. Unmet need for spacing ³ (%)	1.5	2.8
27. Unmet need for limiting ³ (%)	1.8	2.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	83.6	77.2
29. Mothers who had any antenatal care visits (%)	98.8	98.6
30. Mothers who had at least 4 antenatal care visits (%)	60.7	62.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	96.4	86.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	56.7	47.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	40.1	22.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	89.1	95.5
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	87.5	86.4
36. Institutional births in public facility (%)	38.8	40.7
37. Births attended by skilled health personnel ⁵ (%)	91.2	88.0
38. Births delivered by caesarean section (%)	30.1	31.8
39. Births in a private health facility that were delivered by caesarean section (%)	48.4	52.5
40. Births in a public health facility that were delivered by caesarean section (%)	16.8	19.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Ghaziabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	84.2	84.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(4.1)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.3	85.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	71.8	68.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	78.5	(83.8)
46. Children age 12-23 months who have received any vaccine (%)	95.9	93.6
47. Children age 12-23 months who have received BCG (%)	94.1	90.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	71.8	73.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	79.8	79.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	83.1	82.0
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	69.3	(58.2)
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	59.9	78.5
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	82.2	64.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.3	64.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	72.9	(87.8)
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	6.7	(8.6)
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	6.3	4.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.8	0.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(83.3)	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	38.2	18.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	*	(54.5)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	*	(91.9)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	*	(91.3)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	3.3	7.5
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(8.2)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.0	6.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	24.8	28.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.6	17.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.1	3.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	31.8	23.4
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	3.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Ghaziabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	16.0	11.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	38.7	43.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.1	7.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.8	9.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.6	17.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.8	8.0
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.2	11.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.8	20.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.3	10.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.4	6.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.9	19.3
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	15.5
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.4	6.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.9	24.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.9	79.3
88. Women having a mobile phone that they themselves use (%)	72.6	60.0
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	88.8	91.7
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.3	1.7
91. Men age 15 years and above who use any kind of tobacco (%)	28.4	22.3
92. Women age 15 years and above who consume alcohol (%)	0.4	0.1
93. Men age 15 years and above who consume alcohol (%)	16.9	13.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Ghazipur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.9	8.3
2. Population below age 15 years (%)	29.8	29.6
3. Population age 60 years and above (%)	11.9	11.2
4. Population living in households with electricity (%)	96.3	91.8
5. Population living in households with an improved drinking-water source ¹ (%)	99.7	99.3
6. Households using iodized salt (%)	96.2	98.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	32.8	10.6
8. Households with any usual member having a bank account/post office account (%)	98.5	97.4
9. Female population age 6 years and above who ever attended school (%)	71.3	69.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	16.4	11.4
11. Children age 2-4 years who attended pre-school (%)	35.7	13.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	55.3	51.4
13. Men with 10 or more years of schooling (%)	59.6	71.2
14. Women who have ever used the internet (%)	68.6	24.0
15. Men who have ever used the internet (%)	78.0	69.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	14.0	13.5
17. Men age 25-29 years married before age 21 years (%)	*	(16.5)
18. Three or more births in the 5 years preceding the survey (%)	3.9	2.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.9	1.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	62.1	62.9
21. Any modern method ² (%)	37.3	51.2
22. Any traditional method (%)	24.8	11.7
23. Female sterilization (%)	29.4	20.9
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.8	13.4
26. Unmet need for spacing ³ (%)	5.5	4.9
27. Unmet need for limiting ³ (%)	6.3	8.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	68.0	58.1
29. Mothers who had any antenatal care visits (%)	92.3	95.8
30. Mothers who had at least 4 antenatal care visits (%)	39.8	29.2
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.7	93.7
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	22.8	10.6
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.2	2.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.3	97.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.1	88.3
36. Institutional births in public facility (%)	59.9	71.1
37. Births attended by skilled health personnel ⁵ (%)	89.0	90.5
38. Births delivered by caesarean section (%)	16.8	10.9
39. Births in a private health facility that were delivered by caesarean section (%)	43.1	44.3
40. Births in a public health facility that were delivered by caesarean section (%)	7.0	4.6

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Ghazipur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	68.2	71.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(3.4)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	69.3	69.1
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	70.0	60.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	87.8	69.9
46. Children age 12-23 months who have received any vaccine (%)	91.3	96.9
47. Children age 12-23 months who have received BCG (%)	90.0	95.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	71.1	67.6
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	75.3	71.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	81.6	76.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	56.1	53.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	37.4	27.3
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	74.7	33.9
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	60.1	71.8
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.8	97.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.2	2.6
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.4	6.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.7	5.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	46.9	61.8
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	35.6	12.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(54.4)	(60.7)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(95.2)	(97.6)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(85.0)	(77.9)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.6	5.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.9	4.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	22.7	39.3
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	9.7	25.7
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.1	13.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	18.9	38.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	2.1	3.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Ghazipur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	19.8	19.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	25.4	14.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.7	4.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.2	3.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.2	8.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.3	7.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.6	4.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.5	13.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.3	9.0
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.1	3.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.1	14.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	16.6	15.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	9.1	3.5
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	27.1	20.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.5	78.2
88. Women having a mobile phone that they themselves use (%)	60.9	44.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	65.5	77.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.8	4.4
91. Men age 15 years and above who use any kind of tobacco (%)	40.1	37.7
92. Women age 15 years and above who consume alcohol (%)	0.4	0.2
93. Men age 15 years and above who consume alcohol (%)	20.7	10.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Gonda, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.5	9.9
2. Population below age 15 years (%)	34.2	32.4
3. Population age 60 years and above (%)	9.2	10.6
4. Population living in households with electricity (%)	96.1	88.9
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	90.2	85.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	35.3	13.4
8. Households with any usual member having a bank account/post office account (%)	98.8	96.1
9. Female population age 6 years and above who ever attended school (%)	67.1	61.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.5	15.2
11. Children age 2-4 years who attended pre-school (%)	21.0	11.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	36.3	31.5
13. Men with 10 or more years of schooling (%)	37.7	37.1
14. Women who have ever used the internet (%)	75.3	21.1
15. Men who have ever used the internet (%)	84.9	51.4
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	22.0	25.4
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.2	2.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.0	4.2
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	51.6	41.1
21. Any modern method ² (%)	21.3	37.1
22. Any traditional method (%)	30.3	4.0
23. Female sterilization (%)	11.5	8.7
24. Male sterilization (%)	0.0	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	16.6	24.5
26. Unmet need for spacing ³ (%)	6.5	9.2
27. Unmet need for limiting ³ (%)	10.1	15.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.2	44.4
29. Mothers who had any antenatal care visits (%)	96.9	89.3
30. Mothers who had at least 4 antenatal care visits (%)	53.9	41.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.0	90.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	19.2	17.2
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	12.0	12.0
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.4	97.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	86.3	81.8
36. Institutional births in public facility (%)	60.4	66.9
37. Births attended by skilled health personnel ⁵ (%)	84.7	82.6
38. Births delivered by caesarean section (%)	15.0	11.0
39. Births in a private health facility that were delivered by caesarean section (%)	36.3	37.8
40. Births in a public health facility that were delivered by caesarean section (%)	9.2	8.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Gonda, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	67.9	68.6
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.6	3.4
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	70.1	72.1
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	79.5	59.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.8	66.2
46. Children age 12-23 months who have received any vaccine (%)	97.9	91.6
47. Children age 12-23 months who have received BCG (%)	95.6	90.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	81.8	70.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.6	70.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	89.2	71.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	64.5	49.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	59.7	39.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	84.3	21.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	48.5	74.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.0	97.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.0	1.1
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	25.7	15.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.6	11.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	52.3	58.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	38.8	17.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	9.2	(55.9)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	96.5	(89.4)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	78.7	(80.8)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	28.2	5.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	27.4	5.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	27.7	45.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.1	12.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.4	3.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	35.4	28.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	4.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Gonda, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.9	24.6
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	24.8	17.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.1	3.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.9	3.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.6	8.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	12.4	3.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	12.9	5.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	26.0	12.4
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.8	12.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	6.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.1	22.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.1	17.5
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.6	7.5
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.4	28.6
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.5	70.2
88. Women having a mobile phone that they themselves use (%)	60.9	52.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	60.9	61.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	15.7	18.0
91. Men age 15 years and above who use any kind of tobacco (%)	55.4	51.6
92. Women age 15 years and above who consume alcohol (%)	0.2	0.4
93. Men age 15 years and above who consume alcohol (%)	19.1	11.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Gorakhpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.3	9.1
2. Population below age 15 years (%)	26.2	30.5
3. Population age 60 years and above (%)	11.4	10.4
4. Population living in households with electricity (%)	98.6	94.9
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.8
6. Households using iodized salt (%)	89.7	98.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	40.2	17.9
8. Households with any usual member having a bank account/post office account (%)	99.3	97.9
9. Female population age 6 years and above who ever attended school (%)	73.9	67.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.5	15.7
11. Children age 2-4 years who attended pre-school (%)	33.1	12.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	52.5	42.8
13. Men with 10 or more years of schooling (%)	60.3	54.2
14. Women who have ever used the internet (%)	79.4	36.1
15. Men who have ever used the internet (%)	90.9	73.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	6.3	14.6
17. Men age 25-29 years married before age 21 years (%)	*	(17.7)
18. Three or more births in the 5 years preceding the survey (%)	1.5	1.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.1	2.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	63.7	63.8
21. Any modern method ² (%)	35.7	48.9
22. Any traditional method (%)	28.0	14.9
23. Female sterilization (%)	23.7	19.0
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.8	14.7
26. Unmet need for spacing ³ (%)	5.5	4.4
27. Unmet need for limiting ³ (%)	6.3	10.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	77.3	63.9
29. Mothers who had any antenatal care visits (%)	99.3	98.7
30. Mothers who had at least 4 antenatal care visits (%)	70.9	56.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.2	96.3
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	38.6	29.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	24.0	17.0
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.7	98.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	93.2	91.6
36. Institutional births in public facility (%)	55.2	72.5
37. Births attended by skilled health personnel ⁵ (%)	92.3	84.1
38. Births delivered by caesarean section (%)	34.3	15.2
39. Births in a private health facility that were delivered by caesarean section (%)	76.0	57.9
40. Births in a public health facility that were delivered by caesarean section (%)	9.8	5.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Gorakhpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.1	79.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.2	73.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	80.2	67.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.5	69.3
46. Children age 12-23 months who have received any vaccine (%)	94.1	98.9
47. Children age 12-23 months who have received BCG (%)	94.1	97.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	82.7	74.9
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.3	78.3
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.1	85.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	83.0	54.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	63.0	47.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.9	58.3
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	67.2	72.6
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	93.6
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	4.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	18.2	3.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.3	3.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	54.7	(73.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	50.5	38.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(45.7)	(71.2)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(95.1)	(94.2)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(90.8)	(84.0)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(77.7)	(16.8)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	27.9	1.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	29.0	2.3
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	20.5	29.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.4	23.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.8	12.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	29.4	33.7
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.9	1.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Gorakhpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	19.1	18.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	34.1	21.1
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.5	6.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.1	6.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.5	13.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.9	6.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.6	5.1
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.4	11.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.8	10.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.7	4.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.3	16.8
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.9	13.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	7.0	6.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	22.0	20.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	85.9	74.6
88. Women having a mobile phone that they themselves use (%)	68.8	61.9
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	84.2	78.3
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.9	5.4
91. Men age 15 years and above who use any kind of tobacco (%)	42.8	44.6
92. Women age 15 years and above who consume alcohol (%)	0.6	0.1
93. Men age 15 years and above who consume alcohol (%)	24.2	18.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Hamirpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.7	8.7
2. Population below age 15 years (%)	26.6	28.2
3. Population age 60 years and above (%)	13.5	11.3
4. Population living in households with electricity (%)	97.0	94.6
5. Population living in households with an improved drinking-water source ¹ (%)	99.5	99.4
6. Households using iodized salt (%)	61.0	94.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.6	21.9
8. Households with any usual member having a bank account/post office account (%)	96.7	97.4
9. Female population age 6 years and above who ever attended school (%)	67.8	70.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.5	8.2
11. Children age 2-4 years who attended pre-school (%)	45.2	10.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	37.8	37.6
13. Men with 10 or more years of schooling (%)	44.8	50.1
14. Women who have ever used the internet (%)	65.0	27.2
15. Men who have ever used the internet (%)	66.5	52.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	15.3	10.5
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.6	2.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.0	1.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	69.8	61.3
21. Any modern method ² (%)	46.8	47.5
22. Any traditional method (%)	23.0	13.8
23. Female sterilization (%)	27.9	21.9
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.0	14.4
26. Unmet need for spacing ³ (%)	4.4	6.2
27. Unmet need for limiting ³ (%)	4.6	8.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	74.4	77.7
29. Mothers who had any antenatal care visits (%)	96.6	97.4
30. Mothers who had at least 4 antenatal care visits (%)	57.5	43.2
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.8	94.5
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	33.6	27.1
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	17.4	11.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.1	99.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	94.1	96.3
36. Institutional births in public facility (%)	78.6	88.0
37. Births attended by skilled health personnel ⁵ (%)	94.6	96.2
38. Births delivered by caesarean section (%)	9.3	10.7
39. Births in a private health facility that were delivered by caesarean section (%)	43.3	(56.9)
40. Births in a public health facility that were delivered by caesarean section (%)	3.4	6.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Hamirpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	86.0	87.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	88.8	79.0
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	89.6	77.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	86.7	79.8
46. Children age 12-23 months who have received any vaccine (%)	94.7	97.5
47. Children age 12-23 months who have received BCG (%)	94.7	97.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	94.7	84.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	89.6	86.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.1	92.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(79.7)	49.7
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	72.9	63.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.8	45.0
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	71.7	87.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	100.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.9	8.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.2	2.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(59.0)	60.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	69.7	19.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(61.2)	57.2
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(92.4)	87.5
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.1)	65.6
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	14.3	5.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.5	5.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	36.8	48.0
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	33.2	20.6
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	9.1	10.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	44.5	36.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	9.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Hamirpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	33.9	22.5
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	14.9	9.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.4	2.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.2	1.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.8	5.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.3	4.4
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.3	1.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.8	6.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.7	13.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	1.6	2.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	6.4	16.7
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.5	15.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.5	2.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	7.4	18.8
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	85.4	67.6
88. Women having a mobile phone that they themselves use (%)	52.0	41.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	75.2	77.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	28.0	17.0
91. Men age 15 years and above who use any kind of tobacco (%)	68.5	64.5
92. Women age 15 years and above who consume alcohol (%)	0.6	0.2
93. Men age 15 years and above who consume alcohol (%)	20.3	13.3

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Hapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.1	9.7
2. Population below age 15 years (%)	31.6	31.4
3. Population age 60 years and above (%)	9.2	8.0
4. Population living in households with electricity (%)	98.3	98.8
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.9
6. Households using iodized salt (%)	93.3	96.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	39.3	14.9
8. Households with any usual member having a bank account/post office account (%)	97.9	96.3
9. Female population age 6 years and above who ever attended school (%)	72.5	71.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	19.8	9.5
11. Children age 2-4 years who attended pre-school (%)	37.6	16.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	48.2	43.5
13. Men with 10 or more years of schooling (%)	53.2	52.2
14. Women who have ever used the internet (%)	73.0	42.6
15. Men who have ever used the internet (%)	75.9	72.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	9.5	7.1
17. Men age 25-29 years married before age 21 years (%)	*	(26.7)
18. Three or more births in the 5 years preceding the survey (%)	5.3	5.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.0	0.8
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	66.2	70.3
21. Any modern method ² (%)	43.4	47.6
22. Any traditional method (%)	22.8	22.7
23. Female sterilization (%)	16.9	15.3
24. Male sterilization (%)	0.0	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.9	4.8
26. Unmet need for spacing ³ (%)	4.2	2.3
27. Unmet need for limiting ³ (%)	5.7	2.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	81.3	80.1
29. Mothers who had any antenatal care visits (%)	99.3	99.0
30. Mothers who had at least 4 antenatal care visits (%)	65.5	53.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.0	96.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	47.0	29.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	33.3	12.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.3	98.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	83.0	83.3
36. Institutional births in public facility (%)	29.0	34.7
37. Births attended by skilled health personnel ⁵ (%)	86.8	85.6
38. Births delivered by caesarean section (%)	20.1	16.5
39. Births in a private health facility that were delivered by caesarean section (%)	34.3	28.9
40. Births in a public health facility that were delivered by caesarean section (%)	5.5	7.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Hapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.6	83.0
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.4	83.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	83.3	84.4
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	88.7	89.1
46. Children age 12-23 months who have received any vaccine (%)	96.5	98.1
47. Children age 12-23 months who have received BCG (%)	92.2	97.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	86.1	88.3
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.3	94.3
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.7	92.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	77.9	78.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	45.1	65.0
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	89.3	83.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.0	82.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	79.7	93.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	2.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.3	3.9
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.0	2.0
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	79.1	(74.8)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	32.3	38.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	24.1	61.1
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	79.7	95.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	70.9	93.3
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.8	12.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.4	(6.7)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.1	10.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	24.6	30.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	17.9	18.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.6	9.2
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	26.2	26.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.8	5.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Hapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.3	9.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	34.1	32.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.6	4.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.1	3.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.2	7.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.8	4.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.5	4.1
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.9	9.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.1	15.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	6.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.4	22.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.9	20.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.0	5.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.8	25.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.5	80.8
88. Women having a mobile phone that they themselves use (%)	56.6	48.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	79.6	88.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.0	2.4
91. Men age 15 years and above who use any kind of tobacco (%)	30.2	30.8
92. Women age 15 years and above who consume alcohol (%)	0.0	0.1
93. Men age 15 years and above who consume alcohol (%)	14.7	15.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Hardoi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.9	9.9
2. Population below age 15 years (%)	32.6	32.4
3. Population age 60 years and above (%)	10.4	10.2
4. Population living in households with electricity (%)	88.7	71.0
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.9
6. Households using iodized salt (%)	74.7	82.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	34.9	16.0
8. Households with any usual member having a bank account/post office account (%)	99.6	97.1
9. Female population age 6 years and above who ever attended school (%)	66.5	64.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.7	11.8
11. Children age 2-4 years who attended pre-school (%)	30.6	15.4
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	28.7	26.5
13. Men with 10 or more years of schooling (%)	28.9	30.3
14. Women who have ever used the internet (%)	58.0	17.2
15. Men who have ever used the internet (%)	72.7	40.4
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	17.0	19.3
17. Men age 25-29 years married before age 21 years (%)	(23.5)	(44.4)
18. Three or more births in the 5 years preceding the survey (%)	2.3	4.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.2	2.8
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	60.0	58.3
21. Any modern method ² (%)	24.4	46.2
22. Any traditional method (%)	35.6	12.1
23. Female sterilization (%)	7.7	6.9
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.1	13.4
26. Unmet need for spacing ³ (%)	4.5	4.5
27. Unmet need for limiting ³ (%)	6.6	8.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	71.5	68.0
29. Mothers who had any antenatal care visits (%)	96.9	97.0
30. Mothers who had at least 4 antenatal care visits (%)	45.0	39.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	96.4	92.6
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	27.0	23.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.0	13.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.9	96.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	81.2	73.9
36. Institutional births in public facility (%)	60.7	61.3
37. Births attended by skilled health personnel ⁵ (%)	82.6	79.1
38. Births delivered by caesarean section (%)	11.2	7.0
39. Births in a private health facility that were delivered by caesarean section (%)	43.7	43.4
40. Births in a public health facility that were delivered by caesarean section (%)	3.7	2.5

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Hardoi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.2	67.0
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	1.4	5.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.5	62.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	88.6	63.3
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	93.6	62.4
46. Children age 12-23 months who have received any vaccine (%)	96.6	93.2
47. Children age 12-23 months who have received BCG (%)	95.3	88.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	88.6	67.9
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	92.0	76.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.0	79.0
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	76.6	52.4
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	59.1	56.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.9	37.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	65.6	78.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	98.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	14.9	5.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	5.4	3.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	67.3	46.9
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	36.4	20.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(31.3)	53.5
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(83.9)	81.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(80.7)	70.5
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(20.9)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	23.7	8.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	(6.3)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	22.8	7.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	39.0	44.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	23.3	22.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.1	9.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	42.2	33.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	5.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Hardoi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	31.0	22.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	16.1	13.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.5	4.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.7	3.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.5	8.0
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.0	5.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.0	3.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.3	9.8
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.1	11.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.0	4.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.2	18.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.4	14.9
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.7	4.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.0	19.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	83.0	59.6
88. Women having a mobile phone that they themselves use (%)	43.6	31.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	50.0	63.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.3	6.5
91. Men age 15 years and above who use any kind of tobacco (%)	54.2	52.1
92. Women age 15 years and above who consume alcohol (%)	0.1	0.1
93. Men age 15 years and above who consume alcohol (%)	22.7	14.3

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Hathras, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.9	10.4
2. Population below age 15 years (%)	31.7	31.9
3. Population age 60 years and above (%)	10.0	10.4
4. Population living in households with electricity (%)	96.2	94.9
5. Population living in households with an improved drinking-water source ¹ (%)	99.6	99.0
6. Households using iodized salt (%)	90.9	93.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	24.4	10.9
8. Households with any usual member having a bank account/post office account (%)	97.2	95.8
9. Female population age 6 years and above who ever attended school (%)	69.9	67.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	19.1	19.6
11. Children age 2-4 years who attended pre-school (%)	32.1	33.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	35.9	39.1
13. Men with 10 or more years of schooling (%)	47.0	53.4
14. Women who have ever used the internet (%)	56.4	31.6
15. Men who have ever used the internet (%)	76.8	53.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	26.0	28.1
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.6	4.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.5	5.7
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	65.4	57.5
21. Any modern method ² (%)	33.9	37.5
22. Any traditional method (%)	31.5	20.0
23. Female sterilization (%)	15.2	14.6
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.9	13.8
26. Unmet need for spacing ³ (%)	4.3	4.8
27. Unmet need for limiting ³ (%)	5.6	9.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	54.1	49.1
29. Mothers who had any antenatal care visits (%)	87.3	84.6
30. Mothers who had at least 4 antenatal care visits (%)	27.1	27.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.7	87.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	16.0	15.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	10.0	6.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.0	90.0
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	84.1	80.3
36. Institutional births in public facility (%)	48.1	54.0
37. Births attended by skilled health personnel ⁵ (%)	85.3	82.5
38. Births delivered by caesarean section (%)	18.2	13.9
39. Births in a private health facility that were delivered by caesarean section (%)	47.9	40.7
40. Births in a public health facility that were delivered by caesarean section (%)	2.1	5.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Hathras, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	60.8	50.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	1.6
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	64.6	51.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	80.6	71.6
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	88.2	79.5
46. Children age 12-23 months who have received any vaccine (%)	96.9	96.2
47. Children age 12-23 months who have received BCG (%)	91.6	95.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	86.8	76.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	96.9	85.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	95.1	88.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	65.4	48.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	51.7	62.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	87.5	27.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	71.7	61.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.5	92.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.6	4.2
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.9	11.5
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.6	7.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	70.4	54.7
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	44.7	34.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(58.1)	67.4
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(88.8)	92.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.8)	95.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(38.9)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	3.4	5.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(12.9)	(7.3)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.9	5.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	40.3	39.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	12.2	12.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.8	4.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	29.0	24.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.6	3.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Hathras, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	18.3	15.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.5	23.8
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.3	3.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.7	3.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	11.7	7.6
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.6	5.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.5	4.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.6	10.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.0	13.4
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.9	6.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.2	22.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.8	17.5
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	7.0	7.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	19.4	26.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	77.6	67.1
88. Women having a mobile phone that they themselves use (%)	51.3	43.2
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	73.3	66.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.6	8.0
91. Men age 15 years and above who use any kind of tobacco (%)	43.0	43.3
92. Women age 15 years and above who consume alcohol (%)	0.1	0.5
93. Men age 15 years and above who consume alcohol (%)	18.5	12.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Jalaun, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.7	7.6
2. Population below age 15 years (%)	25.9	26.0
3. Population age 60 years and above (%)	14.9	11.8
4. Population living in households with electricity (%)	97.9	96.9
5. Population living in households with an improved drinking-water source ¹ (%)	98.5	99.0
6. Households using iodized salt (%)	77.7	91.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.5	17.6
8. Households with any usual member having a bank account/post office account (%)	98.3	97.5
9. Female population age 6 years and above who ever attended school (%)	69.1	71.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.1	10.0
11. Children age 2-4 years who attended pre-school (%)	26.5	6.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	42.2	38.9
13. Men with 10 or more years of schooling (%)	59.2	60.4
14. Women who have ever used the internet (%)	69.2	21.6
15. Men who have ever used the internet (%)	81.0	51.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	17.7	16.9
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	4.0	2.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.2	2.2
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	67.2	62.4
21. Any modern method ² (%)	41.9	50.5
22. Any traditional method (%)	25.3	11.9
23. Female sterilization (%)	27.1	35.2
24. Male sterilization (%)	0.0	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.4	13.5
26. Unmet need for spacing ³ (%)	5.8	4.6
27. Unmet need for limiting ³ (%)	4.6	8.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	79.6	67.9
29. Mothers who had any antenatal care visits (%)	98.6	97.9
30. Mothers who had at least 4 antenatal care visits (%)	55.8	62.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	97.9	92.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	33.2	20.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	21.4	11.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.0	98.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.0	85.1
36. Institutional births in public facility (%)	66.3	70.0
37. Births attended by skilled health personnel ⁵ (%)	90.2	85.6
38. Births delivered by caesarean section (%)	15.2	12.1
39. Births in a private health facility that were delivered by caesarean section (%)	54.4	60.6
40. Births in a public health facility that were delivered by caesarean section (%)	3.5	4.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Jalaun, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.2	67.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(2.7)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.8	65.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	87.4	66.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(91.2)	64.7
46. Children age 12-23 months who have received any vaccine (%)	96.4	93.6
47. Children age 12-23 months who have received BCG (%)	96.4	90.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	96.4	71.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	87.4	71.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.0	79.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	72.3	58.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	86.6	53.3
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	85.4	32.0
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	65.4	77.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.4	100.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.6	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	15.6	1.2
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.2	0.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	64.3	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	49.6	37.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(17.3)	(58.1)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(90.8)	(100.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(75.8)	(81.3)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.1	3.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.8	2.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	33.5	45.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	28.6	19.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.4	8.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	43.3	36.1
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	2.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Jalaun, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	26.7	22.2
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	21.6	14.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.9	2.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.6	2.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.7	5.2
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	12.0	2.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	12.0	2.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	24.4	6.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.7	11.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	3.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.2	16.8
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.5	14.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.5	4.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.5	20.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.5	77.8
88. Women having a mobile phone that they themselves use (%)	57.0	38.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	74.4	80.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	15.9	13.4
91. Men age 15 years and above who use any kind of tobacco (%)	52.2	49.9
92. Women age 15 years and above who consume alcohol (%)	0.4	0.2
93. Men age 15 years and above who consume alcohol (%)	14.5	11.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Jaunpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.2	9.0
2. Population below age 15 years (%)	30.0	29.3
3. Population age 60 years and above (%)	11.7	11.1
4. Population living in households with electricity (%)	98.7	98.2
5. Population living in households with an improved drinking-water source ¹ (%)	99.7	99.1
6. Households using iodized salt (%)	96.7	97.6
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.8	12.0
8. Households with any usual member having a bank account/post office account (%)	99.4	97.2
9. Female population age 6 years and above who ever attended school (%)	72.9	69.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	19.8	7.3
11. Children age 2-4 years who attended pre-school (%)	25.2	6.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	53.7	52.7
13. Men with 10 or more years of schooling (%)	63.4	61.2
14. Women who have ever used the internet (%)	77.9	45.7
15. Men who have ever used the internet (%)	85.7	77.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	10.3	11.8
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	4.0	3.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.0	0.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	62.1	62.0
21. Any modern method ² (%)	32.8	55.2
22. Any traditional method (%)	29.3	6.8
23. Female sterilization (%)	26.3	23.5
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	13.0	16.2
26. Unmet need for spacing ³ (%)	5.8	5.4
27. Unmet need for limiting ³ (%)	7.2	10.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	70.2	47.9
29. Mothers who had any antenatal care visits (%)	98.5	99.2
30. Mothers who had at least 4 antenatal care visits (%)	46.3	47.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.7	94.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	26.7	14.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.0	7.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.3	98.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	94.9	88.8
36. Institutional births in public facility (%)	51.5	72.4
37. Births attended by skilled health personnel ⁵ (%)	95.2	86.7
38. Births delivered by caesarean section (%)	20.7	7.3
39. Births in a private health facility that were delivered by caesarean section (%)	42.9	32.6
40. Births in a public health facility that were delivered by caesarean section (%)	4.0	2.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Jaunpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.2	82.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.5	79.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	77.0	75.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	86.5	83.6
46. Children age 12-23 months who have received any vaccine (%)	92.5	92.6
47. Children age 12-23 months who have received BCG (%)	91.1	91.4
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	78.4	78.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	85.1	79.9
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	86.3	81.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	69.4	56.9
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	47.7	43.0
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	83.9	47.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	66.3	76.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.5	97.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.5	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.5	5.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.3	4.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	54.9	(45.0)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	38.2	18.0
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(49.3)	54.9
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.6)	97.1
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.4)	77.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.0	4.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	5.2	5.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	21.7	40.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	12.2	14.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.0	6.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	20.4	30.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.7	2.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Jaunpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	18.1	17.5
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	30.6	15.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.8	3.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.6	3.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.8	8.0
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.5	4.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.0	3.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.6	9.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.9	10.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.0	3.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.5	16.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	10.8	13.6
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	6.8	5.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	19.3	19.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.3	83.3
88. Women having a mobile phone that they themselves use (%)	64.8	65.3
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	70.2	83.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	11.2	5.8
91. Men age 15 years and above who use any kind of tobacco (%)	44.7	43.8
92. Women age 15 years and above who consume alcohol (%)	0.2	0.6
93. Men age 15 years and above who consume alcohol (%)	18.3	13.6

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Jhansi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.0	7.7
2. Population below age 15 years (%)	22.4	24.7
3. Population age 60 years and above (%)	12.4	10.7
4. Population living in households with electricity (%)	98.7	95.2
5. Population living in households with an improved drinking-water source ¹ (%)	97.1	96.8
6. Households using iodized salt (%)	74.3	97.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.5	11.4
8. Households with any usual member having a bank account/post office account (%)	92.8	97.1
9. Female population age 6 years and above who ever attended school (%)	71.8	71.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.1	6.0
11. Children age 2-4 years who attended pre-school (%)	32.9	9.4
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	43.4	43.3
13. Men with 10 or more years of schooling (%)	59.3	58.9
14. Women who have ever used the internet (%)	63.9	45.7
15. Men who have ever used the internet (%)	80.8	72.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	17.7	25.1
17. Men age 25-29 years married before age 21 years (%)	*	(16.9)
18. Three or more births in the 5 years preceding the survey (%)	1.3	1.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.6	3.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	76.4	62.0
21. Any modern method ² (%)	58.9	53.6
22. Any traditional method (%)	17.5	8.4
23. Female sterilization (%)	42.3	28.9
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	5.3	13.8
26. Unmet need for spacing ³ (%)	2.2	4.3
27. Unmet need for limiting ³ (%)	3.1	9.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	75.6	70.3
29. Mothers who had any antenatal care visits (%)	91.5	93.3
30. Mothers who had at least 4 antenatal care visits (%)	57.1	36.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	96.0	90.6
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	43.1	24.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	30.4	13.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.7	95.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.5	92.9
36. Institutional births in public facility (%)	59.4	75.6
37. Births attended by skilled health personnel ⁵ (%)	92.2	93.4
38. Births delivered by caesarean section (%)	25.6	12.8
39. Births in a private health facility that were delivered by caesarean section (%)	68.7	47.3
40. Births in a public health facility that were delivered by caesarean section (%)	7.1	6.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Jhansi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	73.8	80.4
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	86.2	76.1
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	(89.0)	44.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(90.2)	53.9
46. Children age 12-23 months who have received any vaccine (%)	(95.8)	95.6
47. Children age 12-23 months who have received BCG (%)	(95.8)	94.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	(95.8)	55.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	(89.0)	69.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	(91.3)	73.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	58.8	44.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	(65.7)	74.3
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	(89.0)	30.3
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	70.8	79.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	(98.1)	98.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	(1.9)	1.5
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.8	4.9
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.2	3.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(85.8)	(40.5)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	59.9	12.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	*	(49.7)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	*	(77.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	*	(61.2)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.4	7.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.3	10.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	26.9	40.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.3	25.2
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.5	10.4
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	38.4	39.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	3.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Jhansi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	23.6	18.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.7	12.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.5	2.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.2	1.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.7	5.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.6	5.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.4	2.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.9	9.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.0	11.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	2.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.7	15.7
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	12.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	3.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.0	17.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.6	80.0
88. Women having a mobile phone that they themselves use (%)	54.3	56.2
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	79.9	82.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	11.5	10.1
91. Men age 15 years and above who use any kind of tobacco (%)	48.3	49.5
92. Women age 15 years and above who consume alcohol (%)	0.3	0.3
93. Men age 15 years and above who consume alcohol (%)	18.2	10.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kannauj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.1	10.1
2. Population below age 15 years (%)	31.7	32.7
3. Population age 60 years and above (%)	10.1	9.3
4. Population living in households with electricity (%)	96.5	87.3
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.9
6. Households using iodized salt (%)	70.4	80.6
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.5	15.3
8. Households with any usual member having a bank account/post office account (%)	98.6	96.6
9. Female population age 6 years and above who ever attended school (%)	72.5	71.3
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.8	11.4
11. Children age 2-4 years who attended pre-school (%)	33.6	25.4
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	41.4	37.3
13. Men with 10 or more years of schooling (%)	39.2	39.3
14. Women who have ever used the internet (%)	62.3	19.5
15. Men who have ever used the internet (%)	75.2	56.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	18.5	10.4
17. Men age 25-29 years married before age 21 years (%)	*	(18.0)
18. Three or more births in the 5 years preceding the survey (%)	2.4	4.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	4.6	1.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	54.6	69.4
21. Any modern method ² (%)	25.5	60.3
22. Any traditional method (%)	29.1	9.1
23. Female sterilization (%)	9.0	7.7
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	15.3	7.2
26. Unmet need for spacing ³ (%)	6.8	2.9
27. Unmet need for limiting ³ (%)	8.5	4.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	62.3	65.6
29. Mothers who had any antenatal care visits (%)	97.6	96.7
30. Mothers who had at least 4 antenatal care visits (%)	49.2	46.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.1	92.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.8	25.4
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.8	10.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.7	96.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	80.6	76.4
36. Institutional births in public facility (%)	49.5	56.5
37. Births attended by skilled health personnel ⁵ (%)	81.7	81.5
38. Births delivered by caesarean section (%)	13.8	11.8
39. Births in a private health facility that were delivered by caesarean section (%)	40.0	48.0
40. Births in a public health facility that were delivered by caesarean section (%)	2.9	4.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kannauj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.6	70.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	1.4	1.5
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.2	66.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	70.9	55.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	82.4	56.8
46. Children age 12-23 months who have received any vaccine (%)	91.4	94.8
47. Children age 12-23 months who have received BCG (%)	86.6	92.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	73.0	62.9
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	76.4	73.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	81.9	82.2
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	67.3	36.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	39.5	56.0
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	75.4	20.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	57.9	72.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	99.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	15.9	5.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.6	3.5
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	74.0	58.2
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	42.5	12.6
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(43.5)	53.9
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.9)	94.0
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(91.4)	66.3
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(59.0)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.2	13.4
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.0	13.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	35.4	43.0
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.3	21.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.6	10.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	41.6	33.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.7	5.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kannauj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.6	20.5
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.4	18.8
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.9	3.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.3	3.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	11.4	7.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.9	2.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.7	3.3
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.3	6.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.4	8.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	3.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.9	14.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.8	10.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.5	2.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.1	14.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.9	63.4
88. Women having a mobile phone that they themselves use (%)	47.1	41.0
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	59.9	66.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.6	8.7
91. Men age 15 years and above who use any kind of tobacco (%)	47.7	50.9
92. Women age 15 years and above who consume alcohol (%)	0.2	0.3
93. Men age 15 years and above who consume alcohol (%)	21.4	15.3

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kanpur Dehat, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.3	8.9
2. Population below age 15 years (%)	26.5	27.9
3. Population age 60 years and above (%)	12.9	11.2
4. Population living in households with electricity (%)	91.3	79.8
5. Population living in households with an improved drinking-water source ¹ (%)	99.6	99.4
6. Households using iodized salt (%)	70.6	81.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	34.8	17.9
8. Households with any usual member having a bank account/post office account (%)	98.8	95.0
9. Female population age 6 years and above who ever attended school (%)	76.5	73.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.5	10.0
11. Children age 2-4 years who attended pre-school (%)	47.2	20.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	46.0	45.4
13. Men with 10 or more years of schooling (%)	47.1	42.6
14. Women who have ever used the internet (%)	69.8	17.8
15. Men who have ever used the internet (%)	74.0	47.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	14.1	12.7
17. Men age 25-29 years married before age 21 years (%)	(15.3)	(14.7)
18. Three or more births in the 5 years preceding the survey (%)	3.1	4.1
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.1	1.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	59.3	60.1
21. Any modern method ² (%)	32.0	41.6
22. Any traditional method (%)	27.3	18.5
23. Female sterilization (%)	11.5	10.7
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	12.8	15.0
26. Unmet need for spacing ³ (%)	5.4	3.9
27. Unmet need for limiting ³ (%)	7.4	11.1
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	74.5	66.8
29. Mothers who had any antenatal care visits (%)	98.3	92.2
30. Mothers who had at least 4 antenatal care visits (%)	56.3	54.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.7	87.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	35.4	24.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	19.9	13.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.1	94.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	81.6	84.3
36. Institutional births in public facility (%)	58.8	69.2
37. Births attended by skilled health personnel ⁵ (%)	84.6	83.8
38. Births delivered by caesarean section (%)	11.5	8.3
39. Births in a private health facility that were delivered by caesarean section (%)	47.9	45.3
40. Births in a public health facility that were delivered by caesarean section (%)	0.9	2.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kanpur Dehat, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.0	67.9
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(10.5)	(2.1)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.8	62.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	79.9	56.3
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(90.9)	68.0
46. Children age 12-23 months who have received any vaccine (%)	94.5	90.2
47. Children age 12-23 months who have received BCG (%)	94.5	85.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	79.9	63.9
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	87.7	69.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	87.7	73.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	77.6	37.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	73.4	38.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	87.7	24.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.5	67.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.3	94.7
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.7	2.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.2	4.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.3	1.5
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	71.3	(47.6)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	53.8	26.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(35.2)	(53.1)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.0)	(97.8)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(83.4)	(73.8)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.7	6.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.6	5.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	37.2	44.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	20.6	12.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.1	6.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	43.4	32.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.3	6.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kanpur Dehat, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	26.1	17.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.3	19.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.0	4.2
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.2	2.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.7	7.6
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	13.3	5.1
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.1	4.1
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	24.0	9.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.0	7.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.6	3.7
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.9	12.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	11.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.9	4.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.5	16.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	85.2	80.4
88. Women having a mobile phone that they themselves use (%)	52.2	40.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	62.8	76.2
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	12.0	10.2
91. Men age 15 years and above who use any kind of tobacco (%)	53.0	50.5
92. Women age 15 years and above who consume alcohol (%)	0.2	0.4
93. Men age 15 years and above who consume alcohol (%)	23.5	16.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kanpur Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	7.1	7.2
2. Population below age 15 years (%)	23.3	24.5
3. Population age 60 years and above (%)	11.5	11.6
4. Population living in households with electricity (%)	98.0	89.3
5. Population living in households with an improved drinking-water source ¹ (%)	99.6	98.8
6. Households using iodized salt (%)	83.8	92.8
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.9	16.1
8. Households with any usual member having a bank account/post office account (%)	98.5	94.7
9. Female population age 6 years and above who ever attended school (%)	82.8	80.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.8	9.7
11. Children age 2-4 years who attended pre-school (%)	48.6	30.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	58.8	58.5
13. Men with 10 or more years of schooling (%)	46.5	55.0
14. Women who have ever used the internet (%)	80.8	43.6
15. Men who have ever used the internet (%)	80.8	58.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.3	10.9
17. Men age 25-29 years married before age 21 years (%)	*	(32.0)
18. Three or more births in the 5 years preceding the survey (%)	2.4	1.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.5	3.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	60.7	73.7
21. Any modern method ² (%)	34.3	57.4
22. Any traditional method (%)	26.4	16.3
23. Female sterilization (%)	10.3	12.8
24. Male sterilization (%)	0.7	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.7	6.8
26. Unmet need for spacing ³ (%)	3.9	2.3
27. Unmet need for limiting ³ (%)	6.8	4.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	78.5	78.4
29. Mothers who had any antenatal care visits (%)	97.6	97.1
30. Mothers who had at least 4 antenatal care visits (%)	59.7	69.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.8	95.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	38.0	26.6
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	25.0	14.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	89.7	95.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	82.4	86.6
36. Institutional births in public facility (%)	47.8	60.3
37. Births attended by skilled health personnel ⁵ (%)	85.7	87.0
38. Births delivered by caesarean section (%)	27.9	19.3
39. Births in a private health facility that were delivered by caesarean section (%)	68.4	46.0
40. Births in a public health facility that were delivered by caesarean section (%)	8.9	11.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kanpur Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.1	71.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(7.6)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	80.2	70.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	(87.1)	72.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(88.8)	(81.2)
46. Children age 12-23 months who have received any vaccine (%)	(100.0)	93.7
47. Children age 12-23 months who have received BCG (%)	(98.6)	93.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	(87.1)	77.4
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	(91.6)	74.7
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	(93.6)	78.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	70.6	57.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	(57.2)	50.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	(91.6)	23.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	62.1	81.6
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	(97.1)	(96.4)
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	(2.9)	(3.6)
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.0	7.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	5.5	3.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	74.0	(46.2)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	52.7	34.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	*	(53.9)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	*	(100.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	*	(68.1)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	20.5	4.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	18.5	3.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	31.1	34.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.0	21.4
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.2	12.6
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	38.9	27.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.8	5.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kanpur Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	21.3	13.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	33.0	25.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.2	5.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.6	4.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.5	10.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.0	3.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.4	3.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.1	8.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.3	10.0
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.7	4.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.8	17.3
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.9	11.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.0	2.8
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.8	16.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	79.8	69.1
88. Women having a mobile phone that they themselves use (%)	69.0	63.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	81.2	87.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	11.5	9.5
91. Men age 15 years and above who use any kind of tobacco (%)	42.6	46.8
92. Women age 15 years and above who consume alcohol (%)	0.1	0.3
93. Men age 15 years and above who consume alcohol (%)	18.0	15.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kasganj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.4	12.1
2. Population below age 15 years (%)	35.6	35.5
3. Population age 60 years and above (%)	9.3	10.1
4. Population living in households with electricity (%)	89.6	87.2
5. Population living in households with an improved drinking-water source ¹ (%)	99.0	99.4
6. Households using iodized salt (%)	84.5	84.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	30.5	15.1
8. Households with any usual member having a bank account/post office account (%)	98.4	93.6
9. Female population age 6 years and above who ever attended school (%)	64.2	62.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.1	14.1
11. Children age 2-4 years who attended pre-school (%)	24.6	27.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	35.4	29.7
13. Men with 10 or more years of schooling (%)	53.2	44.4
14. Women who have ever used the internet (%)	64.0	30.5
15. Men who have ever used the internet (%)	81.6	47.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	19.2	23.2
17. Men age 25-29 years married before age 21 years (%)	(45.2)	(35.5)
18. Three or more births in the 5 years preceding the survey (%)	6.9	5.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	8.3	7.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	68.9	65.2
21. Any modern method ² (%)	23.6	31.3
22. Any traditional method (%)	45.3	33.9
23. Female sterilization (%)	6.7	6.9
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.4	11.8
26. Unmet need for spacing ³ (%)	4.1	3.4
27. Unmet need for limiting ³ (%)	6.3	8.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	61.2	61.2
29. Mothers who had any antenatal care visits (%)	95.6	92.5
30. Mothers who had at least 4 antenatal care visits (%)	30.1	38.1
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	88.0	85.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	22.1	22.4
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	12.4	7.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.0	93.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	85.9	74.8
36. Institutional births in public facility (%)	53.7	46.6
37. Births attended by skilled health personnel ⁵ (%)	87.4	77.2
38. Births delivered by caesarean section (%)	9.4	6.5
39. Births in a private health facility that were delivered by caesarean section (%)	26.1	19.9
40. Births in a public health facility that were delivered by caesarean section (%)	1.8	1.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kasganj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	72.6	62.0
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(4.9)	2.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	73.0	67.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	77.4	69.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	85.8	74.6
46. Children age 12-23 months who have received any vaccine (%)	98.0	94.9
47. Children age 12-23 months who have received BCG (%)	96.9	94.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	80.6	72.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	81.2	82.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	84.4	87.2
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	66.7	55.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	55.3	53.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	83.4	26.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	61.9	59.6
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.4	93.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.6	1.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.1	11.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.9	6.1
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	77.1	74.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	43.1	33.2
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	38.2	67.7
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	89.1	93.0
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	81.0	92.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(42.5)	(51.7)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.1	5.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(16.0)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.3	5.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	38.8	45.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	11.9	19.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.3	9.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	32.0	35.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.6	4.6

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kasganj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	23.2	17.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	23.3	20.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	4.8	3.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	4.6	2.8
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	9.8	6.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.3	6.4
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.4	3.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.0	10.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.7	9.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.7	4.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.7	16.5
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	12.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	5.8
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.7	19.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	85.0	59.8
88. Women having a mobile phone that they themselves use (%)	52.8	45.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	65.1	55.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.5	6.9
91. Men age 15 years and above who use any kind of tobacco (%)	41.4	44.7
92. Women age 15 years and above who consume alcohol (%)	0.1	0.4
93. Men age 15 years and above who consume alcohol (%)	17.1	17.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kaushambi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.1	10.7
2. Population below age 15 years (%)	35.4	34.9
3. Population age 60 years and above (%)	9.6	10.3
4. Population living in households with electricity (%)	83.7	78.7
5. Population living in households with an improved drinking-water source ¹ (%)	99.3	98.3
6. Households using iodized salt (%)	86.8	97.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	31.4	20.2
8. Households with any usual member having a bank account/post office account (%)	98.8	97.5
9. Female population age 6 years and above who ever attended school (%)	65.7	63.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.9	12.7
11. Children age 2-4 years who attended pre-school (%)	18.3	10.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	34.8	31.9
13. Men with 10 or more years of schooling (%)	47.4	59.8
14. Women who have ever used the internet (%)	60.8	26.5
15. Men who have ever used the internet (%)	75.9	59.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	16.0	17.6
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	4.9	3.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.1	2.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	60.0	62.5
21. Any modern method ² (%)	29.7	38.3
22. Any traditional method (%)	30.3	24.2
23. Female sterilization (%)	17.3	19.3
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	14.0	12.5
26. Unmet need for spacing ³ (%)	6.5	4.4
27. Unmet need for limiting ³ (%)	7.5	8.1
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.6	72.2
29. Mothers who had any antenatal care visits (%)	97.8	95.1
30. Mothers who had at least 4 antenatal care visits (%)	61.8	34.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.2	91.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.0	21.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	18.9	11.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.0	96.0
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	85.3	88.6
36. Institutional births in public facility (%)	66.2	72.3
37. Births attended by skilled health personnel ⁵ (%)	86.7	92.8
38. Births delivered by caesarean section (%)	8.9	8.9
39. Births in a private health facility that were delivered by caesarean section (%)	35.4	39.7
40. Births in a public health facility that were delivered by caesarean section (%)	3.2	3.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kaushambi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	76.2	74.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.9	67.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	84.5	65.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	89.1	77.1
46. Children age 12-23 months who have received any vaccine (%)	93.7	94.0
47. Children age 12-23 months who have received BCG (%)	90.1	92.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	93.7	70.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	87.7	73.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	88.1	79.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	74.4	57.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	59.5	52.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	87.7	57.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.3	79.8
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	98.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.4	3.9
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.4	1.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	65.5	(67.8)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	50.2	27.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	6.8	(62.5)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	94.9	(93.2)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	66.4	(83.8)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(80.4)	(34.6)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	36.5	5.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(23.7)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	34.0	6.1
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	44.1	40.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	28.4	18.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.4	4.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	54.6	36.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	0.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kaushambi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.7	24.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	20.1	17.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.4	6.7
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.1	4.5
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.9	11.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.5	7.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.2	8.0
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.3	15.5
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.9	5.0
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.1	3.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.8	9.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.9	6.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.1	2.8
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.9	10.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.3	68.7
88. Women having a mobile phone that they themselves use (%)	55.4	40.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	59.8	67.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	15.0	14.9
91. Men age 15 years and above who use any kind of tobacco (%)	53.8	53.6
92. Women age 15 years and above who consume alcohol (%)	0.6	0.2
93. Men age 15 years and above who consume alcohol (%)	19.9	13.0

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kheri, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.2	10.0
2. Population below age 15 years (%)	32.1	32.4
3. Population age 60 years and above (%)	9.1	8.1
4. Population living in households with electricity (%)	93.9	84.2
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.9
6. Households using iodized salt (%)	78.4	92.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.2	22.0
8. Households with any usual member having a bank account/post office account (%)	99.1	97.2
9. Female population age 6 years and above who ever attended school (%)	62.7	66.3
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.6	13.0
11. Children age 2-4 years who attended pre-school (%)	29.8	5.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	25.8	28.7
13. Men with 10 or more years of schooling (%)	37.9	41.1
14. Women who have ever used the internet (%)	62.3	22.6
15. Men who have ever used the internet (%)	71.3	53.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	17.6	19.7
17. Men age 25-29 years married before age 21 years (%)	*	(28.6)
18. Three or more births in the 5 years preceding the survey (%)	3.3	2.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.3	3.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	53.5	54.9
21. Any modern method ² (%)	34.5	37.9
22. Any traditional method (%)	19.0	17.0
23. Female sterilization (%)	17.7	20.8
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	14.2	16.1
26. Unmet need for spacing ³ (%)	6.8	6.1
27. Unmet need for limiting ³ (%)	7.4	10.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.2	53.9
29. Mothers who had any antenatal care visits (%)	95.2	93.7
30. Mothers who had at least 4 antenatal care visits (%)	56.4	48.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.8	89.9
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	15.5	12.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	2.8	7.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.5	93.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	79.0	82.8
36. Institutional births in public facility (%)	58.2	68.0
37. Births attended by skilled health personnel ⁵ (%)	80.2	81.2
38. Births delivered by caesarean section (%)	13.2	14.2
39. Births in a private health facility that were delivered by caesarean section (%)	58.3	65.8
40. Births in a public health facility that were delivered by caesarean section (%)	1.9	6.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kheri, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	73.1	60.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	3.1	1.6
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.6	55.7
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	72.6	69.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.7	76.2
46. Children age 12-23 months who have received any vaccine (%)	94.6	88.3
47. Children age 12-23 months who have received BCG (%)	89.5	88.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	72.6	69.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	80.8	77.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	84.2	78.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	67.5	63.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	61.6	48.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	80.5	56.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	55.8	65.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	97.0	91.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.0	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.9	3.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.3	1.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	71.2	(72.8)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	43.1	27.0
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(38.8)	65.5
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(94.9)	96.7
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.4)	82.4
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.3	3.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.1	4.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	36.6	47.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	16.4	15.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.1	3.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	37.6	36.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.2	2.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kheri, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	34.9	30.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	15.0	16.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.7	4.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.1	4.0
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.8	8.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.2	4.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.9	4.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.6	9.4
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.4	12.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.6	5.7
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.7	20.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.8	16.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	5.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.5	22.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	84.0	74.0
88. Women having a mobile phone that they themselves use (%)	36.4	29.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	53.8	58.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	19.8	14.8
91. Men age 15 years and above who use any kind of tobacco (%)	60.2	55.2
92. Women age 15 years and above who consume alcohol (%)	0.3	0.2
93. Men age 15 years and above who consume alcohol (%)	19.8	17.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Kushinagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.0	10.3
2. Population below age 15 years (%)	32.8	34.9
3. Population age 60 years and above (%)	10.6	10.1
4. Population living in households with electricity (%)	95.6	91.5
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	100.0
6. Households using iodized salt (%)	81.5	97.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	43.1	15.6
8. Households with any usual member having a bank account/post office account (%)	99.3	96.8
9. Female population age 6 years and above who ever attended school (%)	67.8	63.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.2	11.3
11. Children age 2-4 years who attended pre-school (%)	26.1	16.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	35.6	34.1
13. Men with 10 or more years of schooling (%)	39.6	59.9
14. Women who have ever used the internet (%)	72.2	35.2
15. Men who have ever used the internet (%)	84.3	72.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	13.2	17.7
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.7	3.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.8	2.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	62.5	64.2
21. Any modern method ² (%)	33.9	48.5
22. Any traditional method (%)	28.6	15.7
23. Female sterilization (%)	24.2	23.2
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.6	14.5
26. Unmet need for spacing ³ (%)	5.5	6.5
27. Unmet need for limiting ³ (%)	6.1	8.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	61.4	44.4
29. Mothers who had any antenatal care visits (%)	98.4	96.5
30. Mothers who had at least 4 antenatal care visits (%)	44.4	35.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	91.4	91.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	23.7	17.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.9	9.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.7	96.5
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	88.1	89.4
36. Institutional births in public facility (%)	51.6	71.1
37. Births attended by skilled health personnel ⁵ (%)	90.3	93.1
38. Births delivered by caesarean section (%)	22.5	12.0
39. Births in a private health facility that were delivered by caesarean section (%)	56.4	50.9
40. Births in a public health facility that were delivered by caesarean section (%)	3.7	3.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Kushinagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.9	79.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(3.6)	(2.6)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	76.9	74.7
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	79.5	65.4
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	83.9	82.9
46. Children age 12-23 months who have received any vaccine (%)	93.6	96.0
47. Children age 12-23 months who have received BCG (%)	91.7	91.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	82.5	72.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	83.7	85.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	87.1	87.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	69.9	59.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	44.3	37.1
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.0	44.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	67.9	77.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.8	97.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.2	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.3	4.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.2	5.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	45.7	75.3
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	31.7	16.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(58.2)	63.2
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(98.1)	96.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(85.0)	73.9
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	18.8	3.2
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.9	3.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	22.0	32.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	14.8	24.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.4	9.0
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	24.2	36.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	2.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Kushinagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.9	19.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.7	20.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.9	4.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.2	4.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.0	10.6
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.4	6.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.5	6.3
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.1	12.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.4	9.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.5	4.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.0	16.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	15.0	13.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	8.2	6.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	26.0	21.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.3	82.8
88. Women having a mobile phone that they themselves use (%)	60.4	61.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	79.6	77.3
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.4	4.3
91. Men age 15 years and above who use any kind of tobacco (%)	47.5	45.1
92. Women age 15 years and above who consume alcohol (%)	0.1	0.2
93. Men age 15 years and above who consume alcohol (%)	26.3	16.9

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Lalitpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.1	9.6
2. Population below age 15 years (%)	29.0	31.7
3. Population age 60 years and above (%)	9.9	8.8
4. Population living in households with electricity (%)	97.7	94.1
5. Population living in households with an improved drinking-water source ¹ (%)	97.7	96.6
6. Households using iodized salt (%)	85.1	90.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	31.7	11.7
8. Households with any usual member having a bank account/post office account (%)	98.7	96.5
9. Female population age 6 years and above who ever attended school (%)	67.4	66.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.4	7.0
11. Children age 2-4 years who attended pre-school (%)	22.9	6.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	31.6	23.8
13. Men with 10 or more years of schooling (%)	37.3	37.4
14. Women who have ever used the internet (%)	63.6	17.1
15. Men who have ever used the internet (%)	71.6	50.4
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	40.2	42.5
17. Men age 25-29 years married before age 21 years (%)	(57.9)	(51.6)
18. Three or more births in the 5 years preceding the survey (%)	2.2	3.1
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	9.4	9.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	79.0	73.8
21. Any modern method ² (%)	61.4	65.7
22. Any traditional method (%)	17.6	8.1
23. Female sterilization (%)	53.3	50.0
24. Male sterilization (%)	0.2	0.4
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	6.9	7.4
26. Unmet need for spacing ³ (%)	4.9	3.8
27. Unmet need for limiting ³ (%)	2.0	3.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	63.1	65.7
29. Mothers who had any antenatal care visits (%)	97.2	98.4
30. Mothers who had at least 4 antenatal care visits (%)	38.4	45.7
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.7	89.6
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.9	26.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.8	9.9
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.1	98.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	91.2	90.8
36. Institutional births in public facility (%)	82.9	86.6
37. Births attended by skilled health personnel ⁵ (%)	90.5	84.5
38. Births delivered by caesarean section (%)	8.4	6.4
39. Births in a private health facility that were delivered by caesarean section (%)	(47.9)	*
40. Births in a public health facility that were delivered by caesarean section (%)	5.3	4.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Lalitpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.5	71.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(3.8)	(2.8)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.8	64.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	92.2	70.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.3	73.0
46. Children age 12-23 months who have received any vaccine (%)	98.5	95.7
47. Children age 12-23 months who have received BCG (%)	97.2	94.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	98.5	77.8
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	94.8	80.9
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	94.9	81.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	74.8	44.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	78.7	67.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	92.3	38.9
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	49.6	78.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	100.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	21.1	2.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.7	0.8
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	70.1	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	49.8	29.2
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(57.6)	56.0
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	86.2
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(97.6)	79.5
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(68.5)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.8	4.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	12.3	4.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	29.8	46.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.9	18.7
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.3	7.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	35.2	34.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.6	6.7

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Lalitpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	25.1	22.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	17.1	13.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.1	2.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	4.4	1.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	11.7	4.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.5	3.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.1	1.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.0	6.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.4	10.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.7	4.7
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	7.8	17.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.7	13.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.0	3.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.4	17.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	88.2	66.8
88. Women having a mobile phone that they themselves use (%)	38.9	28.9
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	64.1	63.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	8.7	5.5
91. Men age 15 years and above who use any kind of tobacco (%)	56.8	54.5
92. Women age 15 years and above who consume alcohol (%)	0.2	0.7
93. Men age 15 years and above who consume alcohol (%)	22.1	16.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Lucknow, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	6.8	6.9
2. Population below age 15 years (%)	23.8	23.5
3. Population age 60 years and above (%)	11.9	10.3
4. Population living in households with electricity (%)	98.5	96.8
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.8
6. Households using iodized salt (%)	89.2	97.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	34.3	15.8
8. Households with any usual member having a bank account/post office account (%)	97.8	97.7
9. Female population age 6 years and above who ever attended school (%)	79.7	80.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.0	14.9
11. Children age 2-4 years who attended pre-school (%)	35.3	11.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	52.2	51.9
13. Men with 10 or more years of schooling (%)	54.9	47.0
14. Women who have ever used the internet (%)	79.9	45.7
15. Men who have ever used the internet (%)	88.9	73.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	10.3	9.9
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	1.1	1.2
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.0	2.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	68.1	56.5
21. Any modern method ² (%)	42.8	40.2
22. Any traditional method (%)	25.3	16.3
23. Female sterilization (%)	16.9	16.4
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	6.9	15.7
26. Unmet need for spacing ³ (%)	2.9	5.1
27. Unmet need for limiting ³ (%)	4.0	10.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	82.6	74.8
29. Mothers who had any antenatal care visits (%)	99.0	98.3
30. Mothers who had at least 4 antenatal care visits (%)	73.4	53.1
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	97.0	96.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	42.4	22.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	26.1	15.9
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	90.9	96.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	92.2	91.3
36. Institutional births in public facility (%)	55.9	47.2
37. Births attended by skilled health personnel ⁵ (%)	92.7	87.9
38. Births delivered by caesarean section (%)	37.6	36.7
39. Births in a private health facility that were delivered by caesarean section (%)	71.5	55.9
40. Births in a public health facility that were delivered by caesarean section (%)	20.9	25.6

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Lucknow, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	87.8	78.7
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	92.2	71.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	87.8	68.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(96.7)	(84.5)
46. Children age 12-23 months who have received any vaccine (%)	97.9	91.7
47. Children age 12-23 months who have received BCG (%)	97.9	87.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	87.8	73.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	97.1	78.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	97.1	78.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(89.4)	71.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	80.6	57.0
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	93.1	61.3
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	79.9	70.2
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	97.8	94.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	2.2	5.9
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	9.2	4.9
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.6	1.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	77.3	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	39.9	19.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	*	(71.7)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	*	(90.2)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	*	(83.8)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	23.4	3.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	20.1	2.3
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	31.0	32.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	14.6	11.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.3	1.4
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	27.8	25.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	2.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Lucknow, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.7	15.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	37.8	34.8
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.7	6.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.2	7.8
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.5	14.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.0	5.6
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	13.1	9.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	23.4	16.5
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.5	12.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.1	6.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.0	21.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.0	15.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.8	6.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	18.5	22.8
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.6	73.7
88. Women having a mobile phone that they themselves use (%)	66.7	57.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	85.4	84.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.9	7.7
91. Men age 15 years and above who use any kind of tobacco (%)	39.2	39.2
92. Women age 15 years and above who consume alcohol (%)	0.5	0.2
93. Men age 15 years and above who consume alcohol (%)	18.2	14.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Maharajanj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.7	9.5
2. Population below age 15 years (%)	32.5	32.4
3. Population age 60 years and above (%)	10.3	9.1
4. Population living in households with electricity (%)	95.0	94.8
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	83.6	98.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	41.8	18.2
8. Households with any usual member having a bank account/post office account (%)	98.7	97.9
9. Female population age 6 years and above who ever attended school (%)	65.6	62.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.5	11.2
11. Children age 2-4 years who attended pre-school (%)	23.9	11.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	35.7	33.8
13. Men with 10 or more years of schooling (%)	43.2	39.3
14. Women who have ever used the internet (%)	73.8	29.0
15. Men who have ever used the internet (%)	88.0	74.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	20.6	24.1
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	2.2	1.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.7	3.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	63.1	62.9
21. Any modern method ² (%)	37.9	49.6
22. Any traditional method (%)	25.2	13.3
23. Female sterilization (%)	25.8	27.9
24. Male sterilization (%)	0.3	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.0	13.7
26. Unmet need for spacing ³ (%)	4.9	5.3
27. Unmet need for limiting ³ (%)	6.1	8.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.9	55.5
29. Mothers who had any antenatal care visits (%)	99.0	97.8
30. Mothers who had at least 4 antenatal care visits (%)	57.8	52.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.9	96.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	25.8	26.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.4	12.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.8	97.4
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	87.0	93.3
36. Institutional births in public facility (%)	58.9	77.5
37. Births attended by skilled health personnel ⁵ (%)	88.0	92.0
38. Births delivered by caesarean section (%)	17.4	14.7
39. Births in a private health facility that were delivered by caesarean section (%)	55.1	57.7
40. Births in a public health facility that were delivered by caesarean section (%)	3.2	7.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Maharajanj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.0	77.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.2)	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	76.1	76.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	84.7	76.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	93.2	82.8
46. Children age 12-23 months who have received any vaccine (%)	97.8	100.0
47. Children age 12-23 months who have received BCG (%)	96.7	96.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	85.8	82.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.7	92.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	93.0	91.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	80.4	60.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	60.6	41.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.7	68.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	62.2	81.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	94.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.7	3.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.0	4.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	54.6	63.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	41.2	13.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	54.9	62.1
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	96.0	96.9
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	75.1	90.5
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.3	7.7
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	20.3	8.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	28.9	40.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	23.8	21.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.9	9.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	37.8	37.4
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	3.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Maharajanj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.6	17.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	25.9	19.9
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.0	6.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.5	5.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.9	12.0
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.2	6.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.4	5.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.1	13.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.2	11.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.8	5.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.1	19.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	11.3	16.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.8	6.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	20.1	24.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.6	83.5
88. Women having a mobile phone that they themselves use (%)	61.6	57.0
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	78.9	80.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	6.0	5.9
91. Men age 15 years and above who use any kind of tobacco (%)	49.0	47.8
92. Women age 15 years and above who consume alcohol (%)	0.4	0.3
93. Men age 15 years and above who consume alcohol (%)	22.7	17.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Mahoba, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.4	9.4
2. Population below age 15 years (%)	25.5	28.4
3. Population age 60 years and above (%)	13.6	11.0
4. Population living in households with electricity (%)	97.6	95.9
5. Population living in households with an improved drinking-water source ¹ (%)	97.7	94.3
6. Households using iodized salt (%)	80.2	93.8
7. Households with any usual member covered under a health insurance/financing scheme (%)	31.0	16.2
8. Households with any usual member having a bank account/post office account (%)	97.9	95.4
9. Female population age 6 years and above who ever attended school (%)	64.9	65.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.0	9.8
11. Children age 2-4 years who attended pre-school (%)	32.7	7.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	35.1	30.7
13. Men with 10 or more years of schooling (%)	42.1	37.6
14. Women who have ever used the internet (%)	65.5	24.4
15. Men who have ever used the internet (%)	77.3	63.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	14.8	20.8
17. Men age 25-29 years married before age 21 years (%)	(21.3)	(30.3)
18. Three or more births in the 5 years preceding the survey (%)	3.2	4.2
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.1	4.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	74.7	61.6
21. Any modern method ² (%)	53.8	52.0
22. Any traditional method (%)	20.9	9.6
23. Female sterilization (%)	38.4	32.1
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	6.5	16.6
26. Unmet need for spacing ³ (%)	3.6	6.6
27. Unmet need for limiting ³ (%)	2.9	10.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	79.4	76.0
29. Mothers who had any antenatal care visits (%)	98.5	97.9
30. Mothers who had at least 4 antenatal care visits (%)	61.1	46.2
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.5	92.5
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	43.8	27.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	33.2	13.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.2	98.4
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	93.2	96.3
36. Institutional births in public facility (%)	80.7	87.6
37. Births attended by skilled health personnel ⁵ (%)	93.1	97.0
38. Births delivered by caesarean section (%)	13.4	9.6
39. Births in a private health facility that were delivered by caesarean section (%)	(71.9)	(62.8)
40. Births in a public health facility that were delivered by caesarean section (%)	5.5	4.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Mahoba, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	86.2	86.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	90.0	80.8
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	88.6	70.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	81.1	73.9
46. Children age 12-23 months who have received any vaccine (%)	96.8	90.8
47. Children age 12-23 months who have received BCG (%)	96.8	90.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	96.8	78.8
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	88.6	78.4
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.2	78.2
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	(69.5)	55.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	85.6	73.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	87.3	54.3
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	58.3	88.8
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.5	100.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.5	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	15.8	6.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.1	4.0
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	69.2	(75.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	65.6	20.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(21.0)	67.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	89.9
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(92.7)	82.0
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(14.1)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.5	7.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.2	7.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.5	42.3
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	25.6	25.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.2	11.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	38.3	33.4
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	5.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Mahoba, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	30.3	28.0
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	13.0	10.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.1	3.4
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	4.2	1.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.6	5.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.1	3.1
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.7	2.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.3	6.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.6	10.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	1.7	3.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	6.5	15.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.7	11.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	1.9	2.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	7.8	15.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	89.3	70.1
88. Women having a mobile phone that they themselves use (%)	48.2	35.8
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	78.4	77.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	16.2	13.2
91. Men age 15 years and above who use any kind of tobacco (%)	66.5	63.3
92. Women age 15 years and above who consume alcohol (%)	0.6	0.2
93. Men age 15 years and above who consume alcohol (%)	19.0	12.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Mainpuri, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.9	9.4
2. Population below age 15 years (%)	29.2	30.9
3. Population age 60 years and above (%)	11.4	11.1
4. Population living in households with electricity (%)	96.1	95.2
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.6
6. Households using iodized salt (%)	87.8	74.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.4	16.1
8. Households with any usual member having a bank account/post office account (%)	97.7	95.7
9. Female population age 6 years and above who ever attended school (%)	76.3	74.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	23.1	16.6
11. Children age 2-4 years who attended pre-school (%)	41.1	31.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	45.7	45.8
13. Men with 10 or more years of schooling (%)	48.6	54.7
14. Women who have ever used the internet (%)	65.9	30.2
15. Men who have ever used the internet (%)	76.0	51.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	15.5	16.0
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	4.5	5.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.6	4.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	62.1	80.1
21. Any modern method ² (%)	26.6	46.4
22. Any traditional method (%)	35.5	33.7
23. Female sterilization (%)	10.8	10.8
24. Male sterilization (%)	0.0	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.7	4.4
26. Unmet need for spacing ³ (%)	3.4	1.6
27. Unmet need for limiting ³ (%)	6.3	2.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	68.1	63.7
29. Mothers who had any antenatal care visits (%)	94.7	91.6
30. Mothers who had at least 4 antenatal care visits (%)	51.3	38.1
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.5	92.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	21.6	16.2
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	10.4	3.0
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.9	95.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	82.1	82.7
36. Institutional births in public facility (%)	55.1	59.7
37. Births attended by skilled health personnel ⁵ (%)	82.8	84.7
38. Births delivered by caesarean section (%)	11.9	8.2
39. Births in a private health facility that were delivered by caesarean section (%)	35.9	30.1
40. Births in a public health facility that were delivered by caesarean section (%)	4.1	2.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Mainpuri, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.9	63.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.0)	1.6
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	76.9	66.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	86.2	75.0
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	86.0	87.7
46. Children age 12-23 months who have received any vaccine (%)	100.0	96.7
47. Children age 12-23 months who have received BCG (%)	97.1	94.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	89.1	80.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	100.0	86.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	100.0	89.2
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	65.8	58.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	71.6	74.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	92.9	22.1
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	67.5	68.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.3	93.5
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.7	3.4
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.1	5.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.1	4.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	59.3	49.0
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	36.4	30.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(44.4)	(61.9)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(91.5)	(89.4)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(87.1)	(87.6)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.6	11.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.2	10.1
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	31.7	44.3
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	13.1	14.6
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.4	6.7
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	27.1	33.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.5	4.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Mainpuri, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	19.7	15.3
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.7	21.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.4	4.4
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	4.9	3.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.4	8.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.0	6.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.0	3.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.3	10.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.9	9.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	3.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.8	14.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.7	13.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	3.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.2	17.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.4	75.8
88. Women having a mobile phone that they themselves use (%)	54.3	51.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	67.2	71.3
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	3.3	2.9
91. Men age 15 years and above who use any kind of tobacco (%)	41.2	41.3
92. Women age 15 years and above who consume alcohol (%)	0.1	0.5
93. Men age 15 years and above who consume alcohol (%)	17.5	15.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Mathura, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.0	10.6
2. Population below age 15 years (%)	29.0	32.7
3. Population age 60 years and above (%)	10.7	10.3
4. Population living in households with electricity (%)	98.9	98.2
5. Population living in households with an improved drinking-water source ¹ (%)	97.8	98.4
6. Households using iodized salt (%)	95.4	90.3
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.3	10.8
8. Households with any usual member having a bank account/post office account (%)	98.0	95.4
9. Female population age 6 years and above who ever attended school (%)	67.9	68.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.5	15.9
11. Children age 2-4 years who attended pre-school (%)	31.7	35.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	41.4	34.7
13. Men with 10 or more years of schooling (%)	61.8	57.9
14. Women who have ever used the internet (%)	63.3	24.9
15. Men who have ever used the internet (%)	84.5	50.4
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	25.4	21.3
17. Men age 25-29 years married before age 21 years (%)	(16.2)	*
18. Three or more births in the 5 years preceding the survey (%)	3.0	6.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	7.9	6.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	65.1	58.2
21. Any modern method ² (%)	40.6	43.1
22. Any traditional method (%)	24.5	15.1
23. Female sterilization (%)	29.0	23.5
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.0	12.9
26. Unmet need for spacing ³ (%)	5.0	4.7
27. Unmet need for limiting ³ (%)	6.0	8.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	69.1	59.3
29. Mothers who had any antenatal care visits (%)	97.2	83.4
30. Mothers who had at least 4 antenatal care visits (%)	51.2	39.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.4	87.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	26.6	14.1
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	15.6	3.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.8	89.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	88.6	79.5
36. Institutional births in public facility (%)	45.8	36.9
37. Births attended by skilled health personnel ⁵ (%)	91.3	81.5
38. Births delivered by caesarean section (%)	20.3	15.7
39. Births in a private health facility that were delivered by caesarean section (%)	40.5	32.3
40. Births in a public health facility that were delivered by caesarean section (%)	6.6	5.2

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Mathura, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	71.6	62.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.5	67.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	81.6	73.6
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	81.3	80.5
46. Children age 12-23 months who have received any vaccine (%)	92.1	98.3
47. Children age 12-23 months who have received BCG (%)	90.8	98.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	84.3	75.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	92.1	91.7
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.8	90.3
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	69.9	53.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	56.5	50.3
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	83.4	23.9
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	62.9	69.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.2	91.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.8	3.2
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.8	9.3
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	6.2	6.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	68.4	62.6
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	27.5	24.5
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	44.2	61.0
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	86.4	95.4
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	83.1	82.1
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	2.8	2.4
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(3.3)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	3.0	2.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.8	31.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	10.5	11.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.3	3.2
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	21.7	21.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.8	2.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Mathura, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.6	18.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	27.8	25.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.2	3.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	3.8	2.8
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	10.5	7.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.2	4.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.5	5.0
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.2	10.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.9	12.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	6.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.2	21.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.5	17.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.3	7.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	15.6	25.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	75.6	68.4
88. Women having a mobile phone that they themselves use (%)	48.8	32.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	81.2	74.7
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	7.7	9.3
91. Men age 15 years and above who use any kind of tobacco (%)	39.6	42.9
92. Women age 15 years and above who consume alcohol (%)	0.1	0.1
93. Men age 15 years and above who consume alcohol (%)	13.4	13.0

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Mau, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.4	7.5
2. Population below age 15 years (%)	27.0	27.2
3. Population age 60 years and above (%)	10.9	10.8
4. Population living in households with electricity (%)	98.3	94.2
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.8
6. Households using iodized salt (%)	93.6	98.4
7. Households with any usual member covered under a health insurance/financing scheme (%)	45.0	16.0
8. Households with any usual member having a bank account/post office account (%)	99.1	97.5
9. Female population age 6 years and above who ever attended school (%)	75.7	68.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.8	11.6
11. Children age 2-4 years who attended pre-school (%)	35.7	8.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	58.5	53.1
13. Men with 10 or more years of schooling (%)	50.2	69.8
14. Women who have ever used the internet (%)	74.4	32.6
15. Men who have ever used the internet (%)	81.7	74.2
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	7.0	11.0
17. Men age 25-29 years married before age 21 years (%)	*	(11.4)
18. Three or more births in the 5 years preceding the survey (%)	4.7	2.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.7	0.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	67.9	54.2
21. Any modern method ² (%)	27.9	39.0
22. Any traditional method (%)	40.0	15.2
23. Female sterilization (%)	18.3	11.8
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	7.5	16.8
26. Unmet need for spacing ³ (%)	3.8	7.2
27. Unmet need for limiting ³ (%)	3.7	9.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	68.8	64.4
29. Mothers who had any antenatal care visits (%)	96.1	94.6
30. Mothers who had at least 4 antenatal care visits (%)	35.5	43.2
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	91.0	97.5
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	27.4	27.6
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	17.4	13.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	88.2	98.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	91.9	94.7
36. Institutional births in public facility (%)	37.6	58.9
37. Births attended by skilled health personnel ⁵ (%)	89.5	90.7
38. Births delivered by caesarean section (%)	22.7	14.5
39. Births in a private health facility that were delivered by caesarean section (%)	38.8	33.3
40. Births in a public health facility that were delivered by caesarean section (%)	4.2	4.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Mau, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.9	80.0
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(7.7)	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	80.5	75.7
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	66.7	59.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	80.5	63.3
46. Children age 12-23 months who have received any vaccine (%)	93.5	100.0
47. Children age 12-23 months who have received BCG (%)	90.4	98.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	68.9	73.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	76.9	74.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	82.2	76.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	63.1	47.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	37.5	53.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	75.3	41.3
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	60.5	84.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	98.6	97.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.4	1.6
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.3	4.7
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.2	2.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	68.9	(55.1)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	33.5	37.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	56.2	(65.1)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	90.9	(94.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	78.8	(85.3)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	17.1	3.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(21.3)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	18.1	3.2
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	24.0	25.4
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	9.4	21.2
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	1.2	9.6
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	22.6	30.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.8	1.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Mau, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.4	16.5
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.0	20.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.1	4.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.7	5.4
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.3	11.6
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.8	5.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.8	6.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.2	12.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.6	16.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.7	5.2
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.0	23.9
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	12.9	20.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	5.8	7.9
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	20.6	29.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	88.3	93.3
88. Women having a mobile phone that they themselves use (%)	58.2	61.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	72.1	76.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.9	6.4
91. Men age 15 years and above who use any kind of tobacco (%)	38.4	39.3
92. Women age 15 years and above who consume alcohol (%)	0.4	0.4
93. Men age 15 years and above who consume alcohol (%)	20.5	13.6

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Meerut, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.2	10.0
2. Population below age 15 years (%)	27.7	30.3
3. Population age 60 years and above (%)	10.0	8.6
4. Population living in households with electricity (%)	98.4	98.6
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.7
6. Households using iodized salt (%)	90.9	96.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	40.8	18.4
8. Households with any usual member having a bank account/post office account (%)	97.4	98.3
9. Female population age 6 years and above who ever attended school (%)	75.8	74.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.0	12.7
11. Children age 2-4 years who attended pre-school (%)	38.6	14.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	52.1	47.3
13. Men with 10 or more years of schooling (%)	50.9	59.5
14. Women who have ever used the internet (%)	74.2	44.2
15. Men who have ever used the internet (%)	77.3	71.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	6.4	7.6
17. Men age 25-29 years married before age 21 years (%)	(3.0)	*
18. Three or more births in the 5 years preceding the survey (%)	2.2	5.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.2	2.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	64.3	72.4
21. Any modern method ² (%)	40.1	48.8
22. Any traditional method (%)	24.2	23.6
23. Female sterilization (%)	12.5	14.2
24. Male sterilization (%)	0.3	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.1	5.6
26. Unmet need for spacing ³ (%)	3.0	2.7
27. Unmet need for limiting ³ (%)	5.1	2.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	66.0	73.4
29. Mothers who had any antenatal care visits (%)	95.7	97.6
30. Mothers who had at least 4 antenatal care visits (%)	44.4	49.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.1	92.5
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	36.7	35.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	23.7	13.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.6	95.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.2	80.6
36. Institutional births in public facility (%)	36.5	25.7
37. Births attended by skilled health personnel ⁵ (%)	91.0	88.8
38. Births delivered by caesarean section (%)	29.4	19.9
39. Births in a private health facility that were delivered by caesarean section (%)	41.2	29.9
40. Births in a public health facility that were delivered by caesarean section (%)	21.1	13.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Meerut, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	76.7	81.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	80.9	81.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	78.7	82.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.1	92.2
46. Children age 12-23 months who have received any vaccine (%)	92.3	97.8
47. Children age 12-23 months who have received BCG (%)	92.3	97.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	78.7	83.3
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.7	91.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.3	92.4
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	65.2	79.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	45.5	44.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	81.3	79.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	63.0	79.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	75.9	89.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	4.4	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	14.9	2.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.2	2.0
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	(85.7)	(69.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	29.5	27.2
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(41.5)	(55.4)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(89.1)	(88.1)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(82.4)	(84.7)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(27.6)	(40.1)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.3	6.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(18.0)	(14.1)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	16.5	8.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	22.4	32.1
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.9	10.2
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.8	2.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	27.0	23.7
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.9	1.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Meerut, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	15.4	12.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	37.2	38.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.0	5.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.2	7.8
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.7	14.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.3	7.3
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.3	7.9
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	23.0	16.0
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.1	12.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.8	5.5
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.9	20.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.9	17.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	7.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.3	26.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.0	81.1
88. Women having a mobile phone that they themselves use (%)	62.0	55.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	83.1	83.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	2.4	2.6
91. Men age 15 years and above who use any kind of tobacco (%)	25.2	24.4
92. Women age 15 years and above who consume alcohol (%)	0.3	0.2
93. Men age 15 years and above who consume alcohol (%)	11.4	13.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Mirzapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.3	9.0
2. Population below age 15 years (%)	29.8	31.9
3. Population age 60 years and above (%)	10.7	10.9
4. Population living in households with electricity (%)	98.2	92.7
5. Population living in households with an improved drinking-water source ¹ (%)	93.2	96.9
6. Households using iodized salt (%)	86.1	97.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	42.2	15.0
8. Households with any usual member having a bank account/post office account (%)	99.6	96.5
9. Female population age 6 years and above who ever attended school (%)	72.9	69.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	26.9	10.1
11. Children age 2-4 years who attended pre-school (%)	26.7	6.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	45.8	43.8
13. Men with 10 or more years of schooling (%)	54.0	56.3
14. Women who have ever used the internet (%)	68.8	30.5
15. Men who have ever used the internet (%)	77.8	62.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	14.8	16.6
17. Men age 25-29 years married before age 21 years (%)	(23.1)	*
18. Three or more births in the 5 years preceding the survey (%)	4.3	3.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.0	2.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	72.0	61.5
21. Any modern method ² (%)	45.8	53.3
22. Any traditional method (%)	26.2	8.2
23. Female sterilization (%)	38.8	32.4
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.3	14.5
26. Unmet need for spacing ³ (%)	4.9	6.1
27. Unmet need for limiting ³ (%)	4.4	8.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	71.1	55.7
29. Mothers who had any antenatal care visits (%)	97.9	95.5
30. Mothers who had at least 4 antenatal care visits (%)	67.6	36.5
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	91.8	94.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	27.8	11.6
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	15.4	3.3
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.2	98.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	86.2	91.1
36. Institutional births in public facility (%)	58.9	62.1
37. Births attended by skilled health personnel ⁵ (%)	87.1	88.6
38. Births delivered by caesarean section (%)	17.5	17.6
39. Births in a private health facility that were delivered by caesarean section (%)	43.2	43.7
40. Births in a public health facility that were delivered by caesarean section (%)	9.7	8.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Mirzapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	80.3	69.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(6.9)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.1	69.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	76.8	68.3
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.7	83.3
46. Children age 12-23 months who have received any vaccine (%)	95.8	93.8
47. Children age 12-23 months who have received BCG (%)	94.6	92.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	79.2	74.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	85.1	77.7
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	88.2	80.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	84.6	66.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	42.3	29.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.3	55.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	66.3	72.0
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	100.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	19.3	3.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.1	2.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	56.1	(60.8)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	47.1	8.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(28.8)	71.5
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(92.9)	96.6
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(73.0)	86.1
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.2	4.2
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.2	4.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	37.8	43.4
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.4	12.5
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	3.4	3.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	44.3	27.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.0	2.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Mirzapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.7	20.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	23.4	22.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.3	3.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.4	5.1
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.3	9.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.6	5.4
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.3	5.0
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.5	11.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.5	13.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.8	6.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.3	21.7
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.8	18.6
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.4	8.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.5	29.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	88.7	86.6
88. Women having a mobile phone that they themselves use (%)	55.7	49.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	74.2	75.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	13.0	5.4
91. Men age 15 years and above who use any kind of tobacco (%)	50.1	42.2
92. Women age 15 years and above who consume alcohol (%)	0.3	0.1
93. Men age 15 years and above who consume alcohol (%)	22.0	14.0

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Moradabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.4	9.0
2. Population below age 15 years (%)	29.8	29.4
3. Population age 60 years and above (%)	7.9	8.5
4. Population living in households with electricity (%)	98.0	96.0
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.8
6. Households using iodized salt (%)	93.9	95.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	42.5	17.1
8. Households with any usual member having a bank account/post office account (%)	98.2	96.1
9. Female population age 6 years and above who ever attended school (%)	65.4	67.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	16.9	7.5
11. Children age 2-4 years who attended pre-school (%)	26.9	11.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	40.0	42.6
13. Men with 10 or more years of schooling (%)	45.0	43.5
14. Women who have ever used the internet (%)	67.7	36.2
15. Men who have ever used the internet (%)	78.8	55.0
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	6.4	9.6
17. Men age 25-29 years married before age 21 years (%)	*	(19.9)
18. Three or more births in the 5 years preceding the survey (%)	2.8	4.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.3	1.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.1	69.8
21. Any modern method ² (%)	33.8	47.7
22. Any traditional method (%)	27.3	22.1
23. Female sterilization (%)	11.9	10.2
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.7	6.9
26. Unmet need for spacing ³ (%)	4.8	2.9
27. Unmet need for limiting ³ (%)	3.9	4.0
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	70.8	73.7
29. Mothers who had any antenatal care visits (%)	98.7	97.0
30. Mothers who had at least 4 antenatal care visits (%)	59.1	33.2
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.3	95.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	20.0	17.2
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	11.0	2.9
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.6	97.6
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	88.5	80.3
36. Institutional births in public facility (%)	17.8	40.0
37. Births attended by skilled health personnel ⁵ (%)	91.4	85.6
38. Births delivered by caesarean section (%)	39.7	22.5
39. Births in a private health facility that were delivered by caesarean section (%)	53.4	43.0
40. Births in a public health facility that were delivered by caesarean section (%)	11.3	13.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Moradabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	88.4	76.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(5.6)	3.4
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.9	74.7
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	82.9	84.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	87.2	90.6
46. Children age 12-23 months who have received any vaccine (%)	93.7	97.2
47. Children age 12-23 months who have received BCG (%)	92.8	97.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	83.7	84.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	86.3	90.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.3	89.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	73.8	60.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	47.4	65.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	85.5	61.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	74.8	74.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	84.5	96.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.9	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	12.2	7.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	4.4	2.2
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	63.1	(71.0)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	25.1	17.0
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(27.4)	(56.0)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(91.7)	(100.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(91.7)	(80.4)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(40.2)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	12.4	9.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	12.6	(24.8)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	12.4	12.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	25.9	34.7
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	17.8	19.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.0	9.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	29.0	27.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.7	2.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Moradabad, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.8	13.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	30.9	17.0
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.1	5.2
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.1	3.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.7	9.5
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.7	8.1
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.3	3.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.4	12.9
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.6	13.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.4	6.4
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.3	21.3
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.2	18.3
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.6	6.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.3	25.6
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.9	77.6
88. Women having a mobile phone that they themselves use (%)	48.5	42.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	66.3	78.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.5	3.6
91. Men age 15 years and above who use any kind of tobacco (%)	30.3	32.6
92. Women age 15 years and above who consume alcohol (%)	0.2	0.3
93. Men age 15 years and above who consume alcohol (%)	11.5	13.8

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Muzaffarnagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.6	8.9
2. Population below age 15 years (%)	29.4	30.4
3. Population age 60 years and above (%)	10.2	9.9
4. Population living in households with electricity (%)	99.6	97.8
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.9
6. Households using iodized salt (%)	93.9	96.3
7. Households with any usual member covered under a health insurance/financing scheme (%)	37.3	8.2
8. Households with any usual member having a bank account/post office account (%)	97.4	95.7
9. Female population age 6 years and above who ever attended school (%)	68.4	69.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.3	11.5
11. Children age 2-4 years who attended pre-school (%)	33.7	12.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	40.3	41.8
13. Men with 10 or more years of schooling (%)	50.3	48.6
14. Women who have ever used the internet (%)	74.8	44.6
15. Men who have ever used the internet (%)	89.6	70.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.4	7.3
17. Men age 25-29 years married before age 21 years (%)	(13.9)	(12.2)
18. Three or more births in the 5 years preceding the survey (%)	4.4	4.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.4	1.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.9	73.2
21. Any modern method ² (%)	35.9	44.9
22. Any traditional method (%)	26.0	28.3
23. Female sterilization (%)	12.1	14.4
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	7.8	5.0
26. Unmet need for spacing ³ (%)	3.3	1.9
27. Unmet need for limiting ³ (%)	4.5	3.1
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	75.7	75.5
29. Mothers who had any antenatal care visits (%)	98.0	96.0
30. Mothers who had at least 4 antenatal care visits (%)	46.3	44.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.7	94.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.3	34.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	23.2	8.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	92.1	96.4
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	88.6	87.0
36. Institutional births in public facility (%)	44.7	45.0
37. Births attended by skilled health personnel ⁵ (%)	90.3	89.9
38. Births delivered by caesarean section (%)	20.8	22.4
39. Births in a private health facility that were delivered by caesarean section (%)	35.2	33.3
40. Births in a public health facility that were delivered by caesarean section (%)	11.9	18.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Muzaffarnagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.7	88.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.1	87.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	80.2	69.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	88.2	79.6
46. Children age 12-23 months who have received any vaccine (%)	100.0	92.2
47. Children age 12-23 months who have received BCG (%)	100.0	91.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	81.5	74.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	88.8	75.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.9	77.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	65.3	66.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	58.2	84.1
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	84.2	62.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	65.2	72.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	90.7	93.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.2	4.1
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.4	6.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.5	0.5
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	75.1	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	48.0	18.2
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(14.1)	61.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(91.9)	92.1
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(89.5)	93.3
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(54.8)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	12.3	18.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(4.9)	(10.6)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.0	16.1
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	24.0	29.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	20.6	20.7
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.2	9.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	30.8	29.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.7	4.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Muzaffarnagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.5	11.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	31.0	28.1
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.5	5.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.3	5.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.6	11.5
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.3	8.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.9	5.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.5	14.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.8	14.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.3	5.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.5	21.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.5	16.6
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	6.8
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.5	24.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.1	75.1
88. Women having a mobile phone that they themselves use (%)	51.4	40.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	66.9	78.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.3	3.7
91. Men age 15 years and above who use any kind of tobacco (%)	32.6	31.7
92. Women age 15 years and above who consume alcohol (%)	0.0	0.3
93. Men age 15 years and above who consume alcohol (%)	11.3	13.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Pilibhit, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.8	8.5
2. Population below age 15 years (%)	26.8	29.6
3. Population age 60 years and above (%)	11.2	8.9
4. Population living in households with electricity (%)	92.4	86.8
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	93.5	91.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	45.8	21.4
8. Households with any usual member having a bank account/post office account (%)	98.3	96.6
9. Female population age 6 years and above who ever attended school (%)	63.2	60.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	19.7	11.7
11. Children age 2-4 years who attended pre-school (%)	28.5	13.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	34.3	28.5
13. Men with 10 or more years of schooling (%)	45.9	36.0
14. Women who have ever used the internet (%)	66.2	23.9
15. Men who have ever used the internet (%)	85.8	41.6
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	18.4	16.2
17. Men age 25-29 years married before age 21 years (%)	*	(18.7)
18. Three or more births in the 5 years preceding the survey (%)	2.4	1.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.1	4.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	62.8	74.1
21. Any modern method ² (%)	29.9	45.5
22. Any traditional method (%)	32.9	28.6
23. Female sterilization (%)	12.9	14.4
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.0	4.9
26. Unmet need for spacing ³ (%)	4.2	2.5
27. Unmet need for limiting ³ (%)	4.8	2.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	73.4	67.5
29. Mothers who had any antenatal care visits (%)	98.6	95.7
30. Mothers who had at least 4 antenatal care visits (%)	52.3	43.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	97.1	91.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.1	26.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	18.6	4.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.7	98.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	83.8	76.2
36. Institutional births in public facility (%)	45.2	50.2
37. Births attended by skilled health personnel ⁵ (%)	84.7	80.2
38. Births delivered by caesarean section (%)	23.5	19.0
39. Births in a private health facility that were delivered by caesarean section (%)	55.2	54.2
40. Births in a public health facility that were delivered by caesarean section (%)	4.9	9.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Pilibhit, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	70.6	72.6
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(1.9)	1.6
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	73.6	71.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	82.9	92.3
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	86.0	90.4
46. Children age 12-23 months who have received any vaccine (%)	96.3	97.0
47. Children age 12-23 months who have received BCG (%)	92.8	97.0
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	85.2	92.3
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.9	94.0
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.3	95.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	75.3	69.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	68.1	79.8
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	89.9	70.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	64.6	78.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	97.2
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	1.5
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	14.0	8.6
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.9	3.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	64.1	(69.7)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	43.6	29.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(13.1)	(54.6)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(93.6)	(91.5)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(89.4)	(79.1)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(39.5)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.3	10.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(12.9)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.1	8.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	29.8	38.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	27.4	20.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.1	7.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	39.8	39.4
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.1	1.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Pilibhit, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	25.0	16.4
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	24.7	21.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.0	5.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.2	5.0
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.8	10.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.3	6.4
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.1	6.6
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.6	13.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.7	11.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	5.5
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.0	19.5
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.6	11.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.4	5.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.5	17.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.3	84.7
88. Women having a mobile phone that they themselves use (%)	41.4	38.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	62.6	71.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.9	4.9
91. Men age 15 years and above who use any kind of tobacco (%)	44.8	38.2
92. Women age 15 years and above who consume alcohol (%)	0.1	0.2
93. Men age 15 years and above who consume alcohol (%)	21.4	15.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Pratapgarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.5	9.7
2. Population below age 15 years (%)	27.1	29.0
3. Population age 60 years and above (%)	11.4	12.9
4. Population living in households with electricity (%)	96.7	92.5
5. Population living in households with an improved drinking-water source ¹ (%)	97.6	97.8
6. Households using iodized salt (%)	96.9	95.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.5	14.9
8. Households with any usual member having a bank account/post office account (%)	99.5	97.5
9. Female population age 6 years and above who ever attended school (%)	70.1	67.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	24.9	14.4
11. Children age 2-4 years who attended pre-school (%)	25.6	5.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	51.6	49.7
13. Men with 10 or more years of schooling (%)	49.1	61.4
14. Women who have ever used the internet (%)	80.0	32.4
15. Men who have ever used the internet (%)	84.4	64.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	7.7	11.0
17. Men age 25-29 years married before age 21 years (%)	(17.1)	*
18. Three or more births in the 5 years preceding the survey (%)	2.0	2.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.2	0.7
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	64.2	51.9
21. Any modern method ² (%)	31.8	33.6
22. Any traditional method (%)	32.4	18.3
23. Female sterilization (%)	22.4	24.2
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.0	21.9
26. Unmet need for spacing ³ (%)	5.2	8.7
27. Unmet need for limiting ³ (%)	4.8	13.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	70.7	57.2
29. Mothers who had any antenatal care visits (%)	97.0	97.8
30. Mothers who had at least 4 antenatal care visits (%)	57.5	30.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.6	93.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	31.4	22.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	21.5	8.2
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.1	94.1
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	92.1	90.8
36. Institutional births in public facility (%)	57.7	62.6
37. Births attended by skilled health personnel ⁵ (%)	92.3	93.1
38. Births delivered by caesarean section (%)	19.0	11.8
39. Births in a private health facility that were delivered by caesarean section (%)	41.6	28.3
40. Births in a public health facility that were delivered by caesarean section (%)	8.2	6.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Pratapgarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.7	71.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	82.7	67.3
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	88.3	75.0
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	95.1	92.1
46. Children age 12-23 months who have received any vaccine (%)	95.5	92.6
47. Children age 12-23 months who have received BCG (%)	92.5	89.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	95.5	78.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	92.6	82.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	92.7	87.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	82.6	58.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	34.8	15.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	89.7	42.8
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	65.8	69.3
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	96.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	16.2	2.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.3	1.6
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	55.5	(74.9)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	50.6	13.2
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(19.7)	(51.9)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(87.0)	(90.6)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(70.3)	(84.2)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	19.0	5.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	(0.0)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	18.8	4.2
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	26.3	35.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.9	10.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.5	2.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	39.0	27.7
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.6	1.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Pratapgarh, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.4	19.6
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	25.5	25.0
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.6	6.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.8	6.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.7	14.1
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.3	8.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	12.5	10.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	24.1	19.1
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.7	10.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.3	4.9
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.7	17.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.5	13.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.3	6.1
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.0	22.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	89.7	74.9
88. Women having a mobile phone that they themselves use (%)	61.9	50.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	65.6	69.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	17.4	11.9
91. Men age 15 years and above who use any kind of tobacco (%)	53.6	41.4
92. Women age 15 years and above who consume alcohol (%)	0.2	0.5
93. Men age 15 years and above who consume alcohol (%)	20.6	9.0

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Prayagraj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.9	9.9
2. Population below age 15 years (%)	27.4	31.5
3. Population age 60 years and above (%)	12.4	10.5
4. Population living in households with electricity (%)	97.3	95.0
5. Population living in households with an improved drinking-water source ¹ (%)	97.8	95.8
6. Households using iodized salt (%)	90.2	96.8
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.4	20.6
8. Households with any usual member having a bank account/post office account (%)	99.2	99.1
9. Female population age 6 years and above who ever attended school (%)	71.9	70.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	27.5	14.5
11. Children age 2-4 years who attended pre-school (%)	29.2	13.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	55.3	47.6
13. Men with 10 or more years of schooling (%)	61.3	65.1
14. Women who have ever used the internet (%)	75.6	32.4
15. Men who have ever used the internet (%)	84.4	64.9
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	10.7	13.8
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.4	6.3
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.0	2.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	63.6	63.1
21. Any modern method ² (%)	37.5	45.6
22. Any traditional method (%)	26.1	17.5
23. Female sterilization (%)	26.6	31.3
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.6	11.9
26. Unmet need for spacing ³ (%)	5.5	4.5
27. Unmet need for limiting ³ (%)	5.1	7.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	77.4	58.3
29. Mothers who had any antenatal care visits (%)	98.8	93.6
30. Mothers who had at least 4 antenatal care visits (%)	71.6	45.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.7	88.3
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.8	22.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	20.8	12.0
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.3	95.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	85.3	83.2
36. Institutional births in public facility (%)	43.6	52.6
37. Births attended by skilled health personnel ⁵ (%)	88.0	85.5
38. Births delivered by caesarean section (%)	26.1	15.7
39. Births in a private health facility that were delivered by caesarean section (%)	49.6	43.4
40. Births in a public health facility that were delivered by caesarean section (%)	12.4	4.6

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Prayagraj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.0	67.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(0.0)	(4.2)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	77.7	59.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	75.6	55.8
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.2	67.8
46. Children age 12-23 months who have received any vaccine (%)	97.5	92.2
47. Children age 12-23 months who have received BCG (%)	94.9	85.4
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	96.2	58.4
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	76.8	67.2
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	86.0	70.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	76.5	49.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	46.1	28.6
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	76.8	53.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	56.9	60.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	93.8	95.6
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	6.2	1.7
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	18.5	3.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.7	0.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	59.6	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	42.1	24.8
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(21.9)	(53.4)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	(96.8)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(73.5)	(88.9)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(29.9)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	37.9	0.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	33.8	0.0
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.8	37.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.4	15.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.1	4.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	37.0	32.6
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	1.0

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Prayagraj, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	18.1	17.3
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	30.0	25.5
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.1	6.3
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.6	6.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.0	13.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.2	7.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.3	7.3
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.3	15.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.1	7.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.4	3.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.8	12.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.0	11.9
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	4.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.4	17.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.4	72.4
88. Women having a mobile phone that they themselves use (%)	63.0	50.7
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	74.2	79.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	13.3	11.9
91. Men age 15 years and above who use any kind of tobacco (%)	46.0	44.6
92. Women age 15 years and above who consume alcohol (%)	0.4	0.2
93. Men age 15 years and above who consume alcohol (%)	15.5	12.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Rae Bareli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.0	9.0
2. Population below age 15 years (%)	28.0	30.4
3. Population age 60 years and above (%)	11.1	11.0
4. Population living in households with electricity (%)	94.8	88.9
5. Population living in households with an improved drinking-water source ¹ (%)	99.5	98.1
6. Households using iodized salt (%)	74.0	79.5
7. Households with any usual member covered under a health insurance/financing scheme (%)	38.6	16.8
8. Households with any usual member having a bank account/post office account (%)	99.3	94.7
9. Female population age 6 years and above who ever attended school (%)	72.4	66.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	25.5	11.7
11. Children age 2-4 years who attended pre-school (%)	37.7	19.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	41.1	36.1
13. Men with 10 or more years of schooling (%)	43.2	41.3
14. Women who have ever used the internet (%)	54.8	16.0
15. Men who have ever used the internet (%)	74.6	43.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	11.1	14.3
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.2	2.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.4	1.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	56.1	69.3
21. Any modern method ² (%)	25.2	55.8
22. Any traditional method (%)	30.9	13.5
23. Female sterilization (%)	10.6	14.1
24. Male sterilization (%)	0.3	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	13.4	12.4
26. Unmet need for spacing ³ (%)	6.0	3.6
27. Unmet need for limiting ³ (%)	7.4	8.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	65.7	58.8
29. Mothers who had any antenatal care visits (%)	96.3	87.8
30. Mothers who had at least 4 antenatal care visits (%)	46.4	43.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.3	90.7
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	29.0	16.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	17.7	4.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	97.1	97.3
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.4	89.5
36. Institutional births in public facility (%)	75.9	78.5
37. Births attended by skilled health personnel ⁵ (%)	91.5	90.8
38. Births delivered by caesarean section (%)	13.3	8.8
39. Births in a private health facility that were delivered by caesarean section (%)	59.0	(50.8)
40. Births in a public health facility that were delivered by caesarean section (%)	6.3	4.1

Note: Major indicators are highlighted in grey.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Rae Bareli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.8	60.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(10.4)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.7	61.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	83.2	71.4
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	90.9	77.2
46. Children age 12-23 months who have received any vaccine (%)	97.3	95.2
47. Children age 12-23 months who have received BCG (%)	95.7	93.9
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	86.3	76.6
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.9	81.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.9	82.8
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	70.4	58.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	76.4	41.1
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.0	27.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	72.9	73.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	93.7	97.3
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	3.6	2.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.8	4.0
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	61.4	(37.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	52.5	28.6
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(75.8)	(64.4)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.7)	(93.4)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(94.6)	(80.9)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	7.0	3.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	6.1	3.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	35.5	47.0
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	9.6	13.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	2.4	6.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	29.2	28.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	4.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Rae Bareli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	20.6	18.5
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	22.9	18.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.3	3.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.9	3.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.7	8.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.1	3.1
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.8	3.2
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.3	7.4
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.0	11.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	7.0	5.9
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	16.8	19.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	12.3	11.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	6.8	4.8
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	20.1	17.9
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.4	60.3
88. Women having a mobile phone that they themselves use (%)	57.8	42.4
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	67.1	70.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	14.9	11.8
91. Men age 15 years and above who use any kind of tobacco (%)	57.5	50.4
92. Women age 15 years and above who consume alcohol (%)	0.6	0.5
93. Men age 15 years and above who consume alcohol (%)	17.3	16.5

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Rampur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.5	9.4
2. Population below age 15 years (%)	29.9	31.1
3. Population age 60 years and above (%)	8.0	9.1
4. Population living in households with electricity (%)	96.1	96.2
5. Population living in households with an improved drinking-water source ¹ (%)	98.4	100.0
6. Households using iodized salt (%)	91.7	90.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	34.5	13.1
8. Households with any usual member having a bank account/post office account (%)	98.4	93.6
9. Female population age 6 years and above who ever attended school (%)	60.3	57.6
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	14.0	9.3
11. Children age 2-4 years who attended pre-school (%)	19.4	17.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	31.8	26.9
13. Men with 10 or more years of schooling (%)	24.5	35.1
14. Women who have ever used the internet (%)	63.6	23.8
15. Men who have ever used the internet (%)	74.8	57.2
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	9.0	10.5
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	2.2	3.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.1	3.5
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.4	56.8
21. Any modern method ² (%)	37.7	30.0
22. Any traditional method (%)	23.7	26.8
23. Female sterilization (%)	9.7	9.3
24. Male sterilization (%)	0.3	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.6	12.0
26. Unmet need for spacing ³ (%)	4.4	5.8
27. Unmet need for limiting ³ (%)	4.2	6.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	68.8	73.8
29. Mothers who had any antenatal care visits (%)	98.5	98.4
30. Mothers who had at least 4 antenatal care visits (%)	36.5	50.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.4	94.5
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	25.8	32.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	13.4	12.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.3	97.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.2	81.0
36. Institutional births in public facility (%)	33.6	44.7
37. Births attended by skilled health personnel ⁵ (%)	92.2	81.1
38. Births delivered by caesarean section (%)	30.2	16.7
39. Births in a private health facility that were delivered by caesarean section (%)	50.2	40.0
40. Births in a public health facility that were delivered by caesarean section (%)	5.1	4.8

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Rampur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	75.4	72.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(2.9)	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.5	72.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	82.8	79.4
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.3	81.9
46. Children age 12-23 months who have received any vaccine (%)	98.4	96.1
47. Children age 12-23 months who have received BCG (%)	97.8	96.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	82.8	80.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	98.4	88.7
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	98.4	91.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	66.8	56.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	44.6	54.1
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	84.1	61.8
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	65.1	56.9
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	71.2	97.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	8.6	1.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.7	5.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.2	5.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	90.8	(85.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	41.8	39.6
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	31.0	(43.4)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	94.6	(97.1)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	78.2	(88.4)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(47.8)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.3	7.7
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(1.0)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	7.1	6.9
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	28.0	40.4
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	21.0	17.6
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.3	5.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	34.6	32.1
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	2.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Rampur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.3	13.6
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	31.6	20.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.2	6.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.4	5.5
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	13.1	12.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.3	7.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.5	6.0
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.3	14.4
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.3	10.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.1	5.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.9	17.7
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.9	15.5
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.7	4.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.1	21.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.8	68.9
88. Women having a mobile phone that they themselves use (%)	52.1	35.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	67.0	74.3
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.8	4.9
91. Men age 15 years and above who use any kind of tobacco (%)	32.7	36.1
92. Women age 15 years and above who consume alcohol (%)	0.1	0.1
93. Men age 15 years and above who consume alcohol (%)	12.8	14.6

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Saharanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.8	9.8
2. Population below age 15 years (%)	30.8	28.5
3. Population age 60 years and above (%)	8.1	8.8
4. Population living in households with electricity (%)	98.7	98.3
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.7
6. Households using iodized salt (%)	96.3	96.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	44.7	11.8
8. Households with any usual member having a bank account/post office account (%)	99.1	95.8
9. Female population age 6 years and above who ever attended school (%)	70.8	69.3
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	19.6	9.5
11. Children age 2-4 years who attended pre-school (%)	36.2	10.3
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	37.6	38.2
13. Men with 10 or more years of schooling (%)	43.5	43.3
14. Women who have ever used the internet (%)	77.9	35.7
15. Men who have ever used the internet (%)	86.6	59.2
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	4.3	7.0
17. Men age 25-29 years married before age 21 years (%)	(8.4)	*
18. Three or more births in the 5 years preceding the survey (%)	4.1	4.7
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.5	1.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	61.9	73.7
21. Any modern method ² (%)	33.3	48.9
22. Any traditional method (%)	28.6	24.8
23. Female sterilization (%)	9.2	9.5
24. Male sterilization (%)	0.2	0.3
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	7.8	4.3
26. Unmet need for spacing ³ (%)	3.3	1.6
27. Unmet need for limiting ³ (%)	4.5	2.7
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	82.3	78.7
29. Mothers who had any antenatal care visits (%)	99.3	96.5
30. Mothers who had at least 4 antenatal care visits (%)	73.0	52.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.8	97.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	32.9	25.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.4	7.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	89.9	96.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	82.9	81.7
36. Institutional births in public facility (%)	40.9	53.5
37. Births attended by skilled health personnel ⁵ (%)	86.1	86.9
38. Births delivered by caesarean section (%)	21.7	15.4
39. Births in a private health facility that were delivered by caesarean section (%)	40.9	37.6
40. Births in a public health facility that were delivered by caesarean section (%)	11.1	9.1

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Saharanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	84.7	83.6
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	7.0	(3.7)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	84.6	79.1
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	86.5	89.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	92.2	93.4
46. Children age 12-23 months who have received any vaccine (%)	99.0	97.3
47. Children age 12-23 months who have received BCG (%)	96.8	97.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	87.8	91.0
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	92.7	92.9
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	94.6	93.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	75.9	82.0
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	62.8	82.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	91.5	56.9
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	65.5	82.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	85.2	93.7
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.9	4.5
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	15.3	6.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.1	3.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	72.9	(62.4)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	37.1	26.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(21.2)	67.4
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(93.1)	96.7
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(77.3)	90.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(65.6)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	9.0	5.4
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(14.3)	(6.3)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.4	5.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	22.9	28.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	23.1	22.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.2	12.9
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	31.8	26.7
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.7	1.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Saharanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	21.8	12.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	32.2	16.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.6	4.8
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.0	3.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.9	8.8
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.0	8.7
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.7	2.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.9	11.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.0	15.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.9	6.5
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.7	23.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.2	18.2
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.5	5.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.7	24.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	80.9	80.7
88. Women having a mobile phone that they themselves use (%)	53.3	40.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	72.6	76.7
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.4	2.4
91. Men age 15 years and above who use any kind of tobacco (%)	31.2	30.7
92. Women age 15 years and above who consume alcohol (%)	0.1	0.5
93. Men age 15 years and above who consume alcohol (%)	11.3	14.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Sambhal, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.2	11.2
2. Population below age 15 years (%)	35.1	35.5
3. Population age 60 years and above (%)	6.2	6.6
4. Population living in households with electricity (%)	90.3	91.9
5. Population living in households with an improved drinking-water source ¹ (%)	98.8	100.0
6. Households using iodized salt (%)	88.7	91.1
7. Households with any usual member covered under a health insurance/financing scheme (%)	33.0	13.7
8. Households with any usual member having a bank account/post office account (%)	98.2	96.9
9. Female population age 6 years and above who ever attended school (%)	56.8	56.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	14.4	13.0
11. Children age 2-4 years who attended pre-school (%)	23.4	9.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	23.9	24.1
13. Men with 10 or more years of schooling (%)	39.0	35.1
14. Women who have ever used the internet (%)	56.7	24.8
15. Men who have ever used the internet (%)	69.7	53.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	19.1	21.1
17. Men age 25-29 years married before age 21 years (%)	(27.8)	(39.4)
18. Three or more births in the 5 years preceding the survey (%)	6.1	4.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	7.6	4.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	64.6	65.9
21. Any modern method ² (%)	32.0	40.1
22. Any traditional method (%)	32.6	25.8
23. Female sterilization (%)	8.8	8.6
24. Male sterilization (%)	0.2	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	10.3	8.5
26. Unmet need for spacing ³ (%)	3.0	3.9
27. Unmet need for limiting ³ (%)	7.3	4.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	63.5	60.8
29. Mothers who had any antenatal care visits (%)	96.1	90.9
30. Mothers who had at least 4 antenatal care visits (%)	35.8	32.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.9	90.3
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	20.8	21.3
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	8.6	5.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.1	93.5
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	77.4	74.2
36. Institutional births in public facility (%)	40.7	45.1
37. Births attended by skilled health personnel ⁵ (%)	80.7	78.2
38. Births delivered by caesarean section (%)	11.5	8.1
39. Births in a private health facility that were delivered by caesarean section (%)	28.9	25.5
40. Births in a public health facility that were delivered by caesarean section (%)	2.3	1.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Sambhal, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	62.4	67.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	72.7	67.5
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	88.9	83.0
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	95.9	89.6
46. Children age 12-23 months who have received any vaccine (%)	98.8	96.6
47. Children age 12-23 months who have received BCG (%)	96.7	95.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	91.0	84.7
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	96.0	89.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	97.1	91.3
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	75.8	72.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	56.7	72.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	97.1	73.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	68.0	72.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.1	86.1
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.9	0.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.6	5.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9	2.1
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	88.0	73.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	47.3	31.5
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(18.4)	62.5
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(85.5)	89.8
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(88.9)	84.1
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(44.2)	(25.7)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.7	5.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(3.7)	(3.9)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	7.2	5.6
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	42.3	51.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	24.2	14.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	8.8	5.6
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	47.2	29.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	3.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Sambhal, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	25.5	16.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	21.1	19.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.2	3.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.8	3.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	11.4	7.4
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.1	4.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.5	3.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.0	8.5
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.4	14.1
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.0	4.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.7	19.3
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.8	16.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.9	5.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	7.2	22.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	78.2	75.2
88. Women having a mobile phone that they themselves use (%)	41.7	32.0
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	50.6	60.2
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	4.2	4.9
91. Men age 15 years and above who use any kind of tobacco (%)	40.6	40.3
92. Women age 15 years and above who consume alcohol (%)	0.2	0.2
93. Men age 15 years and above who consume alcohol (%)	14.0	16.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Sant Kabir Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.3	10.3
2. Population below age 15 years (%)	32.3	34.6
3. Population age 60 years and above (%)	10.5	11.4
4. Population living in households with electricity (%)	97.6	94.0
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	99.9
6. Households using iodized salt (%)	93.1	98.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	44.7	20.4
8. Households with any usual member having a bank account/post office account (%)	99.6	98.4
9. Female population age 6 years and above who ever attended school (%)	69.8	63.8
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.3	13.9
11. Children age 2-4 years who attended pre-school (%)	19.5	12.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	42.1	34.8
13. Men with 10 or more years of schooling (%)	43.1	48.3
14. Women who have ever used the internet (%)	77.5	19.0
15. Men who have ever used the internet (%)	84.1	53.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	7.6	16.8
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.1	2.5
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	0.7	1.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	49.4	69.2
21. Any modern method ² (%)	19.5	48.5
22. Any traditional method (%)	29.9	20.7
23. Female sterilization (%)	10.2	13.6
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	16.3	13.0
26. Unmet need for spacing ³ (%)	7.7	6.6
27. Unmet need for limiting ³ (%)	8.6	6.4
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	74.0	62.1
29. Mothers who had any antenatal care visits (%)	96.5	95.6
30. Mothers who had at least 4 antenatal care visits (%)	42.0	43.2
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.5	93.7
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.1	23.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	18.4	9.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	96.4	98.3
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	90.0	90.6
36. Institutional births in public facility (%)	59.0	79.5
37. Births attended by skilled health personnel ⁵ (%)	91.5	89.7
38. Births delivered by caesarean section (%)	17.4	8.3
39. Births in a private health facility that were delivered by caesarean section (%)	47.1	33.0
40. Births in a public health facility that were delivered by caesarean section (%)	4.8	5.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Sant Kabir Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.7	72.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(12.8)	(11.5)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.0	75.1
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	83.4	80.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	94.4	82.9
46. Children age 12-23 months who have received any vaccine (%)	94.1	96.5
47. Children age 12-23 months who have received BCG (%)	94.1	95.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	83.4	85.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.7	87.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	93.0	95.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	75.4	61.8
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	57.3	39.3
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	89.8	37.7
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	69.0	84.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	97.2
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.0	3.5
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.7	4.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	55.5	60.3
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	46.3	12.5
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(21.1)	(61.7)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(97.9)	(96.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(52.2)	(79.3)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	(22.8)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	44.6	7.3
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	44.6	7.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	28.8	42.3
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	24.6	19.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.8	8.8
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	38.6	34.2
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	2.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Sant Kabir Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.0	19.3
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.8	15.9
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.3	4.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.1	3.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.1	8.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.6	5.1
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	13.3	4.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	25.4	10.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.7	12.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.1	5.7
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.8	22.5
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.0	18.0
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.0	7.3
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	14.6	28.2
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.0	84.8
88. Women having a mobile phone that they themselves use (%)	61.2	54.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	67.1	63.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	10.0	9.9
91. Men age 15 years and above who use any kind of tobacco (%)	53.0	51.9
92. Women age 15 years and above who consume alcohol (%)	0.2	0.3
93. Men age 15 years and above who consume alcohol (%)	22.1	14.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Shahjahanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.0	11.6
2. Population below age 15 years (%)	31.3	34.4
3. Population age 60 years and above (%)	8.3	8.3
4. Population living in households with electricity (%)	91.0	84.1
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	100.0
6. Households using iodized salt (%)	90.4	79.8
7. Households with any usual member covered under a health insurance/financing scheme (%)	36.2	14.2
8. Households with any usual member having a bank account/post office account (%)	98.6	97.7
9. Female population age 6 years and above who ever attended school (%)	66.7	63.2
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	21.8	10.6
11. Children age 2-4 years who attended pre-school (%)	27.1	20.0
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	34.7	27.7
13. Men with 10 or more years of schooling (%)	47.1	38.8
14. Women who have ever used the internet (%)	67.0	26.3
15. Men who have ever used the internet (%)	78.7	47.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	18.7	20.9
17. Men age 25-29 years married before age 21 years (%)	(23.0)	(32.9)
18. Three or more births in the 5 years preceding the survey (%)	3.1	4.2
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	7.1	4.6
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	64.7	70.3
21. Any modern method ² (%)	28.5	46.4
22. Any traditional method (%)	36.2	23.9
23. Female sterilization (%)	8.1	10.0
24. Male sterilization (%)	0.1	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.4	6.4
26. Unmet need for spacing ³ (%)	4.0	2.6
27. Unmet need for limiting ³ (%)	5.4	3.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	70.8	61.4
29. Mothers who had any antenatal care visits (%)	97.3	96.6
30. Mothers who had at least 4 antenatal care visits (%)	46.1	35.3
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.0	95.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	29.0	24.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	14.5	5.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.6	98.2
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	72.8	63.3
36. Institutional births in public facility (%)	47.1	47.8
37. Births attended by skilled health personnel ⁵ (%)	76.6	68.3
38. Births delivered by caesarean section (%)	13.9	8.6
39. Births in a private health facility that were delivered by caesarean section (%)	50.0	42.2
40. Births in a public health facility that were delivered by caesarean section (%)	2.4	4.3

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Shahjahanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	61.6	62.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.0	0.7
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	61.8	63.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	85.8	76.5
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	94.7	84.5
46. Children age 12-23 months who have received any vaccine (%)	95.5	97.3
47. Children age 12-23 months who have received BCG (%)	94.5	97.3
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	85.8	80.8
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.6	88.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.6	89.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	77.9	73.1
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	71.7	52.5
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	90.6	48.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	70.2	76.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.1	97.2
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.9	2.8
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	10.4	5.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.9	7.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	75.8	65.8
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	52.6	27.6
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	25.8	55.1
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	87.9	96.4
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	82.9	85.2
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(43.5)	(36.1)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	10.0	5.7
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(2.8)	(9.5)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	8.5	6.7
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.9	44.5
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	24.5	17.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.2	8.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	37.7	34.7
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	1.3	3.2

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Shahjahanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	23.9	17.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	25.7	17.2
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	5.9	4.5
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.5	4.3
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.2	9.1
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.6	3.9
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.4	2.5
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	14.4	6.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.0	11.2
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	4.1
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.1	16.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.5	11.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	3.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.9	14.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.4	80.9
88. Women having a mobile phone that they themselves use (%)	46.6	29.1
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	62.9	59.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	5.3	6.6
91. Men age 15 years and above who use any kind of tobacco (%)	44.1	49.0
92. Women age 15 years and above who consume alcohol (%)	0.1	0.2
93. Men age 15 years and above who consume alcohol (%)	16.9	20.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Shamli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	9.6	10.2
2. Population below age 15 years (%)	29.8	31.3
3. Population age 60 years and above (%)	9.9	9.7
4. Population living in households with electricity (%)	99.2	98.7
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.9
6. Households using iodized salt (%)	97.7	95.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	45.0	14.8
8. Households with any usual member having a bank account/post office account (%)	99.2	97.1
9. Female population age 6 years and above who ever attended school (%)	65.5	62.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	19.4	10.0
11. Children age 2-4 years who attended pre-school (%)	35.4	8.1
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	37.0	32.6
13. Men with 10 or more years of schooling (%)	53.2	41.4
14. Women who have ever used the internet (%)	78.1	32.2
15. Men who have ever used the internet (%)	91.4	59.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	8.2	10.8
17. Men age 25-29 years married before age 21 years (%)	(8.8)	(25.5)
18. Three or more births in the 5 years preceding the survey (%)	6.1	6.6
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.3	2.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	60.1	71.0
21. Any modern method ² (%)	32.3	45.6
22. Any traditional method (%)	27.8	25.4
23. Female sterilization (%)	8.8	10.9
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	9.7	5.6
26. Unmet need for spacing ³ (%)	4.6	2.3
27. Unmet need for limiting ³ (%)	5.1	3.3
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	79.6	82.7
29. Mothers who had any antenatal care visits (%)	98.9	97.9
30. Mothers who had at least 4 antenatal care visits (%)	56.1	41.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	90.8	97.6
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	30.0	25.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	15.1	6.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	91.5	98.3
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	82.7	78.3
36. Institutional births in public facility (%)	46.9	54.3
37. Births attended by skilled health personnel ⁵ (%)	86.3	83.3
38. Births delivered by caesarean section (%)	14.1	11.3
39. Births in a private health facility that were delivered by caesarean section (%)	34.5	35.7
40. Births in a public health facility that were delivered by caesarean section (%)	3.6	5.0

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Shamli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.0	81.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	9.4	1.2
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.1	75.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	77.7	91.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	85.9	95.3
46. Children age 12-23 months who have received any vaccine (%)	94.6	97.5
47. Children age 12-23 months who have received BCG (%)	94.4	97.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	79.9	93.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	84.0	91.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	84.5	94.6
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	85.7	75.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	55.4	92.5
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	84.3	76.9
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	61.3	84.0
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.1	93.0
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.9	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	13.1	1.9
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.9	1.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	80.8	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	29.9	19.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(33.4)	64.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(92.1)	94.3
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(93.7)	85.8
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(42.3)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	16.6	1.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(2.2)	(2.5)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	11.8	1.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	29.1	28.6
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	20.8	24.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.5	13.2
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	36.1	27.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	2.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Shamli, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	17.9	9.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	35.0	22.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.6	4.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.3	3.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.2	8.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.8	7.4
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.9	4.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	19.2	12.5
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.9	12.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.6	7.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.3	21.2
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.6	20.9
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.7	7.2
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.1	28.6
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	76.2	72.6
88. Women having a mobile phone that they themselves use (%)	52.1	34.2
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	65.6	74.1
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	6.0	3.7
91. Men age 15 years and above who use any kind of tobacco (%)	30.8	30.6
92. Women age 15 years and above who consume alcohol (%)	0.3	0.3
93. Men age 15 years and above who consume alcohol (%)	8.7	10.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Shrawasti, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	12.7	13.5
2. Population below age 15 years (%)	39.6	40.2
3. Population age 60 years and above (%)	7.8	8.0
4. Population living in households with electricity (%)	88.8	73.7
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	99.3
6. Households using iodized salt (%)	79.2	81.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	25.4	13.5
8. Households with any usual member having a bank account/post office account (%)	98.0	97.0
9. Female population age 6 years and above who ever attended school (%)	56.2	47.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.3	10.1
11. Children age 2-4 years who attended pre-school (%)	16.2	10.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	15.4	15.9
13. Men with 10 or more years of schooling (%)	24.2	31.1
14. Women who have ever used the internet (%)	65.5	17.4
15. Men who have ever used the internet (%)	79.9	45.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	45.4	51.9
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.8	3.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	5.5	5.3
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	44.9	49.8
21. Any modern method ² (%)	15.7	35.6
22. Any traditional method (%)	29.2	14.2
23. Female sterilization (%)	5.4	6.9
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	18.9	20.1
26. Unmet need for spacing ³ (%)	9.4	8.6
27. Unmet need for limiting ³ (%)	9.5	11.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	62.4	42.2
29. Mothers who had any antenatal care visits (%)	96.9	94.4
30. Mothers who had at least 4 antenatal care visits (%)	74.2	42.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	92.7	94.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	14.3	16.6
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	7.8	8.8
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	90.6	95.9
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	76.2	80.4
36. Institutional births in public facility (%)	63.2	75.3
37. Births attended by skilled health personnel ⁵ (%)	77.4	84.3
38. Births delivered by caesarean section (%)	7.2	3.2
39. Births in a private health facility that were delivered by caesarean section (%)	29.8	(28.0)
40. Births in a public health facility that were delivered by caesarean section (%)	5.2	2.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Shrawasti, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	72.3	68.3
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	8.2	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	71.2	68.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	72.4	59.6
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.5	69.8
46. Children age 12-23 months who have received any vaccine (%)	89.0	93.9
47. Children age 12-23 months who have received BCG (%)	86.2	92.2
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	75.3	65.5
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	79.8	76.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	82.3	78.1
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	73.9	53.7
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	41.7	54.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	78.8	33.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	60.3	81.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.0	98.7
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	1.0	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	15.8	9.8
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.9	6.7
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	39.0	59.1
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	44.1	14.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	22.5	70.2
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	95.9	92.9
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	67.8	82.5
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(90.7)	(25.5)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	33.6	4.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	(0.0)
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	31.0	3.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	42.7	50.9
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	18.2	20.3
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.4	9.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	40.4	40.8
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.3	3.3

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Shrawasti, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	27.8	22.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	20.6	13.7
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.0	3.9
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	6.9	3.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	15.4	8.1
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.4	4.6
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.7	3.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.6	9.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.9	9.6
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.0	4.6
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.9	18.4
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	7.4	13.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.2	3.5
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.8	18.7
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	76.1	79.2
88. Women having a mobile phone that they themselves use (%)	49.6	53.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	50.0	47.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	27.7	24.2
91. Men age 15 years and above who use any kind of tobacco (%)	67.5	64.2
92. Women age 15 years and above who consume alcohol (%)	0.2	0.3
93. Men age 15 years and above who consume alcohol (%)	10.7	8.0

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Siddharth Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	11.8	11.6
2. Population below age 15 years (%)	36.1	41.0
3. Population age 60 years and above (%)	10.0	8.9
4. Population living in households with electricity (%)	94.9	90.7
5. Population living in households with an improved drinking-water source ¹ (%)	99.9	99.9
6. Households using iodized salt (%)	93.2	99.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	28.9	12.2
8. Households with any usual member having a bank account/post office account (%)	99.5	98.7
9. Female population age 6 years and above who ever attended school (%)	62.3	57.7
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.1	13.7
11. Children age 2-4 years who attended pre-school (%)	15.2	12.8
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	27.6	17.0
13. Men with 10 or more years of schooling (%)	45.0	27.3
14. Women who have ever used the internet (%)	77.6	16.9
15. Men who have ever used the internet (%)	86.2	56.3
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	19.2	33.9
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.4	2.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.5	3.7
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	53.5	65.0
21. Any modern method ² (%)	19.2	52.5
22. Any traditional method (%)	34.3	12.5
23. Female sterilization (%)	8.5	6.1
24. Male sterilization (%)	0.1	0.2
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	15.0	14.4
26. Unmet need for spacing ³ (%)	6.8	6.2
27. Unmet need for limiting ³ (%)	8.2	8.2
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	68.5	62.8
29. Mothers who had any antenatal care visits (%)	98.6	93.8
30. Mothers who had at least 4 antenatal care visits (%)	60.7	60.9
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	91.2	93.8
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	17.5	31.7
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	11.9	17.9
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	94.3	98.3
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	77.5	69.7
36. Institutional births in public facility (%)	52.1	62.9
37. Births attended by skilled health personnel ⁵ (%)	79.9	67.6
38. Births delivered by caesarean section (%)	12.0	4.0
39. Births in a private health facility that were delivered by caesarean section (%)	42.5	(21.9)
40. Births in a public health facility that were delivered by caesarean section (%)	2.3	3.9

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Siddharth Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	71.5	61.8
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.9	6.2
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	69.8	57.2
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	83.2	65.1
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	92.6	68.1
46. Children age 12-23 months who have received any vaccine (%)	98.6	97.8
47. Children age 12-23 months who have received BCG (%)	97.3	97.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	86.0	70.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	88.2	77.1
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.8	87.9
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	87.8	68.5
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	41.6	38.7
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	85.7	53.6
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	56.8	82.7
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	99.4	97.4
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.6	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	11.9	4.1
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	0.2	4.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	37.9	51.0
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	46.1	24.9
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	17.1	70.4
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	96.6	94.5
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	72.6	80.6
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(83.0)	(15.1)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	31.5	2.0
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	31.0	3.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	35.4	37.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	23.6	24.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.5	14.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	44.5	36.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.2	2.8

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Siddharth Nagar, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	24.5	25.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	24.9	12.9
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.1	4.0
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.7	4.8
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.3	9.5
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.8	5.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	14.6	5.4
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	26.1	11.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.2	11.7
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.7	5.3
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.5	19.1
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	9.1	14.0
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	6.1	7.4
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	17.4	23.5
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.3	79.4
88. Women having a mobile phone that they themselves use (%)	60.6	61.6
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	61.8	63.0
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	15.8	14.6
91. Men age 15 years and above who use any kind of tobacco (%)	58.3	54.9
92. Women age 15 years and above who consume alcohol (%)	0.2	0.2
93. Men age 15 years and above who consume alcohol (%)	18.9	11.2

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Sitapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.6	10.6
2. Population below age 15 years (%)	33.5	33.6
3. Population age 60 years and above (%)	9.0	8.9
4. Population living in households with electricity (%)	89.9	68.4
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	99.7
6. Households using iodized salt (%)	71.5	96.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	38.5	23.2
8. Households with any usual member having a bank account/post office account (%)	98.5	97.6
9. Female population age 6 years and above who ever attended school (%)	65.8	65.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	17.8	11.5
11. Children age 2-4 years who attended pre-school (%)	27.6	9.9
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	27.2	23.9
13. Men with 10 or more years of schooling (%)	27.5	27.9
14. Women who have ever used the internet (%)	63.3	10.7
15. Men who have ever used the internet (%)	69.7	46.7
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	22.0	20.8
17. Men age 25-29 years married before age 21 years (%)	(21.9)	(42.2)
18. Three or more births in the 5 years preceding the survey (%)	4.5	4.0
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	7.2	3.4
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	50.1	51.1
21. Any modern method ² (%)	33.7	36.5
22. Any traditional method (%)	16.4	14.6
23. Female sterilization (%)	16.8	16.8
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	17.8	17.9
26. Unmet need for spacing ³ (%)	7.0	6.0
27. Unmet need for limiting ³ (%)	10.8	11.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	74.8	66.8
29. Mothers who had any antenatal care visits (%)	98.5	94.9
30. Mothers who had at least 4 antenatal care visits (%)	56.7	35.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	95.9	91.2
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	17.5	18.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	6.5	9.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	90.0	94.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	89.1	84.8
36. Institutional births in public facility (%)	69.3	70.4
37. Births attended by skilled health personnel ⁵ (%)	90.0	87.8
38. Births delivered by caesarean section (%)	10.3	8.4
39. Births in a private health facility that were delivered by caesarean section (%)	38.2	35.7
40. Births in a public health facility that were delivered by caesarean section (%)	4.0	4.6

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Sitapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	78.9	71.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(6.6)	2.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	85.8	67.9
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	85.6	65.9
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	94.0	72.2
46. Children age 12-23 months who have received any vaccine (%)	96.3	97.1
47. Children age 12-23 months who have received BCG (%)	96.3	91.7
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	86.7	71.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	90.1	81.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	91.1	80.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	79.4	60.2
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	77.7	54.9
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	91.1	51.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	60.6	89.4
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	95.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	1.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	21.7	9.5
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.5	5.4
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	53.8	53.9
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	36.1	24.3
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(46.5)	60.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(89.1)	87.1
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(92.1)	73.9
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(64.6)	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	24.0	3.1
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	20.7	4.1
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	45.5	47.8
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	24.0	18.2
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	7.6	6.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	49.9	37.9
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	2.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Sitapur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	34.5	30.1
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	14.8	15.4
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.5	3.7
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	7.0	3.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	16.1	7.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.5	4.5
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.7	4.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	21.8	9.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.6	11.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.6	5.5
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.1	17.8
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.8	16.1
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.2	5.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	7.4	23.0
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.9	67.0
88. Women having a mobile phone that they themselves use (%)	40.2	23.3
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	52.7	55.5
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	14.2	13.8
91. Men age 15 years and above who use any kind of tobacco (%)	55.3	59.7
92. Women age 15 years and above who consume alcohol (%)	0.1	0.1
93. Men age 15 years and above who consume alcohol (%)	19.7	13.8

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Sonbhadra, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	10.8	10.1
2. Population below age 15 years (%)	33.0	31.8
3. Population age 60 years and above (%)	8.6	9.5
4. Population living in households with electricity (%)	95.4	82.9
5. Population living in households with an improved drinking-water source ¹ (%)	91.4	89.6
6. Households using iodized salt (%)	82.4	98.2
7. Households with any usual member covered under a health insurance/financing scheme (%)	44.4	19.7
8. Households with any usual member having a bank account/post office account (%)	99.6	97.1
9. Female population age 6 years and above who ever attended school (%)	67.2	62.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	18.9	9.8
11. Children age 2-4 years who attended pre-school (%)	24.5	6.6
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	36.0	35.7
13. Men with 10 or more years of schooling (%)	56.7	36.4
14. Women who have ever used the internet (%)	69.7	21.3
15. Men who have ever used the internet (%)	81.7	44.1
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	26.5	17.7
17. Men age 25-29 years married before age 21 years (%)	(28.9)	*
18. Three or more births in the 5 years preceding the survey (%)	3.1	4.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	8.8	6.9
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	66.3	67.1
21. Any modern method ² (%)	45.2	53.6
22. Any traditional method (%)	21.1	13.5
23. Female sterilization (%)	37.5	33.7
24. Male sterilization (%)	0.2	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	11.8	11.6
26. Unmet need for spacing ³ (%)	6.1	4.7
27. Unmet need for limiting ³ (%)	5.7	6.9
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	67.0	63.6
29. Mothers who had any antenatal care visits (%)	96.2	93.4
30. Mothers who had at least 4 antenatal care visits (%)	38.8	36.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	88.8	86.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	34.9	28.0
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	17.6	12.5
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.1	95.8
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	72.0	76.8
36. Institutional births in public facility (%)	56.1	56.8
37. Births attended by skilled health personnel ⁵ (%)	74.7	77.1
38. Births delivered by caesarean section (%)	11.2	11.8
39. Births in a private health facility that were delivered by caesarean section (%)	49.2	48.6
40. Births in a public health facility that were delivered by caesarean section (%)	6.1	3.7

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Sonbhadra, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	68.2	59.9
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	0.9	0.0
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	70.1	57.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	75.5	72.7
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.9	82.3
46. Children age 12-23 months who have received any vaccine (%)	90.7	95.7
47. Children age 12-23 months who have received BCG (%)	88.0	93.6
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	80.1	80.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	78.2	83.6
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	80.8	81.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	68.6	57.9
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	53.9	29.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	77.2	51.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	72.8	83.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.6	98.9
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.4	0.0
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	16.2	2.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.6	2.9
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	48.9	(55.5)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	54.5	29.1
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(34.8)	66.8
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(98.2)	98.5
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(83.0)	77.7
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	36.5	1.9
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	36.1	2.5
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	39.0	38.3
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	25.6	26.8
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	6.6	17.4
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	46.6	46.5
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	0.9

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Sonbhadra, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	30.6	20.9
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	15.5	15.3
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	6.5	5.1
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	5.6	2.6
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	12.4	7.9
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	9.7	5.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.3	5.0
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.4	11.3
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	3.9	12.9
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.9	5.0
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	8.4	19.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.8	16.8
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.5	4.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.2	21.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	86.2	82.1
88. Women having a mobile phone that they themselves use (%)	52.3	38.5
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	61.2	68.9
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	13.3	5.8
91. Men age 15 years and above who use any kind of tobacco (%)	64.2	54.1
92. Women age 15 years and above who consume alcohol (%)	4.7	0.6
93. Men age 15 years and above who consume alcohol (%)	44.2	24.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Sultanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.9	8.7
2. Population below age 15 years (%)	28.7	31.4
3. Population age 60 years and above (%)	11.4	11.0
4. Population living in households with electricity (%)	98.5	94.3
5. Population living in households with an improved drinking-water source ¹ (%)	99.8	99.5
6. Households using iodized salt (%)	91.4	97.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	41.9	20.7
8. Households with any usual member having a bank account/post office account (%)	99.4	97.3
9. Female population age 6 years and above who ever attended school (%)	73.5	68.9
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	23.8	17.4
11. Children age 2-4 years who attended pre-school (%)	35.5	6.5
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	46.9	46.6
13. Men with 10 or more years of schooling (%)	47.8	41.9
14. Women who have ever used the internet (%)	77.4	21.5
15. Men who have ever used the internet (%)	85.6	60.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	4.8	7.9
17. Men age 25-29 years married before age 21 years (%)	(12.1)	*
18. Three or more births in the 5 years preceding the survey (%)	2.4	0.9
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	1.6	1.1
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	57.1	47.6
21. Any modern method ² (%)	23.1	24.3
22. Any traditional method (%)	34.0	23.3
23. Female sterilization (%)	12.1	13.6
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	13.7	23.9
26. Unmet need for spacing ³ (%)	6.9	9.1
27. Unmet need for limiting ³ (%)	6.8	14.8
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	74.8	59.6
29. Mothers who had any antenatal care visits (%)	97.4	95.2
30. Mothers who had at least 4 antenatal care visits (%)	44.5	47.0
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	94.5	93.4
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	28.4	27.8
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	13.4	12.4
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	93.8	97.7
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	91.5	87.0
36. Institutional births in public facility (%)	69.0	68.0
37. Births attended by skilled health personnel ⁵ (%)	91.6	83.7
38. Births delivered by caesarean section (%)	16.1	13.0
39. Births in a private health facility that were delivered by caesarean section (%)	53.2	45.1
40. Births in a public health facility that were delivered by caesarean section (%)	6.0	6.5

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Sultanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	80.4	70.5
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(3.9)	(0.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.2	69.4
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	84.5	80.2
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	92.3	92.7
46. Children age 12-23 months who have received any vaccine (%)	95.9	95.7
47. Children age 12-23 months who have received BCG (%)	95.9	94.5
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	87.4	81.3
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	88.9	87.8
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	90.4	89.3
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	81.4	58.6
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	67.0	43.2
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.1	55.5
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	73.5	80.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	100.0	93.8
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	0.0	1.2
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	16.4	2.3
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	1.2	1.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	62.5	(72.3)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	47.7	26.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(11.2)	(64.8)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(100.0)	(97.5)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(65.2)	(83.4)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	(86.1)	(25.4)
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	42.8	2.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	42.5	4.4
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	31.1	33.4
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	18.4	10.7
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	4.3	2.5
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	34.5	28.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.0	1.4

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Sultanpur, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	22.1	20.8
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	26.5	22.9
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.6	5.6
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.5	6.9
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	17.5	13.5
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	11.8	7.8
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	11.9	7.8
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	24.4	16.7
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	4.4	12.8
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.4	5.8
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	9.5	20.6
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	16.4
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.3	6.7
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	10.2	24.3
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	87.9	80.5
88. Women having a mobile phone that they themselves use (%)	58.9	41.9
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	68.5	74.8
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	15.7	14.2
91. Men age 15 years and above who use any kind of tobacco (%)	51.5	47.5
92. Women age 15 years and above who consume alcohol (%)	0.3	0.1
93. Men age 15 years and above who consume alcohol (%)	16.3	10.1

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Unnao, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.6	8.4
2. Population below age 15 years (%)	28.7	28.1
3. Population age 60 years and above (%)	12.2	12.3
4. Population living in households with electricity (%)	95.2	75.4
5. Population living in households with an improved drinking-water source ¹ (%)	100.0	98.5
6. Households using iodized salt (%)	78.4	68.7
7. Households with any usual member covered under a health insurance/financing scheme (%)	43.1	15.3
8. Households with any usual member having a bank account/post office account (%)	98.9	93.6
9. Female population age 6 years and above who ever attended school (%)	71.2	69.0
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	20.3	13.6
11. Children age 2-4 years who attended pre-school (%)	40.8	20.2
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	41.0	37.0
13. Men with 10 or more years of schooling (%)	50.1	35.6
14. Women who have ever used the internet (%)	69.4	20.3
15. Men who have ever used the internet (%)	82.2	51.5
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	9.7	18.9
17. Men age 25-29 years married before age 21 years (%)	*	*
18. Three or more births in the 5 years preceding the survey (%)	3.0	2.4
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	3.7	2.2
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	53.5	48.7
21. Any modern method ² (%)	29.6	46.2
22. Any traditional method (%)	23.9	2.5
23. Female sterilization (%)	11.9	9.6
24. Male sterilization (%)	0.0	0.0
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	13.8	19.1
26. Unmet need for spacing ³ (%)	5.3	4.5
27. Unmet need for limiting ³ (%)	8.5	14.6
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	75.8	44.7
29. Mothers who had any antenatal care visits (%)	98.0	74.9
30. Mothers who had at least 4 antenatal care visits (%)	54.6	19.8
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	93.5	88.1
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	36.5	18.6
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	16.6	9.6
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	95.7	93.0
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	85.9	80.8
36. Institutional births in public facility (%)	56.9	62.6
37. Births attended by skilled health personnel ⁵ (%)	86.1	81.2
38. Births delivered by caesarean section (%)	21.0	18.3
39. Births in a private health facility that were delivered by caesarean section (%)	58.1	64.5
40. Births in a public health facility that were delivered by caesarean section (%)	7.2	10.4

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.

- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.

- Pregnant with an unwanted pregnancy.

- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Unnao, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	79.2	54.2
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	(3.4)	(7.0)
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	81.8	60.6
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	84.2	58.6
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	(95.5)	(62.5)
46. Children age 12-23 months who have received any vaccine (%)	97.4	88.8
47. Children age 12-23 months who have received BCG (%)	97.4	88.8
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	85.7	63.2
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	85.7	71.5
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	86.1	76.7
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	70.4	48.3
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	75.6	54.4
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	86.0	26.2
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	66.4	74.1
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	96.8	88.3
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	3.2	7.4
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	8.6	8.0
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	2.6	2.3
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	67.0	(61.8)
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	59.4	11.7
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(34.6)	(54.7)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(76.0)	(88.2)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(89.4)	(73.5)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.8	5.8
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	*	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	15.3	5.2
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	32.8	39.2
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.6	12.1
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.2	5.1
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	42.1	29.3
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.4	7.5

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Unnao, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	21.4	22.6
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	25.5	20.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	8.9	4.4
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	8.5	3.7
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.0	9.7
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	10.4	4.4
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	9.5	3.7
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	20.2	9.2
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.9	13.5
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.6	5.7
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.7	22.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	6.1	15.7
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	4.0	4.0
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	11.6	21.4
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	81.8	57.3
88. Women having a mobile phone that they themselves use (%)	58.4	45.9
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	65.6	65.4
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	8.5	9.5
91. Men age 15 years and above who use any kind of tobacco (%)	48.2	50.9
92. Women age 15 years and above who consume alcohol (%)	0.5	0.3
93. Men age 15 years and above who consume alcohol (%)	20.7	16.7

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Varanasi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Population and Household Profile	Total	Total
1. Population below age 5 years (%)	8.9	7.3
2. Population below age 15 years (%)	26.2	27.3
3. Population age 60 years and above (%)	10.4	9.6
4. Population living in households with electricity (%)	99.3	97.2
5. Population living in households with an improved drinking-water source ¹ (%)	99.0	98.4
6. Households using iodized salt (%)	96.9	99.9
7. Households with any usual member covered under a health insurance/financing scheme (%)	44.2	12.8
8. Households with any usual member having a bank account/post office account (%)	98.2	95.9
9. Female population age 6 years and above who ever attended school (%)	76.9	74.1
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	22.6	11.7
11. Children age 2-4 years who attended pre-school (%)	43.0	8.7
Characteristics of Adults (age 15-49 years)		
12. Women with 10 or more years of schooling (%)	58.1	55.0
13. Men with 10 or more years of schooling (%)	65.2	61.7
14. Women who have ever used the internet (%)	84.2	38.9
15. Men who have ever used the internet (%)	94.6	65.8
Marriage and Fertility		
16. Women age 20-24 years married before age 18 years (%)	9.6	10.4
17. Men age 25-29 years married before age 21 years (%)	(14.7)	(19.7)
18. Three or more births in the 5 years preceding the survey (%)	2.8	2.2
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.6	0.0
Current Use of Family Planning Methods (currently married women age 15-49 years)		
20. Any method ² (%)	74.3	72.5
21. Any modern method ² (%)	44.4	60.9
22. Any traditional method (%)	29.9	11.6
23. Female sterilization (%)	29.1	23.9
24. Male sterilization (%)	0.3	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)		
25. Total unmet need ³ (%)	8.0	8.7
26. Unmet need for spacing ³ (%)	4.1	4.2
27. Unmet need for limiting ³ (%)	3.9	4.5
Maternal and Child Health		
Antenatal Care (for last birth in the 5 years before the survey)		
28. Mothers who had an antenatal check-up in the first trimester (%)	79.0	74.3
29. Mothers who had any antenatal care visits (%)	97.1	98.3
30. Mothers who had at least 4 antenatal care visits (%)	65.9	51.4
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	89.6	95.0
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	46.8	30.1
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	33.2	10.7
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	86.7	99.0
Delivery Care (for births in the 5 years before the survey)		
35. Institutional births (%)	93.2	95.2
36. Institutional births in public facility (%)	42.0	58.7
37. Births attended by skilled health personnel ⁵ (%)	94.2	93.4
38. Births delivered by caesarean section (%)	39.2	23.0
39. Births in a private health facility that were delivered by caesarean section (%)	62.9	46.3
40. Births in a public health facility that were delivered by caesarean section (%)	16.7	10.5

Note: Major indicators are highlighted in grey. Readers are cautioned when comparing trends and interpreting results of some indicators due to the small sample size.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.

- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.

Varanasi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Postnatal Care (for last birth in the 5 years before the survey)	Total	Total
41. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	84.2	85.1
42. Children born at home who were taken to a health facility for a check-up within 24 hours of birth (%)	*	*
43. Children who received postnatal care from a doctor/nurse/LHV/ANM/midwife/other health personnel within 2 days of delivery* (%)	83.8	82.0
Child Vaccinations and Vitamin A Supplementation		
44. Children age 12-23 months fully vaccinated based on information from either vaccination card or mother's recall ⁶ (%)	74.7	79.3
45. Children age 12-23 months fully vaccinated based on information from vaccination card only ⁷ (%)	84.7	78.6
46. Children age 12-23 months who have received any vaccine (%)	95.5	97.2
47. Children age 12-23 months who have received BCG (%)	94.7	96.1
48. Children age 12-23 months who have received 3 doses of polio vaccine ⁸ (%)	80.4	87.1
49. Children age 12-23 months who have received 3 doses of pentavalent vaccine (%)	78.4	84.3
50. Children age 12-23 months who have received the first dose of measles-containing vaccine (MCV) (%)	84.1	89.5
51. Children age 24-35 months who have received a second dose of measles-containing vaccine (MCV) (%)	69.8	48.9
52. Children age 12-23 months who have received birth dose of hepatitis B vaccine (%)	46.9	45.0
53. Children age 12-23 months who have received 3 doses of rotavirus vaccine (%)	81.3	48.4
54. Children age 9-35 months who received a vitamin A dose in the last 6 months (%)	55.3	80.5
55. Children age 12-23 months who received most of their vaccinations in a public health facility (%)	95.8	97.7
56. Children age 12-23 months who received most of their vaccinations in a private health facility (%)	4.2	2.3
Treatment of Childhood Diseases (children under age 5 years)		
57. Prevalence of diarrhoea in the 2 weeks preceding the survey (%)	17.9	4.4
58. Prevalence of symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey (%)	3.0	2.1
59. Children with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider (%)	70.8	*
Child Feeding Practices and Nutritional Status of Children		
60. Children under age 3 years breastfed within one hour of birth ⁹ (%)	34.5	36.4
61. Children under age 6 months exclusively breastfed ¹⁰ (%)	(28.3)	(47.5)
62. Children under age 6 months currently breastfeeding ¹⁰ (%)	(91.7)	(100.0)
63. Children under age 6 months exclusively breastfed or breastfeeding and consuming plain water/other milk only ¹⁰ (%)	(78.5)	(84.0)
64. Children age 6-8 months receiving solid or semi-solid food and breastmilk ¹⁰ (%)	*	*
65. Breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	31.7	6.6
66. Non-breastfeeding children age 6-23 months receiving an adequate diet ^{10, 11} (%)	(10.7)	*
67. Total children age 6-23 months receiving an adequate diet ^{10, 11} (%)	27.3	5.8
68. Children under 5 years who are stunted (height-for-age) ¹² (%)	25.3	37.4
69. Children under 5 years who are wasted (weight-for-height) ¹² (%)	22.1	21.0
70. Children under 5 years who are severely wasted (weight-for-height) ¹³ (%)	5.6	13.3
71. Children under 5 years who are underweight (weight-for-age) ¹² (%)	33.2	39.0
72. Children under 5 years who are overweight (weight-for-height) ¹⁴ (%)	0.3	3.1

*LHV = Lady Health Visitor; ANM = Auxiliary Nurse Midwife.

⁶Vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁷Among children whose vaccination card was shown to the interviewer, percentage vaccinated with BCG, measles-containing vaccine (MCV)/MR/MMR/Measles, and 3 doses each of polio (excluding polio vaccine given at birth) and DPT or penta vaccine.

⁸Not including polio vaccination given at birth.

⁹Based on the last child born in the 3 years before the survey.

¹⁰Based on the youngest child living with the mother.

¹¹Breastfed children receiving 4 or more food groups and a minimum meal frequency, non-breastfed children fed with a minimum of 3 Infant and Young Child Feeding Practices (fed with other milk or milk products at least twice a day, a minimum meal frequency that is, receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months, and solid or semi-solid foods from at least four food groups not including the milk or milk products food group).

¹²Below -2 standard deviations, based on the WHO standard.

¹³Below -3 standard deviations, based on the WHO standard.

¹⁴Above +2 standard deviations, based on the WHO standard.

Varanasi, Uttar Pradesh - Key Indicators

Indicators	NFHS-6 (2023-24)	NFHS-5 (2019-21)
Nutritional Status of Adults (age 15-49 years)	Total	Total
73. Women whose Body Mass Index (BMI) is below normal (BMI <18.5 kg/m ²) ¹⁵ (%)	18.0	15.7
74. Women who are overweight or obese (BMI ≥25.0 kg/m ²) ¹⁵ (%)	28.6	22.6
Blood Sugar Level among Adults (age 15 years and above)		
Women		
75. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	7.9	4.3
76. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.2	4.2
77. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	18.5	9.3
Men		
78. Blood sugar level - high (141-160 mg/dl) ¹⁶ (%)	12.2	7.2
79. Blood sugar level - very high (>160 mg/dl) ¹⁶ (%)	10.0	5.0
80. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level ¹⁶ (%)	22.9	12.6
Hypertension among Adults (age 15 years and above)		
Women		
81. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	5.0	13.3
82. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	2.3	3.9
83. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	12.0	19.0
Men		
84. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%)	8.7	20.9
85. Moderately or severely elevated blood pressure (Systolic ≥160 mm of Hg and/or Diastolic ≥100 mm of Hg) (%)	3.3	5.6
86. Elevated blood pressure (Systolic ≥140 mm of Hg and/or Diastolic ≥90 mm of Hg) or taking medicine to control blood pressure (%)	13.9	27.1
Women's Empowerment (women age 15-49 years)		
87. Women having a bank or savings account that they themselves use (%)	82.8	89.6
88. Women having a mobile phone that they themselves use (%)	61.8	56.0
89. Women age 15-24 years who use hygienic methods of protection during their menstrual period ¹⁷ (%)	82.8	85.6
Tobacco Use and Alcohol Consumption among Adults (age 15 years and above)		
90. Women age 15 years and above who use any kind of tobacco (%)	10.2	5.5
91. Men age 15 years and above who use any kind of tobacco (%)	43.1	38.4
92. Women age 15 years and above who consume alcohol (%)	0.2	0.4
93. Men age 15 years and above who consume alcohol (%)	22.8	13.4

¹⁵Excludes pregnant women and women with a birth in the preceding 2 months.

¹⁶Random blood sugar measurement.

¹⁷Reusable sanitary pads, disposable sanitary pads, tampons, and menstrual cups are considered to be hygienic methods of protection.

Appendix: State and Districtwise Sample Size

S.No.	District Name	Sample		
		Household	Women	Men
1	Agra	991	1,179	192
2	Aligarh	1,000	1,240	168
3	Ambedkar Nagar	1,023	1,455	153
4	Amethi	1,004	1,355	132
5	Amroha	988	1,209	172
6	Auraiya	1,007	1,119	139
7	Ayodhya	1,012	1,351	169
8	Azamgarh	1,005	1,450	168
9	Baghpat	994	1,282	203
10	Bahraich	1,041	1,365	186
11	Ballia	977	1,387	194
12	Balrampur	1,007	1,436	148
13	Banda	933	1,050	144
14	Barabanki	985	1,237	140
15	Bareilly	986	1,248	170
16	Basti	1,031	1,463	165
17	Bhadohi	1,006	1,485	195
18	Bijnor	1,005	1,256	153
19	Budaun	997	1,273	174
20	Bulandshahr	975	1,198	189
21	Chandauli	998	1,311	166
22	Chitrakoot	953	1,123	145
23	Deoria	984	1,382	154
24	Etah	1,009	1,247	195
25	Etawah	978	1,091	167
26	Farrukhabad	1,000	1,254	175
27	Fatehpur	998	1,131	138
28	Firozabad	1,004	1,269	172
29	Gautam Buddha Nagar	942	993	173
30	Ghaziabad	918	1,028	153
31	Ghazipur	972	1,336	143
32	Gonda	998	1,467	145
33	Gorakhpur	1,005	1,300	163
34	Hamirpur	1,017	1,088	153
35	Hapur	995	1,264	186
36	Hardoi	1,020	1,224	180
37	Hathras	1,020	1,246	180
38	Jalaun	998	1,052	142
39	Jaunpur	964	1,424	129
40	Jhansi	942	983	140
41	Kannauj	1,020	1,238	152
42	Kanpur Dehat	998	1,054	169
43	Kanpur Nagar	938	992	140
44	Kasganj	991	1,206	170
45	Kaushambi	1,013	1,272	147
46	Kheri	984	1,154	130
47	Kushinagar	998	1,389	129
48	Lalitpur	1,008	1,198	194

Appendix: State and Districtwise Sample Size

S.No.	District Name	Sample		
		Household	Women	Men
49	Lucknow	919	918	143
50	Maharajganj	1,028	1,359	143
51	Mahoba	954	978	149
52	Mainpuri	995	1,205	175
53	Mathura	995	1,261	159
54	Mau	988	1,534	164
55	Meerut	978	1,208	165
56	Mirzapur	1,060	1,323	194
57	Moradabad	992	1,358	148
58	Muzaffarnagar	996	1,370	215
59	Pilibhit	1,010	1,199	175
60	Pratapgarh	1,028	1,399	172
61	Prayagraj	1,000	1,217	159
62	Rae Bareli	973	1,154	139
63	Rampur	989	1,324	181
64	Saharanpur	1,012	1,347	191
65	Sambhal	1,006	1,307	187
66	Sant Kabir Nagar	1,039	1,504	155
67	Shahjahanpur	1,004	1,235	193
68	Shamli	1,000	1,315	231
69	Shrawasti	1,049	1,335	157
70	Siddharth Nagar	1,002	1,422	155
71	Sitapur	1,001	1,142	143
72	Sonbhadra	1,018	1,166	158
73	Sultanpur	1,043	1,422	182
74	Unnao	972	1,068	132
75	Varanasi	1,022	1,335	169
	UTTAR PRADESH	74,705	94,159	12,318

For additional information

Director General (Statistics)

Ministry of Health and Family Welfare
Government of India
Room no. 311, 3rd Floor
Indian Red Cross Society Building
Red Cross Road
New Delhi – 110 001 (India)
Landline: 011-2373 6973/2373 6975
Email: dgstats.mohfw@gov.in

Chief Director (Statistics)

Ministry of Health and Family Welfare
Government of India
Room no. 313, 3rd Floor
Indian Red Cross Society Building
Red Cross Road
New Delhi – 110 001 (India)
Landline: 011-2373 6983
Email: cdstats.mohfw@gov.in

Director & Senior Professor/

Principal Investigator (NFHS-6)

International Institute for Population Sciences
Govandi Station Road, B.S. Devashi Marg, Deonar
Mumbai – 400 088 (India)
Telephone: 022-4237 2549
Email: nfhs6pmu@gmail.com
nfhsdata@iipsindia.ac.in
Website: <https://www.iipsindia.ac.in>
<https://www.nfhsiips.in>