

# CALCULUS AND LINEAR ALGEBRA

## UNIT-IV

(Applications of Differential Calculus)

DEPARTMENT OF MATHEMATICS

SRM Institute of Science and  
Technology

## Topics covered

- *Radius of Curvature -Cartesian coordinates.*
- *Radius of Curvature - Polar coordinates.*
- *Centre of curvature.*
- *Circle of curvature.*
- *Evolute of standard curves– Problems*
- *Envelope of standard curves.*
- *Beta Gamma Functions Definitions*
- *Beta Gamma Functions -simple problems.*

## Introduction

The rate of change of the direction of tangent with respect to arc length as the point  $p$  moves along the curve is called curvature. The vector of the curve whose magnitude is called the curvature at  $p$ . Radius of curvature. The reciprocal of the curvature of a curve at any point  $P$  is called the radius of curvature at  $P$  and is denoted by  $\rho$ .

➤ Radius of curvature for Cartesian Curve  $y = f(x)$ , is given by

$$\rho = \frac{[1+y_1^2]^{\frac{3}{2}}}{y_2}, \text{ where } y_1 = dy/dx \text{ and } y_2 = d^2y/dx^2$$

➤ Radius of curvature for parametric equations  $x = f(t)$ ,  $y = g(t)$  is

$$\text{given by } \rho = \frac{(X'^2+Y'^2)^{\frac{3}{2}}}{X'Y''-Y'X''} \text{ where } Z' = dz/dt \text{ and } Z'' = d^2z/dt^2$$

➤ Radius of curvature for polar curve  $r = f(\theta)$  is given by  $\rho$

$$= \frac{(r^2+r_1^2)^{\frac{3}{2}}}{r^2-rr_2+2r_1^2}$$

## Problem-

1 Find the radius of curvature at the point  $\left(\frac{3a}{2}, \frac{3a}{2}\right)$  of the Folium  $x^3 + y^3 = 3axy$ .

**Soln:** Differentiating with respect to  $x$ , we get

$$\begin{aligned} 3x^2 + 3y^2 \frac{dy}{dx} &= 3a \left( y + x \frac{dy}{dx} \right) \\ (y^2 - ax) \frac{dy}{dx} &= ay - x^2 \\ \therefore \frac{dy}{dx} \text{ at } (3a/2, 3a/2) &= -1 \end{aligned} \quad (1)$$

Differentiating (1),

$$\begin{aligned} \left( 2y \frac{dy}{dx} - a \right) \frac{dy}{dx} + (y^2 - ax) \frac{d^2y}{dx^2} &= a \frac{dy}{dx} - 2x \\ \therefore \frac{d^2y}{dx^2} \text{ at } \left( \frac{3a}{2}, \frac{3a}{2} \right) &= \frac{32}{3a} \end{aligned}$$

$$\text{Hence } \rho \text{ at } \left( \frac{3a}{2}, \frac{3a}{2} \right) = \frac{[1 + (\frac{dy}{dx})^2]^{3/2}}{\frac{d^2y}{dx^2}} = \frac{[1 + (-1)^2]^{3/2}}{-32/3a}$$

$$= \frac{3a}{8\sqrt{2}} \text{ (in magnitude)}$$

## Problem-

2

Show that the radius of curvature at any point of the cycloid

$x = a(\theta + \sin\theta)$ ,  $y = a(1 - \cos\theta)$  is  $4a \cos\theta/2$ .

**Soln:** We have  $dy/dx = a(1 + \cos\theta)$ ,  $dy/d\theta = a\sin\theta$

$$\frac{dy}{dx} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dx} = \frac{a\sin\theta}{a(1 + \cos\theta)} = \frac{2\sin\theta/2}{2\cos^2\theta/2} = \frac{\tan\theta}{2}$$

$$\frac{d^2y}{dx^2} = \frac{d}{d\theta} \left( \frac{dy}{dx} \right) \cdot \frac{d\theta}{dx} = \frac{1}{2} \sec^2 \frac{\theta}{2} \cdot \frac{1}{a(1 + \cos\theta)}$$

$$= \frac{1}{2} \sec^2 \frac{\theta}{2} \cdot \frac{1}{2a\cos^2\theta/2} = \frac{1}{4a} \sec^4 \frac{\theta}{2}$$

$$\rho = \frac{\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{3/2}}{d^2y/dx^2} = \frac{4a(1 + \tan^2\theta/2)^{3/2}}{\sec^4 \frac{\theta}{2}}$$

$$= 4a \cdot \left( \frac{\sec^2\theta}{2} \right)^{3/2} \cdot \frac{\cos^4\theta}{2} = 4a \cos\theta/2$$

## Problem-

3 Prove that the radius of curvature at any point of the astroid  $x^{2/3} + y^{2/3} = a^{2/3}$ , is three times the length of the perpendicular from the origin to the tangent at that point.

**Soln:** The parametric equation of the curve is

$$x = a \cos^3 t, y = a \sin^3 t$$

$$x' (= dx/dt) = -3a \cos^2 t \sin t, y' = 3a \sin^2 t \cos t.$$

$$x'' = -3a(\cos^3 t - 2 \cos t \sin^2 t) = 3a \cos t (2 \sin^2 t - \cos^2 t)$$

$$y'' = 3a(2 \sin t \cos t - \sin^3 t)$$

$$= 3a \sin t (2 \cos^2 t - \sin^2 t) x'^2 + y'^2$$

$$= 9a^2(\cos^4 t \sin^2 t + \sin^4 t \cos^2 t) = 9a^2 \sin^2 t \cos^2 t x' y'' - y' x''$$

$$= -9a^2 \cos^2 t \sin^2 t (2 \cos^2 t - \sin^2 t)$$

$$-9a^2 \cos^2 t \sin^2 t (2 \sin^2 t - \cos^2 t) = -9a \sin^2 t \cos^2 t$$

$$\rho = \frac{(x'^2 + y'^2)^{3/2}}{x' y'' - y' x''} = \frac{27 a^3 \sin^3 t \cos^3 t}{-9a^2 \sin^2 t \cos^2 t} = -3a \sin t \cos t.$$

## Problem-3

Cont...

Since  $\frac{dy}{dx} = \frac{y'}{x'} = \tan t$ ,

$\therefore$  Equation of the tangent at  $(a\cos^3 t, a\sin^3 t)$  is

$$y - a\sin^3 t = \tan t (x - a\cos^3 t)$$

$$x \tan t + y - a \sin t = 0$$

length of  $\perp$  from  $(0,0)$  on (2) =  $\frac{0+0-a\sin t}{\sqrt{(\tan^2 t + 1)}} = a \sin t \cos t$ .

Thus  $\rho = 3p$ .

## Problem-

4 Show that the radius of curvature at any point of the cardioid

$r = a(1 - \cos\theta)$  varies as  $\sqrt{r}$ .

**Soln:** Differentiating w.r.t.  $\theta$ , we get

$$r_1 = \frac{dr}{d\theta} = a \sin\theta, r_2 = a \cos\theta$$

$$\therefore (r^2 + r_1^2)^{3/2} = [a^2(1 - \cos\theta)^2 + a^2 \sin^2 \theta]^{3/2} = a^2 [2(1 - \cos\theta)]^{3/2}$$

$$\begin{aligned} r^2 - r r_2 + 2r_1^2 &= a^2(1 - \cos\theta)^2 - a^2(1 - \cos\theta)\cos\theta + 2a^2 \sin^2 \theta \\ &= 3a^2(1 - \cos\theta) \end{aligned}$$

$$\begin{aligned} \text{Thus } \rho &= \frac{(r^2 + r_1^2)^{3/2}}{r^2 - r r_2 + 2r_1^2} = \frac{a^3 2\sqrt{2}(1 - \cos\theta)^{3/2}}{3a^2(1 - \cos\theta)} \\ &= \frac{2\sqrt{2}}{3} a(1 - \cos\theta)^{1/2} = \frac{2\sqrt{2}a}{3} \left(\frac{r}{a}\right)^{1/2} \propto \sqrt{r} \end{aligned}$$



# CENTRE OF CURVATURE

Let  $\Gamma$  be a simple curve having tangent at each point. At any point  $P$  on this curve we can draw a circle having the same curvature at  $P$  as the curve  $\Gamma$ .

This circle is called the circle of curvature and its centre is called the centre of curvature and its radius is the radius of curvature of  $\Gamma$  at  $P$ . Centre of curvature at any point  $P(x, y)$  on the curve  $y = f(x)$  is given

$$\text{by } \bar{x} = x - \frac{y_1(1+y_1^2)}{y_2}$$

$$\bar{y} = y - \frac{1+y_1^2}{y_2}$$

Equation of the circle of curvature at  $P$  is  $(x - \bar{x})^2 + (y - \bar{y})^2 = \rho^2$ .

# EVOLUTE

The locus of centre of curvature of a given curve  $\Gamma$  is called the evolute of the curve. The given curve  $\Gamma$  is called an involute of the evolute. In fact, for the evolute there are many involutes

## Procedure to Find the Evolute

Let  $y = f(x)$  (1) be the equation of the given curve. If  $(\bar{x}, \bar{y})$  is the centre of curvature at any point  $P(x, y)$  on (1), then

$$\bar{x} = x - \frac{y_1(1 + y_1^2)}{y_2} \quad (1)$$

$$\bar{y} = y + \frac{1 + y_1^2}{y_2} \quad (2)$$

Eliminating  $x, y$  using (1), (2) and (3), we get a relation in  $\bar{x}, \bar{y}$ .

Replacing  $\bar{x}$  by  $x$  and  $\bar{y}$  by  $y$ , we get the equation of locus of  $(\bar{x}, \bar{y})$ , which is the evolute of the given curve.

## Problem :1

*Find the centre of curvature at the point of  $P(2,4)$  on the parabola  $y^2 = 4(x + 2)$*

*Given the curve  $y^2 = 4(x + 2)$*   
*tial*

*Differentiating (1) w. r. to  $x$   $2y \frac{dy}{dx} = 4 \Rightarrow \frac{dy}{dx} = \frac{2}{y}$*

$$\frac{d^2y}{dx^2} = -\frac{2}{y^2} \left( \frac{dy}{dx} \right) = \left( \frac{-2}{y^2} \right) \left( \frac{2}{y} \right) = -\frac{4}{y^3}$$

*At  $(2,4)$ ,  $y_1 = \frac{1}{2}$  and  $y_2 = -\frac{1}{16}$*

*Let  $C(\bar{x}, \bar{y})$  be the centre of curvature at the point  $P(2,4)$  on the curve.  $y^2 = 4(x + 2)$ .*

## Problem :1 cont,...

*Then*

$$\bar{x} = x - \frac{y_1}{y_2} (1 + y_1^2) = 2 - \frac{1}{2}(-16) \left[ 1 + \frac{1}{4} \right] = 2 + 8 \left( \frac{5}{4} \right) = 12$$

$$\bar{y} = y + \frac{(1 + y_1^2)}{y_2} = 4 + (-16) \left[ 1 + \frac{1}{4} \right] = 4 - \frac{5}{4}(16) = -16$$

*The centre of curvature is  $C(12, -16)$ .*

## Problem :2

Find the equation to the circle of curvature of the curve  $xy = c^2$  at  $(c, c)$

$xy = c^2$  Differentiating this twice w.r.to  $x$ ,  $\frac{dy}{dx} = -\frac{c^2}{x^2}$

$$\frac{d^2y}{dx^2} = \frac{c^2}{x^3}$$

At the point  $(c, c)$   $y_1 = -1$  and  $y_2 = \frac{2}{c}$

$$\rho = \frac{(1+y_1^2)^{\frac{3}{2}}}{y_2} = \frac{c}{2}(1+1)^{\frac{3}{2}} = c\sqrt{2}$$

Let  $C(\bar{x}, \bar{y})$  be the centre of curvature.

$$\text{Then } \bar{x} = x - \frac{y_1}{y_2}(1+y_1^2) = c + \frac{c}{2}(1+1) = 2c$$

$$\bar{y} = y + \frac{1+y_1^2}{y_2} = c + \frac{c}{2}(1+1) = 2c$$

$$\begin{aligned}\text{Equation of the circle of curvature is } (x - \bar{x})^2 + (y - \bar{y})^2 &= \rho^2 \\ \Rightarrow (x - 2c)^2 + (y - 2c)^2 &= 2c^2\end{aligned}$$

## Problem-

1 Find the coordinates of the center of curvature at any point of the parabola  $y^2 = 4ax$ . Hence show that its evolute is

$$27ay^2 = 4(x - 2a)^3$$

Soln: We have  $2yy_1 = 4a$  i.e.  $y_1 = 2a/y$  and

$$y_2 = -\frac{2a}{y^2} \cdot y_1 = -\frac{4a^2}{y^3}$$

If  $(\bar{x}, \bar{y})$  be the center of curvature, then

$$\begin{aligned}\bar{x} &= x - \frac{y_1(1 + y_1^2)}{y_2} = x - \frac{2a/y(1 + 4a^2/y^2)}{-4a^2/y^3} \\ &= x + \frac{y^2 + 4a^2}{2a} = x + \frac{4ax + 4a^2}{2a} = 3x + 2a\end{aligned}$$

and

## Problem-1

### Cont...

$$\begin{aligned}\bar{y} &= y + \frac{1 + y_1^2}{y_2} = y + \frac{1 + 4a^2/y^2}{-4a^2/y^2} \\ &= y - \frac{y(y^2 + 4a^2)}{4a^2} = \frac{-y^3}{4a^2} = -\frac{2x^{3/2}}{\sqrt{a}}\end{aligned}$$

To find the evolute, we have to eliminate  $x$  from (4) and (5)

$$\therefore (\bar{y})^2 = \frac{4x^3}{a} = \frac{4}{a} \left( \frac{\bar{x} - 2a}{3} \right)^3$$

$$\text{Or } 27a(\bar{y})^2 = 4(\bar{x} - 2a)^3$$

Thus the locus of  $(\bar{x}, \bar{y})$  i.e., evolute is  $27ay^2 = 4(x - 2a)^3$ .

## Problem-

2

Show that the evolute of the cycloid  $x = a(\theta - \sin \theta)$ ,  $y = a(1 - \cos \theta)$  is another equal cycloid.

Soln: We have  $y_1 = \frac{dy}{d\theta} + \frac{dx}{d\theta} = \frac{a \sin \theta}{a(1 - \cos \theta)} = \cot \frac{\theta}{2}$ .

$$\begin{aligned} y_2 &= \frac{d}{dx}(y_1) = \frac{d}{d\theta}(\cot \theta/2) \cdot \frac{d\theta}{dx} \\ &= -\operatorname{cosec}^2 \theta/2 \cdot 1/2 \cdot \frac{1}{a(1 - \cos \theta)} = -\frac{1}{4a \sin^4 \theta/2} \end{aligned}$$

If  $(\bar{x}, \bar{y})$  be the center of curvature, then

$$\bar{x} = x - \frac{y_1(1 + y_1^2)}{y_2} =$$

$$\begin{aligned} &a(\theta - \sin \theta) + \cot \theta/2(-4a \sin^4 \theta/2)(1 + \cot^2 \theta/2) \\ &= a(\theta - \sin \theta) + \frac{\cos \theta/2}{\sin \theta/2} \cdot 4a \sin^4 \theta/2 \cdot \operatorname{cosec}^2 \theta/2 \end{aligned}$$

$$= a(\theta - \sin \theta) + 4a \sin \theta/2 \cos \theta/2 = a(\theta - \sin \theta) + 2a \sin \theta = a(\theta + \sin \theta)$$



## Problem-2

Cont...

$$\bar{y} = y + \frac{1 + y_1^2}{y_2} = a(1 - \cos \theta) + (1 + \cot^2 \theta/2)(-4a\sin^4 \theta/2)$$

$$= a(1 - \cos \theta) - 4a\sin^4 \theta/2 \cdot \operatorname{cosec}^2 \theta/2$$

$$a(1 - \cos \theta) - 4\sin^2 \theta/2$$

$$a(1 - \cos \theta) - 2a(1 - \cos \theta) = -a(1 - \cos \theta)$$

Hence the locus of  $(\bar{x}, \bar{y})$  i.e., the evolute, is given by

$x = a(\theta + \sin \theta), y = -a(1 - \cos \theta)$  which is another equal cycloid.

# ENVELOPE

Consider the system of straight lines  $y = mx + \frac{1}{m}$  (1) where  $m$  is a parameter. For different values of  $m$ , we have different straight lines and so (1) represents a family of straight lines. Each member of this family touches the curve  $y^2 = 4x$ . So, these lines cover the curve  $y^2 = 4x$ . This curve is called the envelope of the family of lines. We shall now define envelope.

## Definition:

Let  $f(x, y, \alpha) = 0$  be a single parameter family of curves, where  $\alpha$  is the parameter. The envelope of this family of curves is a curve which touches every member of the family.

## Problem-

1 Find the envelope of the family of lines  $y = mx + \sqrt{(1 + m^2)}$ ,  $m$  being the parameter.

Soln: We have  $(y - mx)^2 = 1 + m^2$

Differentiating (6) partially with respect to  $m$ ,

$$2(y - mx)(-x) = 2m \text{ or } m = xy/(x^2 - 1)$$

Now eliminate  $m$  from (6) and (7).

Substituting the values of  $m$  in (6), we get

$$\left(y - \frac{x^2 y}{x^2 - 1}\right)^2 = 1 + \left(\frac{xy}{x^2 - 1}\right)^2 \text{ or } y^2 = (x^2 - 1)^2 + x^2 y^2$$

$x^2 + y^2 = 1$  which is the required equation of the envelope

## Problem-

2. Find the envelope of a system of concentric and coaxial ellipses of constant area.

Soln: Taking the common axes of the system of ellipses as the coordinate axes, the equation to an ellipse of the family is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

where  $a$  and  $b$  are the parameters. The area of the ellipse  $= \pi ab$  which is given to be constant, say  $= \pi c^2$

$$ab = c^2 \text{ or } b = c^2/a$$

Substituting in (8),

$$\frac{x^2}{a^2} + \frac{y^2}{(c^2/a)^2} = 1 \text{ or } x^2 a^{-2} + (y^2/c^4) a^2 = 0$$

which is given family of ellipses with  $a$  as the only parameter.

## Problem-2

Cont...

Differentiating partially (10) with respect to  $a$ ,

$$-2x^2a^{-3} + 2(y^2/c^4)a = 0 \text{ or } a^2 = c^2x/y$$

Eliminate  $a$  from (10) and (11)

Substituting the values of  $a^2$  in (10), we get

$$x^2(y/c^2x) + (y^2/c^4)(c^2x/y) = 1 \text{ or } 2xy = c^2$$

which is the required equation of the envelope.

## Problem-

3

Find the evolute of the parabola  $y^2 = 4ax$ .

**Soln:** Any normal to the parabola is  $y = mx - 2am - am^3$

Differentiating it with respect to  $m$  partially,

$$0 = x - 2a - 3am^2 \text{ or } m = [(x - 2a)/3a]^{1/2}$$

Substituting this value of  $m$  in (12),

$$y = \left( \frac{x - 2a}{3a} \right)^{1/2} \left[ x - 2a - a \cdot \frac{x - 2a}{3a} \right]$$

Squaring both sides, we have

$$27ay^2 = 4(x - 2a)^3$$

Which is the evolute of the parabola.

## Beta, Gamma Functions

The beta function is defined as

$$\beta(p, q) = \int_0^1 x^{p-1} (1-x)^{q-1} dx$$

where  $m, n > 0$ . Note that  $\beta(p, q) = \beta(q, p)$ .

The gamma function is defined as

$$\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx$$

where  $n > 0$ .

Note: (1)  $\Gamma(1) = 1$ , (2)  $\Gamma(n+1) = n\Gamma(n) = n!$ , (3)  $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$

The relation between Beta and Gamma functions is

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

## Problems Beta, Gamma Functions

Show that  $\Gamma(n) = \int_0^1 \left(\log \frac{1}{y}\right)^{n-1} dy, (n > 0)$ .

$$\begin{aligned}\text{Soln: } \Gamma(n) &= \int_0^\infty x^{n-1} e^{-x} dx \quad (n > 0) \\ &= \int_0^1 \left(\log \frac{1}{y}\right)^{n-1} y \left(-\frac{1}{y} dy\right)\end{aligned}$$

put  $y = e^{-x}$

i.e.,  $x = \log(1/y)$

so that  $dx = -(1/y)dy$

$$= \int_0^1 \left(\log \frac{1}{y}\right)^{n-1} dy.$$



## Problems based on Beta, Gamma Functions

Show that  $\beta(p, q) = \int_0^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy = \int_0^1 \frac{x^{p-1} + x^{q-1}}{(1+x)^{p+q}} dx$

**Soln:**  $\beta(p, q) = \int_0^1 x^{p-1} (1-x)^{q-1} dx$   
$$= \int_0^1 \frac{1}{(1+y)^{p-1}} \left( \frac{y}{1+y} \right)^{q-1} \frac{-1}{(1+y)^2} dy$$

put  $x = \frac{1}{1+y}$  i.e.,  $y = \frac{1}{x} - 1$

so that  $dx = \frac{-1}{(1+y)^2} dy$   
$$= \int_0^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy = \int_0^1 \frac{y^{q-1}}{(1+y)^{p+q}} dy + \int_1^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy$$

Now substituting  $y = 1/z$  in the second integral, we get

$$\begin{aligned} \int_1^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy &= \int_1^0 \frac{1}{z^q - 1} \cdot \frac{1}{(1 + 1/z)^{p+q}} \left( -\frac{1}{z^2} \right) dz \\ &= \int_0^1 \frac{z^{p-1}}{(1+z)^{p+q}} dz \end{aligned}$$

## Problems based on Beta, Gamma Functions

Express the following integral in terms of gamma function

$$\int_0^1 \frac{dx}{\sqrt{(1-x^4)}}$$

Soln:  $\int_0^1 \frac{dx}{\sqrt{(1-x^4)}}$  Put  $x^2 = \sin \theta$ , i.e.,  $x = \sin^{1/2} \theta$

so that  $dx = 1/2 \sin^{-1/2} \theta \cos \theta d\theta$

$$\begin{aligned} &= \int_0^{\pi/2} \frac{1}{2} \frac{\sin^{-1/2} \theta \cos \theta d\theta}{\sqrt{(1 - \sin^2 \theta)}} \\ &= 1/2 \int_0^{\pi/2} \sin^{-1/2} \theta d\theta \\ &= \frac{1}{2} \cdot \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma\left(-\frac{1}{2} + 1\right)}{\Gamma\left(-\frac{1}{2} + 2\right)} = \frac{\sqrt{\pi}}{4} \frac{\Gamma(1/4)}{\Gamma(3/4)} \end{aligned}$$

## Problems based on Beta, Gamma Functions

Express the following integral in terms of gamma function  $\pi/2$

$$\int_0^{\pi} \sqrt{(\tan \theta)} d\theta$$

$$\begin{aligned}\text{Soln: } \int_0^{\pi/2} \sqrt{(\tan \theta)} d\theta &= \int_0^{\pi/2} \sin^{1/2} \theta \cos^{-1/2} \theta d\theta \\ &= \frac{\Gamma\left(\frac{1}{2} + 1\right) \Gamma\left(\frac{-1}{2} + 1\right)}{2\Gamma\left(\frac{1}{2} - \frac{1}{2} + 2\right)} \\ &= \frac{\Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{1}{4}\right)}{2\Gamma(1)} \\ &= \frac{1}{2} \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right).\end{aligned}$$

## Problems based on Beta, Gamma Functions

Evaluate  $\int_0^{\infty} e^{-ax} x^{m-1} \sin bxdx$  in terms of Gamma functions.

Soln: We have  $\Gamma(m) = \int_0^{\infty} e^{-x} x^{m-1} dx$

$$\begin{aligned} &= \int_0^{\infty} e^{-ay} a^m y^{m-1} dy \\ \int_0^{\infty} e^{-ay} y^{m-1} dy &= \Gamma(m)/a^m \end{aligned}$$

Then  $I = \int_0^{\infty} e^{-ax} x^{m-1} \sin bxdx$

$= \int_0^{\infty} e^{-ax} x^{m-1} (\text{Imaginary part of } e^{ibx}) dx$

$= \text{I.P. of } \int_0^{\infty} e^{-(a-ib)x} x^{m-1} dx$

$= \text{I.P. of } \{\Gamma(m)/(a-ib)^m\} \text{ by (1)}$

## Problems based on Beta, Gamma Functions

= I.P. of  $\{\Gamma(m)/[r^m(\cos \theta - i \sin \theta)^m]$  where  $a = r \cos \theta, b = r \sin \theta$

= I.P. of  $\Gamma(m)/[r^m(\cos m\theta - i \sin m\theta)]$  (Using Demoivre's theorem)

= I.P. of  $\left\{ \frac{\Gamma(m) \cdot (\cos m\theta + i \sin m\theta)}{r^m (\cos m\theta + i \sin m\theta)(\cos m\theta - i \sin m\theta)} \right\}$

=  $\frac{\Gamma(m)}{r^m} \sin m\theta$  Where  $r = \sqrt{a^2 + b^2}, \theta = \tan^{-1} b/a$ .

# Thank You