



PAPER • OPEN ACCESS

Virtual reality for understanding artificial-intelligence-driven scientific discovery with an application in quantum optics

To cite this article: Philipp Schmidt *et al* 2024 *Mach. Learn.: Sci. Technol.* **5** 035045

View the [article online](#) for updates and enhancements.

You may also like

- [Machine-learning strategies for the accurate and efficient analysis of x-ray spectroscopy](#)
Thomas Penfold, Luke Watson, Clelia Middleton et al.
- [Deep learning methods for Hamiltonian parameter estimation and magnetic domain image generation in twisted van der Waals magnets](#)
Woo Seok Lee, Taegeun Song and Kyoung-Min Kim
- [An adaptive particle swarm optimization with information interaction mechanism](#)
Rui Liu, Lisheng Wei and Pinggai Zhang



PAPER

OPEN ACCESS

RECEIVED
8 March 2024REVISED
31 May 2024ACCEPTED FOR PUBLICATION
5 July 2024PUBLISHED
13 August 2024

Original Content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Virtual reality for understanding artificial-intelligence-driven scientific discovery with an application in quantum optics

Philipp Schmidt*, Sören Arlt*, Carlos Ruiz-Gonzalez , Xuemei Gu , Carla Rodríguez and Mario Krenn*

Max Planck Institute for the Science of Light, Erlangen, Germany

* Authors to whom any correspondence should be addressed.

E-mail: ph.s.schmidt@protonmail.com, soeren.arlt@mpl.mpg.de and mario.krenn@mpl.mpg.de

Keywords: generative models, virtual reality, scientific understanding, digital discovery


Abstract

Generative Artificial Intelligence (AI) models can propose solutions to scientific problems beyond human capability. **To truly make conceptual contributions, researchers need to be capable of understanding the AI-generated structures and extracting the underlying concepts and ideas.** When algorithms provide little explanatory reasoning alongside the output, scientists have to reverse-engineer the fundamental insights behind proposals based solely on examples. This task can be challenging as the output is often highly complex and thus not immediately accessible to humans. In this work we show **transferring part of the analysis process into an immersive virtual reality (VR) environment can assist researchers in developing an understanding of AI-generated solutions.** We demonstrate the usefulness of VR in finding interpretable configurations of abstract graphs, representing Quantum Optics experiments. Thereby, we can manually discover new generalizations of AI-discoveries as well as new understanding in experimental quantum optics. Furthermore, it allows us to customize the search space in an informed way—as a human-in-the-loop—to achieve significantly faster subsequent discovery iterations. As concrete examples, with this technology, we discover a new resource-efficient 3-dimensional entanglement swapping scheme, as well as a 3-dimensional 4-particle Greenberger–Horne–Zeilingers-state analyzer. **Our results show the potential of VR to enhance a researcher's ability to derive knowledge from graph-based generative AI.** This type of AI is a widely used abstract data representation in various scientific fields.

1. Introduction

Virtual reality (VR) describes the full immersion of a human in a simulated environment, most commonly via the use of a Head Mounted Display, also known as a VR-Headset or VR-Glasses. Modern headsets feature stereoscopic rendering on top of head- and gaze tracking combined with gesture controls by tracking controllers in all spatial degrees of freedom. The applications of immersive technology are vast and varied [1], including but not limited to entertainment, education [2–4], industry and engineering [5–7], neuro- and medical science and therapy [8–13] and data science [14, 15].

In the natural sciences, VR is used to visualize and analyze complex, often three-dimensional (3D), scientific data, such as molecular or material data [16–20], microscopy data [21–24] or astronomical data [25–27]. **VR is also used to better visualize neural network models [28–30] to address the challenge of making AI explainable [31].** In addition, when the AI output is compatible with existing VR frameworks, it can be studied in an immersive environment such as reviewing drug candidates [32]. Beyond improved visualization, VR offers natural ways of interacting with scientific results via intuitive gestures [33], which can be exploited, for example in interactive molecular simulations [34, 35] or drug discovery [36]. Through this interactivity, the VR platform enables a variety of ingenious methods like using interactive simulations to sample data for machine learning tasks [37, 38] or even manipulating microscopic systems in real-time via in-VR-interaction [39].



In this work, we show that modern VR technology can be effectively applied to enhance the interpretability and conceptualization of scientific results discovered through AI. In particular, we study AI-discovered quantum optics experimental setups. By representing such experiments as abstract colored graphs [40–42], we can map complex setups into networks which in turn benefit greatly in legibility and interpretability when visualized in a true 3D environment [15, 43]. We demonstrate the unlocked potential by showcasing (1) interpretable structures in AI discoveries and (2) generalizations to new experiments derived from AI results by exploiting their graph geometry.

To this end, we built *AriadneVR*¹, a webVR tool that allows us to visualize and edit experimental graphs as well as to dynamically compute their current state. *AriadneVR* is hosted online, and does not require any additional hardware beside of the VR headset and controllers. With it, we discover previously unknown relations in the construction of different state-generation experiments for high-dimensional Greenberger–Horne–Zeilinger (GHZ) states [45] and generalize AI-discovered high-dimensional entanglement swapping experiments to a new setup for high pair counts. Furthermore, we extend to human-in-the-loop AI-discovery by intelligently restricting search space based on observed patterns in a related discovery in the case of a high-dimensional GHZ-state analyzer to achieve the ability to measure a larger state. The methods used for these discoveries are conceptually depicted in figure 1. Our techniques are not limited to quantum optics; they can also be applied to interpret any generative AI outputs involving graph representations.

We find that VR is more practically useful compared to 3-dimensional rendering at a 2-dimensional computer screen for several reasons. First, VR provides full immersion into a virtual world, where the virtual object's orientation follows your natural head movements, allowing for an intuitive and natural observation similar to real-world interaction. Second, the interaction with virtual objects in VR is very natural, using hand gestures to touch and manipulate the objects, enhancing the ease and precision of interactions. Third, VR allows users to stand inside the object and observe structures from within, offering unique internal perspectives that are impossible to achieve with a 2D screen.

¹ In greek mythology Ariadne is Theseus wife, assisting him discovering the way out of the labyrinth. *AriadneVR* assists the *Theseus* algorithm [44] in discovery.

2. AI-driven discovery in quantum physics

AI in natural science - In this work, we leverage the potential of VR to augment the process of extracting scientific understanding from solutions to scientific problems discovered by AI [46, 47]. Among other tasks, AI is used to efficiently search vast configurational spaces to propose testable answers to research questions, such as predicting protein folding patterns [48], developing drug candidates [49], designing new materials and molecules [50–53], quantum circuit design [54] or discovering physics experiments [55, 56].


Even though solutions discovered in this way have value on their own, they might not generally provide scientific understanding if their discovery does not lead to new generalizable insights [57–59]. Achieving such an understanding can be very challenging when algorithms do not provide conceptual reasoning alongside the output. In such cases, the scientist has to deduce the fundamental ideas behind results from examples alone [59].

Discovery in quantum optics - In quantum optics, algorithms are used to efficiently discover new experiments. In its earliest iterations, this is done by assembling a setup out of a pre-defined toolbox of elements to solve a specific task, e.g. designing experiments for quantum state generation, transformation or metrology [55, 60, 61]. However, many setups can fulfill the same task with minor differences between them, which further expands the already vast search space and decreases search efficiency. This challenge is mitigated by choosing a more efficient, abstract representation of quantum optical setups as colored graphs, where a single graph can represent different setups that yield the same result [40–42]. Consequently, the search space for experiments is reduced to the space of graphs. Additionally, such graphs have been shown to be interpretable, allowing the development of conceptual understanding and novel methods from AI solutions [44, 62].

The graph representation - Here, we briefly explain the graph representation of quantum optics for the specific question of finding experimental setups for generating and measuring photonic quantum systems. In this representation, an experimental setup is encoded as a colored, weighted graph by identifying vertices as photon detectors and edges as correlated photon-pair sources [40–42]. An edge linking two vertices corresponds to the pair-photons propagating to the corresponding detectors. The complex edge weight encodes the amplitude and phase of the respective pair. This way, the weights also account for phase shifts and losses in optical elements like beam splitters and mirrors, as well as any occurring interference. The color of the edge is used to specify the internal photon mode, e.g. its polarization, path, spatial mode, time-bin, or frequency.

Experimentally, quantum states are commonly obtained by relying on simultaneous detection events across all detectors, known as *n-fold coincidence detection*. In the graph picture, such events correspond to subsets of edges containing each vertex exactly once, called *perfect matchings*. As an example, a square graph decomposes into two perfect matchings:

a perfect matching in the graph represents an event where every detector (vertex) registers exactly one photon at the same time—this is called an *n-fold coincidence detection*. Each perfect matching connects each input to a unique output such that all vertices are matched, mimicking the situation where all photons are coincidentally detected across all detectors simultaneously.




(1)

One such perfect matching corresponds to one term contributing to the final quantum state. The amplitude and phase associated with the term are computed as the product of all edge weights in the matching. Thereby, a coherent superposition of all the perfect matchings in the graph leads to the final post-selected quantum state.

The colored, weighted graph contains the full information of the quantum optical experiment and can be translated into several different realizations using different technologies. The most straightforward way of translation is using entanglement by path identity, a way of creating superpositions of emission events by perfectly aligning photon paths [63, 64]. Such an example is shown in figure 2. Measurements can be described in the same way using the Choi–Jamiołkowski isomorphism [44, 56]. This technique allows for a native translation of experimental setups to abstract graphs and has been practically implemented in recent experimental works [65–68].

PyTheus - PyTheus is an open-source AI tool for discovering new quantum optics experimental setups [56]. It can discover a variety of different types of experiments, such as state generation setups, communication protocols, measurement devices, or photonic quantum gates. In all cases, the experimental setups are considered ideal (e.g. noiseless and lossless), however additional experimental properties and constraints could be added.

The initial step in searching for an experiment is to define the desired experimental task. Subsequently, one specifies the initial parameters such as the number of detectors, incoming and outgoing photons, and potential topological constraints. With these parameters, PyTheus constructs the largest possible setup in



graph form. The initial graph is then refined in two iterative steps: (1) by minimizing the desired loss by optimizing the graph weights, and (2) by performing topological optimization to remove unnecessary edges. In this way, PyTheus attempts to find the sparsest graph realizing the experiment. The optimization loop is depicted in figure 2.

3. Physical background

We show graphs of experiments for the creation and measurement of GHZ-states as well as high dimensional entanglement swapping. Both topics are highly relevant to the fields of quantum communication and quantum computing but are also used to investigate the fundamental properties of quantum physics [69, 70]. Here, we briefly introduce the quantum physical background of these experiments.

GHZ-states and GHZ-analyzers - GHZ-states are a class of highly entangled multi-particle states originally conceived as a 3-particle generalization of the well-known Bell states [45]. Both classes of states form a multi-particle orthogonal basis set, where each particle is maximally entangled with all others. Since their original proposal, they have been generalized both to particle numbers larger than three and dimensions higher than two [66, 69, 71–73]. In this work, we discuss graphs relating to the GHZ-state of n particles in d dimensions which is given as:

$$\nexists \quad |\text{GHZ}\rangle_n^d = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |j\rangle^{\otimes n}. \quad (2)$$

GHZ-state analyzers are setups distinguishing one or more GHZ-states out of an unknown input state [74]. We discuss binary analyzers distinguishing between the presence or absence of a specific GHZ state given by equation (2).

High dimensional entanglement swapping - In its original form, entanglement swapping refers to the preparation of two Bell pairs and the subsequent projection of one particle of each pair in a joined Bell state. This measurement also collapses the remaining two particles in a joined Bell state of their own, without them ever having interacted [75, 76].

This process can be generalized to multipartite entanglement generation using GHZ states as the initial entangled states [77]. High-dimensional entanglement swapping refers to non-locally creating d -dimensional Bell-pairs of the form:

$$|\Phi_d^+\rangle = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |jj\rangle. \quad (3)$$

We show graphs creating n such pairs simultaneously between two parties and refer to it as n -pair entanglement swapping.




Figure 3. Point of view of an exemplary PyTheus graph analysis inside the environment of AriadneVR. A PyTheus graph is shown with one perfect matching subgraph on the right. A rendered version of this graph is depicted in figure 4(a).

Many experiments implementing the tasks mentioned above require additional photons to realize the desired output. These helper photons are called *ancillae*, opposed state photons.

4. AriadneVR

Here we introduce AriadneVR², our web-based virtual reality application for visualizing and analyzing colored graphs representing quantum optics experiments. It is developed with the open-source framework A-Frame³ using HTML and JavaScript on a Meta (formerly Oculus) Quest 2. The tool runs fully locally in the headset's browser, requiring neither installation nor a tether to a computer for 3D rendering.

For preprocessing graphs, we use the igraph Python library [78] and its pre-implemented **Kamada–Kawai-algorithm [79] to obtain a 3D-mapping of vertices.** This creates an initial arrangement to be perfected manually in VR. The layout-generation step is performed outside the tool and creates a file containing all relevant information to be stored on the host platform (e.g. GitHub for hosting via GitHub pages⁴). When running the tool, A-Frame then uses the THREE.js 3D library⁵ under the hood to construct a model from the graph information. The VR environment is shown in figure 3.

Our tool allows the exploration of graph representations of quantum optics experiments through visualization and interaction, i.e. manual sorting and editing the experimental setup in its graph form. Beyond this, it supports the acceleration of PyTheus' searches by generating instruction files based on constrained geometries defined in VR. Graphs from the library can be spawned alongside each other and investigated simultaneously. **The vertex arrangements are changed by simply grabbing and dragging them around as desired. When a user makes functional changes to a graph, e.g. adding a new edge by drawing with the controller, the changing state is dynamically tracked. The resulting perfect matchings of a graph are available to be spawned as additional 3D models.** To save the current form of a graph, a file can be downloaded to the headset, which can then be placed into the host library to make it available for later sessions. Similarly, the user can download a PyTheus instruction template, where the current graph is set as the initial graph for the search. To streamline this process, edges can be drawn without defining their color, representing a connection where all possible edges are in the initial graph. The framerate for displaying a typical graph is about 40 per second with an Meta (formerly Oculus) Quest 2.


While AriadneVR is built for analyzing quantum optics experiments, its core functionality is visualizing and editing small colored graphs, hence can be applied to other graphs too, assuming they are fed into the pre-processing pipeline in an appropriate format. As AriadneVR is directly processed at the computer of the VR headset, the framerate will decrease for large graphs. For large graphs, e.g. graphs studied in social

² <https://github.com/artificial-scientist-lab/AriadneVR>.

³ <https://aframe.io/>.

⁴ <https://pages.github.com/>.

⁵ <https://threejs.org/>.




science, a powerful external computer is necessary for rendering the structures. In that case, different software such as VRNetzer [15] are better suited.

5. Applications

We use VR in a threefold manner as shown in figure 1. The basis of all our VR analysis is the discovery and interpretation of structure in an experimental graph. We provide interactive 3D figures and basic VR environments for all graphs mentioned in this work at the accompanying website. Through manual sorting, the majority of PyTheus graphs can be transformed into visually clean structures, some of which are immediately interpretable, revealing information on how the respective graph achieves its task. If the mechanism underlying a graph is understood, we can manually generalize it to new experiments. This is done directly in VR by drawing new or extending existing graphs. However, in some cases the underlying structure is not immediately obvious, and it remains unclear how to generalize the graph. Here we can use VR to propose geometries of potential generalizations. We then use PyTheus to search for solutions to confirm or reject our candidates.

Finding interpretable structures in 3D graphs - In the following, we show three selected examples from [56] and [62] where interpretable 3D-representations were found using AriadneVR. For better visual clarity,



we use renderings made with Blender 4.0⁶ instead of screenshots of the VR environment for depictions of 3D-graphs.

Figure 4(a) shows a 3-particle 4-dimensional GHZ-state analyzer. Beyond the striking symmetry of the discovered 3D structure, there is a functional feature exemplified by this highly complex solution. As a measurement graph, it needs to ensure that all coincidence events produce terms belonging to the desired state or interfere destructively. A common way to achieve this in the graph picture is by constructing loops of even edge count with a total negative sign. Such loops were discovered in this particular graph in VR simply by superimposing the 3D models of two interfering perfect matchings.


Figure 4(b) shows (I) $|\text{GHZ}\rangle_3^5$ and (II) $|\text{GHZ}\rangle_4^4$ state creation graphs. Ancillae are described by cube-shaped vertices and state photons by spherical vertices. Through their 3D structure discovered in VR, it is revealed, that both graphs share the same triangular graph structure highlighted in the center of both graphs. The only difference is the central vertex being an ancilla in $|\text{GHZ}\rangle_3^5$ and a state photon in $|\text{GHZ}\rangle_4^4$. In the latter, this core graph creates a 4-particle 3-dimensional GHZ state. The $|\text{GHZ}\rangle_3^5$ -graph declares one photon of this subgraph as an ancilla (vertex 5 in figure 4(b) I) to reduce the state photon count from four to three. Any remaining photon modes are then added by the outer ancilla structure for both graphs. The examples shown here are representative of insights made easily accessible just by moving to stereoscopic rendering and intuitive manual sorting.

In both of these cases the algorithms have discovered the solutions. However, to the best of our knowledge, so far no method exists that can automatically extract the underlying working principles in a similar way as we discovered manually using VR.

Manually discovering new AI-inspired experiments in VR - AriadneVR allows the user to modify and edit graph structures and construct completely new ones. **Geometric understanding obtained through the superior rendering in VR combined with the user's physical understanding of the experimental mechanisms can then be used to manually engineer new graphs.**

This process is shown on an example in figure 5. Here, we show the discovery method in the case of resource-efficient high-dimensional multi-pair entanglement swapping. The discovery is based on the PyTheus-generated solution for 2-pair 3-dimensional entanglement swapping without requiring a 3-dimensional Bell-state measurement (graph 78 in [56]) shown in figure 5(a). By combining it with an AI-inspired construction method (see figure 5(b)) of generating high-dimensional non-locally entangled pairs derived from a different AI-discovered graph for 3-pair 2-dimensional entanglement swapping (graph 77 in [56]) we can generalize to a new, previously unknown 4-pair 3-dimensional entanglement swapping graph. The result is shown in figure 5(c). It effectively creates one whole 3-dimensional pair with no additional resource cost in ancillae by superimposing additional creation processes from single-pair

⁶ <https://www.blender.org/>.



experiments. It is an alternative setup to stacking two of the 2-pair AI solutions showing the same novel decrease in required ancillae. AriadneVR assists this discovery process by firstly providing easy access to graph structures and secondly offering an intuitive method of defining new graphs through manual drawing.

Similar to the examples above, here VR helped us to uncover the underlying working principles, for which we do not know any automated methodology. Applying this new working principle to a new problem is a demonstration that we understand the new working principle.

Accelerate AI discoveries through VR via customized search spaces - Potentially useful generalizations to quantum optical experiments can involve the increase in particle count, dimensionality, or both [69, 70]. Searching for such generalizations from scratch can be inefficient as the corresponding initial graphs, i.e. the search spaces, quickly increase in size. For example, the initial graph for 4-particle 2-dimensional GHZ-state generation has 24 edges. Searching for a 3-dimensional generalization more than doubles this amount to 54 edges, and attempting a 4-dimensional state without additional ancilla raises the size to 96 edges.

We can use AriadneVR to customize our search spaces and decrease the size of the initial graph to accelerate PyTheus to more efficiently discover desired solutions. We achieve this by analyzing existing solutions for lower graph sizes and defining candidate geometries of extensions based on observed patterns. In figure 6 we show this process for the generalization of a 3-particle 3-dimensional to a 4-particle 3-dimensional GHZ-state analyzer. The 3-particle analyzer is highly symmetric. Our candidate for a 4-particle geometry keeps this symmetry and follows the same patterns observed in the smaller solution. By restricting the initial graph to this geometry, the search for the 4-particle analyzer is reduced from 124 edges

down to 74. In this way, we can use human intelligence to assist PyTheus in its discovery task, which in turn can provide the human scientist with the example cases needed for concept extraction.

As a comparison, to find a suitable (high fidelity, low number of edges) solution starting from the complete graph with 124 edges, PyTheus needs 10621.4 ± 881 s (2.9 ± 0.24 h), while the topologically simpler starting graph with 74 edges requires only 343.6 ± 146.4 seconds. Thus, beyond finding an underlying pattern, the pattern has a true advantage in finding solutions.

6. Conclusion and outlook

In this work, we presented ways to leverage the unique potential of immersive technology in the AI-driven discovery process for quantum optics experiments using PyTheus and our purpose-built VR-tool AriadneVR. By interactively exploring the graph representations of quantum optics, a new approach to learning from AI to develop new experiments and insight is opened. Beyond furthering the understanding of existing solutions for GHZ-state creation and measurement, we achieved substantial machine-inspired discoveries with our tool. We engineered a new alternative 4-pair 3-dimensional entanglement swapping setup, where one pair is created at no additional photon cost with knowledge derived from AI discoveries. We also showed the human-assisted AI discovery of a 4-particle 3-dimensional GHZ-state analyzer. To further exploit the potential of VR our prototype could be expanded to include multi-user support for cooperative analysis and direct access to PyTheus within VR. Additionally, a variety of potential visualization features and tools could be added to streamline the analysis process. Examples include computational operations on vertices as well as support for more post-selection conditions. Beyond PyTheus' discoveries, the methods developed here can potentially also prove useful in other AI-discovery fields where graph-based representations are used, such as in quantum computing (quantum circuits [54], especially ZX-graphs [80]), material or protein design (involving crystal structures and molecular graphs [81]) and DNA-origami design [82, 83] in material science and biochemistry. We believe that enhancing human researchers' capabilities to comprehend complex data with technologies like VR is crucial to fully utilize the power of AI for scientific discovery. Beyond task-specific applications, immersive technologies also have the potential to fundamentally transform the scientific workspace by allowing remote collaboration and experimental control [84].

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: <https://github.com/artificial-scientist-lab/AriadneVR>.

ORCID iDs

Carlos Ruiz-Gonzalez  <https://orcid.org/0000-0003-0545-360X>

Xuemei Gu  <https://orcid.org/0000-0002-0734-432X>

Mario Krenn  <https://orcid.org/0000-0003-1620-9207>

References

- [1] Korkut E H and Surer E 2023 Visualization in virtual reality: a systematic review *Virtual Real.* **27** 1447
- [2] Duer Z, Piilonen L and Glasson G 2018 Belle2VR: a Virtual-Reality visualization of subatomic particle physics in the Belle II experiment *IEEE Comput. Graph. Appl.* **38** 33
- [3] Porter C, Smith J, Stagar E, Simmons A, Nieberding M, Orban C, Brown J and Ayers A 2020 Using virtual reality in electrostatics instruction: the impact of training *Phys. Rev. Phys. Educ. Res.* **16** 020119
- [4] Seritan S, Wang Y, Ford J E, Valentini A, Gold T and Martínez T J 2021 InteraChem: Virtual Reality visualizer for reactive interactive molecular dynamics *J. Chem. Educ.* **98** 3486
- [5] Wolfartsberger J 2019 Analyzing the potential of Virtual Reality for engineering design review *Autom. Constr.* **104** 27
- [6] Qin S, Wang Q and Chen X 2020 Application of virtual reality technology in nuclear device design and research *Fusion Eng. Des.* **161** 111906
- [7] Guo Z, Zhou D, Zhou Q, Zhang X, Geng J, Zeng S, Lv C and Hao A 2020 Applications of virtual reality in maintenance during the industrial product lifecycle: a systematic review *J. Manuf. Syst.* **56** 525
- [8] Yuan J, Hassan S S, Wu J, Koger C R, Packard R R S, Shi F, Fei B and Ding Y 2023 Extended reality for biomedicine *Nat. Rev. Methods Primers* **3** 1
- [9] Bohil C J, Alicea B and Biocca F A 2011 Virtual reality in neuroscience research and therapy *Nat. Rev. Neurosci.* **12** 752
- [10] Usher W, Klacansky P, Federer F, Bremer P-T, Knoll A, Yarch J, Angelucci A and Pascucci V 2018 A Virtual Reality visualization tool for neuron tracing *IEEE Trans. Vis. Comput. Graph.* **24** 994
- [11] Wang Y et al 2019 TeraVR empowers precise reconstruction of complete 3-D neuronal morphology in the whole brain *Nat. Commun.* **10** 3474
- [12] Steiniger B S, Pfeffer H, Gaffling S and Lobachev O 2022 The human splenic microcirculation is entirely open as shown by 3D models in virtual reality *Sci. Rep.* **12** 16487

- [13] Glowacki D R, Williams R R, Wonnacott M D, Maynard O M, Freire R, Pike J E and Chatziapostolou M 2022 Group VR experiences can produce ego attenuation and connectedness comparable to psychedelics *Sci. Rep.* **12** 8995
- [14] Donalek C et al 2014 Immersive and collaborative data visualization using virtual reality platforms *IEEE Int. Conf. on Big Data* pp 609–14
- [15] Pirch S et al 2021 The VRNetzer platform enables interactive network analysis in Virtual Reality *Nat. Commun.* **12** 2432
- [16] Kuták D, Vázquez P-P, Isenberg T, Krone M, Baaden M, Byska J, Kozliková B and Miao H 2023 State of the art of molecular visualization in immersive virtual environments *Comput. Graph. Forum* **42** e14738
- [17] O'Connor M et al 2018 Sampling molecular conformations and dynamics in a multiuser virtual reality framework *Sci. Adv.* **4** e2731
- [18] García-Hernández R J and Kranzlmüller D 2019 NOMAD VR: Multiplatform virtual reality viewer for chemistry simulations *Comput. Phys. Commun.* **237** 230
- [19] Kingsley L J, Brunet V, Lelais G, McCloskey S, Milliken K, Leija E, Fuhs S R, Wang K, Zhou E and Spraggon G 2019 Development of a virtual reality platform for effective communication of structural data in drug discovery *J. Mol. Graph. Modelling* **89** 234
- [20] Cassidy K C, Šefčík J, Raghav Y, Chang A, Durrant J D and Dina S-D 2020 ProteinVR: web-based molecular visualization in virtual reality *PLoS Comput. Biol.* **16** e1007747
- [21] Theart R P, Loos B and Niesler T R 2017 Virtual reality assisted microscopy data visualization and colocalization analysis *BMC Bioinform.* **18** 64
- [22] Stefani C, Lacy-Hulbert A and Skillman T 2018 Confocalvr: immersive visualization for confocal microscopy *J. Mol. Biol.* **430** 4028
- [23] Spark A et al 2020 vLUME: 3D virtual reality for single-molecule localization microscopy *Nat. Methods* **17** 1097
- [24] Blanc T, El Beheiry M, Caporal C, Masson J-B and Hajj B 2020 Genuage: visualize and analyze multidimensional single-molecule point cloud data in virtual reality *Nat. Methods* **17** 1100
- [25] Davelaar J, Bronzwaer T, Kok D, Younsi Z, Mo'scibrodzka M and Falcke H 2018 Observing supermassive black holes in virtual reality *Comput. Astrophys. Cosmol.* **5** 1
- [26] Sagrista A, Jordan S, Muller T and Sadlo F 2019 Gaia Sky: navigating the Gaia catalog *IEEE Trans. Vis. Comput. Graph.* **25** 1070
- [27] Baracaglia E and Vogt F P A 2020 E0102-VR: Exploring the scientific potential of Virtual Reality for observational astrophysics *Astron. Comput.* **30** 100352
- [28] Meissler N, Wohlan A, Hochgeschwender N and Schreiber A 2019 Using visualization of convolutional neural networks in virtual reality for machine learning newcomers *IEEE Int. Conf. on Artificial Intelligence and Virtual Reality (AIVR)* pp 152–1526
- [29] Bellgardt M, Scheiderer C and Kühlen T W 2020 An immersive node-link visualization of artificial neural networks for machine learning experts *IEEE Int. Conf. on Artificial Intelligence and Virtual Reality (AIVR)* pp 33–36
- [30] Lyu Z, Li J and Wang B 2021 Alive: interactive visualization and sonification of neural networks in Virtual Reality *IEEE Int. Conf. on Artificial Intelligence and Virtual Reality (AIVR)* pp 251–5
- [31] Minh D, Wang H X, Li Y F and Nguyen T N 2022 Explainable artificial intelligence: a comprehensive review *Artif. Intell. Rev.* **55** 3503
- [32] Zhavoronkov A et al Potential non-covalent SARS-CoV-2 3C-like protease inhibitors designed using generative deep learning approaches and reviewed by human medicinal chemist in virtual reality (2020)
- [33] El Beheiry M, Doutreligne S, Caporal C, Ostertag C, Dahan M and Masson J-B 2019 Virtual reality: beyond visualization *J. Mol. Biol.* **431** 1315
- [34] O'Connor M B, Bennie S J, Deeks H M, Jamieson-Binnie A, Jones A J, Shannon R J, Walters R, Mitchell T J, Mulholland A J and Glowacki D R 2019 Interactive molecular dynamics in virtual reality from quantum chemistry to drug binding: an open-source multi-person framework *J. Chem. Phys.* **150** 220901
- [35] Lanrezac A, Férey N and Baaden M 2022 Wielding the power of interactive molecular simulations *WIREs Comput. Mol. Sci.* **12** e1594
- [36] Walters R K, Gale E M, Barnoud J, Glowacki D R and Mulholland A J 2022 The emerging potential of interactive virtual reality in drug discovery *Exp. Opin. Drug Discovery* **17** 685
- [37] Deeks H M, Zinovjev K, Barnoud J, Mulholland A J, van der Kamp M W and Glowacki D R 2023 Free energy along drug-protein binding pathways interactively sampled in virtual reality *Sci. Rep.* **13** 16665
- [38] Amabilino S, Bratholm L A, Bennie S J, O'Connor M B and Glowacki D R 2020 Training atomic neural networks using fragment-based data generated in virtual reality *J. Chem. Phys.* **153** 154105
- [39] Ferretti S, Bianchi S, Frangipane G and Di Leonardo R 2021 A virtual reality interface for the immersive manipulation of live microscopic systems *Sci. Rep.* **11** 7610
- [40] Krenn M, Gu X and Zeilinger A 2017 Quantum experiments and graphs: multiparty states as Coherent superpositions of perfect matchings *Phys. Rev. Lett.* **119** 240403
- [41] Gu X, Chen L, Zeilinger A and Krenn M 2019 Quantum experiments and graphs. III. high-dimensional and multiparticle entanglement *Phys. Rev. A* **99** 032338
- [42] Gu X, Erhard M, Zeilinger A and Krenn M 2019 Quantum experiments and graphs II: quantum interference, computation and state generation *Proc. Natl Acad. Sci.* **116** 4147
- [43] Hütter C V R, Sin C, Müller F and Menche J 2022 Network cartographs for interpretable visualizations *Nat. Comput. Sci.* **2** 84
- [44] Krenn M, Kottmann J S, Tischler N and Aspuru-Guzik A 2021 Conceptual understanding through efficient automated design of quantum optical experiments *Phys. Rev. X* **11** 031044
- [45] Greenberger D M, Horne M A, Shimony A and Zeilinger A 1990 Bell's theorem without inequalities *Am. J. Phys.* **58** 1131
- [46] Krenn M, Landgraf J, Foesel T and Marquardt F 2023 Artificial intelligence and machine learning for quantum technologies *Phys. Rev. A* **107** 010101
- [47] Wang H et al 2023 Scientific discovery in the age of artificial intelligence *Nature* **620** 47
- [48] Jumper J et al 2021 Highly accurate protein structure prediction with AlphaFold *Nature* **596** 583
- [49] Arnold C 2023 Inside the nascent industry of AI-designed drugs *Nat. Med.* **29** 1292
- [50] Sanchez-Lengeling B and Aspuru-Guzik A 2018 Inverse molecular design using machine learning: generative models for matter engineering *Science* **361** 360
- [51] Elton D C, Boukouvalas Z, Fuge M D and Chung P W 2019 Deep learning for molecular design—a review of the state of the art *Mol. Syst. Des. Eng.* **4** 828
- [52] Nigam A, Pollice R and Aspuru-Guzik A 2022 Parallel tempered genetic algorithm guided by deep neural networks for inverse molecular design *Digit. Discovery* **1** 390

- [53] Merchant A, Batzner S, Schoenholz S S, Aykol M, Cheon G and Cubuk E D 2023 Scaling deep learning for materials discovery *Nature* **624** 80
- [54] Kottmann J S 2023 Molecular quantum circuit design: a graph-based approach *Quantum* **7** 1073
- [55] Krenn M, Erhard M and Zeilinger A 2020 Computer-inspired quantum experiments *Nat. Rev. Phys.* **2** 649
- [56] Ruiz-Gonzalez C, Arlt S, Petermann J, Sayyad S, Jaouni T, Karimi E, Tischler N, Gu X and Krenn M 2023 Digital discovery of 100 diverse quantum experiments with PyTheus *Quantum* **7** 1204
- [57] De Regt H W and Dieks D 2005 A contextual approach to scientific understanding *Synthese* **144** 137
- [58] De Regt H W 2017 *Understanding Scientific Understanding* (Oxford University Press)
- [59] Krenn M et al 2022 On scientific understanding with artificial intelligence *Nat. Rev. Phys.* **4** 761
- [60] Krenn M, Malik M, Fickler R, Lapkiewicz R and Zeilinger A 2016 Automated Search for new Quantum Experiments *Phys. Rev. Lett.* **116** 090405
- [61] Knott P 2016 A search algorithm for quantum state engineering and metrology *New J. Phys.* **18** 073033
- [62] Arlt S Ruiz-Gonzalez C and Krenn M 2022 Digital discovery of a scientific concept at the core of experimental quantum optics (arXiv:2210.09981 [quant-ph])
- [63] Krenn M, Hochrainer A, Lahiri M and Zeilinger A 2017 Entanglement by path identity *Phys. Rev. Lett.* **118** 080401
- [64] Hochrainer A, Lahiri M, Erhard M, Krenn M and Zeilinger A 2022 Quantum indistinguishability by path identity and with undetected photons *Rev. Mod. Phys.* **94** 025007
- [65] Kysela J, Erhard M, Hochrainer A, Krenn M and Zeilinger A 2020 Path identity as a source of high-dimensional entanglement *Proc. Natl Acad. Sci.* **117** 26118
- [66] Bao J et al 2023 Very-large-scale integrated quantum graph photonics *Nat. Photon.* **17** 573
- [67] Qian K, Wang K, Chen L, Hou Z, Krenn M, Zhu S and Ma X-S 2023 Multiphoton non-local quantum interference controlled by an undetected photon *Nat. Commun.* **14** 1480
- [68] Feng L-T, Zhang M, Liu D, Cheng Y-J, Guo G-P, Dai D-X, Guo G-C, Krenn M and Ren X-F 2023 On-chip quantum interference between the origins of a multi-photon state *Optica* **10** 105
- [69] Erhard M, Krenn M and Zeilinger A 2020 Advances in high-dimensional quantum entanglement *Nat. Rev. Phys.* **2** 365
- [70] Hu X-M, Guo Y, Liu B-H, Li C-F and Guo G-C 2023 Progress in quantum teleportation *Nat. Rev. Phys.* **5** 339
- [71] Ryu J, Lee C, Żukowski M and Lee J 2013 Greenberger-horne-zeilinger theorem for n qudits *Phys. Rev. A* **88** 042101
- [72] Lawrence J 2014 Rotational covariance and greenberger-horne-zeilinger theorems for three or more particles of any dimension *Phys. Rev. A* **89** 012105
- [73] Xing W-B, Hu X-M, Guo Y, Liu B-H, Li C-F and Guo G-C 2023 Preparation of multiphoton high-dimensional GHZ state *Opt. Express* **31** 24887
- [74] Pan J-W and Zeilinger A 1998 Greenberger-Horne-Zeilinger-state analyzer *Phys. Rev. A* **57** 2208
- [75] Żukowski M, Zeilinger A, Horne M A and Ekert A K 1993 'event-ready-detectors' bell experiment via entanglement swapping *Phys. Rev. Lett.* **71** 4287
- [76] Pan J-W, Bouwmeester D, Weinfurter H and Zeilinger A 1998 Experimental entanglement swapping: entangling photons that never interacted *Phys. Rev. Lett.* **80** 3891
- [77] Bose S, Vedral V and Knight P L 1998 Multiparticle generalization of entanglement swapping *Phys. Rev. A* **57** 822
- [78] Csardi G and Nepusz T 2005 The Igraph software package for complex network research interjournal complex systems pp 1695
- [79] Kamada T and Kawai S 1989 An algorithm for drawing general undirected graphs *Inf. Process. Lett.* **31** 7–15
- [80] Duncan R, Kissinger A, Perdrix S and Wetering J V D 2020 Graph-theoretic simplification of quantum circuits with the ZX-calculus *Quantum* **4** 279
- [81] Butler K T, Davies D W, Cartwright H, Isayev O and Walsh A 2018 Machine learning for molecular and materials science *Nature* **559** 547
- [82] Rothmund P W K 2006 Folding DNA to create nanoscale shapes and patterns *Nature* **440** 297
- [83] DeLuca M, Sensale S, Lin P-A and Arya G 2023 Prediction and control in DNA nanotechnology *ACS Appl. Bio Mater.* **7** 626–45
- [84] Gómez-Zarà D, Schiffer P and Wang D 2023 The promise and pitfalls of the metaverse for science *Nat. Hum. Behav.* **7** 1237