

Patient Details (or infographic sticker)

Patient Name: _____

Hospital Number: _____

Date of Birth: _____

Whiston Emergency**Department****Open Fracture Management Proforma****Injury Details**

Mechanism of injury?	
Date & Time of injury (<i>not arrival to A&E</i>):/...../..... ..:.....
Fracture location:	Upper limb Lower limb

Clinical History & Examination

--

Neurovascular Status of Affected Limb:

Pre-manipulation			Post-manipulation			2 nd manipulation		
Tick relevant:	Normal	Abnormal	Tick relevant:	Normal	Abnormal	Tick relevant:	Normal	Abnormal
Power			Power			Power		
Sensation			Sensation			Sensation		
Cap Refill			Cap Refill			Cap Refill		
Pulses			Pulses			Pulses		

If any of the above abnormal, please detail below:

--

Whiston Open Fracture Management Proforma

Patient Details (or infographic sticker)

Patient Name: _____

Hospital Number: _____

Date of Birth: _____

Wound Assessment

Is the wound visibly contaminated? *Yes is indication for immediate debridement, contact Orthopaedics (or Plastics for hands) on call.*

Yes

No

Drawing of wound; *please comment on bruising, degloving etc*

Medical photography

Wound photographed?

Yes

No

Mode of photography?

Medical Photography

Careflow Connect

Consent gained and documented?

Yes

No (state reason below)

Wound Management

Saline-soaked gauze with an occlusive film applied?

Yes

No

If no, document reason

Wound irrigation as standard is **NOT** recommended by NICE for open fractures of the long bones, hindfoot or midfoot in ED.

If this has been performed, please document reason

Patient Details (or infographic sticker)

Patient Name: _____

Hospital Number: _____

Date of Birth: _____

Antibiotic Administration **GIVE WITHIN 1 HOUR OF ARRIVAL**	
Patient allergy status	
Antibiotics administered in ED?	Yes (complete below)
	Administered pre-hospital
	No please state reason:
Antibiotic:	
Dose:	
Time/date of administration:/...../..... :.....
Tetanus status?	Up to Date Booster Given Immunoglobulins Given
OPEN FRACTURE ANTIBIOTIC GUIDELINES	
<u>First Line AND Non-Severe Penicillin Allergy</u>	CefUROxime 1.5g Repeated 8 hourly until surgery complete AND at first debridement Gentamicin* 3mg/kg (single dose)
<u>Severe Penicillin Allergy</u>	As above BUT <u>Clindamycin</u> 600mg 6 hourly instead of <u>CefUROxime</u>.
<u>Known MRSA</u>	Vancomycin (IV) (refer to Micro Guide for dosing) AND at first debridement Gentamicin* 3mg/kg (single dose)
Oral antibiotics only in clean wounds <1cm following discussion with specialty team	