



Patient Profile

Name: Mr. Saubhik Bhaumik

Age:

Gender: Male

Summary of Results

All evaluated laboratory parameters fall within accepted reference ranges. The predictive model flags this profile as **Normal**, indicating no immediate biochemical abnormalities requiring urgent clinical intervention.

Detailed Test Explanations

SERUM BILIRUBIN (TOTAL) – 0.9 mg/dl (Normal: 0.2-1.2 mg/dl). Total bilirubin measures overall bilirubin load; values within range suggest normal hepatic processing and no hemolysis.

SERUM BILIRUBIN (DIRECT) – 0.2 mg/dl (Normal: 0-0.3 mg/dl). Direct bilirubin reflects conjugated bilirubin; normal value indicates no cholestatic disease.

SERUM BILIRUBIN (INDIRECT) – 0.7 mg/dl (Normal: 0.1-0.9 mg/dl). Unconjugated bilirubin is within normal limits, further supporting healthy erythrocyte turnover and hepatic conjugation.

SGPT (ALT) – 36 U/L (Normal: 13-40 U/L). ALT is a liver enzyme; level within range indicates no hepatocellular injury.

SGOT (AST) – 32 U/L (Normal: 0-37 U/L). AST is another hepatic enzyme; normal value complements ALT findings.

SERUM ALKALINE PHOSPHATASE – data not provided.

SERUM PROTEIN – 7.2 g/dl (Normal: 6.4-8.3 g/dl). Total protein reflects nutritional status and hepatic synthetic capacity.

SERUM ALBUMIN – 4.7 g/dl (Normal: 3.5-5.2 g/dl). Albumin indicates plasma oncotic pressure and liver synthetic function.

GLOBULIN – 2.5 g/dl (Normal: 1.8-3.6 g/dl). Globulin range is consistent with a healthy immune and metabolic profile.

AIG RATIO – 1.88 (Normal: 1.1-2.4). The albumin-to-globulin ratio falls within expected limits, signifying balanced protein synthesis.

Overall Interpretation

The panel demonstrates a normal hepatic function profile, with bilirubin fractions, transaminases, protein fractions, and albumin/globulin ratio all within their respective reference ranges. The absence of elevated liver enzymes, abnormal bilirubin, or protein derangements suggests no current liver pathology or significant systemic disease.

Recommendations

- Maintain a balanced diet rich in fruits, vegetables, and lean proteins to support ongoing liver health.
- Limit alcohol consumption and avoid hepatotoxic medications where possible.
- Continue regular health screenings; repeat liver function testing if symptoms such as jaundice, abdominal pain, or fatigue arise.

- Discuss with a primary care provider any persistent liver-related concerns or family history of hepatic disease.

Disclaimer

This report is for informational purposes only and does not substitute for professional medical advice, diagnosis, or treatment. Please consult a qualified healthcare professional for personalized guidance.