

Smart AI Medical Report

Generated by Health Navigator

Patient Profile

(Mr. Saubhik Bhaumik)

Summary of Results

All laboratory values fall within established reference ranges. The predictive model has assessed the overall profile as Normal, indicating no evidence of clinically significant abnormalities.

Detailed Test Explanations

SERUM BILIRUBIN (TOTAL) – 0.9/mg/dl

Total bilirubin reflects the liver's ability to conjugate and excrete waste pigments. Levels within 0.2–1.2/mg/dl are considered normal; this result indicates health.

SERUM BILIRUBIN (DIRECT) – 0.2/mg/dl

Direct (conjugated) bilirubin is usually <0.3/mg/dl. The measure suggesting no cholestatic or hepatic cell damage.

SERUM BILIRUBIN (INDIRECT) – 0.70/mg/dl

Indirect (unconjugated) bilirubin should be <1.0/mg/dl. A value of 0.70/mg/dl suggests balanced hemolysis and hepatic processing.

SGPT (ALT) – 36/U

Alanine aminotransferase levels of 13–40/U/L^{1} are considered normal. A value of 36/U does not signal hepatocellular injury; this result demonstrates normal hepatic enzyme activity.

SGOT (AST) – 32/U

Aspartate aminotransferase normal range is 0–37/U/L^{1}. The value of 32/U is within the normal range but does not indicate pathology.

SERUM ALKALINE PHOSPHATASE – Not Provided

With no value available, no interpretation can be made.

SERUM PROTEIN – 7.2/g/dL^{1}

Total serum protein of 6.4–8.3/g/dL^{1} is normal. This measurement reflects liver synthetic function.

SERUM ALBUMIN – 4.7/g/dL^{1}

Normal albumin range is 3.5–5.2/g/dL^{1}, confirming good nutritional status and protein synthesis.

GLOBULIN – 2.5/g/dL^{1}

Globulin values of 1.8–3.6/g/dL^{1} are normal, implying balanced immune response and protein production.

AIG RATIO – 1.88

The albumin-to-globulin ratio normally ranges from 1.1 to 2.4; a ratio of 1.88 falls squarely within this interval.

Overall Interpretation

The composite laboratory profile demonstrates no deviations from normal limits across measured parameters. The absence of elevated liver enzymes, bilirubin species, or protein abnormalities indicates preserved hepatic function and normal systemic protein metabolism. The predictive model, reflecting these findings, flags the profile as Normal, suggesting no immediate clinical concern.

Recommendations

Maintain a balanced diet rich in fruits, vegetables, whole grains, and lean proteins to support hepatic health.

Continue regular physical activity and moderate alcohol consumption.

Schedule routine follow up and complete any pending laboratory phosphatase) to sustain comprehensive monitoring.

Report any new symptoms (jaundice, right upper quadrant pain, fatigue) promptly for reassessment.

Disclaimer

This report is provided for informational purposes only and does not substitute for professional medical evaluation, diagnosis, or treatment. Patients should consult their healthcare provider for personalized medical advice.