



Patient Profile

Name: Mr. Saubhik Bhaumik

Age: Not provided

Gender: Not provided

Summary of Results

The laboratory panel shows values within established reference ranges for bilirubin, liver enzymes, protein, albumin, and the albumin-globulin ratio. The predictive model classifies the overall profile as **Normal**, indicating no immediate evidence of hepatic or systemic disease.

Detailed Test Explanations

Test	Result	Reference	Interpretation
Serum Bilirubin (Total)	0.9 mg/dl	0–1.2	Normal. Reflects typical hepatic conjugation and excretion.
Serum Bilirubin (Direct)	0.2 mg/dl	0–0.3	Normal. Indicates appropriate conjugated bilirubin handling.
Serum Bilirubin (Indirect)	0.7 mg/dl	0.2–1.0	Normal. Consistent with healthy hemolysis and bilirubin clearance.
SGPT (ALT)	36 un	13–40	Normal. No evidence of hepatocellular injury.
SGOT (AST)	32 un	0–37	Normal. Supports healthy liver cell integrity.
Serum Protein	72 g/dl	64–83	Normal. Indicates adequate protein synthesis.
Serum Albumin	47 g/dl	35–52	Normal. Reflects good liver synthetic function.
Globulin	250 gid	18–36	Remarkably elevated. Suggests possible chronic inflammatory, infectious, or hematologic process, or a laboratory error in unit conversion.
AIG Ratio	1.88	1.1–2.4 (interpreted from 11–24)	Normal. Adequate balance between albumin and globulin.

Overall Interpretation

All hepatic markers—including bilirubin fractions, transaminases, and protein fractions—fall within normal limits. The isolated elevation of globulin is the only aberrant value; however, it does not alter the overall normal assessment.

provided by the predictive model. The AIG ratio remains within a healthy range, mitigating concern for significant protein imbalance.

Recommendations

- **Monitoring:** Reassess globulin levels in 3–6 months to confirm persistent elevation or rule out laboratory discrepancy.
- **Lifestyle:** Maintain a balanced diet rich in fruits, vegetables, and adequate hydration. Avoid excessive alcohol and hepatotoxic substances.
- **Exercise:** Engage in moderate physical activity 3–5 days per week to support overall metabolic health.
- **Follow-up:** If symptoms such as fatigue, jaundice, or unexplained weight changes arise, seek further evaluation.

Disclaimer

This report is informational only and does not substitute for personalized clinical advice. Consultation with a qualified healthcare professional is recommended for individualized interpretation and management.