



Patient Profile

Name: Mr. Saubhik Bhaumik

Age:

Gender:

Summary of Results

The comprehensive liver function panel demonstrates values within normal reference ranges. The predictive model flags the overall profile as **Normal**, indicating no immediate abnormality detected.

Detailed Test Explanations

- **SERUM BILIRUBIN (TOTAL)** – 0.9 mg/dl (0–1.2)
Total bilirubin measures combined direct (conjugated) and indirect (unconjugated) bilirubin. Normal values indicate effective hepatic excretion and bilirubin metabolism.
- **SERUM BILIRUBIN (DIRECT)** – 0.2 mg/dl (0–0.3)
Direct bilirubin reflects conjugated bile pigments cleared by the liver. The value is within normal limits, suggesting intact biliary clearance.
- **SERUM BILIRUBIN (INDIRECT)** – 0.7 mg/dl (0.2–1)
Indirect bilirubin represents unconjugated bilirubin. A normal range implies balanced hemolysis and hepatic uptake.
- **SGPT (ALT)** – 36 U/L (13–40)
Alanine aminotransferase is a sensitive marker of hepatocellular injury. The result is normal, indicating no significant liver cell damage.
- **SGOT (AST)** – 32 U/L (0–37)
Aspartate aminotransferase is present in liver and other tissues. A normal value corroborates healthy hepatic function.
- **SERUM ALKALINE PHOSPHATASE** – Not reported
Alkaline phosphatase levels are routinely used to assess cholestasis and bone turnover; absence of data precludes interpretation.
- **SERUM PROTEIN** – 7.2 g/dl (6.4–8.3)
Total protein reflects synthesis of albumin and globulins. The result is within normal limits, supporting adequate hepatic synthetic capacity.
- **SERUM ALBUMIN** – 4.7 g/dl (3.5–5.2)
Albumin indicates liver synthetic function and nutritional status. This value falls comfortably within normal range.
- **GLOBULIN** – 2.5 g/dl (1.8–3.6)
Globulins encompass immunoglobulins and acute-phase proteins. The level is normal, suggesting no abnormal protein synthesis.

- **AIG RATIO** – 1.88 (1.1–2.4)

The albumin-to-globulin ratio evaluates the balance of these major plasma proteins. A value within range indicates no dysregulation.

Overall Interpretation

All assessed biochemical parameters remain within reference intervals, supporting a healthy liver profile. The absence of abnormal elevations in transaminases, bilirubin fractions, and protein synthesis markers suggests no evidence of acute or chronic hepatocellular injury, cholestasis, or synthetic dysfunction at this time.

Recommendations

- **Maintain a balanced diet** rich in fruits, vegetables, and whole grains while limiting saturated fats and excessive alcohol intake to preserve hepatic health.
- **Stay hydrated** and avoid over-use of over-the-counter medications that can be hepatotoxic.
- **Regular monitoring:** Annual liver function tests are advised if no pre-existing liver disease, or sooner if any symptoms develop (jaundice, abdominal pain, fatigue).
- **Lifestyle:** Engage in moderate exercise and maintain a healthy weight to reduce the risk of non-alcoholic fatty liver disease.

Disclaimer

This report is for informational purposes only and does not replace professional medical advice. Consult a qualified healthcare provider for personalized diagnosis and treatment.