

Elementary Differential Geometry

Hassium

1	Calculus on Euclidean Space	6	Geometry of Surfaces in \mathbb{R}^3
2	Frame Fields	7	Riemannian Geometry
3	Euclidean Geometry	8	Global Structure of Surfaces
4	Calculus on a Surface		Exercises and Proofs
5	Shape Operators		Alphabetical Index

1 Calculus on Euclidean Space

Definition 1.1. The *Euclidean 3-space*, denoted \mathbb{R}^3 , is the set of ordered triples of the form $p = (p_1, p_2, p_3)$, where $p_i \in \mathbb{R}$. An element of \mathbb{R}^3 is called a *point*.

Let $p = (p_1, p_2, p_3), q = (q_1, q_2, q_3) \in \mathbb{R}^3$ and let $a \in \mathbb{R}$. Define the addition to be $p + q = (p_i + q_i)$ and define the scalar multiplication to be $ap = (ap_i)$. The additive identity $0 = (0, 0, 0)$ is called the *origin* of \mathbb{R}^3 . It is trivial that \mathbb{R}^3 is a vector space over \mathbb{R} .

Definition 1.2. Let x, y , and z be real-valued functions on \mathbb{R}^3 such that for all $p = (p_1, p_2, p_3) \in \mathbb{R}^3$, $x(p) = p_1$, $y(p) = p_2$, and $z(p) = p_3$. We call x, y , and z the *natural coordinate functions* of \mathbb{R}^3 .

Let x, y , and z be the natural coordinate functions, rewrite $x = x_1$, $y = x_2$, and $z = x_3$. Then we have $p = (p_i) = (x_i(p))$.

Definition 1.3. A real-valued function f on \mathbb{R}^3 is *differetiable* if all partial derivatives exist and continuous.

Definition 1.4. A subset $O \subset \mathbb{R}^3$ is *open* if for all $p \in O$, there exists $\varepsilon > 0$ such that $\{x \in \mathbb{R}^3 \mid \|x - p\| < \varepsilon\} \subset O$.

Let $f : O \rightarrow \mathbb{R}$ be a function defined on an open set. The differetiability of f at p can be determined entirely from values of f on O . This means that differetiation is a local operation. We will give a proof of this later.

Definition 1.5. A *tangent vector* v_p is an ordered pair $v_p = (v, p)$, where $v, p \in \mathbb{R}^3$. Here v is called the *vector part* and p is called its *point of application*. Two tangent vectors are said to be *parallel* if they have the same vector part and different points of application.

Definition 1.6. Let $p \in \mathbb{R}^3$. The *tangent space* at p , denoted $T_p(\mathbb{R}^3)$, is the set of all tangent vectors that have p as point of application.

Fix a tangent space $T_p(\mathbb{R}^3)$.

-
- 2 Frame Fields
 - 3 Euclidean Geometry
 - 4 Calculus on a Surface
 - 5 Shape Operators
 - 6 Geometry of Surfaces in \mathbb{R}^3
 - 7 Riemannian Geometry
 - 8 Global Structure of Surfaces

Exercises and Proofs

Exercise 1.1.1. Let $f = x^2y$ and $g = y \sin z$ be functions on \mathbb{R}^3 . Express the following functions in terms of x , y , and z .

1. fg^2 ;
2. $\frac{\partial f}{\partial x}g + \frac{\partial g}{\partial y}f$;
3. $\frac{\partial^2(fg)}{\partial y \partial z}$;
4. $\frac{\partial}{\partial y}(\sin f)$.

Proof. (i) We have $fg^2 = x^2yy^2\sin^2z = x^2y^3\sin^2z$. (ii) We have $\frac{\partial f}{\partial x} = 2xy$ and $\frac{\partial g}{\partial y} = \sin z$, then $\frac{\partial f}{\partial x}g + \frac{\partial g}{\partial y}f = 2xy^2\sin z + x^2y\sin z$. (iii) We have $fg = x^2y^2\sin z$, then $\frac{\partial^2(fg)}{\partial y \partial z} = 2x^2y\cos z$. (iv) We have $\sin f = \sin(x^2y)$, then $\frac{\partial}{\partial y}(\sin f) = x^2\cos(x^2y)$. \square

Exercise 1.1.3. Express $\frac{\partial f}{\partial x}$ in terms of x , y , and z for the following functions.

1. $f = x \sin(xy) + y \cos(xz)$;
2. $f = \sin g$, $g = e^h$, and $h = x^2 + y^2 + z^2$.

Proof. (i) We have $\frac{\partial f}{\partial x} = \frac{x \sin(xy)}{\partial x} + \frac{\partial y \cos(xz)}{\partial x} = \sin(xy) + xy \cos(xy) - yz \sin(xz)$. (ii) We have $f = \sin(e^{x^2+y^2+z^2})$, then $\frac{\partial f}{\partial x} = 2x \cos(e^{x^2+y^2+z^2})e^{x^2+y^2+z^2}$. \square

Exercise 1.2.1. Let $v = (-2, 1, -1)$ and $w = (0, 1, 3)$. At an arbitrary point p , express the tangent vector $3v_p - 2w_p$ as a linear combination of $U_1(p)$, $U_2(p)$, and $U_3(p)$.

Alphabetical Index

differetiable, 1

Euclidean 3-space, 1

natural coordinate functions, 1

open, 1

origin, 1

parallel, 1

point, 1

point of application, 1

tangent space, 1

tangent vector, 1

vector part, 1