Please check the examination details below before entering your candidate information

Candidate surname	Other names

Pearson Edexcel International Advanced Level

Thursday 14 January 2021

Morning (Time: 1 hour 30 minutes) | Paper Reference **WMA13/01**

Mathematics
International Advanced Level
Pure Mathematics P3

You must have:
Mathematical Formulae and Statistical Tables (Lilac), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions
- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information
- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 30 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice
- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over >

P 6 6 3 8 9 A
www.pearson.com/uk
©2020

---

1. Find

$$\int \frac{x^2-5}{2x^3}\,dx \qquad x>0$$

giving your answer in simplest form.

(3)

1 (a) $\int \frac{x^2-5}{2x^3}\,dx$

$\int \frac{x^2}{2x^3} - \frac{5}{2x^3}\,dx$

$\int \frac{1}{2x} - \frac{5}{2}x^{-3}\,dx$

$I = \frac{\ln|x|}{2} - \frac{\frac{5}{2}x^{-2}}{-2} + c$

$= \frac{1}{2}\ln x + \frac{5}{4x^2} + c$ ✓

2.



Figure 1

Figure 1 shows a sketch of the curve with equation $y = f(x)$, where $x \in \mathbb{R}$ and f(x) is a polynomial.

The curve passes through the origin and touches the x-axis at the point (3, 0)

There is a maximum turning point at (1, 2) and a minimum turning point at (3, 0)

On separate diagrams, sketch the curve with equation

(i) $y = 3f(2x)$

(3)

(ii) $y = f(-x) - 1$

(3)

On each sketch, *show clearly* the coordinates of

- the point where the curve crosses the y-axis
- any maximum or minimum turning points

(2a) 3f(2x)

y: (1,6) (3,3) (0,3)

2x: (3,0) (6,0)

(0,3) (1,6) (3,0) x

Question 2 continued

(ii) f(−x) − 1

x is reflected over y-axis

function shifted down −1

(−3,1) (0,−1) (3,−1)

(Total 6 marks)

Q2

3.

$$f(x) = 3 - \frac{x-2}{x+1} + \frac{5x+26}{2x^2-3x-5} \qquad x>4$$

(a) Show that

$$f(x) = \frac{ax+b}{cx+d} \qquad x>4$$

where a, b, c and d are integers to be found.

(4)

(b) Hence find $f^{-1}(x)$

(2)

(c) Find the domain of f^{-1}

(2)

3(a) $3 - \frac{x-2}{x+1} + \frac{5x+26}{(x+1)(2x-5)}$

$\frac{3(x+1)(2x-5) - (x-2)(2x-5) + (5x+26)}{(x+1)(2x-5)}$

$\frac{3(2x^2-3x-5) - (2x^2-5x-2x+10) + 5x+26}{(x+1)(2x-5)}$

$\frac{6x^2-9x-15-2x^2+7x-10+5x+26}{(x+1)(2x-5)}$

$\frac{4x^2+5x+1}{(x+1)(2x-5)}$

$\frac{(4x+1)(x+1)}{(x+1)(2x-5)} = \frac{4x+1}{2x-5}$ ✓

Question 3 continued

(b) $f^{-1} \Rightarrow$ inverse $x \Leftrightarrow y$

$y = \frac{4x+1}{2x-5}$

$x = \frac{4y+1}{2y-5}$

$2xy - 5x = 4y+1$

$2xy - 4y = 5x+1$

$y(2x-4) = 5x+1$

$y = \frac{5x+1}{2x-4}$ ✓

(c) domain of $y = \frac{5x+1}{2x-4}$

LB $2x-4 \neq 0$

$x \neq 2$

UB $x > 4$ inf

$\therefore y > 4$ inf f^{-1}

Question 3 continued

$4 = \frac{5x+1}{2x-4}$

$8x - 16 = 5x + 1$

$3x = 17$

$x = \frac{17}{3}$

$\therefore 2 < x < \frac{17}{3}$ ✓

4.



Figure 2

Figure 2 shows a sketch of the graph with equation $y = f(x)$, where

$$f(x) = |kx + a| + a$$

and where a is a positive constant.

The graph has a vertex at the point P, as shown in Figure 2.

(a) Find, in terms of a, the coordinates of P.

(2)

(b) Sketch the graph with equation $y = g(x)$, where

$$g(x) = |x + 5a|$$

On your sketch, *show* the coordinates, in terms of a, of each point where the graph cuts or meets the coordinate axes.

(2)

The graph with equation $y = g(x)$ intersects the graph with equation $y = f(x)$ at two points.

(c) Find, in terms of a, the coordinates of the two points.

(5)

4(a) $y = kx + a + a$ or $y = -kx - a + a$

$y = kx + 2a$ $\quad y = -kx$

not $y > 0$ $\therefore kx + 2a = -kx$

$6x = -2a$

$x = -\frac{2a}{6} = -\frac{a}{3}$

Question 4 continued

$\therefore y = -3\left(-\frac{a}{3}\right)$

$= a$

$\therefore \left(-\frac{a}{3}, a\right)$ ✓

(b) $y = x + 5a$ or $y = -x - 5a$

y is always positive

x-int always

$0 = x + 5a$ $\quad 0 = -x - 5a$

$-x = 5a$ $\quad x = -5a$

y-int, let x = 0

$y = 0 + 5a$ $\quad y = 0 - 5a$

$y = 5a$ $\quad y = -5a$ invalid



Question 4 continued

(c)



Figure 2

Only positive y's exist f(x)

$\therefore -x + a = -3x$ or $x + 5a = 3x + 2a$

$4x = -2a$ $\quad 5a = 2x$

$x = -\frac{2a}{4}$ $\quad x = \frac{5a}{2}$

$\therefore y = -3\left(-\frac{2a}{4}\right)$ $\quad \therefore y = 3\left(\frac{5a}{2}\right) + 2a$

$= +\frac{3a}{2}$ $\quad = \frac{15}{2}a + \frac{4}{2}a$

$\therefore \left(\frac{3}{2}a, \frac{3}{2}a\right)$ ✓ $\quad = \frac{19}{2}a$

$\therefore \left(\frac{5}{2}a, \frac{19}{2}a\right)$ ✓

---

5. The temperature, θ°C, inside an oven, t minutes after the oven is switched on, is given by

$$\theta = A - 150e^{-kt}$$

where A and k are positive constants.

Given that the temperature inside the oven is initially 18 °C,

(a) find the value of A.

(2)

The temperature inside the oven, 5 minutes after the oven is switched on, is 90 °C.

(b) Show that $k = p\ln q$ where p and q are rational numbers to be found.

(4)

Hence find

(c) the temperature inside the oven 9 minutes after the oven is switched on, giving your answer to 3 significant figures.

(2)

(d) the rate of increase of the temperature inside the oven 9 minutes after the oven is switched on. Give your answer in °C min⁻¹ to 3 significant figures.

(3)

5. (a) $\theta = 18$ at $t = 0$

$18 = A - 150e^{-k(0)}$

$18 = A - 150 \times 1$

$A = 198$ ✓

(b) $\theta = 90$ when $t = 5$

$90 = 198 - 150e^{-k(5)}$

$-108 = -150e^{-5k}$

$\ln\frac{108}{150} = -5k$

$-5k = \ln\frac{108}{150}$ $\quad k = \frac{\ln\frac{108}{150}}{-5} = -\frac{1}{5}\ln\frac{3}{5}$ ✓

* note: add k is positive constant $\therefore \frac{1}{5}\ln\frac{3}{5}$

Question 5 continued

(c) $t = 9$

$\theta = 198 - 150e^{-\left(-\frac{1}{5}\ln\frac{3}{5}\right)(9)}$

$\theta = 126.2297\ldots$

$= 126$ ✓ (3sf)

(d) rate of increase

$\therefore \frac{d\theta}{dt}$

$\theta = 198 - 150e^{\left(\frac{1}{5}\ln\frac{3}{5}\right)t}$

$\frac{d\theta}{dt} = -150e^{\left(\frac{1}{5}\ln\frac{3}{5}\right)t} \times -\frac{1}{5}\ln\frac{3}{5}$

$t = 9$

$\frac{d\theta}{dt} = -150e^{\left(\frac{1}{5}\ln\frac{3}{5}\right)(9)} \times -\frac{1}{5}\ln\frac{3}{5}$

$= 7.3326\ldots$ ✓

$= 7.33$ °C min⁻¹ (3sf)

6.

$$f(x) = x\cos\left(\frac{x}{3}\right) \qquad x>0$$

(a) Find f′(x)

(2)

(b) Show that the equation $f'(x) = 0$ can be written as

$$x = k\arctan\left(\frac{x}{9}\right)$$

where k is an integer to be found.

(2)

(c) Starting with $x_1 = 2.5$ use the iteration formula

$$x_{n+1} = k\arctan\left(\frac{x_n}{9}\right)$$

with the value of k found in part (b), to calculate the values of x_2 and x_3 giving your answers to 3 decimal places.

(2)

(d) Using a suitable interval and a suitable function that should be stated, show that a root of $f'(x) = 0$ is 2.581 correct to 3 decimal places.

(2)

6. (a) $u = x$ $\quad v = \cos\frac{x}{3}$ $\quad$ PRODUCT RULE

$u' = 1$ $\quad v' = -\sin\frac{x}{3} \times \frac{1}{3}$ $\quad u'v + v'u$

$f'(x) = \cos\frac{x}{3} + x\left(-\frac{1}{3}\sin\frac{x}{3} \times x\right)$

$f'(x) = \cos\frac{x}{3} - \frac{1}{3}x\sin\frac{x}{3}$ ✓

(b) $\cos\frac{x}{3} - \frac{1}{3}x\sin\frac{x}{3} = 0$ $\quad x = k\arctan\left(\frac{x}{9}\right)$

$\cos\frac{x}{3} = \frac{1}{3}x\sin\frac{x}{3}$

$\frac{3}{x} = \frac{\sin\frac{x}{3}}{\cos\frac{x}{3}}$

$\tan\frac{x}{3} = \frac{3}{x}$

Question 6 continued

$\frac{x}{3} = \tan^{-1}\left(\frac{3}{x}\right)$

$x = 3\tan^{-1}\left(\frac{3}{x}\right)$ ✓

$= 3\arctan\left(\frac{3}{x}\right)$

(c) $x_1 = 2.5$

$x_2 = 3\arctan\left(\frac{3}{2.5}\right)$

$x_2 = 2.67817\ldots$

$= 2.678$ ✓ (3dp pt)

Calculator

① [2] [ans] [÷] [9] [=]

(is there)

[3] [tan⁻¹] [3] [÷] [ans] [=]

for x_2

② keep pressing [=] for x_3 etc

$x_3 = 2.55410\ldots$

$x_4 = 2.59652\ldots$

$x_5 = 2.5924\ldots$

$x_6 = 2.5961\ldots$

$= 2.596$ (3dp pt) ✓

Question 6 continued

(d) $x_3 = 2.578$

$x_5 = 2.582\ldots$

$x_9 = 2.58029\ldots$

$x_{10} = 2.581159\ldots$

$x_{11} = 2.58067\ldots$

$x_{12} = 2.581185\ldots$

$x_{13} = 2.58059\ldots$

$x_{14} = 2.58106\ldots$

$x_{15} = 2.5809\ldots$

don't mind this

So, root between 2.5805 and 2.5815

$f'(2.5805) = \cos\left(\frac{2.5805}{3}\right) - \frac{1}{3}(2.5805)\sin\left(\frac{2.5805}{3}\right)$

$= 0.000636\ldots$

$f'(2.5815) = -0.000345\ldots$

There is a change of sign

f′(x) is continuous

therefore 2.581 is a root ✓

---

7. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

(a) Prove that

$$\frac{\sin 2x}{\cos x} + \frac{\cos 2x}{\sin x} = \operatorname{cosec} x \qquad x \neq \frac{n\pi}{2},\ n \in \mathbb{Z}$$

(3)

(b) Hence solve, for $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$,

$$7 + \frac{\sin 4\theta}{\cos 2\theta} + \frac{\cos 4\theta}{\sin 2\theta} = 3\cos^2 2\theta$$

giving your answers in radians to 3 significant figures where appropriate.

(6)

7(a) $\frac{\sin 2x}{\cos x} + \frac{\cos 2x}{\sin x}$ $\quad \frac{\sin 2x}{\cos x} + \frac{\cos 2x}{\sin x} = \operatorname{cosec} x$

$\sin 2x = 2\sin x\cos x$

$\frac{2\sin x\cos x}{\cos x} + \frac{1 - 2\sin^2 x}{\sin x}$ $\quad \cos 2x = \cos^2 x - \sin^2 x$

$= 1 - 2\sin^2 x$

$= 2\cos^2 x - 1$

$2\sin x + \frac{1}{\sin x} - \frac{2\sin^2 x}{\sin x}$

$2\sin x - 2\sin x + \operatorname{cosec} x$

$= RHS$ ✓

(b) $7 + \frac{\sin 4\theta}{\cos 2\theta} + \frac{\cos 4\theta}{\sin 2\theta} = 3\cos^2 2\theta$

$\frac{\sin 2x}{\cos x} + \frac{\cos 2x}{\sin x}$ } compare

So, $\theta = x$ and $x = 2\theta$

Question 7 continued

$\sin^2\theta + \cos^2\theta = 1$

$\sin^2\theta$

$1 + \cos^2\theta = 2\cos^2\theta$

$\cos^2\theta = \cos^2\theta - 1$

$7 + \operatorname{cosec} 2\theta = 3(\operatorname{cosec}^2 2\theta - 1)$

$7 + \operatorname{cosec} 2\theta = 3\operatorname{cosec}^2 2\theta - 3$

$3\operatorname{cosec}^2 2\theta - \operatorname{cosec} 2\theta - 10 = 0$

a b c

$\operatorname{cosec} 2\theta = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$= \frac{-(-1) \pm \sqrt{(-1)^2 - 4(-1)(3)(-10)}}{2(3)}$

$\operatorname{cosec} 2\theta = 2$ or $\operatorname{cosec} 2\theta = -\frac{5}{3}$

$\sin 2\theta = \frac{1}{2}$ or $\sin 2\theta = -\frac{3}{5}$

positive

$2\theta = \sin^{-1}\left(\frac{1}{2}\right)$ $\quad 2\theta = \sin^{-1}\left(-\frac{3}{5}\right)$

$2\theta = \frac{\pi}{6}$ $\quad 2\theta = -0.6435$

S A

T C

S A

T C ✓

Question 7 continued

$2\theta = \frac{\pi}{6}$

$\theta = \frac{\pi}{12} = 2 = \frac{\pi}{12}$ ✓

$= 0.26179\ldots$

$2\theta = \pi - \frac{\pi}{6}$

$\theta = \left(\pi - \frac{\pi}{6}\right) \div 2 = \frac{5\pi}{12}$

$= 1.309\ldots$

CHECK BOUNDARIES $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$

$= -1.57$ $\quad = 1.57$

CHECK REMAINING ANGLE WITHIN RANGE

① $\frac{3}{2} - 1 = \frac{1}{2} + 2.2$

$\frac{1}{2} - 1 = -1.57 - 2.97$

② $1.9106 - 1 = 0.91$ ✓

$= 0.9273$

③ $1.309 - 1 = 0.31$

④ $-1.309 - 1 = -2.31$

$= -2.3562$

$\theta = \frac{\pi}{12}, \frac{5\pi}{12}, -0.322, -1.25$

$\theta = 0.26, 1.31, -0.322, -1.21$ (3sf) ✓

---

8. The percentage, P, of the population of a small country who have access to the internet, is modelled by the equation

$$P = ab^t$$

where a and b are constants and t is the number of years after the start of 2005

Using the data for the years between the start of 2005 and the start of 2010, a graph is plotted of $\log_{10} P$ against t.

The points are found to lie approximately on a straight line with gradient 0.09 and intercept 0.68 on the $\log_{10} P$ axis.

(a) Find, according to the model, the value of a and the value of b, giving your answers to 2 decimal places.

(4)

(b) In the context of the model, give a practical interpretation of the constant a.

(1)

(c) Use the model to estimate the percentage of the population who had access to the internet at the start of 2015

(2)

8a $P = ab^t$ $\quad$ log both sides

$\log P = \log a + t\log b$ $\quad \log b^t = \log a + \log b^t = \log a + t\log b$

$\log b = 0.09$

$\log a = 0.68$

$b = 10^{0.09} = 1.2302\ldots = 1.23$ (2dp pt) ✓

$a = 10^{0.68} = 4.786\ldots = 4.79$ (2dp pt) ✓

$a^x = c$

$\log_a c = b$

$\log a = \log_a a$

Question 8 continued

8 (b) a is where t = 0

$ab^0 = a(1) = a$

∴ a is percentage population with internet access at start ∴ 2005 ✓

(c) $P = 4.79 \times 1.23^t$

in 2015 t = 10 (10 years after start)

$P = 4.79 \times 1.23^{10}$

$P = 37.9652\ldots$

$P = 38$ ✓

(Total 7 marks)

Q8

---

9. Find

(i) $\int \frac{3x-2}{3x^2-4x+5}\,dx$

(2)

(ii) $\int \frac{e^{2x}}{(e^{2x}-1)^3}\,dx \qquad x \neq 0$

(2)

9(i) $\int \frac{3x-2}{3x^2-4x+5}\,dx$

check if $\frac{f'(x)}{f(x)} \times k$

$y = 3x^2 - 4x - 5$

$\frac{dy}{dx} = 6x - 4$

$= 2(3x - 2)$

$k = 2$

then $\ln|fx| \times \frac{1}{k}$

$I = \ln|3x^2 - 4x - 5| \times \frac{1}{2} + c$

$= \frac{1}{2}\ln(3x^2 - 4x + 5) + c$

Question 9 continued

(ii) $\int \frac{e^{2x}}{(e^{2x}-1)^3}$

$= \int e^{2x}(e^{2x}-1)^{-3}$

check if $[f(x)]^n f'(x)$

then $\frac{[f(x)]^{n+1}}{n+1} \times \frac{1}{k}$

$y = e^{2x} - 1$

$\frac{dy}{dx} = 2e^{2x}$ $\quad k = 2$

$I = (e^{2x}-1)^{-2} \times \frac{1}{2} \times \frac{1}{2} + c$

$I = -\frac{(e^{2x}-1)^{-2}}{2 \times 2} + c$

$I = -\frac{1}{4}(e^{2x}-1)^{-2} + c$ ✓

(Total 4 marks)

Q9

---

10. The curve C has equation

$$x = 3\sec^2 2y \qquad x > 3 \qquad 0 < y < \frac{\pi}{4}$$

(a) Find $\frac{dy}{dx}$ in terms of y.

(2)

(b) Hence show that

$$\frac{dy}{dx} = \frac{p}{q\sqrt{x-3}}$$

where p is irrational and q is an integer, stating the values of p and q.

(3)

(c) Find the equation of the normal to C at the point where $y = \frac{\pi}{12}$, giving your answer in the form $y = mx + c$, giving m and c as exact irrational numbers.

(5)

10(a) $x = 3\sec^2 2y$ $\quad$ if $y = \sec\theta$ then $\frac{dy}{dx} = \sec\theta\tan\theta$

$x = 3(\sec 2y)^2$ $\quad$ use chain rule (double)

$\frac{dx}{dy} = 3(\sec 2y)^1 \times \sec 2y \tan 2y \times 2$

$= 12\sec^2 2y\tan 2y$ ✓

(b) $\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$

① find $\frac{dy}{dx}$ in terms of y

② Rearrange / manipulate to get in terms of x

Question 10 continued

$\frac{dy}{dx} = \frac{1}{12\sec^2 2y\tan 2y}$

sub $x = 3\sec^2 2y$

$\therefore \frac{x}{3} = \sec^2 2y$

$\tan 2y = \sqrt{\sec^2 2y - 1}$

$= \sqrt{\frac{x}{3} - 1}$

$\frac{\cos^2\theta + \sin^2\theta = 1}{\cos^2\theta}$

$1 + \tan^2\theta = \sec^2\theta$

$\tan^2\theta = \sec^2\theta - 1$

$\tan\theta = \sqrt{\sec^2\theta - 1}$

$\therefore \frac{dy}{dx} = \frac{1}{12\left(\frac{x}{3}\right)\left(\sqrt{\frac{x}{3}-1}\right)}$

$\frac{dy}{dx} = \frac{1}{4x\sqrt{\frac{x}{3}-1}}$

$\frac{dy}{dx} = \frac{1}{4x\sqrt{\frac{x-3}{3}}}$

$\frac{dy}{dx} = \frac{1}{4x\sqrt{\frac{x-3}{3}}}$

$\frac{dy}{dx} = \frac{\sqrt{3}}{4x\sqrt{x-3}}$ ✓

Question 10 continued

(c) Normal to curve $m_{tangent} \times m_{normal} = -1$

$y = mx + c$

$m_{tangent} = \frac{dy}{dx}$ of C at $y = \frac{\pi}{12}$

① find x when $y = \frac{\pi}{12}$

sub $y = \frac{\pi}{12}$ into c

$x = 3\sec^2 2\left(\frac{\pi}{12}\right)$

$= 3\sec^2\left(\frac{\pi}{6}\right)$

$= 3\left(\frac{1}{\cos\frac{\pi}{6}}\right)^2$

$= 4$

② sub x = 4 into $\frac{dy}{dx}$ for $m_{tangent}$

$m = \frac{\sqrt{3}}{4(4)\sqrt{4-3}}$

$= \frac{\sqrt{3}}{16}$

$\therefore m_{normal} = -\frac{16}{\sqrt{3}}$

③ Sub $\left(4, \frac{\pi}{12}\right)$ into $y = mx + c$ $\quad y - y_1 = m(x - x_1)$

$y - \frac{\pi}{12} = -\frac{16}{\sqrt{3}}(x - 4)$

$y = \left(-\frac{16}{\sqrt{3}} + \frac{16(4)}{\sqrt{3}}\right) + \frac{\pi}{12}$

$y = -\frac{16\sqrt{3}}{3}x + \frac{64\sqrt{3}}{3} + \frac{\pi}{12}$ ✓

(Total 10 marks)

Q10

TOTAL FOR PAPER IS 75 MARKS

END