

TRIGONOMETRY

1

Including

Brief note on trig graphs.

p.2

Brief introduction to the unit circle

p.3

Pythagorus

p.3

Special angles in 90 degree triangles

p.3

Area rule

p.4

Cos rule

p.4

Sin rule

p.4

Reduction formulae & co-ratios

p5

Quadrants in degrees and radians

p5

Identities

- Quotient identity
- Pyth. Identity
- Double angle formulae
- Compound angle formulae

p6


General solution (trig equations)

p7

TRIG. GRAPHS

2

$\sin \theta$




degrees | radians

Convert:

radians to degrees

$$\theta \times \frac{180}{\pi}$$


$\cos \theta$



degrees to radians

$$\theta \times \frac{\pi}{180}$$

$\tan \theta$



shift $\uparrow +$
 $\downarrow -$

General form: $a \cdot \sin bx + q$

reflect
over x-axis

amplitude

max to min

horizontal stretch/squish
intuitive

$a > 1$ longer

$-1 < a < 1$ $a \neq 0$ shorter

vertical
squish/stretch
Counter intuitive

$b > 1$

shorter

$-1 < b < 1$


$b \neq 0$

longer

UNIT CIRCLE

A circle with radius = 1

$$x^2 + y^2 = 1$$




$$\begin{aligned}\cos \theta &= x \\ \sin \theta &= y \\ \tan \theta &= \frac{y}{x}\end{aligned}$$

OTHER USEFUL NOTES:

$$S_H^O C_H^A T_A^O$$



$$S_r^y C_r^x T_x^y$$

Pythagorus



$$a^2 + b^2 = c^2$$

SPECIAL TRIANGLES




ANGLES OF INCLINATION

$$\tan \theta = m$$

AREA RULE

In $\triangle ABC$




$$\text{Area} = \frac{1}{2} \cdot a \cdot b \cdot \sin \hat{C}$$

$$= \frac{1}{2} \cdot a \cdot c \cdot \sin \hat{B}$$

$$= \frac{1}{2} \cdot b \cdot c \cdot \sin \hat{A}$$

Cos rule


In triangle ABC



$$a^2 = b^2 + c^2 - 2(b)(c) \cos \hat{C}$$

Sin rule

In triangle ABC




$$\frac{a}{\sin \hat{A}} = \frac{b}{\sin \hat{B}} = \frac{c}{\sin \hat{C}}$$

REDUCTION FORMULAE ($180^\circ / 360^\circ$)

5

$$\begin{matrix} \sin \\ \cos \\ \tan \end{matrix} \begin{pmatrix} \text{obtuse or} \\ \text{reflex} \\ \text{angle} \end{pmatrix} \Leftrightarrow \begin{matrix} \pm \sin \\ \pm \cos \\ \pm \tan \end{matrix} (\text{acute } <)$$

the CAST diagram shows where each will be positive: degrees or radians



CO-RATIOS (for 90°)

$$\begin{aligned} \sin(90^\circ + \theta) &\Leftrightarrow \cos \theta \\ \sin(90^\circ - \theta) &\Leftrightarrow \cos \theta \\ \cos(90^\circ + \theta) &\Leftrightarrow -\sin \theta \\ \cos(90^\circ - \theta) &\Leftrightarrow \sin \theta \end{aligned}$$

if you remember this one, the others just switch cos/sin

in radians:

$$\begin{aligned} \sin\left(\frac{\pi}{2} + \theta\right) &\Leftrightarrow \cos \theta \\ \sin\left(\frac{\pi}{2} - \theta\right) &\Leftrightarrow \cos \theta \\ \cos\left(\frac{\pi}{2} + \theta\right) &\Leftrightarrow -\sin \theta \\ \cos\left(\frac{\pi}{2} - \theta\right) &\Leftrightarrow \sin \theta \end{aligned}$$

TRIG IDENTITIES

Quotient identity

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

Pyth. Identity

$$\cos^2 \theta + \sin^2 \theta = 1$$

DOUBLE ANGLE FORMULAE

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\begin{aligned} \cos 2\theta &= \cos^2 \theta - \sin^2 \theta \\ &= 2 \cos^2 \theta - 1 \\ &= 1 - 2 \sin^2 \theta \end{aligned}$$

COMPOUND ANGLE FORMULAE

$$\begin{aligned} \sin(A+B) &= \sin A \cos B + \cos A \sin B \\ \sin(A-B) &= \sin A \cos B - \cos A \sin B \end{aligned}$$

$$\begin{aligned} \cos(A+B) &= \cos A \cos B - \sin A \sin B \\ \cos(A-B) &= \cos A \cos B + \sin A \sin B \end{aligned}$$

GENERAL SOLUTION

Gives every possible answer considering that trig graphs are repeating.

- STEPS:**
- ① Determine the positive reference angle
 - ② Apply the reference angle to the relevant quadrants
 - ③ Simplify and don't forget $n \in \mathbb{Z}$

Example 1

given $\tan x \sin x + \cos x \tan x = 0$

Determine the general solution.

Answer:

① $\tan x \sin x + \cos x \tan x = 0$

$\tan x (\sin x + \cos x) = 0$

$\therefore \tan x = 0$

OR

$\sin x + \cos x = 0$

$x = \tan^{-1}(0)$
 $= 0$

$\sin x = -\cos x$

$\frac{\sin x}{\cos x} = -1$

$\tan x = -1$


$x = \tan^{-1}(-1)$

$= (-)\frac{\pi}{4} / (-)45$

\therefore reference angles :

$$x = 0 \text{ or } x = 45 \mid \frac{\pi}{4}$$

② tan is **negative** in quadrant 2 and 4



tan period of 180

③ $\therefore x = 180 - 0 + n180$
 $= 180 + n180 \quad (n \in \mathbb{Z})$
 or $x = 360 - 0 + n180$
 $= 360 + n180 \quad (n \in \mathbb{Z})$

or

$$x = 180 - 45 + n180$$

$$= 135 + n180 \quad (n \in \mathbb{Z})$$

or

$$x = 360 - 45 + n180$$

$$= 315 + n180 \quad (n \in \mathbb{Z})$$