

# Important Notice to Authors

*No further publication processing will occur until we receive your response to this proof.*

Attached is a PDF proof of your forthcoming article in *Physical Review Letters*. The article accession code is LZ15969.

Your paper will be in the following section of the journal: LETTERS — Nuclear Physics

Please note that as part of the production process, APS converts all articles, regardless of their original source, into standardized XML that in turn is used to create the PDF and online versions of the article as well as to populate third-party systems such as Portico, Crossref, and Web of Science. We share our authors' high expectations for the fidelity of the conversion into XML and for the accuracy and appearance of the final, formatted PDF. This process works exceptionally well for the vast majority of articles; however, please check carefully all key elements of your PDF proof, particularly any equations or tables.

Figures submitted electronically as separate files containing color appear in color in the online journal.

However, all figures will appear as grayscale images in the print journal unless the color figure charges have been paid in advance, in accordance with our policy for color in print (<https://journals.aps.org/authors/color-figures-print>).

## Specific Questions and Comments to Address for This Paper

The numbered items below correspond to numbers in the margin of the proof pages pinpointing the source of the question and/or comment. The numbers will be removed from the margins prior to publication.

- 1 PRL prefers to refer to the current article as “Letter.” Please indicate when terms such as “study” and/or “work” can be changed to “Letter” throughout the article or when the terms “study” and “work” refer to the larger body of research behind the Letter.
- 2 Journal style dictates that acronyms not appear in the abstract unless they occur more than once.
- 3 Copyediting changes have been made to your article to conform to APS style and improve readability. Please check to make sure changes have not altered your intended meaning.
- 4 Please supply a definition for “EMC” as it is not on our list of acronyms not needing definition. Journal style dictates that acronyms be defined at their first occurrence in the body of the article.
- 5 Please check the sentence beginning “The previous measurements of DVCS...” to be sure your meaning has not been changed.
- 6 Please note the change to outer square brackets throughout the article to avoid nested fences of the same size and type per PRL style.
- 7 Please check all equations carefully to be sure errors were not introduced during conversion.
- 8 Please supply a definition for “PDF” as it is not on our list of acronyms not needing to be defined and there is more than one possible definition. Journal style dictates that acronyms used only once in the body of the article be spelled out and the acronym deleted.
- 9 Journal style dictates that full words, abbreviations, and acronyms with more than one letter appearing in the inferior or superior positions be set roman.
- 10 PRL prefers italic font for single-letter acronyms and single-letter variables and abbreviations in text and in math.
- 11 Please review the funding information section of the proof’s cover letter and respond as appropriate. We must receive confirmation that the funding agencies have been properly identified before the article can publish.
- 12 This query was generated by an automatic reference checking system. Reference [18] could not be located in the databases used by the system. While the references may be correct, we ask that you check them so we can provide as many links to the referenced articles as possible.

- 13** NOTE: External links, which appear as blue text in the reference section, are created for any reference where a Digital Object Identifier (DOI) can be found. Please confirm that the links created in this PDF proof, which can be checked by clicking on the blue text, direct the reader to the correct references online. If there is an error, correct the information in the reference or supply the correct DOI for the reference. If no correction can be made or the correct DOI cannot be supplied, the link will be removed.
- 14** APS now requires that journal references with authors numbering 10 or fewer display all author names. Please verify author names in expanded author list for Ref. [8].
- 15** Please provide publication information for Ref. [41] if available.

## Titles in References

The editors now encourage insertion of article titles in references to journal articles and e-prints. This format is optional, but if chosen, authors should provide titles for *all* eligible references. If article titles remain missing from eligible references, the production team will remove the existing titles at final proof stage.

## Funding Information

Information about an article's funding sources is now submitted to Crossref to help you comply with current or future funding agency mandates. Crossref's Open Funder Registry (<https://www.crossref.org/services/funder-registry/>) is the definitive registry of funding agencies. Please ensure that your acknowledgments include all sources of funding for your article following any requirements of your funding sources. Where possible, please include grant and award ids. Please carefully check the following funder information we have already extracted from your article and ensure its accuracy and completeness:

- Comisión Nacional de Investigación Científica y Tecnológica, FundRef ID <http://dx.doi.org/10.13039/501100002848> (Republic of Chile/CL)
- Istituto Nazionale di Fisica Nucleare, FundRef ID <http://dx.doi.org/10.13039/501100004007> (Repubblica Italiana/IT)
- Centre National de la Recherche Scientifique, FundRef ID <http://dx.doi.org/10.13039/501100004794> (Republic of France/FR)
- Commissariat à l'Énergie Atomique et aux Énergies Alternatives, FundRef ID <http://dx.doi.org/10.13039/501100006489> (Republic of France/FR)
- U.S. Department of Energy, FundRef ID <http://dx.doi.org/10.13039/100000015> (United States/US)
- Science and Technology Facilities Council, FundRef ID <http://dx.doi.org/10.13039/501100000271> (United Kingdom of Great Britain and Northern Ireland/GB)
- Scottish Universities Physics Alliance, FundRef ID <http://dx.doi.org/10.13039/501100000708> (United Kingdom of Great Britain and Northern Ireland/GB)
- National Research Foundation of Korea, FundRef ID <http://dx.doi.org/10.13039/501100003725> (Republic of Korea/KR)
- Mississippi State University, FundRef ID <http://dx.doi.org/10.13039/100007250> (United States/US)
- Consulat Général de France à Jérusalem
- Thomas Jefferson National Accelerator Facility, FundRef ID <http://dx.doi.org/10.13039/100011665> (United States/US)

## Other Items to Check

- Please note that the original manuscript has been converted to XML prior to the creation of the PDF proof, as described above. Please carefully check all key elements of the paper, particularly the equations and tabular data.
- Title: Please check; be mindful that the title may have been changed during the peer-review process.
- Author list: Please make sure all authors are presented, in the appropriate order, and that all names are spelled correctly.

- Please make sure you have inserted a byline footnote containing the email address for the corresponding author, if desired. Please note that this is not inserted automatically by this journal.
- Affiliations: Please check to be sure the institution names are spelled correctly and attributed to the appropriate author(s).
- Receipt date: Please confirm accuracy.
- Acknowledgments: Please be sure to appropriately acknowledge all funding sources.
- References: Please check to ensure that titles are given as appropriate.
- Hyphenation: Please note hyphens may have been inserted in word pairs that function as adjectives when they occur before a noun, as in “x-ray diffraction,” “4-mm-long gas cell,” and “*R*-matrix theory.” However, hyphens are deleted from word pairs when they are not used as adjectives before nouns, as in “emission by x rays,” “was 4 mm in length,” and “the *R* matrix is tested.”  
Note also that Physical Review follows U.S. English guidelines in that hyphens are not used after prefixes or before suffixes: superresolution, quasiequilibrium, nanoprecipitates, resonancelike, clockwise.
- Please check that your figures are accurate and sized properly. Make sure all labeling is sufficiently legible. Figure quality in this proof is representative of the quality to be used in the online journal. To achieve manageable file size for online delivery, some compression and downsampling of figures may have occurred. Fine details may have become somewhat fuzzy, especially in color figures. The print journal uses files of higher resolution and therefore details may be sharper in print. Figures to be published in color online will appear in color on these proofs if viewed on a color monitor or printed on a color printer.
- Overall, please proofread the entire *formatted* article very carefully. The redlined PDF should be used as a guide to see changes that were made during copyediting. However, note that some changes to math and/or layout may not be indicated.

## Ways to Respond

- **Web:** If you accessed this proof online, follow the instructions on the web page to submit corrections.
- **Email:** Send corrections to [aps-robot@luminad.com](mailto:aps-robot@luminad.com). Include the accession code LZ15969 in the subject line.
- **Fax:** Return this proof with corrections to +1.855.808.3897.

## If You Need to Call Us

You may leave a voicemail message at +1.855.808.3897. Please reference the accession code and the first author of your article in your voicemail message. We will respond to you via email.

## Exploring the Structure of the Bound Proton with Deeply Virtual Compton Scattering

M. Hattawy,<sup>1,22,32</sup> N. A. Baltzell,<sup>1,32,38</sup> R. Dupré,<sup>1,22</sup> S. Bültmann,<sup>32</sup> R. De Vita,<sup>19</sup> A. El Alaoui,<sup>1,39</sup> L. El Fassi,<sup>1,28</sup> H. Egiyan,<sup>38</sup> F. X. Girod,<sup>38</sup> M. Guidal,<sup>22</sup> K. Hafidi,<sup>1</sup> D. Jenkins,<sup>42</sup> S. Liuti,<sup>43</sup> Y. Perrin,<sup>27</sup> S. Stepanyan,<sup>38</sup> B. Torayev,<sup>32</sup> E. Voutier,<sup>22,27</sup> S. Adhikari,<sup>12</sup> Giovanni Angelini,<sup>15</sup> C. Ayerbe Gayoso,<sup>44</sup> L. Barion,<sup>17</sup> M. Battaglieri,<sup>19</sup> I. Bedlinskiy,<sup>23</sup> A. S. Biselli,<sup>10</sup> F. Bossù,<sup>6</sup> W. Brooks,<sup>39</sup> F. Cao,<sup>8</sup> D. S. Carman,<sup>38</sup> A. Celentano,<sup>19</sup> P. Chatagnon,<sup>22</sup> T. Chetry,<sup>31</sup> G. Ciullo,<sup>17,11</sup> L. Clark,<sup>40</sup> P. L. Cole,<sup>26,16</sup> M. Contalbrigo,<sup>17</sup> V. Crede,<sup>13</sup> A. D'Angelo,<sup>20,34</sup> N. Dashyan,<sup>45</sup> E. De Sanctis,<sup>18</sup> M. Defurne,<sup>6</sup> A. Deur,<sup>38</sup> S. Diehl,<sup>8</sup> C. Djalali,<sup>31,36</sup> M. Ehrhart,<sup>22</sup> P. Eugenio,<sup>13</sup> S. Fegan,<sup>40,\*</sup> A. Filippi,<sup>21</sup> T. A. Forest,<sup>16</sup> A. Fradi,<sup>22,†</sup> M. Garçon,<sup>6</sup> G. Gavalian,<sup>38,32</sup> N. Gevorgyan,<sup>45</sup> G. P. Gilfoyle,<sup>33</sup> K. L. Giovanetti,<sup>24</sup> E. Golovatch,<sup>35</sup> R. W. Gothe,<sup>36</sup> K. A. Griffioen,<sup>44</sup> N. Harrison,<sup>38</sup> F. Hauenstein,<sup>32</sup> T. B. Hayward,<sup>44</sup> D. Heddle,<sup>7,38</sup> K. Hicks,<sup>31</sup> M. Holtrop,<sup>29</sup> Y. Ilieva,<sup>36</sup> D. G. Ireland,<sup>40</sup> E. L. Isupov,<sup>35</sup> H. S. Jo,<sup>25</sup> S. Johnston,<sup>1</sup> D. Keller,<sup>43,31</sup> G. Khachatryan,<sup>45</sup> M. Khachatryan,<sup>32</sup> A. Khanal,<sup>12</sup> M. Khandaker,<sup>30,‡</sup> C. W. Kim,<sup>15</sup> W. Kim,<sup>25</sup> F. J. Klein,<sup>5</sup> V. Kubarovsky,<sup>38</sup> S. E. Kuhn,<sup>32</sup> L. Lanza,<sup>20</sup> M. L. Kabir,<sup>28</sup> P. Lenisa,<sup>17</sup> K. Livingston,<sup>40</sup> I. J. D. MacGregor,<sup>40</sup> D. Marchand,<sup>22</sup> N. Markov,<sup>8</sup> M. Mayer,<sup>32</sup> B. McKinnon,<sup>40</sup> Z. E. Meziani,<sup>37</sup> T. Mineeva,<sup>39</sup> M. Mirazita,<sup>18</sup> R. A. Montgomery,<sup>40</sup> C. Munoz Camacho,<sup>22</sup> P. Nadel-Turonski,<sup>38,5</sup> S. Niccolai,<sup>22</sup> A. I. Ostrovidov,<sup>13</sup> L. L. Pappalardo,<sup>17</sup> R. Paremuzyan,<sup>29,45</sup> E. Pasyuk,<sup>38,2</sup> O. Pogorelko,<sup>23</sup> J. Poudel,<sup>32</sup> Y. Prok,<sup>32,43</sup> D. Protopopescu,<sup>40</sup> M. Ripani,<sup>19</sup> D. Riser,<sup>8</sup> A. Rizzo,<sup>20,34</sup> G. Rosner,<sup>40</sup> P. Rossi,<sup>38,18</sup> F. Sabatié,<sup>6</sup> C. Salgado,<sup>30</sup> R. A. Schumacher,<sup>4</sup> Y. G. Sharabian,<sup>38</sup> Iu. Skorodumina,<sup>36,35</sup> D. Sokhan,<sup>40</sup> O. Soto,<sup>39</sup> N. Sparveris,<sup>37</sup> S. Strauch,<sup>36</sup> M. Taiuti,<sup>14,§</sup> J. A. Tan,<sup>25</sup> N. Tyler,<sup>36</sup> M. Ungaro,<sup>38,8</sup> H. Voskanyan,<sup>45</sup> R. Wang,<sup>22</sup> D. P. Watts,<sup>41</sup> X. Wei,<sup>38</sup> L. B. Weinstein,<sup>32</sup> M. H. Wood,<sup>3</sup> N. Zachariou,<sup>41</sup> J. Zhang,<sup>43,32</sup> and Z. W. Zhao<sup>9,36</sup>

(CLAS Collaboration)

<sup>1</sup>Argonne National Laboratory, Argonne, Illinois 60439, USA<sup>2</sup>Arizona State University, Tempe, Arizona 85287-1504, USA<sup>3</sup>Canisius College, Buffalo, New York, USA<sup>4</sup>Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA<sup>5</sup>Catholic University of America, Washington, DC 20064, USA<sup>6</sup>IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France<sup>7</sup>Christopher Newport University, Newport News, Virginia 23606, USA<sup>8</sup>University of Connecticut, Storrs, Connecticut 06269, USA<sup>9</sup>Duke University, Durham, North Carolina 27708-0305, USA<sup>10</sup>Fairfield University, Fairfield Connecticut 06824, USA<sup>11</sup>Università di Ferrara, 44121 Ferrara, Italy<sup>12</sup>Florida International University, Miami, Florida 33199, USA<sup>13</sup>Florida State University, Tallahassee, Florida 32306, USA<sup>14</sup>Università di Genova, 16146 Genova, Italy<sup>15</sup>The George Washington University, Washington, DC 20052, USA<sup>16</sup>Idaho State University, Pocatello, Idaho 83209, USA<sup>17</sup>INFN, Sezione di Ferrara, 44100 Ferrara, Italy<sup>18</sup>INFN, Laboratori Nazionali di Frascati, 00044 Frascati, Italy<sup>19</sup>INFN, Sezione di Genova, 16146 Genova, Italy<sup>20</sup>INFN, Sezione di Roma Tor Vergata, 00133 Rome, Italy<sup>21</sup>INFN, Sezione di Torino, 10125 Torino, Italy<sup>22</sup>Institut de Physique Nucléaire, IN2P3-CNRS, Université Paris-Sud, Université Paris-Saclay, F-91406 Orsay, France<sup>23</sup>Institute of Theoretical and Experimental Physics, Moscow, 117259, Russia<sup>24</sup>James Madison University, Harrisonburg, Virginia 22807, USA<sup>25</sup>Kyungpook National University, Daegu 41566, Republic of Korea<sup>26</sup>Lamar University, 4400 MLK Boulevard, P.O. Box 10009, Beaumont, Texas 77710, USA<sup>27</sup>LPSC, Université Grenoble-Alpes, CNRS/IN2P3, 38026 Grenoble, France<sup>28</sup>Mississippi State University, Mississippi State, Mississippi 39762-5167, USA<sup>29</sup>University of New Hampshire, Durham, New Hampshire 03824-3568, USA<sup>30</sup>Norfolk State University, Norfolk, Virginia 23504, USA<sup>31</sup>Ohio University, Athens, Ohio 45701, USA<sup>32</sup>Old Dominion University, Norfolk, Virginia 23529, USA<sup>33</sup>University of Richmond, Richmond, Virginia 23173, USA

<sup>34</sup>*Universita' di Roma Tor Vergata, 00133 Rome, Italy*

<sup>35</sup>*Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, 119234 Moscow, Russia*

<sup>36</sup>*University of South Carolina, Columbia, South Carolina 29208, USA*

<sup>37</sup>*Temple University, Philadelphia, Pennsylvania 19122, USA*

<sup>38</sup>*Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA*

<sup>39</sup>*Universidad Técnica Federico Santa María, Casilla 110-V Valparaíso, Chile*

<sup>40</sup>*University of Glasgow, Glasgow G12 8QQ, United Kingdom*

<sup>41</sup>*University of York, York YO10 5DD, United Kingdom*

<sup>42</sup>*Virginia Tech, Blacksburg, Virginia 24061-0435, USA*

<sup>43</sup>*University of Virginia, Charlottesville, Virginia 22901, USA*

<sup>44</sup>*College of William and Mary, Williamsburg, Virginia 23187-8795, USA*

<sup>45</sup>*Yerevan Physics Institute, 375036 Yerevan, Armenia*

(Received 19 December 2018; revised manuscript received 12 March 2019)

In the past two decades, deeply virtual Compton scattering of electrons has been successfully used to advance our knowledge of the partonic structure of the free proton and investigate correlations between the transverse position and the longitudinal momentum of quarks inside the nucleon. Meanwhile, the structure of bound nucleons in nuclei has been studied in inclusive deep-inelastic lepton scattering experiments off nuclear targets, showing a significant difference in longitudinal momentum distribution of quarks inside the bound nucleon, known as the EMC effect. In this work, we report the first beam spin asymmetry (BSA) measurement of exclusive deeply virtual Compton scattering off a proton bound in  $^4\text{He}$ . The data used here were accumulated using a 6 GeV longitudinally polarized electron beam incident on a pressurized  $^4\text{He}$  gaseous target placed within the CLAS spectrometer in Hall-B at the Thomas Jefferson National Accelerator Facility. The azimuthal angle ( $\phi$ ) dependence of the BSA was studied in a wide range of virtual photon and scattered proton kinematics. The  $Q^2$ ,  $x_B$ , and  $t$  dependencies of the BSA on the bound proton are compared with those on the free proton. In the whole kinematical region of our measurements, the BSA on the bound proton is smaller by 20% to 40%, indicating possible medium modification of its partonic structure.

DOI:

**3** Electromagnetic probes have played a major role in advancing our knowledge about the structure of the nucleon. While lepton-nucleon elastic scattering measurements have taught us about the spatial charge and magnetization distributions [1,2], deep-inelastic scattering experiments have uncovered the partonic structure of the nucleon and the longitudinal momentum distributions of the constituent partons, i.e., quarks and gluons [3]. With nuclear targets, deeply inelastic lepton scattering measurements have revealed that the distribution of quarks in a nucleus is not a simple convolution of their distributions within nucleons, an observation known as the “EMC effect” [4] (for reviews on the topic, see [5–8]).

A wealth of information on the structure of hadrons lies in the correlations between the momentum and spatial degrees of freedom of the partons. These correlations can be revealed through deeply virtual Compton scattering (DVCS), i.e., the hard exclusive lepton production of a real photon, which provides access to a three-dimensional (3D) imaging of partons within the generalized parton distributions (GPDs) framework [9–13]. The measurement of free proton DVCS has been the focus of a worldwide effort [14–26] involving several accelerator facilities such as Jefferson Lab, DESY, and CERN. These measurements

now enable the extractions of GPDs and a 3D tomography of the free proton [27,28]. New measurements of DVCS from the  $^4\text{He}$  nucleus are a critical step towards providing a similar 3D picture of the quark structure of the nucleus [29]. In the nuclear case, however, two channels are available, the coherent channel where the scattering is off the entire nucleus, which is left intact in the final state [30], and the incoherent channel where the DVCS occurs on a nucleon, which is ejected from the nucleus. The latter is the focus of this Letter and provides a unique access to the modification of the partonic structure of the bound nucleons [31–33]. The  $^4\text{He}$  nucleus is an ideal experimental target for this measurement as it is characterized by a strong binding energy, a relatively high nuclear core density, and a large EMC effect [34]. Moreover, it remains simple enough that precise calculation of its structure can be performed, making this nucleus the perfect target for our investigation of the medium modifications of the nucleon’s partonic structure. The previous measurements of DVCS off nuclei, and in particular off  $^4\text{He}$ , performed at HERMES [35], yielded results with both “coherent enriched” and “incoherent enriched” event samples; hence they are not fully exclusive, but significant enough to be compared with our results below.






FIG. 1. Representation of the leading-order, twist-2, handbag diagram of the incoherent DVCS process off  $^4\text{He}$ , where the four vectors of the electrons, photons, and protons are denoted by  $k/k'$ ,  $q/q'$ , and  $p/p'$ , respectively.  $x + \xi$  is the nucleon longitudinal momentum fraction carried by the struck quark,  $-2\xi$  is the longitudinal momentum fraction of the momentum transfer  $\Delta (= q - q')$ , and  $t (= \Delta^2)$  is the squared momentum transfer between the initial and the final state nucleon.

In this Letter, we present the first exclusive measurement of the beam-spin asymmetry (BSA) in deeply virtual electroproduction of a real photon off a bound proton in  $^4\text{He}$ . Figure 1 illustrates the leading-twist handbag diagram for the DVCS process. In the Bjorken regime, i.e., large virtual photon four-momentum squared [ $Q^2 = -q^2 = -(k - k')^2$ ], and at small invariant momentum transfer [ $t = (q - q')^2$ ], the DVCS scattering process can be factorized, leaving the nonperturbative structure of the nucleon to be parametrized in terms of four chirally even GPDs:  $H$ ,  $E$ ,  $\tilde{H}$ , and  $\tilde{E}$ , representing the four helicity-spin combinations of the quark-nucleon states [36,37]. Experimentally, we measure the squared sum of the Bethe-Heitler (BH) and the DVCS amplitudes. The BH process, where the real photon is emitted by the incident or the scattered electron rather than the nucleon, dominates the cross section at our kinematics. The BSA arises from the interference of these two terms and is directly sensitive to the DVCS amplitude that contains the information on the GPDs. Using a longitudinally polarized electron beam ( $L$ ) and an unpolarized target ( $U$ ), the BSA is defined as

$$A_{LU} = \frac{d^5\sigma^+ - d^5\sigma^-}{d^5\sigma^+ + d^5\sigma^-}, \quad (1)$$

where  $d^5\sigma^+ (d^5\sigma^-)$  is the virtual photoproduction differential cross section for a positive (negative) beam helicity. Following the cross section decomposition provided in [38], the different components can be expressed in terms of Fourier coefficients associated with  $\phi$  harmonics, where  $\phi$  is the angle between the leptonic and the hadronic planes of the reaction. At leading twist, the BSA can be parametrized as

$$A_{LU}(\phi) = \frac{a_0 \sin(\phi)}{1 + a_1 \cos(\phi) + a_2 \cos(2\phi)}, \quad (2)$$

where the parameters  $a_{0,1,2}$  are combinations of the aforementioned Fourier coefficients. The  $\sin(\phi)$  harmonic is dominant in  $A_{LU}$  and is proportional to the following combination of Compton form factors (CFFs)  $\mathcal{H}$ ,  $\mathcal{E}$ , and  $\tilde{\mathcal{H}}$  as [27]

$$a_0 \propto \text{Im} \left[ F_1 \mathcal{H} - \frac{t}{4M^2} F_2 \mathcal{E} + \frac{x_B}{2} (F_1 + F_2) \tilde{\mathcal{H}} \right], \quad (3)$$

where  $F_1$  and  $F_2$  are the Dirac and Pauli form factors, respectively, and  $x_B$  the Bjorken scaling variable. The real and the imaginary parts of the CFF  $\mathcal{H}$  relate to the GPD  $H$  as

$$\Re(\mathcal{H}) = \mathcal{P} \int_0^1 dx [H(x, \xi, t) - H(-x, \xi, t)] C^+(x, \xi), \quad (4)$$

$$\Im(\mathcal{H}) = -\pi [H(\xi, \xi, t) - H(-\xi, \xi, t)], \quad (5)$$

with  $\mathcal{P}$  the Cauchy principal value integral and  $C^+$  a coefficient function defined as  $[1/(x - \xi) + 1/(x + \xi)]$ , where  $\xi$  is the skewing factor and can be related to  $x_B$  by  $\xi \approx (x_B/2 - x_B)$ . Similar expressions apply for the GPDs  $E$ ,  $\tilde{H}$ , and  $\tilde{E}$  [27]. At the forward limit,  $\xi \rightarrow 0$  and  $t \rightarrow 0$ , the GPD  $H$  reduces to quark, antiquark PDFs, and its zeroth moment in  $x$  represents the elastic Dirac form factor  $F_1$ .

The experiment (E08-024 [39]) took place in Hall-B of Jefferson Lab using the nearly 100% duty factor, longitudinally polarized electron beam (83% polarization) from the Continuous Electron Beam Accelerator Facility (CEBAF) at an energy of 6.064 GeV. The data were accumulated over 40 days using a 6-atm-pressure, 292-mm-long, and 6-mm-diameter gaseous  $^4\text{He}$  target centered 64 cm upstream of the CEBAF Large Acceptance Spectrometer (CLAS) coordinate center. For DVCS experiments, the CLAS baseline design [40] was supplemented with an inner calorimeter (IC) and a solenoid magnet. The IC extended the photon detection acceptance of CLAS down to a polar angle of  $4^\circ$ . The 5-Tesla solenoid magnet, in the center of which the target was located, prevented the high-rate low-energy Møller electrons from reaching the CLAS drift chambers by guiding these electrons inside a tungsten shield placed around the beam line.

Incoherent DVCS events were selected by requiring an electron, a proton, and at least one photon in the final state using the standard particle identification framework of the CLAS event reconstruction (see [41] for additional details on the particle identification). Note that, even though the DVCS reaction has only one real photon in the final state, events with more than one photon were not discarded at this stage. These extra photons were

mostly soft photons from accidental coincidence which, as will be discussed below, the DVCS exclusivity cuts easily eliminated. In the following stage, the most energetic photon was considered as the DVCS photon candidate.

Further requirements were applied to clean the identified initial set of incoherent DVCS events from accidental and physics background events. First, events were selected with  $Q^2$  greater than 1 GeV<sup>2</sup> and the  $\gamma^*p$  invariant mass [ $W = \sqrt{(q+p)^2}$ , assuming that the initial nucleon is at rest] greater than 2 GeV. This is a commonly accepted region of kinematics used by the previous DVCS experiments and avoids the nucleon resonance region. The squared transferred momentum to the recoil proton  $t$ , calculated from the four-momentum vectors of the incoming and outgoing photons, was required to be greater than a minimum kinematically allowed value ( $t_{\min}$ ) at given  $Q^2$  and  $W$  defined as

$$t_{\min} = -Q^2 \frac{2(1-x_B)(1-\sqrt{1+\epsilon^2}) + \epsilon^2}{4x_B(1-x_B) + \epsilon^2}, \quad (6)$$

where  $\epsilon^2 = (4M_p^2 x_B^2 / Q^2)$  and  $M_p$  is the proton mass. This cut was applied to avoid accepting events that appear in unphysical regions of kinematics due to detector resolution and radiative effects. We specifically use the kinematics of the photons to determine  $t$  because the initial proton kinematics is unknown due to Fermi motion.

In the final sample, the exclusivity of the incoherent DVCS events was ensured by imposing a series of constraints based on the four-momentum conservation in the reaction  $ep \rightarrow e'p'\gamma$ . These kinematical variables are the coplanarity angle  $\Delta\phi$  between the  $(\gamma, \gamma^*)$  and  $(\gamma^*, p')$  planes, the missing energy, mass, and transverse momentum of the  $e'\gamma$  and  $e'p'\gamma$  systems, the missing mass squared of the  $e'p'$  system, and the angle  $\theta$  between the measured photon and the missing momentum of the  $e'p'$  system. The experimental distributions for the most relevant exclusivity variables are shown in Fig. 2. Because of the Fermi motion of the nucleons in the helium nucleus, the cuts indicated by the dashed lines are slightly wider than those previously used for free proton experiments [21]. After the corrections discussed below, the asymmetries appear to be stable as a function of cut width, and we saw no sizable effect that could be related to the initial momentum of the nucleons. We also rejected events where a  $\pi^0$  was identified by the invariant mass of two photons. At the end of this selection process, about 30k events passed all the requirements.

The two main backgrounds that contributed to the event sample after the exclusivity cuts are due to accidental coincidences and exclusive  $\pi^0$  production, where one of the photons from the  $\pi^0$  decay escapes detection. The contribution from accidental events, i.e.,  $e'p'\gamma$  collections with particles originating from different electron scatterings, was




FIG. 2. The distributions from left to right and from top to bottom are  $\Delta\phi$ , missing energy, missing mass squared, and the cone angle ( $\theta$ ) between the measured and the calculated photons in the  $e'p'$  final-state system. The incoherent DVCS exclusivity cuts are represented by the vertical red-dashed lines. The black distributions represent the incoherent DVCS event candidates before the exclusivity cuts. The shaded distributions represent the incoherent DVCS events that passed all of these cuts except the quantity plotted.

evaluated to be 6.5% by selecting events passing all our selection cuts but originating from different vertices. The  $\pi^0$  contamination was estimated and subtracted using detector simulation and experimental data. From simulation, we calculated the ratio ( $R = N_{\text{sim}}^{1\gamma} / N_{\text{sim}}^{2\gamma}$ ) of the number of  $\pi^0$  events that were wrongly identified as exclusive  $ep \rightarrow e'p'\gamma$  events ( $N_{\text{sim}}^{1\gamma}$ ) to the number of events correctly identified as exclusive  $ep \rightarrow e'p'\pi^0$  ( $N_{\text{sim}}^{2\gamma}$ ). Then, in each kinematical bin and for each beam-helicity state, the  $\pi^0$ -subtracted experimental DVCS events were calculated as  $N = N_{\text{exp}}^{ep \rightarrow e'p'\gamma} - RN_{\text{exp}}^{ep \rightarrow e'p'\pi^0}$ , where  $N_{\text{exp}}^{ep \rightarrow e'p'\gamma}$  ( $N_{\text{exp}}^{ep \rightarrow e'p'\pi^0}$ ) is the number of the experimentally identified  $ep \rightarrow e'p'\gamma$  ( $ep \rightarrow e'p'\pi^0$ ) events. Depending on the kinematics, we subtracted between 8% and 10% of the data due to the  $\pi^0$  contamination.

Experimentally,  $A_{LU}$  is defined as

$$A_{LU} = \frac{1}{P_B(1-C)} \frac{N^+ - N^-}{N^+ + N^-}, \quad (7)$$

where  $N^+$  and  $N^-$  are the number of DVCS events for the positive and negative beam-helicity states,  $P_B$  is the longitudinal beam polarization, and  $C$  stands for the contamination percentage of the accidental coincidences.

In the kinematical phase space of our experiment, the  $\phi$  dependence of  $A_{LU}$  is most sensitive to the imaginary part




FIG. 3. The incoherent  $A_{LU}$  as a function of  $\phi$  for different  $t$  bins. The error bars represent the statistical uncertainties. The gray bands represent the systematic uncertainties, including the normalization uncertainties. The black curves are the results of our fits with the form  $[a_0 \sin(\phi)/1 + a_1 \cos(\phi)]$ .

of the CFFs through the  $a_0$  term of Eq. (2), as confirmed by high statistics measurements on the free proton [21,26]. In the determination of  $a_0$  in Eq. (3), the CFF  $\mathcal{E}$  and  $\tilde{\mathcal{H}}$  are suppressed due to form factors and the smallness of the coefficients. Therefore, the dominant contribution to the BSA comes from the CFF  $\mathcal{H}$  and, hence, the GPD  $H$ . Because of limited statistics, the data were binned two dimensionally into 36 bins. That is, four bins in one of the kinematical variables of interest ( $Q^2$ ,  $x_B$ , or  $t$ ) and then nine bins in the azimuthal angle ( $\phi$ ). Figure 3 presents the measured incoherent  $A_{LU}$  as a function of  $\phi$  in bins of  $t$  (integrated over the full  $Q^2$  and  $x_B$  ranges). The curves on the plots are fits of the form  $[a_0 \sin(\phi)/1 + a_1 \cos(\phi)]$ . The main contributions to systematic uncertainties on these fits are from the choice of the DVCS exclusivity cuts (6%) and the large bin size (7%). The systematic uncertainties sum up to less than 10% for all data points and, thus, always remain significantly smaller than the statistical uncertainties.

Figure 4 presents the dependence of the fitted  $A_{LU}$  values at  $\phi = 90^\circ$  ( $a_0$  parameter from the individual fits in Fig. 3) on the kinematical variables  $Q^2$ ,  $x_B$ , and  $t$ . Within the given uncertainties,  $A_{LU}$  does not show a strong dependence on  $Q^2$ . The  $x_B$  and  $t$  dependencies are compared to the theoretical calculations performed by

S. Liuti and K. Taneja [31]. Their model uses a nuclear spectral function and mainly considers the effect of the nucleon off shellness. The calculations are carried out at slightly different kinematics than our data but still provide important guidance. The experimental results appear to have smaller asymmetries especially at small  $x_B$  than the calculations. These differences may arise from nuclear effects that are not taken into account in the model, such as long-range interactions and final state interactions of the knocked-out proton. On the graph for the  $-t$  dependence, we show previous measurements by HERMES Collaboration [35], in which only electrons and photons were measured. Because of the large experimental uncertainties of the HERMES points, the two measurements are completely compatible.

One can use the nuclear DVCS to measure a “generalized” EMC effect in order to see if significant nuclear effects are also visible within the GPD framework. To explore this idea, we constructed the ratio of  $A_{LU}$  for bound protons to that on a free proton target. Figure 5 presents the BSA ratio based on interpolation of the free proton asymmetries from CLAS [21] as a function of the kinematical variable  $t$ . The  $A_{LU}$  ratios show 25%–40% lower asymmetries that are independent of  $t$  for a bound proton compared to the free proton. The measurements disagree




FIG. 4. The  $Q^2$  (left),  $x_B$  (middle), and  $t$  dependencies (right) of the fitted  $A_{LU}$  at  $\phi = 90^\circ$  (black squares). The error bars represent the statistical uncertainties, while the gray bands represent the systematic uncertainties. On the middle plot, the curves are theoretical calculations from [31]. On the right plot, the solid (empty) green circles are the HERMES –  $A_{LU}$  (a positron beam was used) inclusive measurements for the incoherent (coherent) enriched region [35]; the curves represent theoretical calculations from [31].






FIG. 5. The  $A_{LU}$  ratio of the bound to the free proton at  $\phi = 90^\circ$  as a function of  $t$ . The black squares are from this work, the green circle is the HERMES measurement [35]. The error bars represent the statistical uncertainties, while the gray band represents the systematic uncertainties. The blue and red curves are results of off-shell calculations [31]. The solid and dashed black curves are from on-shell calculations [33].

with the off-shell [31] and the on-shell calculations that use the medium-modified GPDs as calculated from the quark-meson coupling model [33]. Our results show that an important nuclear effect is missing from the existing models in order to explain this strong quenching of the BSA. More theoretical developments will be needed to identify the origin of this quenching, in particular, it will be important to differentiate initial from final state effects and how they affect the DVCS asymmetries.

In summary, we have presented the first BSA measurement associated with bound proton DVCS off  $^4\text{He}$  using an upgraded setup of the CLAS spectrometer at Jefferson Lab. Our results are compared to model calculations based on different assumptions of the nuclear medium effects at the partonic level. The bound-proton BSA is largely suppressed compared to the free proton BSA. This result is a first step in using a novel experimental method of understanding the properties of bound nucleons directly from the basic degrees of freedom of QCD, quarks, and gluons. Planned experiments at Jefferson Lab will continue and extend these studies of the bound nucleon structure using DVCS. We have an experimental program called ALERT using the CLAS12 detector in Hall-B of Jefferson Lab. These experiments will improve the DVCS measurements with the detection of nuclear fragments to better control the final state interactions and the initial state kinematics of the bound nucleon.

The authors acknowledge the staff of the Accelerator and Physics Divisions at the Thomas Jefferson National Accelerator Facility who made this experiment possible. This work was supported in part by the Chilean Comisión Nacional de Investigación Científica y Tecnológica (CONICYT), by CONICYT PIA Grant No. ACT1413, the Italian Istituto Nazionale di Fisica Nucleare, the French

Centre National de la Recherche Scientifique, the French Commissariat à l'Énergie Atomique, the U.S. Department of Energy under Contract No. DE-AC02-06CH11357, the United Kingdom Science and Technology Facilities Council (STFC), the Scottish Universities Physics Alliance (SUPA), the National Research Foundation of Korea, and the Office of Research and Economic Development at Mississippi State University. M. H. also acknowledges the support of the Consulat Général de France à Jérusalem. The Southeastern Universities Research Association operates the Thomas Jefferson National Accelerator Facility for the United States Department of Energy under Contract No. DE-AC05-06OR23177.

\*Present address: The George Washington University, Washington, DC 20052, USA.

†Present address: Imam Abdulrahman Bin Faisal University, Industrial Jubail 31961, Saudi Arabia.

‡Present address: Idaho State University, Pocatello, Idaho 83209, USA.

§Present address: INFN, Sezione di Genova, 16146 Genova, Italy.

- [1] R. Hofstadter and R. W. McAllister, *Phys. Rev.* **98**, 217 (1955).
- [2] C. F. Perdrisat, V. Punjabi, and M. Vanderhaeghen, *Prog. Part. Nucl. Phys.* **59**, 694 (2007).
- [3] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
- [4] J. J. Aubert *et al.*, *Phys. Lett.* **123B**, 275 (1983).
- [5] M. Arneodo, *Phys. Rep.* **240**, 301 (1994).
- [6] D. F. Geesaman, K. Saito, and A. W. Thomas, *Annu. Rev. Nucl. Part. Sci.* **45**, 337 (1995).
- [7] P. R. Norton, *Rep. Prog. Phys.* **66**, 1253 (2003).
- [8] O. Hen, G. A. Miller, E. Piasetzky, and L. B. Weinstein, *Rev. Mod. Phys.* **89**, 045002 (2017).
- [9] D. Mueller, D. Robaschik, B. Geyer, F. M. Dittes, and J. Horejsi, *Fortschr. Phys.* **42**, 101 (1994).
- [10] X. D. Ji, *Phys. Rev. Lett.* **78**, 610 (1997).
- [11] X. D. Ji, *Phys. Rev. D* **55**, 7114 (1997).
- [12] A. V. Radyushkin, *Phys. Lett. B* **380**, 417 (1996).
- [13] A. V. Radyushkin, *Phys. Rev. D* **56**, 5524 (1997).
- [14] S. Stepanyan *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **87**, 182002 (2001).
- [15] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. Lett.* **87**, 182001 (2001).
- [16] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. D* **75**, 011103 (2007).
- [17] S. Chekanov *et al.* (ZEUS Collaboration), *Phys. Lett. B* **573**, 46 (2003).
- [18] A. Aktas *et al.* (H1 Collaboration), *Eur. Phys. J. C* **44**, 1 (2005).
- [19] S. Chen *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **97**, 072002 (2006).
- [20] M. Defurne *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. C* **92**, 055202 (2015).
- [21] F. X. Girod *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **100**, 162002 (2008).

---

425	[22] M. Mazouz <i>et al.</i> (Jefferson Lab Hall A Collaboration),	[32] V. Guzey and T. Teckentrup, <i>Phys. Rev. D</i> <b>74</b> , 054027 (2006).	444
426	<i>Phys. Rev. Lett.</i> <b>99</b> , 242501 (2007).	[33] V. Guzey, A. W. Thomas, and K. Tsushima, <i>Phys. Lett. B</i>	445
427	[23] G. Gavalian <i>et al.</i> (CLAS Collaboration), <i>Phys. Rev. C</i> <b>80</b> ,	<b>673</b> , 9 (2009).	446
428	035206 (2009).	[34] J. Seely <i>et al.</i> , <i>Phys. Rev. Lett.</i> <b>103</b> , 202301 (2009).	447
429	[24] E. Seder <i>et al.</i> (CLAS Collaboration), <i>Phys. Rev. Lett.</i> <b>114</b> ,	[35] A. Airapetian <i>et al.</i> (HERMES Collaboration), <i>Phys. Rev. C</i>	448
430	032001 (2015).	<b>81</b> , 035202 (2010).	449
431	[25] S. Pisano <i>et al.</i> (CLAS Collaboration), <i>Phys. Rev. D</i> <b>91</b> ,	[36] J. C. Collins and A. Freund, <i>Phys. Rev. D</i> <b>59</b> , 074009	450
432	052014 (2015).	(1999).	451
433	[26] H. S. Jo <i>et al.</i> (CLAS Collaboration), <i>Phys. Rev. Lett.</i> <b>115</b> ,	[37] X.-D. Ji and J. Osborne, <i>Phys. Rev. D</i> <b>58</b> , 094018 (1998).	452
434	212003 (2015).	[38] A. V. Belitsky, D. Mueller, and A. Kirchner, <i>Nucl. Phys.</i>	453
435	[27] M. Guidal, H. Moutarde, and M. Vanderhaeghen, <i>Rep. Prog.</i>	<b>B629</b> , 323 (2002); Updated in: A. V. Belitsky and D.	454
436	<i>Phys.</i> <b>76</b> , 066202 (2013).	Mueller, <i>Phys. Rev. D</i> <b>82</b> , 074010 (2010).	455
437	[28] R. Dupré, M. Guidal, and M. Vanderhaeghen, <i>Phys. Rev. D</i>	[39] K. Hafidi <i>et al.</i> , proposal PR-08-024 to JLab PAC33, <a href="https://www.jlab.org/exp_prog/proposals/08/PR-08-024.pdf">https://</a>	456
438	<b>95</b> , 011501(R) (2017).	<a href="https://www.jlab.org/exp_prog/proposals/08/PR-08-024.pdf">www.jlab.org/exp_prog/proposals/08/PR-08-024.pdf</a> .	457
439	[29] R. Dupré and S. Scopetta, <i>Eur. Phys. J. A</i> <b>52</b> , 159 (2016).	[40] B. A. Mecking <i>et al.</i> , <i>Nucl. Instrum. Methods Phys. Res.,</i>	458
440	[30] M. Hattawy <i>et al.</i> (CLAS Collaboration), <i>Phys. Rev. Lett.</i>	<b>Sect. A</b> <b>503</b> , 513 (2003).	459
441	<b>119</b> , 202004 (2017).	[41] M. Hattawy, Ph. D. thesis, Université Paris Sud—Paris	460
442	[31] S. Liuti and S. K. Taneja, <i>Phys. Rev. C</i> <b>72</b> , 032201(R)	XI [Institution Report No. 2015PA112161, 2015 (to be	461
443	(2005).	published)].	462
			463