

Important Notice to Authors

Attached is a PDF proof of your forthcoming article in *Physical Review Letters*. The article accession code is LG16705. Your paper will be in the following section of the journal: LETTERS — Elementary Particles and Fields

Please note that as part of the production process, APS converts all articles, regardless of their original source, into standardized XML that in turn is used to create the PDF and online versions of the article as well as to populate third-party systems such as Portico, Crossref, and Web of Science. We share our authors' high expectations for the fidelity of the conversion into XML and for the accuracy and appearance of the final, formatted PDF. This process works exceptionally well for the vast majority of articles; however, please check carefully all key elements of your PDF proof, particularly any equations or tables.

Figures submitted electronically as separate PostScript files containing color appear in color in the online journal. However, all figures will appear as grayscale images in the print journal unless the color figure charges have been paid in advance, in accordance with our policy for color in print (<http://journals.aps.org/authors/color-figures-print>).

No further publication processing will occur until we receive your response to this proof.

Specific Questions and Comments to Address for This Paper

The numbered items below correspond to numbers in the margin of the proof pages pinpointing the source of the question and/or comment. The numbers will be removed from the margins prior to publication.

- 1 Please check that the author list is correct.
- 2 We have defined CEBAF as Continuous Electron Beam Accelerator Facility. Is this correct?
- 3 Please provide postal code for author affiliation 22 and 26.
- 4 Please note the change in Eq. (2) to outer square brackets to avoid nested fences of the same size and type per journal style.
- 5 Text footnotes are prohibited and have been incorporated into the text as parenthetical remarks. Please check.
- 6 Please note the change in Eq. (5) to outer square brackets to avoid nested fences of the same size and type per journal style.
- 7 The center dot in the last equation in Eq. (6) has been removed. Please see <http://journals.aps.org/authors/multiplication-signs-h11> for more information. Also, please note the change in the third equation to square brackets to avoid the use of nested fences.
- 8 We have defined VGG as Vanderhaeghen-Guichon-Guidal. Is this correct?
- 9 Please review the funding information section of the proof's cover letter and respond as appropriate. We must receive confirmation that the funding agencies have been properly identified before the article can publish.
- 10 This query was generated by an automatic reference checking system. References [11, 15, 30] could not be located in the databases used by the system. While the references may be correct, we ask that you check them so we can provide as many links to the referenced articles as possible.
- 11 NOTE: External links, which appear as blue text in the reference section, are created for any reference where a Digital Object Identifier (DOI) can be found. Please confirm that the links created in this PDF proof, which can be checked by clicking on the blue text, direct the reader to the correct references online. If there is an error, correct the information in the reference or supply the correct DOI for the reference. If no correction can be made or the correct DOI cannot be supplied, the link will be removed.
- 12 We have updated Ref. [36]. Please check.

Titles in References

The editors now encourage insertion of article titles in references to journal articles and e-prints. This format is optional, but if chosen, authors should provide titles for *all* eligible references. If article titles remain missing from eligible references, the

production team will remove the existing titles at final proof stage.

Funding Information

Information about an article's funding sources is now submitted to Crossref to help you comply with current or future funding agency mandates. Crossref's Open Funder Registry (<https://www.crossref.org/services/funder-registry/>) is the definitive registry of funding agencies. Please ensure that your acknowledgments include all sources of funding for your article following any requirements of your funding sources. Where possible, please include grant and award ids. Please carefully check the following funder information we have already extracted from your article and ensure its accuracy and completeness:

- Comisión Nacional de Investigación Científica y Tecnológica, FundRef ID <http://dx.doi.org/10.13039/501100002848> (Republic of Chile)
- Istituto Nazionale di Fisica Nucleare, FundRef ID <http://dx.doi.org/10.13039/501100004007> (Repubblica Italiana)
- Centre National de la Recherche Scientifique, FundRef ID <http://dx.doi.org/10.13039/501100004794> (Republic of France)
- Commissariat à l'Énergie Atomique et aux Énergies Alternatives, FundRef ID <http://dx.doi.org/10.13039/501100006489> (Republic of France)
- U.S. Department of Energy, FundRef ID <http://dx.doi.org/10.13039/100000015> (United States/District of Columbia)
- Science and Technology Facilities Council, FundRef ID <http://dx.doi.org/10.13039/501100000271> (United Kingdom of Great Britain and Northern Ireland)
- Scottish Universities Physics Alliance, FundRef ID <http://dx.doi.org/10.13039/501100000708> (United Kingdom of Great Britain and Northern Ireland)
- National Research Foundation of Korea, FundRef ID <http://dx.doi.org/10.13039/501100003725> (Republic of Korea)
- Office of Research and Economic Development, University of California, Riverside, FundRef ID <http://dx.doi.org/10.13039/100011115> (United States/California)
- Consulat Général de France à Jérusalem

Other Items to Check

- Please note that the original manuscript has been converted to XML prior to the creation of the PDF proof, as described above. Please carefully check all key elements of the paper, particularly the equations and tabular data.
- Title: Please check; be mindful that the title may have been changed during the peer review process.
- Author list: Please make sure all authors are presented, in the appropriate order, and that all names are spelled correctly.
- Please make sure you have inserted a byline footnote containing the email address for the corresponding author, if desired. Please note that this is not inserted automatically by this journal.
- Affiliations: Please check to be sure the institution names are spelled correctly and attributed to the appropriate author(s).
- Receipt date: Please confirm accuracy.
- Acknowledgments: Please be sure to appropriately acknowledge all funding sources.
- References: Please check to ensure that titles are given as appropriate.
- Hyphenation: Please note hyphens may have been inserted in word pairs that function as adjectives when they occur before a noun, as in “x-ray diffraction,” “4-mm-long gas cell,” and “*R*-matrix theory.” However, hyphens are deleted from word pairs when they are not used as adjectives before nouns, as in “emission by x rays,” “was 4 mm in length,” and “the *R* matrix is tested.”
Note also that Physical Review follows U.S. English guidelines in that hyphens are not used after prefixes or before suffixes: superresolution, quasiequilibrium, nanoprecipitates, resonancelike, clockwise.

- Please check that your figures are accurate and sized properly. Make sure all labeling is sufficiently legible. Figure quality in this proof is representative of the quality to be used in the online journal. To achieve manageable file size for online delivery, some compression and downsampling of figures may have occurred. Fine details may have become somewhat fuzzy, especially in color figures. The print journal uses files of higher resolution and therefore details may be sharper in print. Figures to be published in color online will appear in color on these proofs if viewed on a color monitor or printed on a color printer.
- Overall, please proofread the entire *formatted* article very carefully. The redlined PDF should be used as a guide to see changes that were made during copyediting. However, note that some changes to math and/or layout may not be indicated.

Ways to Respond

- **Web:** If you accessed this proof online, follow the instructions on the web page to submit corrections.
- **Email:** Send corrections to aps-robot@luminad.com. Include the accession code LG16705 in the subject line.
- **Fax:** Return this proof with corrections to +1.855.808.3897.

If You Need to Call Us

You may leave a voicemail message at +1.855.808.3897. Please reference the accession code and the first author of your article in your voicemail message. We will respond to you via email.

First Exclusive Measurement of Deeply Virtual Compton Scattering off ^4He : Toward the 3D Tomography of Nuclei

M. Hattawy,^{1,22} N. A. Baltzell,^{1,37} R. Dupré,^{1,22,*} K. Hafidi,¹ S. Stepanyan,³⁷ S. Bültmann,³¹ R. De Vita,¹⁹ A. El Alaoui,^{1,38} L. El Fassi,²⁷ H. Egiyan,³⁷ F. X. Girod,²² M. Guidal,²² D. Jenkins,⁴³ S. Liuti,⁴² Y. Perrin,²⁶ B. Torayev,³¹ E. Voutier,^{26,22} K. P. Adhikari,²⁷ S. Adhikari,¹² D. Adikaram,^{31,†} Z. Akbar,¹³ M. J. Amarian,³¹ S. Anefalos Pereira,¹⁸ Whitney R. Armstrong,¹ H. Avakian,³⁷ J. Ball,⁷ M. Bashkanov,³⁹ M. Battaglieri,¹⁹ V. Batourine,³⁷ I. Bedlinskiy,²³ A. S. Biselli,¹⁰ S. Boiarinov,³⁷ W. J. Briscoe,¹⁵ W. K. Brooks,³⁸ V. D. Burkert,³⁷ Frank Thanh Cao,⁹ D. S. Carman,³⁷ A. Celentano,¹⁹ G. Charles,³¹ T. Chetry,³⁰ G. Ciullo,^{17,11} L. Clark,⁴⁰ L. Colaneri,²² P. L. Cole,¹⁶ M. Contalbrigo,¹⁷ O. Cortes,¹⁶ V. Crede,¹³ A. D'Angelo,^{20,33} N. Dashyan,⁴⁵ E. De Sanctis,¹⁸ A. Deur,³⁷ C. Djalali,³⁵ L. Elouadrhiri,³⁷ P. Eugenio,¹³ G. Fedotov,^{35,34} S. Fegan,^{40,‡} R. Fersch,^{8,44} A. Filippi,²¹ J. A. Fleming,³⁹ T. A. Forest,¹⁶ A. Fradi,^{22,§} M. Garçon,⁷ N. Gevorgyan,⁴⁵ Y. Ghandilyan,⁴⁵ G. P. Gilfoyle,³² K. L. Giovanetti,²⁴ C. Gleason,³⁵ W. Gohn,^{9,||} E. Golovatch,³⁴ R. W. Gothe,³⁵ K. A. Griffioen,⁴⁴ L. Guo,^{12,37} H. Hakobyan,^{38,45} C. Hanretty,^{37,13} N. Harrison,³⁷ D. Heddle,^{8,37} K. Hicks,³⁰ M. Holtrop,²⁸ S. M. Hughes,³⁹ D. G. Ireland,⁴⁰ B. S. Ishkhanov,³⁴ E. L. Isupov,³⁴ H. Jiang,³⁵ K. Joo,⁹ S. Joosten,³⁶ D. Keller,^{42,30} G. Khachatryan,⁴⁵ M. Khachatryan,³¹ M. Khandaker,^{29,¶} A. Kim,⁹ W. Kim,²⁵ A. Klein,³¹ F. J. Klein,⁶ V. Kubarovskiy,³⁷ S. E. Kuhn,³¹ S. V. Kuleshov,^{38,23} L. Lanza,²⁰ P. Lenisa,¹⁷ K. Livingston,⁴⁰ H. Y. Lu,³⁵ I. J. D. MacGregor,⁴⁰ N. Markov,⁹ M. Mayer,³¹ M. E. McCracken,⁵ B. McKinnon,⁴⁰ C. A. Meyer,⁵ Z. E. Meziani,³⁶ T. Mineeva,^{38,9} M. Mirazita,¹⁸ V. Moiseev,³⁷ R. A. Montgomery,⁴⁰ H. Moutarde,⁷ A. Movsisyan,¹⁷ C. Munoz Camacho,²² P. Nadel-Turonski,³⁷ L. A. Net,³⁵ S. Niccolai,²² G. Niculescu,²⁴ I. Niculescu,²⁴ M. Osipenko,¹⁹ A. I. Ostrovidov,¹³ M. Paolone,³⁶ R. Paremuzyan,^{28,45} K. Park,^{37,35} E. Pasyuk,^{37,2} E. Phelps,³⁵ W. Phelps,¹² S. Pisano,^{18,22} O. Pogorelko,²³ J. W. Price,³ Y. Prok,^{31,42} D. Protopopescu,⁴⁰ M. Ripani,¹⁹ B. G. Ritchie,² A. Rizzo,^{20,33} G. Rosner,⁴⁰ P. Rossi,^{37,18} F. Sabatié,⁷ C. Salgado,²⁹ R. A. Schumacher,⁵ E. Seder,⁹ Y. G. Sharabian,³⁷ A. Simonyan,⁴⁵ Iu. Skorodumina,^{35,34} G. D. Smith,³⁹ D. Sokhan,^{40,39} N. Sparveris,³⁶ S. Strauch,³⁵ M. Taiuti,^{14,‡} M. Ungaro,^{37,9} H. Voskanyan,⁴⁵ N. K. Walford,⁶ D. P. Watts,³⁹ X. Wei,³⁷ L. B. Weinstein,³¹ M. H. Wood,⁴ N. Zachariou,³⁹ L. Zana,^{39,28} J. Zhang,⁴² and Z. W. Zhao^{31,35}

(CLAS Collaboration)

¹Argonne National Laboratory, Argonne, Illinois 60439, USA

²Arizona State University, Tempe, Arizona 85287-1504, USA

³California State University, Dominguez Hills, Carson, California 90747, USA

⁴Canisius College, Buffalo, New York, USA

⁵Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

⁶Catholic University of America, Washington, DC 20064, USA

⁷IRFU/SPHn, CEA, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

⁸Christopher Newport University, Newport News, Virginia 23606, USA

⁹University of Connecticut, Storrs, Connecticut 06269, USA

¹⁰Fairfield University, Fairfield, Connecticut 06824, USA

¹¹Università di Ferrara, 44121 Ferrara, Italy

¹²Florida International University, Miami, Florida 33199, USA

¹³Florida State University, Tallahassee, Florida 32306, USA

¹⁴Università di Genova, 16146 Genova, Italy

¹⁵The George Washington University, Washington, DC 20052, USA

¹⁶Idaho State University, Pocatello, Idaho 83209, USA

¹⁷INFN, Sezione di Ferrara, 44100 Ferrara, Italy

¹⁸INFN, Laboratori Nazionali di Frascati, 00044 Frascati, Italy

¹⁹INFN, Sezione di Genova, 16146 Genova, Italy

²⁰INFN, Sezione di Roma Tor Vergata, 00133 Rome, Italy

²¹INFN, Sezione di Torino, 10125 Torino, Italy

²²Institut de Physique Nucléaire, CNRS/IN2P3 and Université Paris Sud, Orsay, France

²³Institute of Theoretical and Experimental Physics, Moscow 117259, Russia

²⁴James Madison University, Harrisonburg, Virginia 22807, USA

²⁵Kyungpook National University, Daegu 41566, Republic of Korea

²⁶LPSC, Université Grenoble-Alpes, CNRS/IN2P3, Grenoble, France

²⁷Mississippi State University, Mississippi State, Mississippi 39762-5167, USA

²⁸University of New Hampshire, Durham, New Hampshire 03824-3568, USA

²⁹Norfolk State University, Norfolk, Virginia 23504, USA

³⁰Ohio University, Athens, Ohio 45701, USA

³¹Old Dominion University, Norfolk, Virginia 23529, USA

³²University of Richmond, Richmond, Virginia 23173, USA

³³Universita' di Roma Tor Vergata, 00133 Rome, Italy

³⁴Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, 119234 Moscow, Russia

³⁵University of South Carolina, Columbia, South Carolina 29208, USA

³⁶Temple University, Philadelphia, Pennsylvania 19122, USA

³⁷Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

³⁸Universidad Técnica Federico Santa María, Casilla 110-V Valparaíso, Chile

³⁹Edinburgh University, Edinburgh EH9 3JZ, United Kingdom

⁴⁰University of Glasgow, Glasgow G12 8QQ, United Kingdom

⁴¹Virginia Tech, Blacksburg, Virginia 24061-0435, USA

⁴²University of Virginia, Charlottesville, Virginia 22901, USA

⁴³Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA

⁴⁴College of William and Mary, Williamsburg, Virginia 23187-8795, USA

⁴⁵Yerevan Physics Institute, 375036 Yerevan, Armenia

(Received 13 July 2017; revised manuscript received 11 September 2017)

We report on the first measurement of the beam-spin asymmetry in the exclusive process of coherent deeply virtual Compton scattering off a nucleus. The experiment uses the 6 GeV electron beam from the Continuous Electron Beam Accelerator Facility (CEBAF) accelerator at Jefferson Lab incident on a pressurized ^4He gaseous target placed in front of the CEBAF Large Acceptance Spectrometer (CLAS). The scattered electron is detected by CLAS and the photon by a dedicated electromagnetic calorimeter at forward angles. To ensure the exclusivity of the process, a specially designed radial time projection chamber is used to detect the recoiling ^4He nuclei. We measure beam-spin asymmetries larger than those observed on the free proton in the same kinematic domain. From these, we are able to extract, in a model-independent way, the real and imaginary parts of the only ^4He Compton form factor, \mathcal{H}_A . This first measurement of coherent deeply virtual Compton scattering on the ^4He nucleus, with a fully exclusive final state via nuclear recoil tagging, leads the way toward 3D imaging of the partonic structure of nuclei.

DOI:

The generalized parton distribution (GPD) framework offers the opportunity to obtain information about the momentum and spatial degrees of freedom of the quarks and gluons inside hadrons [1–5]. In impact parameter space, the GPDs are indeed interpreted as a tomography of the transverse plane for partons carrying a given fraction of the proton longitudinal momentum [6–9]. The most promising way to access GPDs experimentally is through the measurement of deeply virtual Compton scattering (DVCS), i.e., the hard exclusive electroproduction of a real photon on a hadron. While other processes are also known to be sensitive to GPDs, the measurement of DVCS is considered the cleanest probe and has been the focus of efforts at Jefferson Lab, HERA, and CERN [10–25]. The vast majority of these measurements focused on the study of the proton and allowed for an extraction of its three-dimensional image (for reviews of the field, see [26–31]). These recent developments could also be applicable to nuclei, giving access to novel information about nuclear structure in terms of quarks and gluons [32–35]. Such a study of the 3D structure of nuclei appears to be especially interesting in light of the large, yet unresolved, nuclear

effects observed in nuclear parton distribution functions [36–38]. The results presented in this Letter demonstrate the feasibility of such an approach and constitute the first step toward a tomography of nuclei.

Figure 1 illustrates the handbag diagram for coherent DVCS on ^4He , where the four-vectors of the electrons, photons, and ^4He are denoted by k/k' , q/q' , and p/p' , respectively. For large virtual photon 4-momentum squared, $Q^2 = -(k - k')^2$, and small squared momentum




FIG. 1. Representation of the leading-order handbag diagram of the DVCS process off ^4He .

transfer, $t = (p - p')^2$, the DVCS handbag diagram can be factorized into two parts [39,40]. The hard part includes the photon-quark interaction and is calculable in perturbative QED. The nonperturbative part is parametrized in terms of GPDs, which embed the partonic structure of the hadron. The GPDs depend on the three variables x , ξ , and t . ξ relates to the Bjorken variable x_A : $\xi \approx (x_A/2 - x_A)$, where $x_A = (Q^2/2M_A\nu)$, ν is the energy of the virtual photon, and M_A is the nuclei mass. x is the quark's internal loop momentum fraction and cannot be accessed experimentally in DVCS. We in fact measure Compton form factors (CFFs), which are complex quantities defined as

$$\text{Re}(\mathcal{H}_A) = \mathcal{P} \int_0^1 dx [H_A(x, \xi, t) - H_A(-x, \xi, t)] C^+(x, \xi), \quad (1)$$

$$\text{Im}(\mathcal{H}_A) = -\pi [H_A(\xi, \xi, t) - H_A(-\xi, \xi, t)], \quad (2)$$

with H_A a GPD, \mathcal{P} the Cauchy principal value integral, and a coefficient function $C^+ = (1/x - \xi) + (1/x + \xi)$.

Until now, the only available data on nuclear DVCS were from the HERMES experiment [12]. In this experiment, the exclusivity of the reaction was obtained through kinematic cuts using only the measured scattered electron and real photon. This measurement was performed on a large set of nuclei (^4He , ^{14}N , ^{20}Ne , ^{85}Kr , and ^{131}Xe), but contamination from incoherent processes can be suspected to influence the results significantly [41]. The direct detection of the recoil nucleus can, however, guarantee that the nucleus remains intact.

The ^4He nucleus is an ideal experimental target for nuclear DVCS, as it is light enough to be detected by our experimental setting, while it is subject to interesting nuclear effects [42]. Its zero spin also leads to an important simplification, as a spin-zero hadron is parametrized by only one chiral even GPD [$H_A(x, \xi, t)$] at leading twist, while four GPDs arise for the spin- $\frac{1}{2}$ nucleon. This significantly simplifies the interpretation of the data and allows a model-independent extraction of the ^4He CFF (\mathcal{H}_A) presented at the end of this Letter.

The Continuous Electron Beam Accelerator Facility (CEBAF) Large Acceptance Spectrometer (CLAS) in Hall B at Jefferson Lab [43] has been previously supplemented with an inner calorimeter (IC) and a solenoid magnet to measure DVCS observables on the nucleon [18,20,21,23,24]. The IC extended the photon detection acceptance of CLAS to polar angles as low as 4° . The 5-T solenoid magnet acted as a guiding field for the low-energy Møller electrons that were absorbed in a heavy shield placed around the beam line.

In the kinematic range of the present experiment, the recoil ^4He nuclei have low momentum, averaging 300 MeV. CLAS could not detect such low-energy α particles, so, in

order to ensure the exclusivity of the measurement, we built a small and light radial time projection chamber (RTPC). The RTPC was a 20-cm-long cylinder with a diameter of 15 cm, positioned in the solenoid magnet. In the center of the RTPC was the target cell, a 25-cm-long and 6-mm-diameter Kapton tube with 27- μm -thick walls filled with gaseous ^4He at 6 atm (see [44] for a detailed description of the RTPC and its performances). The experiment (E08-024) [45] collected data over 40 days at the end of 2009 using a nearly 100% duty factor, longitudinally polarized electron beam ($83.7 \pm 3.5\%$ polarization [46]) at an energy of 6.064 GeV. The RTPC was calibrated specifically for the detection of ^4He nuclei using elastic scattering ($e^4\text{He} \rightarrow e'^4\text{He}'$) with a 1.2 GeV electron beam.

To identify coherent DVCS events, we first selected events where one electron, one ^4He , and at least one photon were detected in the final state. Electrons were identified using their measured momentum, light yield, time, and energy obtained from the CLAS drift chambers, Čerenkov counters, scintillator counters, and electromagnetic calorimeters, respectively. The recoiling ^4He nuclei were identified in the RTPC using time and energy-loss cuts for tracks in the fiducial region [47]. In addition, we applied a vertex-matching cut to ensure that the electron and helium nucleus originated from a common reaction vertex in the target cell. The photons were detected in either the IC or the CLAS electromagnetic calorimeters. Note that, even though the DVCS reaction has only one real photon in the final state, events with more than one good photon were not discarded at this stage. These were mainly caused by accidental coincidences of soft photons and did not directly affect this measurement, as only the most energetic photon of an event was considered a DVCS photon candidate. This prescription, however, slightly increased the corrections associated with the π^0 and the accidental backgrounds discussed below.

We selected events with Q^2 greater than 1 GeV^2 for which the DVCS handbag diagram is believed to be dominant. Then the exclusivity of the reaction was ensured by applying a set of cuts on the following kinematic variables: the coplanarity angle $\Delta\phi$ between the (γ, γ^*) and $(\gamma^*, ^4\text{He}')$ planes, the missing energy, mass, and transverse momentum of the $e'^4\text{He}'$ system, the missing mass squared of the $e'^4\text{He}'$ system, and the angle θ between the measured photon and the missing momentum of the $e'^4\text{He}'$ system. The experimental data for the most relevant exclusivity variables and applied cuts are shown in Fig. 2 (see [47] for additional details). We also rejected events where a π^0 was identified by the invariant mass of two photons. About 3200 events passed all these requirements; their kinematic distributions are shown in Fig. 3. (We use here and for other results $x_B = (Q^2/2M_N\nu)$ with M_N the proton mass instead of x_A . This makes it easier to compare these results with the proton DVCS data available in the literature).




FIG. 2. Four of the six coherent DVCS exclusivity cuts. The black distributions represent the initial candidate events, while the shaded distributions represent those that passed all of the exclusivity cuts except the quantity plotted. The vertical red lines represent the applied cuts. The distributions from left to right and from top to bottom are coplanarity angle $\Delta\phi$, missing energy E_X , missing mass squared M_X^2 , and the cone angle θ between the measured photon and the missing momentum of the $e'^4\text{He}'$ system.

We identified two main backgrounds: accidental coincidences and exclusive coherent π^0 production. The accidentals have particles originating from different events, and we estimated their contribution to be 4.1% of the data sample. We evaluated this contribution by selecting events passing all the cuts but with the scattered electron and ^4He originating from different vertices. The π^0 production can be mistaken for DVCS when one of the two photons from the π^0 decay is produced at low energy in the laboratory frame and remains undetected. To estimate the effect of this contamination, we developed an event generator tuned on the experimental yield of exclusive π^0 with two photons measured. We used this generator together with a GEANT3




FIG. 3. Coherent DVCS event distributions for Q^2 after exclusivity cuts. The distributions are shown as a function of Bjorken variable x_B (left) and as a function of squared-momentum transfer $-t$ (right).

simulation of our detectors to estimate the ratio of the number of π^0 events where the two photons are detected to those that are misidentified as DVCS events. This ratio is then multiplied by the measured yield of exclusive π^0 events to correct the DVCS data. Depending on the kinematics, we found contaminations of 2%–4%.

In this work, the physics observable extracted using coherent DVCS events is the beam-spin asymmetry A_{LU} . On an unpolarized target, A_{LU} is defined as the difference of cross sections for the reaction with opposite beam helicities normalized to the total cross section:

$$A_{LU} = \frac{d^4\sigma^+ - d^4\sigma^-}{d^4\sigma^+ + d^4\sigma^-}, \quad (3)$$

where $d^4\sigma^\pm$ is the DVCS differential cross section for positive (negative) beam helicity.


In this ratio, luminosity normalization and detector efficiencies largely cancel, and A_{LU} can be extracted from the reaction yields for the two helicities (N^\pm):

$$A_{LU} = \frac{1}{P_B} \frac{N^+ - N^-}{N^+ + N^-}, \quad (4)$$

where P_B is the degree of longitudinal polarization of the incident electron beam.

There is an additional process contributing to the same final state as the DVCS, the so-called Bethe-Heitler (BH) process, where the real photon is emitted by the incoming or the outgoing lepton. The DVCS and BH processes are indistinguishable experimentally, and the amplitude of the electroproduction of a real photon includes a sum of the amplitudes of these two processes. The BH amplitude depends on the target elastic form factors, which are well known in this kinematic region, while the DVCS amplitude depends on the GPDs. In our kinematics, the cross section of the real photon electroproduction is dominated by the BH contribution, which varies strongly with ϕ , the azimuthal angle between the (e, e') and $(\gamma^*, ^4\text{He}')$ planes. The DVCS contribution is smaller by about a factor of 2 but independent of ϕ at twist-2 [48], and thus it is possible to separate these contributions with a cross section measurement. However, the DVCS-BH interference term offers an easier experimental access by generating spin asymmetries. We have, in particular, for a spin-zero target the beam spin asymmetry A_{LU} , which can be expressed at leading order and leading twist [49,50] as

$$A_{LU}(\phi) = \frac{\alpha_0(\phi)\text{Im}(\mathcal{H}_A)}{\alpha_1(\phi) + \alpha_2(\phi)\text{Re}(\mathcal{H}_A) + \alpha_3(\phi)[\text{Re}(\mathcal{H}_A)^2 + \text{Im}(\mathcal{H}_A)^2]}. \quad (5)$$



F4:1 FIG. 4. A_{LU} as a function of azimuthal angle ϕ . Results are
 F4:2 presented for different Q^2 bins (top panel), x_B bins (middle
 F4:3 panel), and t bins (bottom panel). The error bars represent the
 F4:4 statistical uncertainties. The gray bands represent the systematic
 F4:5 uncertainties, including the normalization uncertainties. The red
 F4:6 curves are the results of fits with Eq. (5).

276 Explicit expressions of the kinematic factors α_i are derived
 278 7 from expressions in Ref. [50]:


$$\begin{aligned}
 \alpha_0(\phi) &= \frac{x_A(1+\epsilon^2)^2}{y} S_{++}(1) \sin(\phi), \\
 \alpha_1(\phi) &= c_0^{\text{BH}} + c_1^{\text{BH}} \cos(\phi) + c_2^{\text{BH}} \cos(2\phi), \\
 \alpha_2(\phi) &= \frac{x_A(1+\epsilon^2)^2}{y} [C_{++}(0) + C_{++}(1) \cos(\phi)], \\
 \alpha_3(\phi) &= \frac{x_A^2 t(1+\epsilon^2)^2}{y} \mathcal{P}_1(\phi) \mathcal{P}_2(\phi) 2 \frac{2-2y+y^2+\frac{\epsilon^2}{2}y^2}{1+\epsilon^2}, \quad (6)
 \end{aligned}$$

where $S_{++}(1)$ and $C_{++}(0, 1)$ are the Fourier harmonics of
 the interference amplitude in the leptonic tensor, $c_{0,1,2}^{\text{BH}}$ the
 Fourier harmonics of the BH amplitude, and finally $\mathcal{P}_{1,2}(\phi)$
 the BH propagators, which include $\cos(\phi)$ dependencies.
 (The explicit expression of all these terms can be found in
 Appendix A of Ref. [51].) We observe that, using the
 different $\sin(\phi)$, $\cos(\phi)$, and $\cos(2\phi)$ contributions, one
 can extract unambiguously both the imaginary and real
 parts of \mathcal{H}_A with a fit of the $A_{LU}(\phi)$ distribution.

We present in Fig. 4 A_{LU} as a function of azimuthal angle ϕ
 and the kinematical variables Q^2 , x_B , and t . Because of limited
 statistics, these latter variables are studied separately with a
 two-dimensional data binning. The curves on the plots are fits
 using the function presented in Eq. (5), where the real and
 imaginary parts of the CFF \mathcal{H}_A are the only free parameters.

Studies of systematic uncertainties showed that the main
 contributions come from the choice of DVCS exclusivity
 cuts (8% systematic uncertainty) and the large binning size
 (5.1%). These values are relative and quoted for A_{LU} at
 $\phi = 90^\circ$. Added quadratically, the total systematic uncer-
 tainty is about 10% at 90° (or 0.03, absolute), which is
 significantly smaller than the statistical uncertainties at all
 kinematical bins.

In Fig. 5, the Q^2 , x_B , and t dependencies of the fitted A_{LU}
 at $\phi = 90^\circ$ are shown. The comparison to HERMES data
 shows that we obtain the same sign, but the size of the error
 bars and the difference of kinematics do not permit us to
 say much more. The x_B and t dependencies are also
 compared to theoretical calculations by Liuti and Taneja
 [52]. The model accounts for the effect of the nucleon
 virtuality (off-shellness) on the quark distribution. The
 calculations are at slightly different kinematics than the
 data but still allow us to draw some conclusions. The model
 appears to predict smaller asymmetries than observed. The
 difference may arise from the theoretical uncertainty in the
 determination of the crossing point where the parton
 nuclear distribution becomes larger than the nucleon one
 and reverses the sign of the nuclear effect.



F5:1 FIG. 5. The Q^2 (left), x_B (middle), and t dependencies (right) of A_{LU} at $\phi = 90^\circ$ (black squares). In the middle plot, the full red and the
 F5:2 dashed blue curves are theoretical calculations from Ref. [52]. On the right, the green circles are the HERMES $-A_{LU}$ (a positron beam
 F5:3 was used) inclusive measurements [12], and the curve is the theoretical calculation from Ref. [52]. The gray bands represent the
 F5:4 systematic errors.




FIG. 6. The leading-twist model-independent extraction of the imaginary (top panel) and real (bottom panel) parts of the ^4He CFF \mathcal{H}_A as a function of Q^2 (left panel), x_B (middle panel), and t (right panel). The full red curves are the calculations based on the convolution model from Ref. [53]. The black dashed curves are calculations from the same model using different GPDs for the nucleons [Vanderhaeghen-Guichon-Guidal (VGG) model [55]]. The blue long-dashed curve on the top-right plot is from the off-shell model as described in Ref. [57].

The Q^2 , x_B , and t dependencies of the ^4He CFF \mathcal{H}_A extracted from the fit to the azimuthal dependence of A_{LU} are shown in Fig. 6. The curves on the graphs are model calculations, labeled *convolution* and *off-shell*. In the convolution model [53], the nucleus is assumed to be composed of nonrelativistic nucleons, each interacting independently with the probe. The convolution-dual model is based on nucleon GPDs from the dual parametrization [54], where the convolution VGG uses nucleon GPDs from the VGG model [55] and is based on the double distributions ansatz [56]. The off-shell model is the same as in Fig. 5 using a more recent GPD model for the nucleon [57].

The results in Fig. 6 show that the extraction of the CFF from the A_{LU} is possible without model-dependent assumptions beyond leading-twist and leading-order dominance. The amplitude and the dependencies observed as a function of Q^2 , x_B , and t are in agreement with the theoretical expectations. One can see a difference between the precision of the extracted imaginary and real parts, which is due to α_2 being much smaller than α_1 in Eq. (5). While the precision of this measurement is not at a sufficient level to discriminate between the models, these results demonstrate the possibility of extracting the CFF of a spin-0 target directly from a A_{LU} measurement.

In summary, we have presented the first measurement of the beam-spin asymmetry of exclusive coherent DVCS off ^4He using the CLAS spectrometer supplemented with a RTPC. This setup allowed detection of the low-energy ^4He recoils in order to ensure an exclusive measurement of the

coherent DVCS process. The azimuthal dependence of the measured A_{LU} has been used to extract, in a model-independent way, the real and the imaginary parts of the ^4He CFF, \mathcal{H}_A . The extracted CFF is in agreement with predictions of the available models. This first fully exclusive experiment opens new perspectives for studying nuclear structure with the GPD framework and paves the way for future measurements at JLab using 12 GeV CEBAF and upgraded equipment [51].

The authors thank the staff of the Accelerator and Physics Divisions at the Thomas Jefferson National Accelerator Facility who made this experiment possible. This work was supported in part by the Chilean Comisión Nacional de Investigación Científica y Tecnológica (CONICYT), the Italian Istituto Nazionale di Fisica Nucleare, the French Centre National de la Recherche Scientifique, the French Commissariat à l'Energie Atomique, the U.S. Department of Energy under Contract No. DE-AC02-06CH11357, the United Kingdom Science and Technology Facilities Council (STFC), the Scottish Universities Physics Alliance (SUPA), the National Research Foundation of Korea, and the Office of Research and Economic Development at Mississippi State University. M. Hattawy also acknowledges the support of the Consulat Général de France à Jérusalem. The Southeastern Universities Research Association operates the Thomas Jefferson National Accelerator Facility for the United States Department of Energy under Contract No. DE-AC05-06OR23177.

*Corresponding author.
dupre@ipno.in2p3.fr

[†]Present address: Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA.

[‡]Present address: INFN, Sezione di Genova, 16146 Genova, Italy.

[§]Present address: Faculty of Sciences of Gabes, Department of Physics, 6072-Gabes, Tunisia.

^{||}Present address: University of Kentucky, Lexington, KY 40506, USA.

[¶]Present address: Idaho State University, Pocatello, ID 83209, USA.

- [1] D. Mueller, D. Robaschik, B. Geyer, F.M. Dittes, and J. Horejsi, *Fortschr. Phys.* **42**, 101 (1994).
- [2] X. D. Ji, *Phys. Rev. Lett.* **78**, 610 (1997).
- [3] X. D. Ji, *Phys. Rev. D* **55**, 7114 (1997).
- [4] A. V. Radyushkin, *Phys. Lett. B* **380**, 417 (1996).
- [5] A. V. Radyushkin, *Phys. Rev. D* **56**, 5524 (1997).
- [6] M. Burkardt, *Phys. Rev. D* **62**, 071503 (2000); **66**, 119903 (E) (2002).
- [7] M. Diehl, *Eur. Phys. J. C* **25**, 223 (2002); **31**, 277(E) (2003).
- [8] A. V. Belitsky and D. Mueller, *Nucl. Phys.* **A711**, 118 (2002).
- [9] M. Burkardt, *Phys. Rev. D* **72**, 094020 (2005).
- [10] S. Stepanyan *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **87**, 182002 (2001).

- [11] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. Lett.* **87**, 182001 (2001); *Phys. Rev. D* **75**, 011103 (2007); *J. High Energy Phys.* 06 (2008) 066; **11** (2009) 083; **06** (2010) 019; *Phys. Lett. B* **704**, 15 (2011); *J. High Energy Phys.* 07 (2012) 032; **10** (2012) 042.
- [12] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. C* **81**, 035202 (2010).
- [13] C. Adloff *et al.* (H1 Collaboration), *Phys. Lett. B* **517**, 47 (2001).
- [14] S. Chekanov *et al.* (ZEUS Collaboration), *Phys. Lett. B* **573**, 46 (2003).
- [15] A. Aktas *et al.* (H1 Collaboration), *Eur. Phys. J. C* **44**, 1 (2005).
- [16] S. Chen *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **97**, 072002 (2006).
- [17] C. Muñoz Camacho *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. Lett.* **97**, 262002 (2006).
- [18] F. X. Girod *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **100**, 162002 (2008).
- [19] M. Mazouz *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. Lett.* **99**, 242501 (2007).
- [20] G. Gavalian *et al.* (CLAS Collaboration), *Phys. Rev. C* **80**, 035206 (2009).
- [21] E. Seder *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **114**, 032001 (2015).
- [22] M. Defurne *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. C* **92**, 055202 (2015).
- [23] S. Pisano *et al.* (CLAS Collaboration), *Phys. Rev. D* **91**, 052014 (2015).
- [24] H. S. Jo *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **115**, 212003 (2015).
- [25] P. Joerg (COMPASS Collaboration), *Proc. Sci.*, DIS2016 (2016) 235.
- [26] K. Goeke, M. V. Polyakov, and M. Vanderhaeghen, *Prog. Part. Nucl. Phys.* **47**, 401 (2001).
- [27] M. Diehl, *Phys. Rep.* **388**, 41 (2003).
- [28] X. D. Ji, *Annu. Rev. Nucl. Part. Sci.* **54**, 413 (2004).
- [29] A. V. Belitsky and A. V. Radyushkin, *Phys. Rep.* **418**, 1 (2005).
- [30] S. Boffi and B. Pasquini, *Riv. Nuovo Cimento Soc. Ital. Fis.* **30**, 387 (2007).
- [31] M. Guidal, H. Moutarde, and M. Vanderhaeghen, *Rep. Prog. Phys.* **76**, 066202 (2013).
- [32] E. R. Berger, F. Cano, M. Diehl, and B. Pire, *Phys. Rev. Lett.* **87**, 142302 (2001).
- [33] F. Cano and B. Pire, *Eur. Phys. J. A* **19**, 423 (2004).
- [34] V. Guzey and M. Siddikov, *J. Phys. G* **32**, 251 (2006).
- [35] R. Dupré and S. Scopetta, *Eur. Phys. J. A* **52**, 159 (2016).
- [36] O. Hen, G. A. Miller, E. Piasetzky, and L. B. Weinstein, *arXiv:1611.09748* [Rev. Mod. Phys. (to be published)].
- [37] P. R. Norton, *Rep. Prog. Phys.* **66**, 1253 (2003).
- [38] D. F. Geesaman, K. Saito, and A. W. Thomas, *Annu. Rev. Nucl. Part. Sci.* **45**, 337 (1995).
- [39] J. C. Collins and A. Freund, *Phys. Rev. D* **59**, 074009 (1999).
- [40] X.-D. Ji and J. Osborne, *Phys. Rev. D* **58**, 094018 (1998).
- [41] V. Guzey and M. Strikman, *Phys. Rev. C* **68**, 015204 (2003).
- [42] J. Seely *et al.*, *Phys. Rev. Lett.* **103**, 202301 (2009).
- [43] B. A. Mecking *et al.* (CLAS Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **503**, 513 (2003).
- [44] R. Dupré *et al.*, *arXiv:1706.10160*.
- [45] K. Hafidi *et al.*, proposal PR-08-024 to JLab PAC33 (unpublished).
- [46] Y. Perrin, Ph.D. thesis, Université de Grenoble, France, 2012 [Institution Report No. 2012GRENY092].
- [47] M. Hattawy, Ph.D. thesis, Université Paris Sud—Paris XI, France, 2015 [Institution Report No. 2015PA112161].
- [48] A. V. Belitsky, D. Mueller, and A. Kirchner, *Nucl. Phys. B* **629**, 323 (2002).
- [49] A. Kirchner and D. Mueller, *Eur. Phys. J. C* **32**, 347 (2003).
- [50] A. V. Belitsky and D. Mueller, *Phys. Rev. D* **79**, 014017 (2009).
- [51] W. Armstrong *et al.*, *arXiv:1708.00888*.
- [52] S. Liuti and S. K. Taneja, *Phys. Rev. C* **72**, 032201 (2005).
- [53] V. Guzey (private communication); *Phys. Rev. C* **78**, 025211 (2008).
- [54] V. Guzey and T. Teckentrup, *Phys. Rev. D* **74**, 054027 (2006).
- [55] M. Vanderhaeghen, P. A. M. Guichon, and M. Guidal, *Phys. Rev. Lett.* **80**, 5064 (1998); *Phys. Rev. D* **60**, 094017 (1999); K. Goeke, M. V. Polyakov, and M. Vanderhaeghen, *Prog. Part. Nucl. Phys.* **47**, 401 (2001); M. Guidal, M. V. Polyakov, A. V. Radyushkin, and M. Vanderhaeghen, *Phys. Rev. D* **72**, 054013 (2005).
- [56] I. V. Musatov and A. V. Radyushkin, *Phys. Rev. D* **61**, 074027 (2000).
- [57] J. O. Gonzalez-Hernandez, S. Liuti, G. R. Goldstein, and K. Kathuria, *Phys. Rev. C* **88**, 065206 (2013).