Vaccine	Purpose	Govt. Scheme
B.C.G.	For Tuberculosis (TB)	Yes
O.P.V. (1 st Dose) (Oral Polio Vaccine)	For Polio	Yes
Hepatitis B (1 st Dose) (Within 24 hrs)	For Hepatitis B	Yes
DTWP/DtaP (1st Dose)	For Diphtheria, Tetanus, and Pertussis	Yes
I.P.V. (1 st dose)/O.P.V. (2 nd Dose)	For Polio	Yes
Hepatitis B (2 nd Dose)	For Hepatitis B	Yes
H.I.B. (1 st Dose)	For H. Influenzae type B infection	Yes
Penta-1	For Diphtheria, Tetanus, Hepatitis B, Pertussis & Haemophilus influenzae Type B	Yes
Rota Virus (1 st Dose)	For rotavirus infections (severe diarrhea)	May be
Pnemococcal/PCV (1 st Dose)	To prevent pneumococcal disease	No
	B.C.G. O.P.V. (1 st Dose) (Oral Polio Vaccine) Hepatitis B (1 st Dose) (Within 24 hrs) DTWP/DtaP (1 st Dose) I.P.V. (1 st dose)/O.P.V. (2 nd Dose) Hepatitis B (2 nd Dose) H.I.B. (1 st Dose) Penta-1 Rota Virus (1 st Dose) Pnemococcal/PCV	B.C.G. For Tuberculosis (TB) O.P.V. (1st Dose) (Oral Polio Vaccine) Hepatitis B (1st Dose) (Within 24 hrs) DTWP/DtaP (1st Dose) I.P.V. (1st dose)/O.P.V. (2nd Dose) Hepatitis B (2nd Dose) Hepatitis B (2nd Dose) For Hepatitis B H.I.B. (1st Dose) For Hepatitis B Rota Virus (1st Dose) For Diphtheria, Tetanus, and Pertussis For Hepatitis B Pertussis & Haemophilus influenzae Type B Rota Virus (1st Dose) For rotavirus infections (severe diarrhea) Pnemococcal/PCV To prevent pneumococcal disease

Age	Vaccine	Purpose	Govt. Scheme
10 Weeks	DTWP/DtaP (2 nd Dose)	For Diphtheria, Tetanus, and Pertussis	Yes
	I.P.V. (2 nd dose) /O.P.V. (3 rd Dose)	For Polio	Yes
	Hepatitis B (3 rd Dose)	For Hepatitis B	Yes
	H.I.B. (2 nd Dose)	For H. Influenzae type B infection	Yes
	Penta-2	For Diphtheria, Tetanus, Hepatitis B, Pertussis & Haemophilus influenzae Type B	Yes
	Rota Virus (2 nd Dose)	For rotavirus infections (severe diarrhea)	May be
	Pnemococcal/PCV (2 nd Dose)	To prevent pneumococcal disease	No

Age	Vaccine	Purpose	Govt. Scheme
14 Weeks	DTWP/DtaP (3 rd Dose)	For Diphtheria, Tetanus, and Pertussis	Yes
	I.P.V. (3 rd dose)/ O.P.V. (4 th Dose)	For Polio	Yes
	Hepatitis B (4 th Dose)	For Hepatitis B	Yes
	H.I.B. (3 rd Dose)	For H. Influenzae type B infection	Yes
	Penta-3	For Diphtheria, Tetanus, Hepatitis B, Pertussis & Haemophilus influenzae Type B	Yes
	Rota Virus (3 rd Dose)	For rotavirus infections (severe diarrhea)	May be
	Pnemococcal/PCV (3 rd Dose)	To prevent pneumococcal disease	No

Age	Vaccine	Purpose	Govt. Scheme
6 Months	Flu Vaccine	To prevent infection by influenza viruses	No
	O.P.V. (5 th Dose)	For Polio	Yes
9 Months	M.M.R. (1 st Dose)	For Measles, Mumps & Rubella	Yes
	Vitamin A (1 st Dose)	For more protection against measles	Yes
	Typhoid Conjugate Vaccine (9 to 12 Months)	For Typhoid	No
12 Months	Hepatitis A (Only Dose if Live Vaccine)	For Hepatitis A	No
	Japanese Encephalitis	For Viral Brain Infection (Mandatory for endemic states)	Optional
15 Months	M.M.R. (2 nd Dose)	For Measles, Mumps & Rubella	Yes
	Varicella (1 st Dose)	For Chickenpox	No
	Pnemococcal/PCV (4 th Dose)	To prevent pneumococcal disease	No

Age	Vaccine	Purpose	Govt. Scheme
16-18 Months	DTWP/DtaP (Booster)	For Diphtheria, Tetanus, and Pertussis	Yes
	H.I.B. (Booster)	For H. Influenzae type B infection	Yes
18 to 24 Months	Typhoid Conjugate Vaccine (Booster)	For Typhoid	No
4 to 6 Years	DTWP/DtaP (Booster)	For Diphtheria, Tetanus, and Pertussis	Yes
	M.M.R. (Booster)	For Measles, Mumps & Rubella	Yes
	Varicella (Booster)	For Chickenpox	No
10 Years	Tdap/Td	For Tetanus, Diphtheria, Pertussis	Yes
	HPV (Human Papilloma Virus) (Girls Only)	To prevent cancer of the cervix (3 Dosage)	No