

# In-EVM Mina State Verification Circuit Description

Cherniaeva Alisa

[a.cherniaeva@nil.foundation](mailto:a.cherniaeva@nil.foundation)

=nil; Crypto3 (<https://crypto3.nil.foundation>)

Shirobokov Ilia

[i.shirobokov@nil.foundation](mailto:i.shirobokov@nil.foundation)

=nil; Crypto3 (<https://crypto3.nil.foundation>)

November 16, 2021

## 1 Introduction

### WIP

High level description according to RfP<sup>1</sup>

1. Computing several hash values from the data of the proof. This involves using the Poseidon hash function with 55 full rounds both over  $\mathbb{F}_p$  and  $\mathbb{F}_q$  with round constants and MDS matrix specified for  $\mathbb{F}_p$ <sup>2</sup> and for  $\mathbb{F}_q$ <sup>3</sup>.
2. Checking arithmetic equations.
3. Performing one multi-scalar multiplication (MSM) of size  $2n_2 + 4 + (2 + 25) = 63$ , for which some of the bases are fixed and some are variable.
4. For each  $i \in \{1, 2\}$ , performing a multi-scalar multiplication over  $\mathbb{G}_i$  of size  $2^{n_i}$  with a fixed array of bases, and with scalars that can be very efficiently computed from the proof.

Note that for MSM in Step 4:

$$\sum_{i=0}^{2^{n_k}-1} s_i \cdot G_i = H$$
$$s_i := \prod_{\substack{0 \leq j \leq n_k \\ \text{bits}(i)[j]=1}} \phi(c_j),$$

where:

- $\phi: \{0, 1\}^{128} \rightarrow \mathbb{F}$  is defined as `to_field` in the implementation<sup>4</sup>.
- Given an integer  $i < 2^{n_k}$ ,  $\text{bits}(i)$  is defined as the little-endian bit array of length  $n$  representing the binary expansion of  $i$ .
- $G_0, \dots, G_{2^{n_k}-1} \in \mathbb{G}_k$  is a fixed sequence of group elements<sup>5</sup>.
- $c_0, \dots, c_{n_k-1} \in \{0, 1\}^{128}$  is a sequence of challenges.

We use the same 15-wires PLONK circuits that are designed for Mina.<sup>6</sup>

## 2 Preliminaries

### WIP

<sup>1</sup>[https://hackmd.io/u\\_2Ygx8XS5Ss1a0bgOFjkA](https://hackmd.io/u_2Ygx8XS5Ss1a0bgOFjkA)

<sup>2</sup><https://github.com/o1-labs/proof-systems/blob/master/oracle/src/pasta/fp.rs>

<sup>3</sup><https://github.com/o1-labs/proof-systems/blob/master/oracle/src/pasta/fq.rs>

<sup>4</sup><https://github.com/o1-labs/proof-systems/blob/49f81edc9c86e5907d26ea791fa083640ad0ef3e/oracle/src/sponge.rs#L33>

<sup>5</sup><https://github.com/o1-labs/proof-systems/blob/master/dlog/commitment/src/srs.rs#L70>

<sup>6</sup>[https://o1-labs.github.io/mina-book/specs/15\\_wires/15\\_wires.html](https://o1-labs.github.io/mina-book/specs/15_wires/15_wires.html)

## 2.1 Pasta Curves

Let  $n_1 = 17$ ,  $n_2 = 16$ . Pasta curves parameters:

- $p = 2^{254} + 45560315531419706090280762371685220353$
- $q = 2^{254} + 45560315531506369815346746415080538113$
- Pallas:

$$\mathbb{G}_1 = \{(x, y) \in \mathbb{F}_p | y^2 = x^3 + 5\}$$

$$|\mathbb{G}_1| = q$$

- Vesta:

$$\mathbb{G}_2 = \{(x, y) \in \mathbb{F}_q | y^2 = x^3 + 5\}$$

$$|\mathbb{G}_2| = p$$

## 2.2 Verification Algorithm

Notations:

$N_{\text{wires}}$	Number of wires ('advice columns')
$N_{\text{perm}}$	Number of wires that are included in the permutation argument
$N_{\text{prev}}$	Number of previous challenges
$S_{\sigma_i}(X)$	Permutation polynomials for $0 \leq i < N_{\text{perm}}$
$pub(X)$	Public input polynomial
$w_i(X)$	Witness polynomials for $0 \leq i < N_{\text{wires}}$
$\eta_i(X)$	Previous challenges polynomials for $0 \leq i < N_{\text{prev}}$
$\omega$	$n$ -th root of unity

Proof  $\pi$  constains (here  $\mathbb{F}_r$  is a scalar field of  $\mathbb{G}$ ):

- Commitments:
  - Witness polynomials:  $w_{0,\text{comm}}, \dots, w_{N_{\text{wires}},\text{comm}} \in \mathbb{G}$
  - Permutation polynomial:  $z_{\text{comm}} \in \mathbb{G}$
  - Quotient polynomial:  $t_{\text{comm}} = (t_{1,\text{comm}}, t_{2,\text{comm}}, \dots, t_{N_{\text{perm}},\text{comm}}) \in (\mathbb{G}^{N_{\text{perm}}} \times \mathbb{G})$
- Evaluations:
  - $w_0(\zeta), \dots, w_{N_{\text{wires}}}(\zeta) \in \mathbb{F}_r$
  - $w_0(\zeta\omega), \dots, w_{N_{\text{wires}}}(\zeta\omega) \in \mathbb{F}_r$
  - $z(\zeta), z(\zeta\omega) \in \mathbb{F}_r$
  - $S_{\sigma_0}(\zeta), \dots, S_{\sigma_{N_{\text{perm}}}}(\zeta) \in \mathbb{F}_r$
  - $S_{\sigma_0}(\zeta\omega), \dots, S_{\sigma_{N_{\text{perm}}}}(\zeta\omega) \in \mathbb{F}_r$
  - $\bar{L}(\zeta\omega) \in \mathbb{F}_r$
- Opening proof  $o_\pi$ :
  - $(L_i, R_i) \in \mathbb{G} \times \mathbb{G}$  for  $0 \leq i < \text{lr\_rounds}$
  - $\delta, \hat{G} \in \mathbb{G}$
  - $z_1, z_2 \in \mathbb{F}_r$
- previous challenges:
  - $\{\eta_i(\xi_j)\}_{j, \eta_{i,\text{comm}}}$ , for  $0 \leq i < \text{prev}$

Denote multi-scalar multiplication  $\sum_{s_i \in \mathbf{s}, G_i \in \mathbf{G}} [s_i]G_i$  by  $\text{MSM}(\mathbf{s}, \mathbf{G})$  for  $l_{\mathbf{s}} = l_{\mathbf{G}}$  where  $l_{\mathbf{s}} = |\mathbf{s}|$ ,  $l_{\mathbf{G}} = |\mathbf{G}|$ .

If  $l_{\mathbf{s}} < l_{\mathbf{G}}$ , then we use only first  $l_{\mathbf{s}}$  elements of  $\mathbf{G}$

**Remark:** For simplicity, we do not use distinct proofs index  $i$  for each element in the algorithm below. For instance, we write  $pub_{\text{comm}}$  instead of  $pub_{i,\text{comm}}$ .

---

**Algorithm 1** Verification

---

**Input:**  $\pi_0, \dots, \pi_{\text{batch\_size}}$  (see 2.2)**Output:** acc or rej

1. for each  $\pi_i$ :
    - 1.1  $\text{pub}_{\text{comm}} = \text{MSM}(\mathbf{L}, \text{pub}) \in \mathbb{G}$ , where  $\mathbf{L}$  is Lagrange bases vector
    - 1.2 **random\_oracle**( $p_{\text{comm}}, \pi_i$ ):
      - 1.2.1  $H_{\mathbb{F}_q}.\text{absorb}(\text{pub}_{\text{comm}} || w_{0,\text{comm}} || \dots || w_{N_{\text{wires}},\text{comm}})$
      - 1.2.2  $\beta, \gamma = H_{\mathbb{F}_q}.\text{squeeze}()$
      - 1.2.3  $H_{\mathbb{F}_q}.\text{absorb}(z_{\text{comm}})$
      - 1.2.4  $\alpha = \phi(H_{\mathbb{F}_q}.\text{squeeze}())$
      - 1.2.5  $H_{\mathbb{F}_q}.\text{absorb}(t_{1,\text{comm}} || \dots || t_{N_{\text{perm}},\text{comm}} || \dots || \infty ||)$
      - 1.2.6  $\zeta = \phi(H_{\mathbb{F}_q}.\text{squeeze}())$
      - 1.2.7 Transform  $H_{\mathbb{F}_q}$  to  $H_{\mathbb{F}_r}$
      - 1.2.8  $H_{\mathbb{F}_r}.\text{absorb}(\text{pub}(\zeta) || w_0(\zeta) || \dots || w_{N_{\text{wires}}}(\zeta) || S_0(\zeta) || \dots || S_{N_{\text{perm}}}(\zeta))$
      - 1.2.9  $H_{\mathbb{F}_r}.\text{absorb}(\text{pub}(\zeta\omega) || w_0(\zeta\omega) || \dots || w_{N_{\text{wires}}}(\zeta\omega) || S_0(\zeta\omega) || \dots || S_{N_{\text{perm}}}(\zeta\omega))$
      - 1.2.10  $H_{\mathbb{F}_r}.\text{absorb}(\bar{L}(\zeta\omega))$
      - 1.2.11  $v = \phi(H_{\mathbb{F}_r}.\text{squeeze}())$
      - 1.2.12  $u = \phi(H_{\mathbb{F}_r}.\text{squeeze}())$
      - 1.2.13 Compute evaluation of  $\eta_i(\zeta), \eta_i(\zeta\omega)$  for  $0 \leq i < N_{\text{prev}}$
      - 1.2.14 Compute evaluation of  $\bar{L}(\zeta)$
    - 1.3  $\mathbf{f}_{\text{base}} := \{S_{\sigma_{N_{\text{perm}}-1},\text{comm}}, \text{gate}_{\text{mult},\text{comm}}, w_{0,\text{comm}}, w_{1,\text{comm}}, w_{2,\text{comm}}, q_{\text{const},\text{comm}}, \text{gate}_{\text{psdn},\text{comm}}, \text{gate}_{\text{rc},\text{comm}}, \text{gate}_{\text{ec\_add},\text{comm}}, \text{gate}_{\text{ec\_dbl},\text{comm}}, \text{gate}_{\text{ec\_endo},\text{comm}}, \text{gate}_{\text{ec\_vbase},\text{comm}}\}$
    - 1.4  $s_{\text{perm}} := (w_0(\zeta) + \gamma + \beta \cdot S_{\sigma_0}(\zeta)) \cdot \dots \cdot (w_5(\zeta) + \gamma + \beta \cdot S_{\sigma_{N_{\text{perm}}}}(\zeta))$
    - 1.5  $\mathbf{f}_{\text{scalars}} := \{-z(\zeta\omega) \cdot \beta \cdot \alpha_0 \cdot zkp(\zeta) \cdot s_{\text{perm}}, w_0(\zeta) \cdot w_1(\zeta), w_0(\zeta), w_1(\zeta), 1, s_{\text{psdn}}, s_{\text{rc}}, s_{\text{ec\_add}}, s_{\text{ec\_dbl}}, s_{\text{ec\_endo}}, s_{\text{ec\_vbase}}\}$
    - 1.6  $f_{\text{comm}} = \text{MSM}(\mathbf{f}_{\text{base}}, \mathbf{f}_{\text{scalars}})$
    - 1.7  $\bar{L}_{\text{comm}} = f_{\text{comm}} - t_{\text{comm}} \cdot (\zeta^n - 1)$
    - 1.8 **PE** is a set of elements of the form  $(f_{\text{comm}}, f(\zeta), f(\zeta\omega))$  for the following polynomials:  
 $\eta_0, \dots, \eta_{N_{\text{prev}}}, \text{pub}, w_0, \dots, w_{N_{\text{wires}}}, z, S_{\sigma_0}, \dots, S_{\sigma_{N_{\text{perm}}}}, \bar{L}$
    - 1.9  $\mathcal{P}_i = \{H_{\mathbb{F}_q}, \zeta, v, u, \mathbf{PE}, o_{\pi_i}\}$
  2. **final\_check**( $\mathcal{P}_0, \dots, \mathcal{P}_{\text{batch\_size}}$ )
-

---

**Algorithm 2** Final Check

---

**Input:**  $\pi_0, \dots, \pi_{\text{batch\_size}}$ , where  $\pi_i = \{H_{i, \mathbb{F}_q}, \zeta_i, \zeta_i \omega, v_i, u_i, \mathbf{PE}_i, o_{\pi_i}\}$

**Output:** acc or rej

1.  $\rho_1 \rightarrow \mathbb{F}_r$
  2.  $\rho_2 \rightarrow \mathbb{F}_r$
  3.  $r_0 = r'_0 = 1$
  4. for  $0 \leq i < \text{batch\_size}$ :
    - 4.1  $\text{cip}_i = \text{combined\_inner\_product}(\zeta_i, \zeta_i \omega, v_i, u_i, \mathbf{PE}_i)$
    - 4.2  $H_{i, \mathbb{F}_q}.\text{absorb}(\text{cip}_i - 2^{255})$
    - 4.3  $U_i = (H_{i, \mathbb{F}_q}.\text{squeeze}()).\text{to\_group}()$
    - 4.4 Calculate opening challenges  $\xi_{i,j}$  from  $o_{\pi_i}$
    - 4.5  $h_i(X) := \prod_{k=0}^{\log(d+1)-1} (1 + \xi_{\log(d+1)-k} X^{2^k})$ , where  $d = \text{lr\_rounds}$
    - 4.6  $b_i = h_i(\zeta) + u_i \cdot h_i(\zeta \omega)$
    - 4.7  $C_i = \sum_j v_i^j (\sum_k r_i^k f_{j, \text{comm}})$ , where  $f_{j, \text{comm}}$  from  $\mathbf{PE}_i$ .
    - 4.8  $Q_i = \sum (\xi_{i,j} \cdot L_{i,j} + \xi_{i,j}^{-1} \cdot R_j) + \text{cip}_i \cdot U_i + C_i$
    - 4.9  $c_i = \phi(H_{i, \mathbb{F}_q}.\text{squeeze}())$
    - 4.10  $r_i = r_{i-1} \cdot \rho_1$
    - 4.11  $r'_i = r'_{i-1} \cdot \rho_2$
    - 4.12 Check  $\hat{G}_i = \langle s, G \rangle$ , where  $s$  is set of  $h(X)$  coefficients.  
**Remark:** This check can be done inside the MSM below using  $r'_i$ .
  5.  $\text{res} = \sum_i r^i (c_i Q_i + \text{delta}_i - (z_{i,1}(\hat{G}_i + b_i U_i) + z_{i,2} H))$
  6. return  $\text{res} == 0$
- 

---

**Algorithm 3** Combined Inner Product

---

**Input:**  $\xi, r, f_0(\zeta_1), \dots, f_k(\zeta_1), f_0(\zeta_2), \dots, f_k(\zeta_2)$

**Output:**  $s$

1.  $s = \sum_{i=0}^k \xi^i \cdot (f_i(\zeta_1) + r \cdot f_i(\zeta_2))$
- 

## 3 Elliptic Curve Arithmetic

WIP

### 3.1 Addition

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i$	$x_1$	$y_1$	$x_2$	$y_2$	$x_3$	$y_3$	$r$	...	...	...	...	...	...	...	...

Constraints:

- $(x_2 - x_1) \cdot (y_3 + y_1) - (y_1 - y_2) \cdot (x_1 - x_3)$
- $(x_1 + x_2 + x_3) \cdot (x_1 - x_3) \cdot (x_1 - x_3) - (y_3 + y_1) \cdot (y_3 + y_1)$
- $(x_2 - x_1) \cdot r = 1$

### 3.2 Doubling and Tripling

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i$	$x_1$	$y_1$	$x_2$	$y_2$	$x_3$	$y_3$	$r_1$	$r_2$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$

Constraints:

- Doubling:
  - $4 \cdot y_1^2 \cdot (x_2 + 2 \cdot x_1) = 9 \cdot x_1^4$
  - $2 \cdot y_1 \cdot (y_2 + y_1) = (3 \cdot x_1^2) \cdot (x_1 - x_2)$
  - $y_1 \cdot r_1 = 1$
- Addition (for tripling):
  - $(x_2 - x_1) \cdot (y_3 + y_1) - (y_1 - y_2) \cdot (x_1 - x_3)$
  - $(x_1 + x_2 + x_3) \cdot (x_1 - x_3) \cdot (x_1 - x_3) - (y_3 + y_1) \cdot (y_3 + y_1)$
  - $(x_2 - x_1) \cdot r_2 = 1$

### 3.3 Variable Base Scalar Multiplication

For  $S = [r]T$ , where  $r = 2^n + k$  and  $k = [k_n \dots k_0]$ ,  $k_i \in \{0, 1\}$ : <sup>7</sup>

1.  $S = [2]T$
2. for  $i$  from  $n - 1$  to 0:
  - 2.1  $Q = k_{i+1} ? T : -T$
  - 2.2  $R = S + Q$
  - 2.3  $S = R + S$
3.  $S = k_0 ? S - T : S$

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i$	$x_T$	$y_T$	$x_S$	$y_S$	$x_P$	$y_P$	$n = 0$	$x_R$	$y_R$	$s_1$	$s_2$	$b_1$	$s_3$	$s_4$	$b_2$
$i + 1$	$s_5$	$b_3$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$x_V$	$y_V$	$s_1$	$b_1$	$s_3$	$b_2$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$i + 100$	$x_T$	$y_T$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$s_1$	$s_2$	$b_1$	$s_3$	$s_4$	$b_2$
$i + 101$	$s_5$	$b_3$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$x_V$	$y_V$	$s_1$	$b_1$	$s_3$	$b_2$

Constraints for  $i + z$ , where  $z \bmod 2 = 0$ :

- $b_1 \cdot (b_1 - 1) = 0$
- $b_2 \cdot (b_2 - 1) = 0$
- $(x_P - x_T) \cdot s_1 = y_P - (2b_1 - 1) \cdot y_T$
- $s_1^2 - s_2^2 = x_T - x_R$
- $(2 \cdot x_P + x_T - s_1^2) \cdot (s_1 + s_2) = 2y_P$
- $(x_P - x_R) \cdot s_2 = y_R + y_P$
- $(x_R - x_T) \cdot s_3 = y_R - (2b_2 - 1) \cdot y_T$
- $s_3^2 - s_4^2 = x_T - x_S$
- $(2 \cdot x_R + x_T - s_3^2) \cdot (s_3 + s_4) = 2 \cdot y_R$
- $(x_R - x_S) \cdot s_4 = y_S + y_R$
- $n = 32 \cdot \text{next}(n) + 16 \cdot b_1 + 8 \cdot b_2 + 4 \cdot \text{next}(b_1) + 2 \cdot \text{next}(b_2) + \text{next}(b_3)$

Constraints for  $i + z$ , where  $z \bmod 2 = 1$ :

<sup>7</sup>Using the results from <https://arxiv.org/pdf/math/0208038.pdf>

- $b_1 \cdot (b_1 - 1) = 0$
- $b_2 \cdot (b_2 - 1) = 0$
- $b_3 \cdot (b_3 - 1) = 0$
- $(x_P - x_T) \cdot s_1 = y_P - (2b_1 - 1) \cdot y_T$
- $(2 \cdot x_P + x_T - s_1^2) \cdot ((x_P - x_R) \cdot s_1 + y_R + y_P) = (x_P - x_R) \cdot 2y_P$
- $(y_R + y_P)^2 = (x_P - x_R)^2 \cdot (s_1^2 - x_T + x_R)$
- $(x_T - x_R) \cdot s_3 = (2b_2 - 1) \cdot y_T - y_R$
- $(2x_R - s_3^2 + x_T) \cdot ((x_R - x_V) \cdot s_3 + y_V + y_R) = (x_R - x_V) \cdot 2y_R$
- $(y_V + y_R)^2 = (x_R - x_V)^2 \cdot (s_3^2 - x_T + x_V)$
- $(x_T - x_V) \cdot s_5 = (2b_3 - 1) \cdot y_T - y_V$
- $(2x_V - s_5^2 + x_T) \cdot ((x_V - x_S) \cdot s_5 + y_S + y_V) = (x_V - x_S) \cdot 2y_V$
- $(y_S + y_V)^2 = (x_V - x_S)^2 \cdot (s_5^2 - x_T + x_S)$

### 3.4 Variable Base Endo-Scalar Multiplication

For  $S = [r]T$ , where  $r = [r_n \dots r_0]$  and  $r_i \in \{0, 1\}$ :<sup>8</sup>

1.  $S = [2](\phi(T) + T)$
2. for  $i$  from  $\frac{\lambda}{2} - 1$  to 0:
  - 2.1  $Q = r_{2i+1} ? \phi([2r_{2i} - 1]T) : [2r_{2i} - 1]T$
  - 2.2  $R = S + Q$
  - 2.3  $S = R + S$

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i$	$x_T$	$y_T$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$s_1$	$s_3$	$b_1$	$b_2$	$b_3$	$b_4$
$i + 1$	$s_5$	$b_3$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$s_1$	$s_3$	$b_1$	$b_2$	$b_3$	$b_4$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$i + 62$	$x_T$	$y_T$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$s_1$	$s_3$	$b_1$	$b_2$	$b_3$	$b_4$
$i + 63$	$s_5$	$b_3$	$x_S$	$y_S$	$x_P$	$y_P$	$n$	$x_R$	$y_R$	$s_1$	$s_3$	$b_1$	$b_2$	$b_3$	$b_4$

Constraints:

- $b_1 \cdot (b_1 - 1) = 0$
- $b_2 \cdot (b_2 - 1) = 0$
- $b_3 \cdot (b_3 - 1) = 0$
- $b_4 \cdot (b_4 - 1) = 0$
- $((1 + (\text{endo} - 1) \cdot b_2) \cdot x_T - x_P) \cdot s_1 = (2 \cdot b_1 - 1) \cdot y_T - y_P$
- $(2 \cdot x_P - s_1^2 + (1 + (\text{endo} - 1) \cdot b_2) \cdot x_T) \cdot ((x_P - x_R) \cdot s_1 + y_R + y_P) = (x_P - x_R) \cdot 2 \cdot y_P$
- $(y_R + y_P)^2 = (x_P - x_R)^2 \cdot (s_1^2 - (1 + (\text{endo} - 1) \cdot b_2) \cdot x_T + x_R)$
- $((1 + (\text{endo} - 1) \cdot b_2) \cdot x_T - x_R) \cdot s_3 = (2 \cdot b_3 - 1) \cdot y_T - y_R$
- $(2 \cdot x_R - s_3^2 + (1 + (\text{endo} - 1) \cdot b_4) \cdot x_T) \cdot ((x_R - x_S) \cdot s_3 + y_S + y_R) = (x_R - x_S) \cdot 2 \cdot y_R$
- $(y_S + y_R)^2 = (x_R - x_S)^2 \cdot (s_3^2 - (1 + (\text{endo} - 1) \cdot b_4) \cdot x_T + x_S)$
- $n = 16 \cdot \text{next}(n) + 8 \cdot b_1 + 4 \cdot b_2 + 2 \cdot b_3 + b_4$

### 3.5 Fixed-base scalar multiplication circuit

We precompute all values  $w(B, s, k) = (k_i + 2) \cdot 8^s B$ , where  $k_i \in \{0, \dots, 7\}$ ,  $s \in \{0, \dots, 83\}$  and  $w(B, s, k) = (k_i \cdot 8^s - \sum_{j=0}^{84} 8^{j+1}) \cdot B$ , where  $k_i \in \{0, \dots, 7\}$ ,  $s = 84$ .

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i$	$b_0$	$b_1$	$b_2$	$b_3$	$b_4$	$b_5$	$u_0$	$u_1$	$v_0$	$v_1$	$x_1$	$y_1$	$x_2$	$y_2$	$acc$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$i + 42$	$b_0$	$b_1$	$b_2$	$u_0$	$v_0$	$x_w$	$y_w$	$\alpha$	$\beta$	$\gamma$	$\delta$	$\lambda$	$-$	$-$	$b$

<sup>8</sup>Using the results from <https://eprint.iacr.org/2019/1021.pdf>

Define the following functions:

1.  $\phi_1 : (x_1, x_2, x_3, x_4) \mapsto$   
 $x_3 \cdot (-u'_0 \cdot x_2 \cdot x_1 + u'_0 \cdot x_1 + u'_0 \cdot x_2 - u'_0 + u'_2 \cdot x_1 \cdot x_2 - u'_2 \cdot x_2 + u'_4 \cdot x_1 \cdot x_2 - u'_4 \cdot x_2 - u'_6 \cdot x_1 \cdot x_2 +$   
 $u'_1 \cdot x_2 \cdot x_1 - u'_1 \cdot x_1 - u'_1 \cdot x_2 + u'_1 - u'_3 \cdot x_1 \cdot x_2 + u'_3 \cdot x_2 - u'_5 \cdot x_1 \cdot x_2 + u'_5 \cdot x_2 + u'_7 \cdot x_1 \cdot x_2) - (x_4 -$   
 $u'_0 \cdot x_2 \cdot x_1 + u'_0 \cdot x_1 + u'_0 \cdot x_2 - u'_0 + u'_2 \cdot x_1 \cdot x_2 - u'_2 \cdot x_2 + u'_4 \cdot x_1 \cdot x_2 - u'_4 \cdot x_2 - u'_6 \cdot x_1 \cdot x_2)$
2.  $\phi_2 : (x_1, x_2, x_3, x_4) \mapsto$   
 $x_3 \cdot (-v'_0 \cdot x_2 \cdot x_1 + v'_0 \cdot x_1 + v'_0 \cdot x_2 - v'_0 + v'_2 \cdot x_1 \cdot x_2 - v'_2 \cdot x_2 + v'_4 \cdot x_1 \cdot x_2 - v'_4 \cdot x_2 - v'_6 \cdot x_1 \cdot x_2 + v'_1 \cdot$   
 $x_2 \cdot x_1 - v'_1 \cdot x_1 - v'_1 \cdot x_2 + v'_1 - v'_3 \cdot x_1 \cdot x_2 + v'_3 \cdot x_2 - v'_5 \cdot x_1 \cdot x_2 + v'_5 \cdot x_2 + v'_7 \cdot x_1 \cdot x_2) - (x_4 - v'_0 \cdot$   
 $x_2 \cdot x_1 + v'_0 \cdot x_1 + v'_0 \cdot x_2 - v'_0 + v'_2 \cdot x_1 \cdot x_2 - v'_2 \cdot x_2 + v'_4 \cdot x_1 \cdot x_2 - v'_4 \cdot x_2 - v'_6 \cdot x_1 \cdot x_2)$

Constraints:

- For  $i + 0$ :
  - $b_i \cdot (b_i - 1) = 0$ , where  $i \in \{0, \dots, 5\}$
  - $\phi_1(b_0, b_1, b_2, u_0) = 0$ , where  $(u'_i, v'_i) = w(B, 0, i)$
  - $\phi_1(b_3, b_4, b_5, u_1) = 0$ , where  $(u'_i, v'_i) = w(B, 1, i)$
  - $\phi_2(b_0, b_1, b_2, v_0) = 0$ , where  $(u'_i, v'_i) = w(B, 0, i)$
  - $\phi_2(b_3, b_4, b_5, v_1) = 0$ , where  $(u'_i, v'_i) = w(B, 1, i)$
  - $acc = b_0 + b_1 \cdot 2 + b_2 \cdot 2^2 + b_3 \cdot 2^3 + b_4 \cdot 2^4 + b_5 \cdot 2^5$
  - $(x_1, y_1) = (u_0, v_0)$
  - $(x_2, y_2) = (x_1, y_1) + (u_1, v_1)$  incomplete addition, where  $x_1 \neq u_1$
- For  $i + z$ ,  $z \in 1, \dots, 41$ :
  - $b_i \cdot (b_i - 1) = 0$ , where  $i \in \{0, \dots, 5\}$
  - $\phi_1(b_0, b_1, b_2, u_0) = 0$ , where  $(u'_i, v'_i) = w(B, z \cdot 2, i)$
  - $\phi_1(b_3, b_4, b_5, u_1) = 0$ , where  $(u'_i, v'_i) = w(B, z \cdot 2 + 1, i)$
  - $\phi_2(b_0, b_1, b_2, v_0) = 0$ , where  $(u'_i, v'_i) = w(B, z \cdot 2, i)$
  - $\phi_2(b_3, b_4, b_5, v_1) = 0$ , where  $(u'_i, v'_i) = w(B, z \cdot 2 + 1, i)$
  - $acc = b_0 + b_1 \cdot 2 + b_2 \cdot 2^2 + b_3 \cdot 2^3 + b_4 \cdot 2^4 + b_5 \cdot 2^5 + acc_{prev} \cdot 2^6$
  - $(x_1, y_1) = (u_0, v_0) + (x_2, y_2)_{prev}$  incomplete addition, where  $u_0 \neq x_2$
  - $(x_2, y_2) = (x_1, y_1) + (u_1, v_1)$  incomplete addition, where  $x_1 \neq u_1$
- For  $i + 42$ :
  - $b_i \cdot (b_i - 1) = 0$ , where  $i \in \{0, \dots, 2\}$
  - $\phi_1(b_0, b_1, b_2, u_0) = 0$ , where  $(u'_i, v'_i) = w(B, 84, i)$
  - $\phi_2(b_0, b_1, b_2, v_0) = 0$ , where  $(u'_i, v'_i) = w(B, 84, i)$
  - $b = b_0 + b_1 \cdot 2 + b_2 \cdot 2^2 + acc_{prev} \cdot 2^3$
  - $(x_w, y_w) = (u_0, v_0) + (x_2, y_2)_{prev}$  complete addition from [Orchard](#)

## 4 Multi-Scalar Multiplication Circuit

**WIP**

Input:  $G_0, \dots, G_{k-1} \in \mathbb{G}, s_0, \dots, s_{k-1} \in \mathbb{F}_r$ , where  $\mathbb{F}_r$  is scalar field of  $\mathbb{G}$ .

Output:  $S = \sum_{i=0}^k s_i \cdot G_i$

### 4.1 Naive Algorithm

Using endomorphism:

1.  $A = \infty$
2. for  $j$  from 0 to  $k - 1$ :
  - 2.1  $r := s_j, T := G_j$
  - 2.2  $S = [2](\phi(T) + T)$
  - 2.3 for  $i$  from  $\frac{\lambda}{2} - 1$  to 0:

- 2.3.1  $Q = r_{2i+1} ? \phi([2r_{2i} - 1]T) : [2r_{2i} - 1]T$
- 2.3.2  $R = S + Q$
- 2.3.3  $S = R + S$
- 2.4  $A = A + S$

$$\text{rows} \approx k \cdot (\text{sm\_rows} + 1 + 2) \approx 67k,$$

where **sm\_rows** is the number of rows in the scalar multiplication circuit.

#### Without endomorphism:

- 1.  $A = \infty$
- 2. for  $j$  from 0 to  $k - 1$ :
  - 2.1  $r := s_j, T := G_j$
  - 2.2  $S = [2]T$
  - 2.3 for  $i$  from  $n - 1$  to 0:
    - 2.3.1  $Q = k_{i+1} ? T : -T$
    - 2.3.2  $R = S + Q$
    - 2.3.3  $S = R + S$
  - 2.4  $S = k_0 ? S - T : S$
  - 2.5  $A = A + S$

$$\text{rows} \approx k \cdot (\text{sm\_rows} + 1 + 1) \approx 105k,$$

where **sm\_rows** is the number of rows in the scalar multiplication circuit.

## 4.2 Simultaneous Doubling

#### Using endomorphism:

- 1.  $A = \sum_{j=0}^k [2](\phi(G_j) + G_j)$
- 2. for  $i$  from  $\frac{\lambda}{2} - 1$  to 0:
  - 2.1 for  $j$  from 0 to  $k - 1$ :
    - 2.1.1  $r := s_j, T := G_j$
    - 2.1.2  $Q = r_{2i+1} ? \phi([2r_{2i} - 1]T) : [2r_{2i} - 1]T$
    - 2.1.3  $A = A + Q$
  - 2.2 if  $i \neq 0$ :
    - 2.2.1  $A = 2 \cdot A$

$$\text{rows} \approx \frac{\lambda}{2} \cdot (k \cdot \text{add\_rows} + \text{dbl\_rows}) + 2k \approx 64 \cdot (k + 1) \approx 66k + 64,$$

where

- **add\_rows** is the number of rows in the addition circuit.
- **dbl\_rows** is the number of rows in the doubling circuit.



**Without endomorphism:**

1.  $A = \sum_{j=0}^k [2]G_j$
2. for  $i$  from  $n - 1$  to 0:
  - 2.1 for  $j$  from 0 to  $k - 1$ :
    - 2.1.1  $r := s_j, T := G_j$
    - 2.1.2  $Q = k_{i+1} ? T : -T$
    - 2.1.3  $A = A + Q$
  - 2.2 if  $i \neq 0$ :
    - 2.2.1  $A = 2 \cdot A$
3.  $A = A + \sum_{j=0}^k [1 - s_{j,0}]G_j$

$$\text{rows} \approx \frac{2}{3}n \cdot (k \cdot \text{add\_rows} + \text{dbl\_rows}) + k \approx 103 \cdot (k + 1) + 2k \approx 104k + 103,$$

where

- **add\_rows** is the number of rows in the addition circuit.
- **dbl\_rows** is the number of rows in the doubling circuit.

## 5 Poseidon Circuit

**WIP**

Mina uses Poseidon hash with width = 3. Therefore, each permutation state is represented by 3 elements and each row contains 5 states.

Denote  $i$ -th permutation state by  $T_i = (T_{i,0}, T_{i,1}, T_{i,2})$ .

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i$	$T_{0,0}$	$T_{0,1}$	$T_{0,2}$	$T_{4,0}$	$T_{4,1}$	$T_{4,2}$	$T_{1,0}$	$T_{1,1}$	$T_{1,2}$	$T_{2,0}$	$T_{2,1}$	$T_{2,2}$	$T_{3,0}$	$T_{3,1}$	$T_{3,2}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$i + 10$	$T_{50,0}$	$T_{50,1}$	$T_{50,2}$	$T_{54,0}$	$T_{54,1}$	$T_{54,2}$	$T_{51,0}$	$T_{51,1}$	$T_{51,2}$	$T_{52,0}$	$T_{52,1}$	$T_{52,2}$	$T_{53,0}$	$T_{53,1}$	$T_{53,2}$
$i + 11$	$T_{55,0}$	$T_{55,1}$	$T_{55,2}$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$

State change constraints:

$$\text{STATE}(i + 1) = \text{STATE}(i)^\alpha \cdot \text{MDS} + \text{RC}$$

Denote the index of the first state in the row by **start** (e.g. **start** = 50 for 10-th row). We can expand the previous formula to:

- For  $i$  from **start** to **start** + 5:
  - $T_{i+1,0} = T_{i,0}^5 \cdot \text{MDS}[0][0] + T_{i,1}^5 \cdot \text{MDS}[0][2] + T_{i,2}^5 \cdot \text{MDS}[0][2] + \text{RC}_{i+1,0}$
  - $T_{i+1,1} = T_{i,0}^5 \cdot \text{MDS}[1][0] + T_{i,1}^5 \cdot \text{MDS}[1][2] + T_{i,2}^5 \cdot \text{MDS}[1][2] + \text{RC}_{i+1,1}$
  - $T_{i+1,2} = T_{i,0}^5 \cdot \text{MDS}[2][2] + T_{i,1}^5 \cdot \text{MDS}[2][2] + T_{i,2}^5 \cdot \text{MDS}[2][2] + \text{RC}_{i+1,2}$

Notice that the constraints above include the state from the next row (**start** + 5).

## 6 Other Circuits

**WIP**

## 6.1 Combined Inner Product

$$\sum_{i=0}^k \xi^i \cdot (f_i(\zeta_1) + r \cdot f_i(\zeta_2))$$

Row	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$i + 0$	$f_1$	$f'_1$	$f_2$	$f'_2$	<b>acc</b>	$\xi$	$\xi_{\text{acc}}$	$s_1$	$s_2$	$\xi'_{\text{acc}}$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$
$i + 1$	$f_3$	$f'_3$	$f_4$	$f'_4$	<b>acc</b>	$r$	$\xi_{\text{acc}}$	$s_1$	$s_2$	$\xi'_{\text{acc}}$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$i + \lceil \frac{k}{2} \rceil - 1$	$f_{k-3}$	$f'_{k-3}$	$f_{k-2}$	$f'_{k-2}$	<b>acc</b>	$\xi$	$\xi_{\text{acc}}$	$s_1$	$s_2$	$\xi'_{\text{acc}}$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$
$i + \lceil \frac{k}{2} \rceil$	$f_{k-1}$	$f'_{k-1}$	$f_k$	$f'_k$	<b>acc</b>	$r$	$\xi_{\text{acc}}$	$s_1$	$s_2$	$\xi'_{\text{acc}}$	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$

Constraints for  $i + z$ , where  $z \bmod 2 = 0$ :

- $(w_0 + w_1 \cdot \text{next}(w_5)) \cdot w_6 = w_7$
- $(w_2 + w_3 \cdot \text{next}(w_5)) \cdot w_9 = w_8$
- $w_5 \cdot w_6 = w_9$
- $w_5 \cdot w_9 = \text{next}(w_9)$
- $w_5 \cdot \text{next}(w_9) = \text{next}(w_5)$
- $w_4 + w_7 + w_8 + \text{next}(w_7) + \text{next}(w_8) = \text{next}(w_4)$

Constraints for  $i + z$ , where  $z \bmod 2 = 1$ :

- $(w_0 + w_1 \cdot w_5) \cdot w_9 = w_7$
- $(w_2 + w_3 \cdot w_5) \cdot w_6 = w_8$

## References