

令和 6 年度
修士学位論文

論文用テンプレート

〇〇所属

〇〇課程 〇〇専攻

〇〇分野

指導教員 〇〇 〇〇教授

令和〇年入学

学籍番号 82313206

氏名 八木颯仁

概要

We study a family of elliptic curves $y^2 = x(x - a^2)(x + b^2)$, where (a, b, c) are Pythagorean triples. This is the family of the Frey curves of degree 2. We can 1-parameterize Pythagorean triples by rational numbers and consider the family as an elliptic curve over a function field.

$$E_{1,s} : y^2 = x(x - 4s^2)(x + (s^2 - 1)^2) \quad (1)$$

It is known that the generic rank of the Mordell-Weil group of $E_{1,s}$ over $\overline{\mathbb{Q}}(s)$ is 0. We found an infinite subfamily of $E_{1,s}$ whose Mordell-Weil rank over $\overline{\mathbb{Q}}(s)$ is 1, which means that there are infinitely many $s \in \overline{\mathbb{Q}}$ such that the Mordell-Weil group of $E_{1,s}$ has positive rank over $\overline{\mathbb{Q}}$.

We use the theory of elliptic surfaces to prove it. Each elliptic curve over a function field corresponds to an elliptic surface. The Shioda-Tate formula gives the relation between the Mordell-Weil rank and the Néron-Severi rank of elliptic surfaces. We compute the types of special fibers of the elliptic surfaces and consider the upper bound of the rank of the Néron-Severi group.

目次

1.	Introduction	1
2.	Preliminaries	2
3.	Proof of Theorem 1.0.1	3
4.	The Generic Rank of $E_{2,t}$	4
5.	Reductions	7
5.1	$E_{0,u}^{(1+3u)}$	7
参考文献		8

1. Introduction

Theorem 1.0.1. Let

$$E_{1,s} : y^2 = x(x - 4s^2)(x + (s^2 - 1)^2) \quad (1.1)$$

be an elliptic curve over $\overline{\mathbb{Q}}(s)$. Then, the Mordell-Weil group

$$E_{1,s}(\overline{\mathbb{Q}}(s)) \cong \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}, \quad (1.2)$$

especially the rank is 0. The torsion subgroup is generated by

$$T_1 := (2s(s+1)^2, 2s(s+1)^2(s^2+1)), \quad (1.3)$$

$$T_2 := (2\sqrt{-1}s(s^2-1), 2\sqrt{-1}s(s+\sqrt{-1})^2(s^2-1)). \quad (1.4)$$

Corollary 1.0.2.

$$E_{1,s}(\mathbb{Q}(s)) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z} \quad (1.5)$$

is generated by T_1 and $2T_2 = (0, 0)$.

Remark 1.0.3. 課題研究では、多項式の解の非存在から背理法を用いて 8-torsion point が存在しないことを示した。

Theorem 1.0.4. Let

$$E_{2,t} : y^2 = x(x - 4s^2)(x + (s^2 - 1)^2), \quad s = \frac{2t}{t^2 - 3} \quad (1.6)$$

be an elliptic curve over $\overline{\mathbb{Q}}(t)$. Then, the Mordell-Weil group

$$E_{2,t}(\overline{\mathbb{Q}}(t)) \cong \mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}, \quad (1.7)$$

especially the rank is 1. The torsion subgroup is generated by T_1 and T_2 in Theorem 1.0.1 and the free part is generated by

$$\left(s^2 - 1, \sqrt{-1}s(s^2 - 1) \frac{t^2 + 3}{t^2 - 3} \right). \quad (1.8)$$

2. Preliminaries

In order to get the lower bound of the rank of the Mordell-Weil group, finding generators is enough. It is more difficult to get the upper bound of the rank. The following theorem behaves a key role in the proof of the main theorem.

Theorem 2.0.1. (Shioda-Tate formula, [1, Corollary 5.3]) Let C be a smooth irreducible projective curve over an algebraically closed field k and E an elliptic curve over a function field $k(C)$. Let $\mathcal{E} \rightarrow C$ be the Néron model of E . Let $R \subset C$ be the set of points where the special fiber of \mathcal{E} is singular. For each $v \in R$, let m_v be the number of components of the special fiber of \mathcal{E} at v . Let $\rho(\mathcal{E})$ denote the rank of the Néron-Severi group of \mathcal{E} . Then, we have

$$\rho(\mathcal{E}) = 2 + \sum_{v \in R} (m_v - 1) + \text{rank}(E(k(C))). \quad (2.1)$$

We can calculate R and m_v by Tate's algorithm, but it is still difficult to determine $\rho(\mathcal{E})$. We have the following theorem to get the upper bound of $\rho(\mathcal{E})$.

Theorem 2.0.2.

$$\rho(\mathcal{E}) \leq \frac{5}{6}e(\tilde{S}) + 2, \quad (2.2)$$

$$e(\tilde{S}) := \sum_{v \in R} e(F_v). \quad (2.3)$$

where $e(\tilde{S})$ is the Euler number, $e(F_v)$ is the local Euler number of the special fiber of \mathcal{E} at v for each $v \in R$ and

$$e(F_v) = \begin{cases} m_v & \text{if the fiber has multiplicative reduction,} \\ m_v + 1 & \text{if the fiber has additive reduction.} \end{cases} \quad (2.4)$$

証明 TODO: Naskrencki の PhD の 2.2.19(ii), 2.2.9, 2.2.10 (やその引用元) を引用するオイラー数については [2, pp. 136-137 付録 2], [3, p.14 Table II] □

Theorem 2.0.3. ([4, Lem.3.5])

$$E(\overline{\mathbb{Q}}(s))_{\text{tors}} \hookrightarrow \prod_{v \in R} G(F_v) \quad (2.5)$$

where $G(F_v)$ is the group generated by all simple components of the fiber at v . If F_v is of type I_n in Kodaira's symbol, then $G(F_v) \cong \mathbb{Z}/n\mathbb{Z}$.

証明 TODO

□

3. Proof of Theorem 1.0.1

証明 of Theorem 1.0.1

$$\Delta_{E_{1,s}} = 256s^4(s+1)^4(s-1)^4(s^2+1)^4 \quad (3.1)$$

Table 3.1 Singular fibers of $E_{1,s}$

Place	Type	m_v
$s = 0$	I_4	4
$s = \pm 1$	I_4	4
$s = \pm\sqrt{-1}$	I_4	4
$s = \infty$	I_4	4

$$e(\mathcal{E}_{1,s}) = 24 \quad (3.2)$$

したがって $\mathcal{E}_{1,s}$ は K3 曲面であり． $\rho(\mathcal{E}_{1,s}) \leq 20$ である． Theorem 2.0.1 より

$$\text{rank}(E_{1,s}) = 0 \quad (3.3)$$

As for the torsion subgroup, we have

$$E_{1,s}(\overline{\mathbb{Q}}(s))[2] = \{\mathcal{O}, (0, 0), (4s^2, 0), (-(s^2 - 1)^2, 0)\} \quad (3.4)$$

$$2T_1 = (4s^2, 0) \quad (3.5)$$

$$2T_2 = (0, 0) \quad (3.6)$$

Theorem 2.0.3 より

$$E_{2,t}(\overline{\mathbb{Q}}(t))_{\text{tors}} \hookrightarrow (\mathbb{Z}/4\mathbb{Z})^{12} \quad (3.7)$$

なので位数 8 の点は存在しない.

□

4. The Generic Rank of $E_{2,t}$

In order to prove Theorem 1.0.4, Theorem 2.0.2 is not enough to get the exact upper bound of the ranks of the Néron-Severi group. Actually, we get $\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) \leq 2$ from Theorem 2.0.1 and Theorem 2.0.2.

Table 4.1 Singular fibers of $E_{2,t}$

Place	Type	m_v
$t = 0$	I_4	4
$t = \pm 1$	I_4	4
$t = \pm 3$	I_4	4
$t = \pm\sqrt{3}$	I_4	4
$t^4 - 2t^2 + 9 = 0$	I_4	4
$t = \infty$	I_4	4

証明

$$\Delta_{E_{2,t}} = 4096t^4(t-1)^4(t+1)^4(t-3)^4(t+3)^4(t^2-3)^4(t^4-2t^2+9)^4 \quad (4.1)$$

$$e(\mathcal{E}_{2,t}) = 48 \quad (4.2)$$

TODO: $\rho(\mathcal{E}_{2,t}) \leq 40$ である. Theorem 2.0.1 より

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) \leq 2 \quad (4.3)$$

□

On the other hand, we have only one point of infinite order in $E_{2,t}(\overline{\mathbb{Q}}(t))$. Now, our goal is to show the upper bound of the rank of $E_{2,t}(\overline{\mathbb{Q}}(t))$ is 1.

We use another method to estimate the upper bound of the rank of Néron-Severi group, which we will explain in Chapter 5. Beforehand, we express the rank of $E_{2,t}(\overline{\mathbb{Q}}(t))$ in terms of ranks of elliptic curves with lower order coefficients in the Weierstrass equations to make the later computation feasible.

Definition 4.0.1. Let C be a smooth curve over an algebraically closed field k . Let E be an elliptic curve over a function field $k(C)$ given by the Weierstrass equation

$$E : y^2 = x^3 + a_2x^2 + a_4x + a_6 \quad (4.4)$$

where $a_2, a_4, a_6 \in k(C)$. For a fixed $u \in k(C)^*$, we denote

$$E^{(u)} : uy^2 = x^3 + a_2x^2 + a_4x + a_6 \quad (4.5)$$

to be the quadratic twist of E by u .

Theorem 4.0.2. ([5, Exercise 10.16]) Let E be an elliptic curve over a function field $k(C)$ and $u \in k(C)^*$. Then, the following equation holds

$$\text{rank } E(k(C)(\sqrt{u})) = \text{rank } E(k(C)) + \text{rank } E^{(u)}(k(C)). \quad (4.6)$$

Theorem 4.0.3. Let

$$E_{0,u} : y^2 = x(x - 4u)(x + (u - 1)^2) \quad (4.7)$$

be an elliptic curve over $\overline{\mathbb{Q}}(u)$. Then, we have

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) = \text{rank } E_{1,s}(\overline{\mathbb{Q}}(s)) + \text{rank } E_{1,s}^{(1+3s^2)}(\overline{\mathbb{Q}}(s)), \quad (4.8)$$

$$\text{rank } E_{1,s}^{(1+3s^2)}(\overline{\mathbb{Q}}(s)) = \text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) + \text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)). \quad (4.9)$$

Therefore, we have

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) = \text{rank } E_{1,s}(\overline{\mathbb{Q}}(s)) + \text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) + \text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)). \quad (4.10)$$

証明 Since solving $s = \frac{2t}{t^2-3}$ for t yields $t = \frac{1 \pm \sqrt{1+3s^2}}{s}$, we have

$$E_{2,t}(\overline{\mathbb{Q}}(t)) = E_{1,s}(\overline{\mathbb{Q}}(s)(\sqrt{1+3s^2})) \quad (4.11)$$

By Theorem 4.0.2, we get (4.8). Similarly, $E_{1,s}$ is obtained by substituting $u = s^2$ into $E_{0,u}$, so we have

$$E_{1,s}^{(1+3s^2)}(\overline{\mathbb{Q}}(s)) = E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)(\sqrt{u})), \quad (4.12)$$

then we get (4.9). □

Theorem 4.0.4. TODO

$$\text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) \leq 1 \quad (4.13)$$

証明

$$\Delta(E_{0,u}^{(1+3u)}) = 256u^2(u-1)^4(u+1)^4(3u+1)^6 \quad (4.14)$$

Table 4.2 Singular fibers of $E_{0,u}^{(1+3u)}$

Place	Type	m_v
$u = 0$	I_2	2
$u = \pm 1$	I_4	4
$u = -\frac{1}{3}$	I_0^*	5
$u = \infty$	I_2^*	7

$$e(\mathcal{E}_{0,u}^{(1+3u)}) = 24 \quad (4.15)$$

Theorem 2.0.1 からは

$$\text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) \leq 1 \quad (4.16)$$

□

Theorem 4.0.5.

$$\text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)) = 1 \quad (4.17)$$

証明

$$(u-1, \sqrt{-1}(u-1)) \in E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)) \quad (4.18)$$

より rank は正である.

$$\Delta(E_{0,u}^{(u(1+3u))}) = 256u^8(u-1)^4(u+1)^4(3u+1)^6 \quad (4.19)$$

上と同様に

$$\text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)) \leq 1 \quad (4.20)$$

□

Table 4.3 Singular fibers of $E_{0,u}^{(u(1+3u))}$

Place	Type	m_v
$u = 0$	I_2^*	7
$u = \pm 1$	I_4	4
$u = -\frac{1}{3}$	I_0^*	5
$u = \infty$	I_2	2

5. Reductions

5.1 $E_{0,u}^{(1+3u)}$

We denote by $\mathcal{E}_{0,u}^{(1+3u)} \rightarrow \mathbb{P}^1$ the elliptic surface with the generic fiber $E_{0,u}^{(1+3u)}$.

Table 4.2 より K3 なので

$$\dim_{\mathbb{Q}_l} H_{\text{ét}}^2(\tilde{S}, \mathbb{Q}_l) = 22 \quad (5.1)$$

である. Let V be the subspace of $\text{NS}(\tilde{S})$ generated by the singular fibers and the zero section. Then V is of rank 19, on which the Frobenius automorphism acts by multiplication by p .

$$\text{char}(\Phi_{\tilde{S}}^*|V) = (x - 5)^{19} \quad (5.2)$$

Note that all the multiplicative fibers are split.

$$t_m := \text{Tr}((\Phi_{\tilde{S}, H_{\text{ét}}^2/V}^*)^m) = \#\tilde{S}(\mathbb{F}_{5^m}) - 1 - 5^{2m} - 19 \cdot 5^m \quad (5.3)$$

m	1	2	3
$\#\tilde{S}(\mathbb{F}_{5^m})$	120	1080	18264
t_m	-1	-21	263

$$\text{char}(\Phi_{\tilde{S}, H_{\text{ét}}^2/V}^*) = x^3 + x^2 + 11x - 77 \quad (5.4)$$

参考文献

- [1] T. Shioda. On the Mordell-Weil Lattices. *Commentarii Mathematici Universitatis Sancti Pauli* 39, (1990).
- [2] 徹. 塩田. Mordell-Weil lattice の理論とその応用. *jpn. 東京大学数理科学セミナーノート ; 1.* 東京: Graduate school of mathematical sciences, 1993.
- [3] K. Kodaira. On Compact Analytic Surfaces, III. *Annals of Mathematics* 78.1, pp. 1–40, (1963). URL: <http://www.jstor.org/stable/1970500> (accessed on 2025-01-12).
- [4] B. Naskręcki. Mordell-Weil ranks of families of elliptic curves associated to Pythagorean triples. *eng. Acta Arithmetica* 160.2, pp. 159–183, (2013). URL: <http://eudml.org/doc/279803>.
- [5] J. H. Silverman. The arithmetic of elliptic curves. *eng. 2nd ed. Graduate texts in mathematics ; 106.* New York: Springer, 2009.