

令和6年度
修士学位論文

論文用テンプレート

〇〇所属

〇〇課程 〇〇専攻

〇〇分野

指導教員 〇〇 〇〇教授

令和〇年入学

学籍番号 82313206

氏名 八木颯仁

概要

We study a family of elliptic curves $y^2 = x(x - a^2)(x + b^2)$, where (a, b, c) are Pythagorean triples. This is the family of the Frey curves of degree 2. We can one-parameterize Pythagorean triples by rational numbers and consider the family as an elliptic curve over a function field.

$$E_{1,s} : y^2 = x(x - 4s^2)(x + (s^2 - 1)^2) \quad (1)$$

It is known that the generic rank of the Mordell-Weil group of $E_{1,s}$ over $\overline{\mathbb{Q}}(s)$ is 0. We found an infinite subfamily of $E_{1,s}$ whose Mordell-Weil rank over $\overline{\mathbb{Q}}(s)$ is 1, which means that there are infinitely many $s \in \overline{\mathbb{Q}}$ such that the Mordell-Weil group of $E_{1,s}$ has positive rank over $\overline{\mathbb{Q}}$.

We use the theory of elliptic surfaces to prove it. Each elliptic curve over a function field corresponds to an elliptic surface. The Shioda-Tate formula gives the relation between the Mordell-Weil rank and the Néron-Severi rank of elliptic surfaces. We compute the types of special fibers of the elliptic surfaces and consider an upper bound of the rank of the Néron-Severi group.

It is relatively easy to prove that the generic rank of the subfamily is less than 2. In order to prove that the generic rank is exactly 1, we calculate the characteristic polynomial of the action of the Frobenius automorphism on the second l -adic étale cohomology group to get the tight upper bound of the rank of the Néron-Severi group.

目次

1.	Introduction	1
2.	Preliminaries	2
3.	Proof of Theorem 1.0.1	4
4.	The Generic Rank of $E_{2,t}$	5
5.	Reductions	8
	Acknowledgments	11
	参考文献	12

1. Introduction

Let (a, b, c) be a Pythagorean triple, i.e., $a^2 + b^2 = c^2$, $abc \neq 0$, and $a, b, c \in \mathbb{Z}$, and consider the elliptic curve defined by a Weierstrass equation

$$y^2 = x(x - a^2)(x + b^2), \quad (1.1)$$

which we call the Frey curve of degree 2. There are several examples of this form with rank 0 and 1.

We can parameterize the Frey curves of degree 2 by $s \in \mathbb{Q}$, then the family of the Frey curves of degree 2 is equivalent to a family of elliptic curves with the following Weierstrass equation

$$E_{1,s} : y^2 = x(x - 4s^2)(x + (s^2 - 1)^2). \quad (1.2)$$

We consider $E_{1,s}$ as an elliptic curve over a function field $\overline{\mathbb{Q}}(s)$. First, we determine the Mordell-Weil group of $E_{1,s}$.

Theorem 1.0.1. The Mordell-Weil group of $E_{1,s}$ over $\overline{\mathbb{Q}}(s)$ satisfies

$$E_{1,s}(\overline{\mathbb{Q}}(s)) \cong \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}, \quad (1.3)$$

especially the rank is 0. The torsion subgroup is generated by

$$T_1 := (2s(s+1)^2, 2s(s+1)^2(s^2+1)), \quad (1.4)$$

$$T_2 := (2\sqrt{-1}s(s^2-1), 2\sqrt{-1}s(s+\sqrt{-1})^2(s^2-1)). \quad (1.5)$$

Corollary 1.0.2.

$$E_{1,s}(\mathbb{Q}(s)) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z} \quad (1.6)$$

is generated by T_1 and $2T_2 = (0, 0)$.

Remark 1.0.3. 課題研究では、多項式の解の非存在から背理法を用いて 8-torsion point が存在しないことを示した。TODO: specialization theorem は injectivity しか言ってないので、各 special fiber について 8-torsion point が存在しないことを示した課題研究は上の系より真に強い？

Inspite of the fact that the generic rank of $E_{1,s}$ is 0, Yoshida ([1, Corollary 4.7.]) showed that there are infinitely many $s \in \mathbb{Q}$ such that the Mordell-Weil group of $E_{1,s}$ has positive rank over \mathbb{Q} . However, the

infinite family is not explicitly parameterized. In this paper, we find the subset of $s \in \overline{\mathbb{Q}}$ with the positive rank of the Mordell-Weil group of $E_{1,s}$ parameterized by an rational function of one variable $t \in \overline{\mathbb{Q}}$. In order to prove it, it is enough to find a subfamily of $E_{1,s}$ whose generic rank is 1, since the following theorem about the relation between an elliptic curve over a function field and its special fibers holds.

Theorem 1.0.4. (Specialization Theorem, [2, Theorem 11.4.]) Let E be an elliptic curve over a function field $k(C)$ of a smooth projective curve C over an algebraically closed field k . Let $\pi : \mathcal{E} \rightarrow C$ be the Néron model of E and $F_v := \pi^{-1}(v)$ be the special fiber for $v \in C(k)$. Then for all but finitely many $v \in C(k)$, a map

$$E(k(C)) \rightarrow F_v(k), \quad (1.7)$$

called the specialization map at $v \in C(k)$, is injective

By substituting $s = \frac{2t}{t^2-3}$ into $E_{1,s}$, we get a new family of elliptic curves

$$E_{2,t} : y^2 = x \left(x - 4 \left(\frac{2t}{t^2-3} \right)^2 \right) \left(x + \left(\left(\frac{2t}{t^2-3} \right)^2 - 1 \right)^2 \right), \quad (1.8)$$

which is a subfamily of $E_{1,s}$.

The following is our main result.

Theorem 1.0.5. The Mordell-Weil group of $E_{2,t}$ over $\overline{\mathbb{Q}}(t)$ satisfies

$$E_{2,t}(\overline{\mathbb{Q}}(t)) \cong \mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}, \quad (1.9)$$

especially the rank is 1. We denote $s = \frac{2t}{t^2-3}$. The torsion subgroup is generated by T_1 and T_2 in Theorem 1.0.1 and the free part is generated by

$$\left(s^2 - 1, \sqrt{-1}s(s^2 - 1) \frac{t^2 + 3}{t^2 - 3} \right). \quad (1.10)$$

The important point is that we prove that the generic rank of $E_{2,t}$ is exactly 1, not only the existence of a point of infinite order. Our proof is based on the method of Naskręcki in [3].

2. Preliminaries

For any elliptic curves E over a function field $k(C)$ of a projective curve C over an algebraically closed field k , there is an elliptic surface $\mathcal{E} \rightarrow \mathbb{P}^1$ with the generic fiber E called the Néron model, and we

can use some theorems in the theory of surfaces. In order to get the lower bound of the rank of the Mordell-Weil group, finding generators is enough. It is more difficult to get the upper bound of the rank. The following theorem behaves a key role in the proof of the main theorem.

Theorem 2.0.1. (Shioda-Tate formula, [4, Corollary 5.3]) Let C be a smooth irreducible projective curve over an algebraically closed field k and E an elliptic curve over a function field $k(C)$. Let $\mathcal{E} \rightarrow C$ be the Néron model of E . Let $R \subset C$ be the set of points where the special fiber of \mathcal{E} is singular. For each $v \in R$, let m_v be the number of components of the special fiber of \mathcal{E} at v . Let $\rho(\mathcal{E})$ denote the rank of the Néron-Severi group of \mathcal{E} . Then, we have

$$\rho(\mathcal{E}) = 2 + \sum_{v \in R} (m_v - 1) + \text{rank}(E(k(C))). \quad (2.1)$$

We can calculate R and m_v by Tate's algorithm, but it is still difficult to determine $\rho(\mathcal{E})$. We have the following theorem to get the upper bound of $\rho(\mathcal{E})$.

Theorem 2.0.2.

$$\rho(\mathcal{E}) \leq \frac{5}{6}e(\tilde{\mathcal{S}}), \quad (2.2)$$

$$e(\tilde{\mathcal{S}}) := \sum_{v \in R} e(F_v). \quad (2.3)$$

where $e(\tilde{\mathcal{S}})$ is the Euler number, $e(F_v)$ is the local Euler number of the special fiber of \mathcal{E} at v for each $v \in R$ and

$$e(F_v) = \begin{cases} m_v & \text{if the fiber has multiplicative reduction,} \\ m_v + 1 & \text{if the fiber has additive reduction.} \end{cases} \quad (2.4)$$

証明 TODO: Naskrencki の PhD の 2.2.19(ii), 2.2.9, 2.2.10 (やその引用元) を引用するオイラー数については [5, pp. 136-137 付録 2], [6, p.14 Table II] □

Theorem 2.0.3. ([3, Lem.3.5])

$$E(\overline{\mathbb{Q}}(s))_{\text{tors}} \hookrightarrow \prod_{v \in R} G(F_v) \quad (2.5)$$

where $G(F_v)$ is the group generated by all simple components of the fiber at v . If F_v is of type I_n in Kodaira's symbol, then $G(F_v) \cong \mathbb{Z}/n\mathbb{Z}$.

証明 TODO □

3. Proof of Theorem 1.0.1

証明 of Theorem 1.0.1

$$\Delta_{E_{1,s}} = 256s^4(s+1)^4(s-1)^4(s^2+1)^4 \quad (3.1)$$

Table 3.1 Singular fibers of $E_{1,s}$

Place	Type	m_v
$s = 0$	I_4	4
$s = \pm 1$	I_4	4
$s = \pm\sqrt{-1}$	I_4	4
$s = \infty$	I_4	4

$$e(\mathcal{E}_{1,s}) = 24 \quad (3.2)$$

したがって $\mathcal{E}_{1,s}$ は K3 曲面であり. $\rho(\mathcal{E}_{1,s}) \leq 20$ である. Theorem 2.0.1 より

$$\text{rank}(E_{1,s}) = 0 \quad (3.3)$$

As for the torsion subgroup, we have

$$E_{1,s}(\overline{\mathbb{Q}}(s))[2] = \{\mathcal{O}, (0, 0), (4s^2, 0), (-(s^2 - 1)^2, 0)\}, \quad (3.4)$$

and we can check by calculation that

$$2T_1 = (4s^2, 0), \quad (3.5)$$

$$2T_2 = (0, 0). \quad (3.6)$$

By Theorem 2.0.3, we have

$$E_{1,s}(\overline{\mathbb{Q}}(s))_{\text{tors}} \hookrightarrow (\mathbb{Z}/4\mathbb{Z})^6. \quad (3.7)$$

Therefore, we have

$$E_{1,s}(\overline{\mathbb{Q}}(s))_{\text{tors}} \cong \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}. \quad (3.8)$$

□

4. The Generic Rank of $E_{2,t}$

In order to prove Theorem 1.0.5, Theorem 2.0.2 is not enough to get the tight upper bound of the ranks of the Néron-Severi group. Actually, we get $\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) \leq 2$ from Theorem 2.0.1 and Theorem 2.0.2.

Table 4.1 Singular fibers of $E_{2,t}$

Place	Type	m_v
$t = 0$	I_4	4
$t = \pm 1$	I_4	4
$t = \pm 3$	I_4	4
$t = \pm\sqrt{3}$	I_4	4
$t^4 - 2t^2 + 9 = 0$	I_4	4
$t = \infty$	I_4	4

証明

$$\Delta_{E_{2,t}} = 4096t^4(t-1)^4(t+1)^4(t-3)^4(t+3)^4(t^2-3)^4(t^4-2t^2+9)^4 \quad (4.1)$$

$$e(\mathcal{E}_{2,t}) = 48 \quad (4.2)$$

TODO: $\rho(\mathcal{E}_{2,t}) \leq 40$ である. Theorem 2.0.1 より

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) \leq 2 \quad (4.3)$$

□

Lemma 4.0.1.

$$E_{2,t}(\overline{\mathbb{Q}}(t))_{\text{tors}} \cong \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}. \quad (4.4)$$

証明 By Theorem 2.0.3, we have

$$E_{2,t}(\overline{\mathbb{Q}}(t))_{\text{tors}} \hookrightarrow (\mathbb{Z}/4\mathbb{Z})^{12}. \quad (4.5)$$

Obviously, we have

$$E_{1,s}(\overline{\mathbb{Q}}(s))_{\text{tors}} \subset E_{2,t}(\overline{\mathbb{Q}}(t))_{\text{tors}}. \quad (4.6)$$

□

On the other hand, we have only one point of infinite order in $E_{2,t}(\overline{\mathbb{Q}}(t))$.

Lemma 4.0.2.

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) \geq 1 \quad (4.7)$$

Now, our goal is to show the upper bound of the rank of $E_{2,t}(\overline{\mathbb{Q}}(t))$ is 1.

We use another method to estimate the upper bound of the rank of Néron-Severi group, which we will explain in Chapter 5. Beforehand, we express the rank of $E_{2,t}(\overline{\mathbb{Q}}(t))$ in terms of ranks of elliptic curves with lower order coefficients in the Weierstrass equations to make the later computation feasible.

Definition 4.0.3. Let C be a smooth curve over an algebraically closed field k . Let E be an elliptic curve over a function field $k(C)$ given by the Weierstrass equation

$$E : y^2 = x^3 + a_2x^2 + a_4x + a_6 \quad (4.8)$$

where $a_2, a_4, a_6 \in k(C)$. For a fixed $u \in k(C)^*$, we denote

$$E^{(u)} : uy^2 = x^3 + a_2x^2 + a_4x + a_6 \quad (4.9)$$

to be the quadratic twist of E by u .

Theorem 4.0.4. ([7, Exercise 10.16]) Let E be an elliptic curve over a function field $k(C)$ and $u \in k(C)^*$. Then, the following equation holds

$$\text{rank } E(k(C)(\sqrt{u})) = \text{rank } E(k(C)) + \text{rank } E^{(u)}(k(C)). \quad (4.10)$$

Theorem 4.0.5. Let

$$E_{0,u} : y^2 = x(x - 4u)(x + (u - 1)^2) \quad (4.11)$$

be an elliptic curve over $\overline{\mathbb{Q}}(u)$. Then, we have

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) = \text{rank } E_{1,s}(\overline{\mathbb{Q}}(s)) + \text{rank } E_{1,s}^{(1+3s^2)}(\overline{\mathbb{Q}}(s)), \quad (4.12)$$

$$\text{rank } E_{1,s}^{(1+3s^2)}(\overline{\mathbb{Q}}(s)) = \text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) + \text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)). \quad (4.13)$$

Therefore, we have

$$\text{rank } E_{2,t}(\overline{\mathbb{Q}}(t)) = \text{rank } E_{1,s}(\overline{\mathbb{Q}}(s)) + \text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) + \text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)). \quad (4.14)$$

証明 Since solving $s = \frac{2t}{t^2-3}$ for t yields $t = \frac{1 \pm \sqrt{1+3s^2}}{s}$, we have

$$E_{2,t}(\overline{\mathbb{Q}}(t)) = E_{1,s}(\overline{\mathbb{Q}}(s)(\sqrt{1+3s^2})) \quad (4.15)$$

By Theorem 4.0.4, we get (4.12). Similarly, $E_{1,s}$ is obtained by substituting $u = s^2$ into $E_{0,u}$, so we have

$$E_{1,s}^{(1+3s^2)}(\overline{\mathbb{Q}}(s)) = E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)(\sqrt{u})), \quad (4.16)$$

then we get (4.13). \square

Theorem 4.0.6. TODO

$$\text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) \leq 1 \quad (4.17)$$

証明

$$\Delta(E_{0,u}^{(1+3u)}) = 256u^2(u-1)^4(u+1)^4(3u+1)^6 \quad (4.18)$$

Table 4.2 Singular fibers of $E_{0,u}^{(1+3u)}$

Place	Type	m_v
$u = 0$	I_2	2
$u = \pm 1$	I_4	4
$u = -\frac{1}{3}$	I_0^*	5
$u = \infty$	I_2^*	7

$$e(\mathcal{E}_{0,u}^{(1+3u)}) = 24 \quad (4.19)$$

Theorem 2.0.1 からは

$$\text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) \leq 1 \quad (4.20)$$

\square

Theorem 4.0.7.

$$\text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)) = 1 \quad (4.21)$$

証明

$$(u - 1, \sqrt{-1}(u - 1)) \in E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)) \quad (4.22)$$

より rank は正である.

$$\Delta(E_{0,u}^{(u(1+3u))}) = 256u^8(u - 1)^4(u + 1)^4(3u + 1)^6 \quad (4.23)$$

上と同様に

Table 4.3 Singular fibers of $E_{0,u}^{(u(1+3u))}$

Place	Type	m_v
$u = 0$	I_2^*	7
$u = \pm 1$	I_4	4
$u = -\frac{1}{3}$	I_0^*	5
$u = \infty$	I_2	2

$$\text{rank } E_{0,u}^{(u(1+3u))}(\overline{\mathbb{Q}}(u)) \leq 1 \quad (4.24)$$

□

5. Reductions

Let A be a discrete valuation ring of a number field K with maximal ideal \mathfrak{m} , whose residue field k has $q = p^r$ elements with p prime. Let S be an integral scheme with a morphism $S \rightarrow \text{Spec } A$ that is projective and smooth of relative dimension 2. Then the projective surface $\overline{S} = S_{\overline{\mathbb{Q}}}$ and $\tilde{S} = S_{\overline{k}}$ are smooth over the algebraically closed field $\overline{\mathbb{Q}}$ and \overline{k} , respectively. We will assume that \overline{S} and \tilde{S} are integrals, i.e., they are irreducible, nonsingular, projective surfaces.

For $l \neq p$ be a prime number, we denote by $H_{\text{ét}}^2(\tilde{S}, \mathbb{Q}_l)$ the l -adic étale cohomology group of X and by $H_{\text{ét}}^2(\tilde{S}, \mathbb{Q}_l)(1)$ the Tate twist of it.

Theorem 5.0.1. ([8, Proposition 6.2.])

There are natural injective homomorphisms

$$\mathrm{NS}(\bar{S}) \otimes_{\mathbb{Z}} \mathbb{Q}_l \hookrightarrow \mathrm{NS}(\tilde{S}) \otimes_{\mathbb{Z}} \mathbb{Q}_l \hookrightarrow H_{\mathrm{\acute{e}t}}^2(\tilde{S}, \mathbb{Q}_l)(1) \quad (5.1)$$

of finite-dimensional vector spaces over \mathbb{Q}_l .

Let $F : S_k \rightarrow S_k$ denote the absolute Frobenius, which acts as the identity on the points and by $f \mapsto f^p$ on the structure sheaf. Set $\varphi := F^r$ and let $\varphi^{(i)}$ denote the automorphism on $H_{\mathrm{\acute{e}t}}^i(\tilde{S}, \mathbb{Q}_l)$ induced by $\varphi \times 1$ acting on $S_k \times_{\mathrm{Spec} k} \mathrm{Spec} \bar{k} \cong \tilde{S}$.

Corollary 5.0.2. ([8, Corollary 6.4.]) The ranks of $\mathrm{NS}(\bar{S})$ and $\mathrm{NS}(\tilde{S})$ are bounded from above by the number of eigenvalues λ of $\varphi^{(2)}$ for which λ/q is a root of unity, counted with multiplicity.

Remark 5.0.3. ([8, Remark 6.5.]) Tate's conjecture states that the upper bound mentioned in Corollary 5.0.2 is actually equal to the rank of $\mathrm{NS}(\tilde{S})$. Tate's conjecture has been proven for elliptic K3 surfaces.

Now we want to calculate the characteristic polynomial $\mathrm{char}(\varphi^{(2)})$. Beforehand, we recall the Lefschetz trace formula.

Theorem 5.0.4.

$$\#\tilde{S}(\mathbb{F}_{q^m}) = \sum_{i=0}^n (-1)^i \mathrm{Tr}((\varphi^{(i)})^m) \quad (5.2)$$

Corollary 5.0.5.

$$\mathrm{Tr}((\varphi^{(2)})^m) = \#\tilde{S}(\mathbb{F}_{q^m}) - 1 - q^{2m} \quad (5.3)$$

証明

$$\dim H_{\mathrm{\acute{e}t}}^1(\tilde{S}, \mathbb{Q}_l) = \dim H_{\mathrm{\acute{e}t}}^3(\tilde{S}, \mathbb{Q}_l) = 0 \quad (5.4)$$

and $\varphi^{(4)}$ acts on $H_{\mathrm{\acute{e}t}}^4(\tilde{S}, \mathbb{Q}_l) \cong \mathbb{Q}_l$ by multiplication by q^2 . \square

Let V be the linear subspace of $H_{\mathrm{\acute{e}t}}^2(\tilde{S}, \mathbb{Q}_l)$ generated by the components of the singular fibers and by the zero section and $W = H_{\mathrm{\acute{e}t}}^2(\tilde{S}, \mathbb{Q}_l)/V$, then

$$\dim V = \sum_{v \in R} (m_v - 1) + 2. \quad (5.5)$$

By the multiplicativity of the characteristic polynomial, we have

$$\mathrm{char}(\varphi^{(2)}) = \mathrm{char}(\varphi^{(2)}|V) \cdot \mathrm{char}(\varphi_W^{(2)}) \quad (5.6)$$

and

$$\mathrm{Tr}((\varphi^{(2)})^m) = \mathrm{Tr}((\varphi^{(2)}|V)^m) + \mathrm{Tr}((\varphi_W^{(2)})^m) \quad (5.7)$$

for any $m \in \mathbb{Z}$, where $\varphi_W^{(2)} : W \rightarrow W$ is induced by $\varphi^{(2)}$. Since $\varphi^{(2)}$ acts on V by multiplication by q , we have

$$\text{char}(\varphi^{(2)}|V) = (x - q)^{\dim V}. \quad (5.8)$$

As for the characteristic polynomial of $\varphi_W^{(2)}$, let $t_m := \text{Tr}((\varphi_W^{(2)})^m)$, then $\text{char}(\varphi_W^{(2)})$ is the polynomial part of

$$\frac{x^{\dim W}}{\exp\left(\sum_{m=1}^{\infty} \frac{t_m}{m} x^{-m}\right)} = x^{\dim W} \left(1 + t_1 x^{-1} + \frac{t_1^2 - t_2}{2} x^{-2} + \frac{-t_1^3 + 3t_1 t_2 - 2t_3}{6} x^{-3} + \dots\right). \quad (5.9)$$

Here, by (5.7) and Corollary 5.0.5, we have

$$t_m = \#\tilde{S}(\mathbb{F}_{q^m}) - 1 - q^{2m} - \dim V \cdot q^m. \quad (5.10)$$

Lemma 5.0.6. ([9, Theorem 4, Part III]) If \tilde{S} is a K3 surface, then the second Betti number of \tilde{S} is 22.

Theorem 5.0.7.

$$\text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) = 0 \quad (5.11)$$

証明 We denote by $S = \mathcal{E}_{0,u}^{(1+3u)} \rightarrow \mathbb{P}^1$ the elliptic surface with the generic fiber $E_{0,u}^{(1+3u)}$.

Table 4.2 より K3 なので Lemma 5.0.6 より

$$\dim_{\mathbb{Q}_l} H_{\text{ét}}^2(\tilde{S}, \mathbb{Q}_l) = 22 \quad (5.12)$$

である. V is of rank 19, on which the Frobenius automorphism acts by multiplication by p .

$$\text{char}(\varphi^{(2)}|V) = (x - 5)^{19} \quad (5.13)$$

Note that all the multiplicative fibers are split in \mathbb{F}_{5^m} for $m = 1, 2, 3$.

$$t_m = \#\tilde{S}(\mathbb{F}_{5^m}) - 1 - 5^{2m} - 19 \cdot 5^m \quad (5.14)$$

m	1	2	3
$\#\tilde{S}(\mathbb{F}_{5^m})$	120	1080	18264
t_m	-1	-21	263

$$\text{char}(\varphi_W^{(2)}) = x^3 + x^2 + 11x - 77 \quad (5.15)$$

If $\text{char}(\varphi_W^{(2)})$ has a root of the form $x = 5\zeta$ for some root of unity ζ , then ζ is a root of the polynomial

$$125x^3 + 25x^2 + 55x - 77, \quad (5.16)$$

which is irreducible over \mathbb{Q} . It contradicts the fact that ζ is an algebraic integer. By Corollary 5.0.2, $\rho(\mathcal{E}_{0,u}^{(1+3u)}) \leq 19$. Then by Theorem 2.0.1, we have

$$\text{rank } E_{0,u}^{(1+3u)}(\overline{\mathbb{Q}}(u)) = 0. \quad (5.17)$$

□

Acknowledgments

I would like to express my deepest appreciation to my professor Masato Kurihara for his guidance and encouragement. I also could not have undertaken this journey without the support of Dr. Bartosz Naskręcki. He has politely answered my many emails with questions and given me valuable advice.

参考文献

- [1] T. Yoshida. The relationship between face cuboids and elliptic curves. 2024. arXiv: 2407.09825 [math.NT]. URL: <https://arxiv.org/abs/2407.09825>.
- [2] J. H. Silverman. Advanced topics in the arithmetic of elliptic curves. eng. Graduate texts in mathematics ; 151. New York: Springer, 1994.
- [3] B. Naskręcki. Mordell-Weil ranks of families of elliptic curves associated to Pythagorean triples. eng. Acta Arithmetica 160.2, pp. 159–183, (2013). URL: <http://eudml.org/doc/279803>.
- [4] T. Shioda. On the Mordell-Weil Lattices. Commentarii Mathematici Universitatis Sancti Pauli 39, (1990).
- [5] 徹. 塩田. Mordell-Weil lattice の理論とその応用. jpn. 東京大学数理科学セミナーノート ; 1. 東京: Graduate school of mathematical sciences, 1993.
- [6] K. Kodaira. On Compact Analytic Surfaces, III. Annals of Mathematics 78.1, pp. 1–40, (1963). URL: <http://www.jstor.org/stable/1970500> (accessed on 2025-01-12).
- [7] J. H. Silverman. The arithmetic of elliptic curves. eng. 2nd ed. Graduate texts in mathematics ; 106. New York: Springer, 2009.
- [8] R. van Luijk. An elliptic K3 surface associated to Heron triangles. Journal of Number Theory 123.1, pp. 92–119, (2007). DOI: <https://doi.org/10.1016/j.jnt.2006.06.006>. URL: <https://www.sciencedirect.com/science/article/pii/S0022314X06001326>.
- [9] D. Mumford. Selected papers on the classification of varieties and moduli spaces. eng. New York: Springer, 2004.