

Supplementary Material to:

Title: Stock delineation of Striped Snakehead, *Channa striata* using multivariate generalised linear models with otolith shape and chemistry data

Authors: Salman Khan^{1#}, Hayden T. Schilling^{2,3#}, Mohammad Afzal Khan¹, Devendra Kumar Patel⁴, Ben Maslen⁵, Kaish Miyan¹

1. Section of Fishery Science and Aquaculture, Department of Zoology, Aligarh Muslim University, Aligarh -202 002, India
2. Centre for Marine Science & Innovation, UNSW Australia, Sydney 2052, Australia
3. Sydney Institute of Marine Science, Chowder Bay Road, Mosman 2088, Australia
4. Indian Institute of Toxicology Research, Council of Scientific and Industrial Research, Lucknow-226 001, India
5. Mark Wainwright Analytical Centre, UNSW Australia, Sydney 2052, Australia

SK and HTS are joint first authors.

Corresponding author: Hayden T. Schilling, h.schilling@unsw.edu.au , +614 02 9435 4600
(no fax)

19 **Table S1** Details of the samples used in the otolith chemistry and shape analysis of *C. striata*

Site	Sample	Mean	Minimum	Maximum	Mean	Minimum	Maximum
	Size	Fish	Fish Total	Fish Total	Otolith	Otolith	Otolith
		Total	Length	Length	Weight	weight	weight
		Length	(cm)	(cm)	(mg)	(mg)	(mg)
		(cm)					
Agra	18	41.0	32.0	48.0	191.1	151.1	264.2
Lucknow	18	32.8	24.5	44.0	107.0	56.1	180.4
Narora	18	31.2	21.5	39.0	103.8	46.2	156.6

20

21