

# Problem 7-4 Random failures in uncorrelated networks

1. Poisson d.:  $p_k = e^{-\mu} \frac{\mu^k}{k!}$

From literature:  $\langle k \rangle = \mu$

(1)

$\langle k^2 \rangle = \mu(1+\mu)$

(2)

critical threshold:

$$f_c = 1 - \frac{1}{\frac{\langle k^2 \rangle}{\langle k \rangle} - 1} \stackrel{(1),(2)}{=} 1 - \frac{1}{\frac{\mu(1+\mu)}{\mu} - 1} = 1 - \frac{1}{1+\mu-1}$$

$$= 1 - \frac{1}{\mu} \quad (= 1 - \frac{1}{\langle k \rangle} = f_c^{ER})$$

2. Discrete exponential d.:  $p_k = (1 - e^{-\lambda}) e^{-\lambda k}$

From literature:  $\langle k \rangle = (e^{\lambda} - 1)^{-1}$

(3)

$\langle k^2 \rangle = (e^{\lambda} + 1)(e^{\lambda} - 1)^{-2}$

(4)

critical threshold:

$$f_c = 1 - \frac{1}{\frac{\langle k^2 \rangle}{\langle k \rangle} - 1} \stackrel{(3),(4)}{=} 1 - \frac{1}{(e^{\lambda} + 1)(e^{\lambda} - 1)^{-2} (e^{\lambda} - 1) - 1}^{-1}$$

$$= 1 - \frac{1}{(e^{\lambda} + 1)(e^{\lambda} - 1) - 1} = 1 - \frac{1}{e^{\lambda} + 1 - (e^{\lambda} - 1)}$$

$$= 1 - \frac{1}{(e^{\lambda} - 1)^2} = \frac{2 - (e^{\lambda} - 1)}{2} = \frac{1}{2}(e^{\lambda} + 3)$$

$$= \frac{1}{2}(3 - e^{\lambda})$$

3. Dirac delta d.:  $p_k = \delta_{k,k_0} = \begin{cases} 1 & k=k_0 \\ 0 & \text{otherwise} \end{cases}$

From literature:  $\langle k \rangle = k_0$

$\langle k^2 \rangle = k_0^2$

critical threshold:

$$f_c = 1 - \frac{1}{\frac{\langle k^2 \rangle}{\langle k \rangle} - 1} = 1 - \frac{1}{\frac{k_0^2}{k_0} - 1} = \frac{(k_0 - 1) - 1}{k_0 - 1} = \frac{k_0 - 2}{k_0 - 1}$$

Consequences for the network robustness:

1. The network becomes more robust <sup>(towards random failures)</sup> if  $\mu \rightarrow \infty$ , i.e. if  $\langle k \rangle \rightarrow \infty$ ,

which is the same result as for random networks. A larger average degree  $\langle k \rangle$  improves the robustness, a small  $\langle k \rangle$  worsens it.

2. The network becomes more robust towards random failures if  $\lambda \rightarrow 0$  ( $\lim_{\lambda \rightarrow 0} f_c = \lim_{\lambda \rightarrow 0} \frac{1}{2}(3 - e^{\lambda}) = 1$ ), which leads to  $\langle k \rangle \rightarrow \infty$  ( $\lim_{\lambda \rightarrow 0} \frac{1}{e^{\lambda} - 1} = \infty$ ), i.e. a large average degree. The network becomes less robust if  $\lambda \rightarrow \ln(3)$ , or if  $\langle k \rangle \rightarrow \frac{1}{2}$ .

3. The network becomes more robust towards random failures for  $k_0 = \langle k \rangle \rightarrow \infty$  and less robust if  $k_0 \rightarrow 2$ .