



Interpretable Machine Learning

Shapley Values



Learning goals


- Learn what game theory is
- Understand the concept behind cooperative games
- Understand the Shapley value in game theory



COOPERATIVE GAMES IN GAME THEORY

► Shapley (1951)


- Game theory is the study of strategic games between players, “game” refers to any series of interactions between actors / agents with gains and losses of quantifiable utility value



COOPERATIVE GAMES IN GAME THEORY

► Shapley (1951)


- Game theory is the study of strategic games between players, “game” refers to any series of interactions between actors / agents with gains and losses of quantifiable utility value
- Cooperative games: For all possible players $P = \{1, \dots, p\}$, each subset of players $S \subseteq P$ forms a coalition – each coalition S achieves a certain payout



COOPERATIVE GAMES IN GAME THEORY

► Shapley (1951)


- Game theory is the study of strategic games between players, “game” refers to any series of interactions between actors / agents with gains and losses of quantifiable utility value
- Cooperative games: For all possible players $P = \{1, \dots, p\}$, each subset of players $S \subseteq P$ forms a coalition – each coalition S achieves a certain payout
- A value function $v : 2^{|P|} \mapsto \mathbb{R}$ maps all $2^{|P|}$ possible coalitions to their payout (or gain)



COOPERATIVE GAMES IN GAME THEORY

► Shapley (1951)


- Game theory is the study of strategic games between players, “game” refers to any series of interactions between actors / agents with gains and losses of quantifiable utility value
- Cooperative games: For all possible players $P = \{1, \dots, p\}$, each subset of players $S \subseteq P$ forms a coalition – each coalition S achieves a certain payout
- A value function $v : 2^{|P|} \mapsto \mathbb{R}$ maps all $2^{|P|}$ possible coalitions to their payout (or gain)
- $v(S)$ is the payout of coalition $S \subseteq P$ (payout of empty coalition must be zero: $v(\emptyset) = 0$)



COOPERATIVE GAMES IN GAME THEORY

► Shapley (1951)


- Game theory is the study of strategic games between players, “game” refers to any series of interactions between actors / agents with gains and losses of quantifiable utility value
- Cooperative games: For all possible players $P = \{1, \dots, p\}$, each subset of players $S \subseteq P$ forms a coalition – each coalition S achieves a certain payout
- A value function $v : 2^{|P|} \mapsto \mathbb{R}$ maps all $2^{|P|}$ possible coalitions to their payout (or gain)
- $v(S)$ is the payout of coalition $S \subseteq P$ (payout of empty coalition must be zero: $v(\emptyset) = 0$)
- As some players contribute more than others, we want to fairly divide the total achievable payout $v(P)$ among the players according to a player’s individual contribution






COOPERATIVE GAMES IN GAME THEORY

► Shapley (1951)



- Game theory is the study of strategic games between players, “game” refers to any series of interactions between actors / agents with gains and losses of quantifiable utility value
- Cooperative games: For all possible players $P = \{1, \dots, p\}$, each subset of players $S \subseteq P$ forms a coalition – each coalition S achieves a certain payout
- A value function $v : 2^{|P|} \mapsto \mathbb{R}$ maps all $2^{|P|}$ possible coalitions to their payout (or gain)
- $v(S)$ is the payout of coalition $S \subseteq P$ (payout of empty coalition must be zero: $v(\emptyset) = 0$)
- As some players contribute more than others, we want to fairly divide the total achievable payout $v(P)$ among the players according to a player’s individual contribution
- We call the individual payout per player $\phi_j, j \in P$ (later: Shapley value)




COOPERATIVE GAMES WITHOUT INTERACTIONS





COOPERATIVE GAMES WITHOUT INTERACTIONS




⇒ Fair Payouts are Trivial Without Interactions



COOPERATIVE GAMES WITH INTERACTIONS




⇒ Unclear how to fairly distribute payouts when players interact




COOPERATIVE GAMES WITH INTERACTIONS

Question: What is a fair payout for player “yellow”?

Idea: Compute marginal contribution of the player of interest across different coalitions




- Compute the total payout of each coalition
- Compute difference in payouts for each coalition with and without player “yellow” (= marginal contribution)
- Average marginal contributions using appropriate weights



COOPERATIVE GAMES WITH INTERACTIONS

Question: What is a fair payout for player “yellow”?



Idea: Compute marginal contribution of the player of interest across different coalitions



- Compute the total payout of each coalition
- Compute difference in payouts for each coalition with and without player “yellow” (= marginal contribution)
- Average marginal contributions using appropriate weights

Note: Each marginal contribution is weighted w.r.t. number of possible orders of its coalition


\rightsquigarrow More players in $S \Rightarrow$ more orderings of S



SHAPLEY VALUE - SET DEFINITION

This idea refers to the **Shapley value** which assigns a payout value to each player according to its marginal contribution in all possible coalitions.


- Let $v(S \cup \{j\}) - v(S)$ be the marginal contribution of player j to coalition S
 \rightsquigarrow measures how much a player j increases the value of a coalition S



SHAPLEY VALUE - SET DEFINITION

This idea refers to the **Shapley value** which assigns a payout value to each player according to its marginal contribution in all possible coalitions.

- Let $v(S \cup \{j\}) - v(S)$ be the marginal contribution of player j to coalition S
 \rightsquigarrow measures how much a player j increases the value of a coalition S
- Average marginal contributions for all possible coalitions $S \subseteq P \setminus \{j\}$
 \rightsquigarrow order of how players join the coalition matters \Rightarrow different weights depending on size of S




SHAPLEY VALUE - SET DEFINITION

This idea refers to the **Shapley value** which assigns a payout value to each player according to its marginal contribution in all possible coalitions.

- Let $v(S \cup \{j\}) - v(S)$ be the marginal contribution of player j to coalition S
 \rightsquigarrow measures how much a player j increases the value of a coalition S
- Average marginal contributions for all possible coalitions $S \subseteq P \setminus \{j\}$
 \rightsquigarrow order of how players join the coalition matters \Rightarrow different weights depending on size of S
- Shapley value via **set definition** (weighting via multinomial coefficient):

$$\phi_j = \sum_{S \subseteq P \setminus \{j\}} \frac{|S|!(|P| - |S| - 1)!}{|P|!} (v(S \cup \{j\}) - v(S))$$




SHAPLEY VALUE - ORDER DEFINITION

The Shapley value was introduced as summation over sets $S \subseteq P \setminus \{j\}$, but it can be equivalently defined as a summation of all orders of players:

$$\phi_j = \frac{1}{|P|!} \sum_{\tau \in \Pi} (v(S_j^\tau \cup \{j\}) - v(S_j^\tau))$$

- Π : All possible orders of players (we have $|P|!$ in total)



SHAPLEY VALUE - ORDER DEFINITION

The Shapley value was introduced as summation over sets $S \subseteq P \setminus \{j\}$, but it can be equivalently defined as a summation of all orders of players:

$$\phi_j = \frac{1}{|P|!} \sum_{\tau \in \Pi} (v(S_j^\tau \cup \{j\}) - v(S_j^\tau))$$

- Π : All possible orders of players (we have $|P|!$ in total)
- S_j^τ : Set of players before player j in order $\tau = (\tau^{(1)}, \dots, \tau^{(p)})$ where $\tau^{(i)}$ is i -th element


⇒ Example: Players 1, 2, 3 ⇒

$$\Pi = \{(1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1)\}$$

↪ For order $\tau = (2, 1, 3)$ and player of interest $j = 3 \Rightarrow S_j^\tau = \{2, 1\}$

↪ For order $\tau = (3, 1, 2)$ and player of interest $j = 1 \Rightarrow S_j^\tau = \{3\}$

↪ For order $\tau = (3, 1, 2)$ and player of interest $j = 3 \Rightarrow S_j^\tau = \emptyset$



SHAPLEY VALUE - ORDER DEFINITION

The Shapley value was introduced as summation over sets $S \subseteq P \setminus \{j\}$, but it can be equivalently defined as a summation of all orders of players:

$$\phi_j = \frac{1}{|P|!} \sum_{\tau \in \Pi} (v(S_j^\tau \cup \{j\}) - v(S_j^\tau))$$

- Π : All possible orders of players (we have $|P|!$ in total)
- S_j^τ : Set of players before player j in order $\tau = (\tau^{(1)}, \dots, \tau^{(p)})$ where $\tau^{(i)}$ is i -th element

⇒ Example: Players 1, 2, 3 ⇒

$$\Pi = \{(1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1)\}$$


↪ For order $\tau = (2, 1, 3)$ and player of interest $j = 3 \Rightarrow S_j^\tau = \{2, 1\}$

↪ For order $\tau = (3, 1, 2)$ and player of interest $j = 1 \Rightarrow S_j^\tau = \{3\}$

↪ For order $\tau = (3, 1, 2)$ and player of interest $j = 3 \Rightarrow S_j^\tau = \emptyset$

- Order definition: Marginal contribution of orders that yield set $S = \{1, 2\}$ is summed twice

↪ In set definition, it has the weight $\frac{2!(3-2-1)!}{3!} = \frac{2 \cdot 0!}{6} = \frac{2}{6}$





SHAPLEY VALUE - COMMENTS ON ORDER DEFINITION

- Order and set definition are equivalent
- Reason: The number of orders which yield the same coalition S is

$$|S|!(|P| - |S| - 1)!$$

\Rightarrow There are $|S|!$ possible orders of players within coalition S

\Rightarrow There are $(|P| - |S| - 1)!$ possible orders of players without S and j




SHAPLEY VALUE - COMMENTS ON ORDER DEFINITION

- Order and set definition are equivalent
- Reason: The number of orders which yield the same coalition S is

$$|S|!(|P| - |S| - 1)!$$

\Rightarrow There are $|S|!$ possible orders of players within coalition S

\Rightarrow There are $(|P| - |S| - 1)!$ possible orders of players without S and j




- Relevance of the order definition: Approximate Shapley values by sampling permutations

\leadsto randomly sample a fixed number of M permutations and average them:



$$\phi_j = \frac{1}{M} \sum_{\tau \in \Pi_M} (v(S_j^\tau \cup \{j\}) - v(S_j^\tau))$$

where $\Pi_M \subset \Pi$ is a random subset of Π containing only M orders of players




WEIGHTS FOR MARGINAL CONTRIBUTION - ILLUSTRATION

$|P|! = 6$ orders



WEIGHTS FOR MARGINAL CONTRIBUTION - ILLUSTRATION

$|P|! = 6$ orders



via orders

(players before "yellow" player)

$|S| = 2$
weight = $1/6$



$|S| = 2$
weight = $1/6$



$|S| = 1$
weight = $1/6$




$|S| = 1$
weight = $1/6$



$|S| = 0$
weight = $1/6$




$|S| = 0$
weight = $1/6$



WEIGHTS FOR MARGINAL CONTRIBUTION - ILLUSTRATION

$|P|! = 6$ orders



via orders
(players before "yellow" player)

$|S| = 2$
weight = $1/6$



$|S| = 2$
weight = $1/6$



$|S| = 1$
weight = $1/6$



$|S| = 1$
weight = $1/6$



$|S| = 0$
weight = $1/6$



$|S| = 0$
weight = $1/6$



via sets

$|S| = 2$
weight = $2/6$




$|S| = 1$
weight = $1/6$



$|S| = 1$
weight = $1/6$





$|S| = 0$
weight = $2/6$





SHAPLEY VALUES - ILLUSTRATION

- Shapley value of player j is the marginal contribution to the value when it enters any coalition
- Produce all possible joining orders of player coalitions





SHAPLEY VALUES - ILLUSTRATION

- Shapley value of player j is the marginal contribution to the value when it enters any coalition
- Produce all possible joining orders of player coalitions
- Measure and average the difference in payout after player 1 enters the coalition





SHAPLEY VALUES - ILLUSTRATION

- Shapley value of player j is the marginal contribution to the value when it enters any coalition
- Produce all possible joining orders of player coalitions
- Measure and average the difference in payout after player 2 enters the coalition





SHAPLEY VALUES - ILLUSTRATION

- Shapley value of player j is the marginal contribution to the value when it enters any coalition
- Produce all possible joining orders of player coalitions
- Measure and average the difference in payout after player 3 enters the coalition



SHAPLEY VALUES - ILLUSTRATION

- Shapley value of player j is the marginal contribution to the value when it enters any coalition
- Produce all possible joining orders of player coalitions




AXIOMS OF FAIR PAYOUTS

Why is this a fair payout solution?

One possibility to define fair payouts are the following axioms for a given value function v :

- **Efficiency:** Player contributions add up to the total payout of the game:

$$\sum_{j=1}^p \phi_j = v(P)$$



AXIOMS OF FAIR PAYOUTS

Why is this a fair payout solution?


One possibility to define fair payouts are the following axioms for a given value function v :

- **Efficiency:** Player contributions add up to the total payout of the game:

$$\sum_{j=1}^p \phi_j = v(P)$$

- **Symmetry:** Players $j, k \in P$ who contribute the same to any coalition get the same payout:

If $v(S \cup \{j\}) = v(S \cup \{k\})$ for all $S \subseteq P \setminus \{j, k\}$, then $\phi_j = \phi_k$




AXIOMS OF FAIR PAYOUTS

Why is this a fair payout solution?

One possibility to define fair payouts are the following axioms for a given value function v :

- **Efficiency:** Player contributions add up to the total payout of the game:
$$\sum_{j=1}^p \phi_j = v(P)$$
- **Symmetry:** Players $j, k \in P$ who contribute the same to any coalition get the same payout:
If $v(S \cup \{j\}) = v(S \cup \{k\})$ for all $S \subseteq P \setminus \{j, k\}$, then $\phi_j = \phi_k$
- **Dummy/Null Player:** Payout is 0 for players who don't contribute to the value of any coalition:
If $v(S \cup \{j\}) = v(S) \quad \forall \quad S \subseteq P \setminus \{j\}$, then $\phi_j = 0$



AXIOMS OF FAIR PAYOUTS

Why is this a fair payout solution?

One possibility to define fair payouts are the following axioms for a given value function v :

- **Efficiency:** Player contributions add up to the total payout of the game:
$$\sum_{j=1}^p \phi_j = v(P)$$
- **Symmetry:** Players $j, k \in P$ who contribute the same to any coalition get the same payout:
If $v(S \cup \{j\}) = v(S \cup \{k\})$ for all $S \subseteq P \setminus \{j, k\}$, then $\phi_j = \phi_k$
- **Dummy/Null Player:** Payout is 0 for players who don't contribute to the value of any coalition:
If $v(S \cup \{j\}) = v(S) \quad \forall \quad S \subseteq P \setminus \{j\}$, then $\phi_j = 0$
- **Additivity:** For a game v with combined payouts $v(S) = v_1(S) + v_2(S)$, the payout is the sum of payouts: $\phi_{j,v} = \phi_{j,v_1} + \phi_{j,v_2}$

