



# Introduction to Machine Learning

## Bernoulli Loss



### Learning goals

- Know the Bernoulli loss and related losses (log-loss, logistic loss, Binomial loss)
- Derive the risk minimizer
- Derive the optimal constant model
- Understand the connection between log-loss and entropy splitting




# BERNOULLI LOSS

$$L(y, f(\mathbf{x})) = \log(1 + \exp(-y \cdot f(\mathbf{x}))) \quad \text{for } y \in \{-1, +1\}$$

$$L(y, f(\mathbf{x})) = -y \cdot f(\mathbf{x}) + \log(1 + \exp(f(\mathbf{x}))) \quad \text{for } y \in \{0, 1\}$$

- Two equivalent formulations for different label encodings
- Negative log-likelihood of Bernoulli model, e.g., logistic regression
- Convex, differentiable




# BERNOULLI LOSS ON PROBABILITIES

If scores are transformed into probabilities by the logistic function

$\pi(\mathbf{x}) = (1 + \exp(-f(\mathbf{x})))^{-1}$  (or equivalently if  $f(x) = \log\left(\frac{\pi(\mathbf{x})}{1-\pi(\mathbf{x})}\right)$  are the log-odds of  $\pi(\mathbf{x})$ ), we arrive at another equivalent formulation of the loss, where  $y$  is again encoded as  $\{0, 1\}$ :

$$L(y, \pi(\mathbf{x})) = -y \log(\pi(\mathbf{x})) - (1 - y) \log(1 - \pi(\mathbf{x})).$$



# BERNOULLI LOSS: RISK MINIMIZER

The risk minimizer for the Bernoulli loss defined for probabilistic classifiers  $\pi(\mathbf{x})$  and on  $y \in \{0, 1\}$  is

$$\pi^*(\mathbf{x}) = \eta(\mathbf{x}) = \mathbb{P}(y = 1 \mid \mathbf{x} = \mathbf{x}).$$

**Proof:** We can write the risk for binary  $y$  as follows:

$$\mathcal{R}(f) = \mathbb{E}_{\mathbf{x}} [L(1, \pi(\mathbf{x})) \cdot \eta(\mathbf{x}) + L(0, \pi(\mathbf{x})) \cdot (1 - \eta(\mathbf{x}))],$$

with  $\eta(\mathbf{x}) = \mathbb{P}(y = 1 \mid \mathbf{x} = \mathbf{x})$  (see chapter on the 0-1-loss for more details).


For a fixed  $\mathbf{x}$  we compute the point-wise optimal value  $c$  by setting the derivative to 0:

$$\frac{\partial}{\partial c} (-\log c \cdot \eta(\mathbf{x}) - \log(1 - c) \cdot (1 - \eta(\mathbf{x}))) = 0$$

$$-\frac{\eta(\mathbf{x})}{c} + \frac{1 - \eta(\mathbf{x})}{1 - c} = 0$$

$$\frac{-\eta(\mathbf{x}) + \eta(\mathbf{x})c + c - \eta(\mathbf{x})c}{c(1 - c)} = 0$$

$$c = \eta(\mathbf{x}).$$



# BERNOULLI LOSS: RISK MINIMIZER


The risk minimizer for the Bernoulli loss defined on  $y \in \{-1, 1\}$  and scores  $f(\mathbf{x})$  is the point-wise log-odds:

$$f^*(\mathbf{x}) = \log\left(\frac{\mathbb{P}(y = 1 \mid \mathbf{x} = \mathbf{x})}{1 - \mathbb{P}(y = 1 \mid \mathbf{x} = \mathbf{x})}\right).$$

The function is undefined when  $P(y = 1 \mid \mathbf{x} = \mathbf{x}) = 1$  or  $P(y = 1 \mid \mathbf{x} = \mathbf{x}) = 0$ , but predicts a smooth curve which grows when  $P(y = 1 \mid \mathbf{x} = \mathbf{x})$  increases and equals 0 when  $P(y = 1 \mid \mathbf{x} = \mathbf{x}) = 0.5$ .

**Proof:** As before we minimize

$$\begin{aligned}\mathcal{R}(f) &= \mathbb{E}_{\mathbf{x}} [L(1, f(\mathbf{x})) \cdot \eta(\mathbf{x}) + L(-1, f(\mathbf{x})) \cdot (1 - \eta(\mathbf{x}))] \\ &= \mathbb{E}_{\mathbf{x}} [\log(1 + \exp(-f(\mathbf{x})))\eta(\mathbf{x}) + \log(1 + \exp(f(\mathbf{x})))(1 - \eta(\mathbf{x}))]\end{aligned}$$



# BERNOULLI LOSS: RISK MINIMIZER

For a fixed  $\mathbf{x}$  we compute the point-wise optimal value  $c$  by setting the derivative to 0:

$$\frac{\partial}{\partial c} \log(1 + \exp(-c))\eta(\mathbf{x}) + \log(1 + \exp(c))(1 - \eta(\mathbf{x})) = 0$$


$$-\frac{\exp(-c)}{1 + \exp(-c)}\eta(\mathbf{x}) + \frac{\exp(c)}{1 + \exp(c)}(1 - \eta(\mathbf{x})) = 0$$

$$-\frac{\exp(-c)}{1 + \exp(-c)}\eta(\mathbf{x}) + \frac{1}{1 + \exp(-c)}S(1 - \eta(\mathbf{x})) = 0$$

$$-\eta(\mathbf{x}) + \frac{1}{1 + \exp(-c)} = 0$$

$$\eta(\mathbf{x}) = \frac{1}{1 + \exp(-c)}$$

$$c = \log\left(\frac{\eta(\mathbf{x})}{1 - \eta(\mathbf{x})}\right)$$



## BERNOULLI: OPTIMAL CONSTANT MODEL

The optimal constant probability model  $\pi(\mathbf{x}) = \theta$  w.r.t. the Bernoulli loss for labels from  $\mathcal{Y} = \{0, 1\}$  is:

$$\hat{\theta} = \arg \min_{\theta} \mathcal{R}_{\text{emp}}(\theta) = \frac{1}{n} \sum_{i=1}^n y^{(i)}$$

Again, this is the fraction of class-1 observations in the observed data. We can simply prove this again by setting the derivative of the risk to 0 and solving for  $\theta$ .





# BERNOULLI-LOSS: NAMING CONVENTION

We have seen three loss functions that are closely related. In the literature, there are different names for the losses:

$$L(y, f(\mathbf{x})) = \log(1 + \exp(-yf(\mathbf{x}))) \quad \text{for } y \in \{-1, +1\}$$


$$L(y, f(\mathbf{x})) = -y \cdot f(\mathbf{x}) + \log(1 + \exp(f(\mathbf{x}))) \quad \text{for } y \in \{0, 1\}$$

are referred to as Bernoulli, Binomial or logistic loss.

$$L(y, \pi(\mathbf{x})) = -y \log(\pi(\mathbf{x})) - (1 - y) \log(1 - \pi(\mathbf{x})) \quad \text{for } y \in \{0, 1\}$$

is referred to as cross-entropy or log-loss.

We usually refer to all of them as **Bernoulli loss**, and rather make clear whether they are defined on labels  $y \in \{0, 1\}$  or  $y \in \{-1, +1\}$  and on scores  $f(\mathbf{x})$  or probabilities  $\pi(\mathbf{x})$ .





## BERNOULLI LOSS MIN = ENTROPY SPLITTING

$$\begin{aligned}\mathcal{R}(\mathcal{N}) &= \sum_{(\mathbf{x}, y) \in \mathcal{N}} \left( - \sum_{k=1}^g [y = k] \log \pi_k(\mathbf{x}) \right) \\ &\stackrel{(*)}{=} - \sum_{k=1}^g \sum_{(\mathbf{x}, y) \in \mathcal{N}} [y = k] \log \pi_k^{(\mathcal{N})} \\ &= - \sum_{k=1}^g \log \pi_k^{(\mathcal{N})} \underbrace{\sum_{(\mathbf{x}, y) \in \mathcal{N}} [y = k]}_{n_{\mathcal{N}} \cdot \pi_k^{(\mathcal{N})}} \\ &= - n_{\mathcal{N}} \sum_{k=1}^g \pi_k^{(\mathcal{N})} \log \pi_k^{(\mathcal{N})} = n_{\mathcal{N}} \text{Imp}(\mathcal{N}),\end{aligned}$$