

Solution 1: Multiclass Classification with 0-1-Loss

- (a) As seen in the 0-1-Loss presentation, slide 2, the discrete classifier that minimizes the risk h^* (The Bayes optimal classifier) is:

$$\begin{aligned} h^*(\mathbf{x}) &= \arg \max_{l \in \mathcal{Y}} \underbrace{\mathbb{P}(y = l \mid \mathbf{x}=\mathbf{x})}_{\sim \text{Unif}\{1, \dots, x\}} \\ &= \arg \max_x \frac{1}{x} \cdot \mathbb{I}_{[1 \leq l \leq x]} \end{aligned}$$

As the distribution of y given x is uniform, any value between 1 and x is optimal.

$$h^*(\mathbf{x}) = \{1, \dots, x\}$$

- (b) The Bayes risk for the 0-1-loss, also known as the Bayes error rate, is defined as :

$$\begin{aligned} \mathcal{R}^* &= 1 - \mathbb{E}_x \left[\max_{l \in \mathcal{Y}} \mathbb{P}(y = l \mid \mathbf{x}=\mathbf{x}) \right] \\ &= 1 - \underbrace{\mathbb{E}_x \left[\frac{1}{x} \right]}_{p_x \sim \text{Unif}\{1, \dots, 10\}} \\ &= 1 - \sum_{x=1}^{10} \frac{1}{x} \frac{1}{10} \\ &= 1 - \frac{7381}{25200} \end{aligned}$$

- (c) The point-wise optimizer for the 0-1 loss over all discrete classifiers $h^*(\mathbf{x})$ is:

$$h^*(\mathbf{x}) = \arg \max_{l \in \mathcal{Y}} \mathbb{P}(y = l \mid \mathbf{x}=\mathbf{x})$$

The optimal constant model can be obtained by forgetting the conditioning on \mathbf{x} , leading to:

$$\hat{h}(\mathbf{x}) = \arg \max_{l \in \mathcal{Y}} \mathbb{P}(y = l)$$

Using the law of total probability:

$$\begin{aligned}
\hat{h}(\mathbf{x}) &= \arg \max_{l \in \mathcal{Y}} \sum_{x=1}^{10} \mathbb{P}(y = l \mid \mathbf{x} = \mathbf{x}) \cdot \mathbb{P}(\mathbf{x} = \mathbf{x}) \\
&= \arg \max_{l \in \mathcal{Y}} \sum_{x=1}^{10} \frac{1}{x} \cdot \mathbb{I}_{[1 \leq l \leq x]} \cdot \frac{1}{10} \\
&= \arg \max_{l \in \mathcal{Y}} \begin{cases} \frac{7381}{25200}, & l = 1 \\ \frac{7381}{25200} - \frac{1}{10}, & l = 2 \\ \frac{7381}{25200} - \frac{1}{10} - \frac{1}{20}, & l = 3 \\ \vdots & \vdots \\ \frac{7381}{25200} - \sum_{z=1}^{l-1} \frac{1}{10 \cdot z}, & l = 10 \end{cases}
\end{aligned}$$

As the probability is monotonically decreasing with l , we can conclude that the optimal constant model is :

$$\hat{h}(\mathbf{x}) = 1$$

(d) The Risk is calculated by:

$$\begin{aligned}
\mathbb{R}_L(\hat{h}) &= 1 - \max \mathbb{P}(y = l) \\
&= 1 - \mathbb{P}(y = 1) \\
&= 1 - \frac{7381}{25200}
\end{aligned}$$