

Random Variables

Hayk Aprikyan, Hayk Tarkhanyan

Random Variables

Recap:

During the previous lecture we had a problem like this:

Example

Both the bus and you get to the bus stop at random times between 12 pm and 1 pm. When the bus arrives, it waits for 5 minutes before leaving. When you arrive, you wait for 20 minutes before leaving if the bus doesn't come. What is the probability that you catch the bus?

Here we did not know the **exact values** of the arrival times, so in order to compare them, we **denoted them** by y and b – and calculated their probabilities.

In this case, we say that y and b are **random variables**.

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4.

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] =$$

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] = \mathbb{P}[4] =$$

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] = \mathbb{P}[4] = \frac{1}{6}$$

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] = \mathbb{P}[4] = \frac{1}{6} \qquad \mathbb{P}[X = 1] =$$

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] = \mathbb{P}[4] = \frac{1}{6} \qquad \mathbb{P}[X = 1] = \mathbb{P}[\{1, 5\}] =$$

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] = \mathbb{P}[4] = \frac{1}{6} \qquad \mathbb{P}[X = 1] = \mathbb{P}[\{1, 5\}] = \frac{2}{6}$$

Random Variables

In general, we often deal with situations where some quantity is unknown:

- i.e. we don't know its exact value,

but we know the **probabilistic distribution** of its values, i.e.

- what possible values it can take, and
- how probable each value is.

Example

You throw a fair dice. The outcome is unknown, but we know it can be 1, 2, 3, 4, 5, or 6, each with probability $\frac{1}{6}$.

Let X be the remainder of the outcome when divided by 4. Then,

- the possible values of X are 0, 1, 2, 3,
- the probabilities of these values are:

$$\mathbb{P}[X = 0] = \mathbb{P}[4] = \frac{1}{6}$$

$$\mathbb{P}[X = 1] = \mathbb{P}[\{1, 5\}] = \frac{2}{6}$$

etc.

Definition

A quantity whose value is unknown, but we can describe the probabilities of its possible values, is called a *random variable*.

Definition

A quantity whose value is unknown, but we can describe the probabilities of its possible values, is called a *random variable*.

Formally, any unknown X for which we can find the probability that

- $\mathbb{P}[X = 4]$,
- $\mathbb{P}[X \leq 10]$,
- $\mathbb{P}[6 > X > 3.1]$,
- etc.,

is a random variable.

Random Variables

Definition

A quantity whose value is unknown, but we can describe the probabilities of its possible values, is called a *random variable*.

Formally, any unknown X for which we can find the probability that

- $\mathbb{P}[X = 4]$,
- $\mathbb{P}[X \leq 10]$,
- $\mathbb{P}[6 > X > 3.1]$,
- etc.,

is a random variable. In other words, any quantity X about which we can answer questions like *"What is the probability that X is less than or equal to k ?"* or *"What is the probability that X equals k ?"* is a random variable.

Random Variables (optional)

Technically speaking, this means there should be a way to measure the probabilities

$$\mathbb{P}[X \leq k] \quad \text{or} \quad \mathbb{P}[X = k] \quad \text{or} \quad \mathbb{P}[X > k]$$

for any number k .

Random Variables (optional)

Technically speaking, this means there should be a way to measure the probabilities

$$\mathbb{P}[X \leq k] \quad \text{or} \quad \mathbb{P}[X = k] \quad \text{or} \quad \mathbb{P}[X > k]$$

for any number k .

Recall: The sets A for which $\mathbb{P}[A]$ can be measured are called *events*, and we write it as $A \in \mathcal{F}$.

Random Variables (optional)

Technically speaking, this means there should be a way to measure the probabilities

$$\mathbb{P}[X \leq k] \quad \text{or} \quad \mathbb{P}[X = k] \quad \text{or} \quad \mathbb{P}[X > k]$$

for any number k .

Recall: The sets A for which $\mathbb{P}[A]$ can be measured are called *events*, and we write it as $A \in \mathcal{F}$.

In other words, if X is a random variable, the set of outcomes

$$\{\omega \text{ for which } X \leq k\}$$

should be an event.

Random Variables (optional)

Technically speaking, this means there should be a way to measure the probabilities

$$\mathbb{P}[X \leq k] \quad \text{or} \quad \mathbb{P}[X = k] \quad \text{or} \quad \mathbb{P}[X > k]$$

for any number k .

Recall: The sets A for which $\mathbb{P}[A]$ can be measured are called *events*, and we write it as $A \in \mathcal{F}$.

In other words, if X is a random variable, the set of outcomes

$$\{\omega \text{ for which } X \leq k\}$$

should be an event.

Should we worry about this technicality? Fortunately, no:
In practice, *every unknown thing* is a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

What is not a random variable?

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

What is not a random variable?

- Who threw the dice,

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

What is not a random variable?

- Who threw the dice,
- What is the capital of Armenia,

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

What is not a random variable?

- Who threw the dice,
- What is the capital of Armenia,
- What would show up if we threw a third die

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

What is not a random variable?

- Who threw the dice,
- What is the capital of Armenia,
- What would show up if we threw a third die

Example

Two fair dice are thrown.

- The number X on the first die is a random variable.
- The number Y on the second die is a random variable.
- The sum $X + Y$ is also a random variable.
- The product $X \cdot Y$ is also a random variable.
- The maximum of two numbers is a random variable.

What is not a random variable?

- Who threw the dice,
- What is the capital of Armenia,
- What would show up if we threw a third die

are **not** random variables.

Example

We throw a fair die and win \$1 if it is prime, lose \$1 if it is composite, and stay even otherwise. If we denote the outcome of the die by ω ,

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega \in \{1\} \end{cases}$$

Example

We throw a fair die and win \$1 if it is prime, lose \$1 if it is composite, and stay even otherwise. If we denote the outcome of the die by ω ,

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega \in \{1\} \end{cases}$$

Example

Two teams play football.

- The number of goals they score is a random variable.
- The difference between their scores is also a random variable.

Random Variables

Example

We throw a fair die and win \$1 if it is prime, lose \$1 if it is composite, and stay even otherwise. If we denote the outcome of the die by ω ,

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega \in \{1\} \end{cases}$$

Example

Two teams play football.


- The number of goals they score is a random variable.
- The difference between their scores is also a random variable.

Example

The number of letters in a randomly selected word is a random variable.

Suppose we toss a fair coin 3 times, and we are interested in:


- X = the number of heads
- Y = the number of tails



PMF

Suppose we toss a fair coin 3 times, and we are interested in:

- X = the number of heads
- Y = the number of tails



Question

What are the possible values of X and Y , with their probabilities?

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{TTT\} \\ 1 & \text{if } \omega \in \{HTT, THT, TTH\} \\ 2 & \text{if } \omega \in \{HHT, THH, HTH\} \\ 3 & \text{if } \omega \in \{HHH\} \end{cases}$$

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{TTT\} \\ 1 & \text{if } \omega \in \{HTT, THT, TTH\} \\ 2 & \text{if } \omega \in \{HHT, THH, HTH\} \\ 3 & \text{if } \omega \in \{HHH\} \end{cases}$$

with probabilities:

$$\mathbb{P}[X = 0] =$$

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{\text{TTT}\} \\ 1 & \text{if } \omega \in \{\text{HTT}, \text{THT}, \text{TTH}\} \\ 2 & \text{if } \omega \in \{\text{HHT}, \text{THH}, \text{HTH}\} \\ 3 & \text{if } \omega \in \{\text{HHH}\} \end{cases}$$

with probabilities:

$$\mathbb{P}[X = 0] = \frac{1}{8}$$

$$\mathbb{P}[X = 1] =$$

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{TTT\} \\ 1 & \text{if } \omega \in \{HTT, THT, TTH\} \\ 2 & \text{if } \omega \in \{HHT, THH, HTH\} \\ 3 & \text{if } \omega \in \{HHH\} \end{cases}$$

with probabilities:

$$\mathbb{P}[X = 0] = \frac{1}{8}$$

$$\mathbb{P}[X = 1] = \frac{3}{8}$$

$$\mathbb{P}[X = 2] =$$

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{TTT\} \\ 1 & \text{if } \omega \in \{HTT, THT, TTH\} \\ 2 & \text{if } \omega \in \{HHT, THH, HTH\} \\ 3 & \text{if } \omega \in \{HHH\} \end{cases}$$

with probabilities:

$$\mathbb{P}[X = 0] = \frac{1}{8}$$

$$\mathbb{P}[X = 1] = \frac{3}{8}$$

$$\mathbb{P}[X = 2] = \frac{3}{8}$$

$$\mathbb{P}[X = 3] =$$

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{TTT\} \\ 1 & \text{if } \omega \in \{HTT, THT, TTH\} \\ 2 & \text{if } \omega \in \{HHT, THH, HTH\} \\ 3 & \text{if } \omega \in \{HHH\} \end{cases}$$

with probabilities:

$$\mathbb{P}[X = 0] = \frac{1}{8}$$

$$\mathbb{P}[X = 1] = \frac{3}{8}$$

$$\mathbb{P}[X = 2] = \frac{3}{8}$$

$$\mathbb{P}[X = 3] = \frac{1}{8}$$

The possible values of X are 0, 1, 2, 3:

$$X = \begin{cases} 0 & \text{if } \omega \in \{\text{TTT}\} \\ 1 & \text{if } \omega \in \{\text{HTT, THT, TTH}\} \\ 2 & \text{if } \omega \in \{\text{HHT, THH, HTH}\} \\ 3 & \text{if } \omega \in \{\text{HHH}\} \end{cases}$$

with probabilities:

$$\mathbb{P}[X = 0] = \frac{1}{8} = p_X(0)$$

$$\mathbb{P}[X = 1] = \frac{3}{8} = p_X(1)$$

$$\mathbb{P}[X = 2] = \frac{3}{8} = p_X(2)$$

$$\mathbb{P}[X = 3] = \frac{1}{8} = p_X(3)$$

A convenient way to represent this information:

"What is the probability that $X = k$?" i.e. $\mathbb{P}[X = k]$

is to denote it as:

$$p_X(k)$$

and call it the PMF of X .

Definition

The function which takes k and returns the probability that $X = k$:

$$p_X(k) = \mathbb{P}[X = k]$$

is called the *probability mass function* or *PMF* of X .

A convenient way to represent this information:

"What is the probability that $X = k$?" i.e. $\mathbb{P}[X = k]$

is to denote it as:

$$p_X(k)$$

and call it the PMF of X .

Definition

The function which takes k and returns the probability that $X = k$:

$$p_X(k) = \mathbb{P}[X = k]$$

is called the *probability mass function* or *PMF* of X .

If you know the PMF of X , you know everything about the *distribution* of X (i.e. how likely each value is).

In our example, the PMF of X is:

k	$p_X(k)$
0	1/8
1	3/8
2	3/8
3	1/8

PMF

In our example, the PMF of X is:

k	$p_X(k)$
0	1/8
1	3/8
2	3/8
3	1/8

Question

What is $p_X(4)$?

PMF

In our example, the PMF of X is:

k	$p_X(k)$
0	1/8
1	3/8
2	3/8
3	1/8

Question

What is $p_X(4)$?

Question

What is the PMF of Y ?

PMF

In our example, the PMF of X is:

k	$p_X(k)$
0	1/8
1	3/8
2	3/8
3	1/8

Question

What is $p_X(4)$?

Question

What is the PMF of Y ?

Property

The sum of all probabilities in the PMF is

PMF

In our example, the PMF of X is:

k	$p_X(k)$
0	1/8
1	3/8
2	3/8
3	1/8

Question

What is $p_X(4)$?

Question

What is the PMF of Y ?

Property

The sum of all probabilities in the PMF is 1.

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam.

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] =$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15}.$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] =$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}[X = 2] =$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}[X = 2] = \frac{4}{15} \cdot \frac{3}{14}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}[X = 2] = \frac{4}{15} \cdot \frac{3}{14} = \frac{6}{105} \quad (\text{or just } 1 - \frac{11}{21} - \frac{44}{105})$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.


$$\mathbb{P}[X = 0] = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}[X = 1] = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$


$$\mathbb{P}[X = 2] = \frac{4}{15} \cdot \frac{3}{14} = \frac{6}{105} \quad (\text{or just } 1 - \frac{11}{21} - \frac{44}{105})$$

k (right answers)	$p_X(k)$
0	0.52
1	0.42
2	0.06

Let's look at the PMFs of X and Y , showing average travel times (in minutes) for two US states:




Let's look at the PMFs of X and Y , showing average travel times (in minutes) for two US states:



This is too much information! What if we just want to know the probability that the travel time is under 10 minutes:

$$\mathbb{P}[X \leq 10] = ?$$

Let's look at the PMFs of X and Y , showing average travel times (in minutes) for two US states:



This is too much information! What if we just want to know the probability that the travel time is under 10 minutes:

$$F_X(10) = \mathbb{P}[X \leq 10] = ?$$

The function that takes k and returns the probability that X is less than or equal to k :

$$\mathbb{P}[X \leq k]$$

is called the *cumulative distribution function* or *CDF* of X .

The function that takes k and returns the probability that X is less than or equal to k :

$$\mathbb{P}[X \leq k]$$

is called the *cumulative distribution function* or *CDF* of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}[X \leq k]$$

The function that takes k and returns the probability that X is less than or equal to k :

$$\mathbb{P}[X \leq k]$$

is called the *cumulative distribution function* or *CDF* of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}[X \leq k]$$

Properties

- For any $x \in \mathbb{R}$, $0 \leq F_X(x) \leq 1$

The function that takes k and returns the probability that X is less than or equal to k :

$$\mathbb{P}[X \leq k]$$

is called the *cumulative distribution function* or *CDF* of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}[X \leq k]$$

Properties

- For any $x \in \mathbb{R}$, $0 \leq F_X(x) \leq 1$
- $F_X(x)$ is non-decreasing

The function that takes k and returns the probability that X is less than or equal to k :

$$\mathbb{P}[X \leq k]$$

is called the *cumulative distribution function* or *CDF* of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}[X \leq k]$$

Properties

- For any $x \in \mathbb{R}$, $0 \leq F_X(x) \leq 1$
- $F_X(x)$ is non-decreasing
- $\lim_{x \rightarrow -\infty} F_X(x) = 0$ and $\lim_{x \rightarrow +\infty} F_X(x) = 1$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) =$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) =$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

$$F_X(5.9) =$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

$$F_X(5.9) = \frac{5}{6}$$

Example

Let X indicate the number on a fair die. Then


$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

$$F_X(5.9) = \frac{5}{6}$$

$$F_X(x) = \begin{cases} 0, & \text{if } x < 1, \\ \frac{1}{6}, & \text{if } 1 \leq x < 2, \\ \frac{2}{6}, & \text{if } 2 \leq x < 3, \\ \frac{3}{6}, & \text{if } 3 \leq x < 4, \\ \frac{4}{6}, & \text{if } 4 \leq x < 5, \\ \frac{5}{6}, & \text{if } 5 \leq x < 6, \\ 1, & \text{if } x \geq 6 \end{cases}$$

Graphs of CDFs usually look like this:



identically distributed RVs

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

identically distributed RVs

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- The students may be asked different questions and get different grades, so $X \neq Y$.

identically distributed RVs

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- The students may be asked different questions and get different grades, so $X \neq Y$.
- However, Y has the same PMF and CDF as X (check it!)

identically distributed RVs

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- The students may be asked different questions and get different grades, so $X \neq Y$.
- However, Y has the same PMF and CDF as X (check it!)

identically distributed RVs

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- The students may be asked different questions and get different grades, so $X \neq Y$.
- However, Y has the same PMF and CDF as X (check it!)

In this case, we say that X and Y are identically distributed:

Definition

Two random variables are *identically distributed* if their PMFs/CDFs are equal.

identically distributed RVs

In some sense, identically distributed random variables are "similar" to each other, **but not necessarily** equal.

Example

Say we toss a fair coin. X and Y are RVs such that:

$$X = \begin{cases} 0, & \text{if } \omega = H, \\ 1, & \text{if } \omega = T, \end{cases} \quad Y = \begin{cases} 1, & \text{if } \omega = H, \\ 0, & \text{if } \omega = T, \end{cases}$$

They both have the same PMFs:

$$\mathbb{P}[X = 0] = \mathbb{P}[X = 1] = \mathbb{P}[Y = 0] = \mathbb{P}[Y = 1] = \frac{1}{2}$$

identically distributed RVs

In some sense, identically distributed random variables are "similar" to each other, **but not necessarily** equal.

Example

Say we toss a fair coin. X and Y are RVs such that:

$$X = \begin{cases} 0, & \text{if } \omega = H, \\ 1, & \text{if } \omega = T, \end{cases} \quad Y = \begin{cases} 1, & \text{if } \omega = H, \\ 0, & \text{if } \omega = T, \end{cases}$$

They both have the same PMFs:

$$\mathbb{P}[X = 0] = \mathbb{P}[X = 1] = \mathbb{P}[Y = 0] = \mathbb{P}[Y = 1] = \frac{1}{2} \quad \text{but } X \neq Y$$

Random Variables

So far we have only seen random variables that take a **finite** or **countable** number of values, like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$ – because they showed some quantities like the number of heads, number of goals, etc.

Random Variables

So far we have only seen random variables that take a **finite** or **countable** number of values, like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$ – because they showed some quantities like the number of heads, number of goals, etc. However, random variables showing e.g. time, distance, coordinates, etc. can take **uncountably** many values as well – i.e. values in intervals like $(0, +\infty)$. For example, the time until the bus arrives can be any real number between 0 and 60 minutes, so the possible values form an interval.

Random Variables

So far we have only seen random variables that take a **finite** or **countable** number of values, like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$ – because they showed some quantities like the number of heads, number of goals, etc. However, random variables showing e.g. time, distance, coordinates, etc. can take **uncountably** many values as well – i.e. values in intervals like $(0, +\infty)$. For example, the time until the bus arrives can be any real number between 0 and 60 minutes, so the possible values form an interval.

Definition

If the values of X can be represented as a list (say, x_1, x_2, x_3, \dots with probabilities 0.2, 0.1, 0.4, \dots), it is called a *discrete random variable*.

Random Variables

So far we have only seen random variables that take a **finite** or **countable** number of values, like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$ – because they showed some quantities like the number of heads, number of goals, etc. However, random variables showing e.g. time, distance, coordinates, etc. can take **uncountably** many values as well – i.e. values in intervals like $(0, +\infty)$. For example, the time until the bus arrives can be any real number between 0 and 60 minutes, so the possible values form an interval.

Definition

If the values of X can be represented as a list (say, x_1, x_2, x_3, \dots with probabilities 0.2, 0.1, 0.4, \dots), it is called a *discrete random variable*.

Otherwise,

Definition

If the values of X cannot be represented as a list (e.g. they are an interval), it is called a *continuous random variable*.

Example

Examples of discrete random variables:

- the sum of two consecutive dice rolls,
- the number of goals in a football match,
- the difference between the scores of two teams,
- the number of children in a randomly selected family,
- the number of customers that enter a shop on a given day.

Random Variables

Example

Examples of discrete random variables:

- the sum of two consecutive dice rolls,
- the number of goals in a football match,
- the difference between the scores of two teams,
- the number of children in a randomly selected family,
- the number of customers that enter a shop on a given day.

Example

Examples of continuous random variables:

- the time until a bus arrives,
- the height of a randomly selected person,
- the stocks of Apple tomorrow,
- the x -coordinate of a randomly selected point on the (x, y) -plane.

Okay, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}[X = 0.5]$?

Okay, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}[X = 0.5]$? What about $\mathbb{P}[X = 0.589]$?

Okay, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}[X = 0.5]$? What about $\mathbb{P}[X = 0.589]$? As previously, it is 0.

Okay, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}[X = 0.5]$? What about $\mathbb{P}[X = 0.589]$? As previously, it is 0.

What can we say about the PMF of X ?

Okay, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}[X = 0.5]$? What about $\mathbb{P}[X = 0.589]$? As previously, it is 0.

What can we say about the PMF of X ? It is always zero!

Okay, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}[X = 0.5]$? What about $\mathbb{P}[X = 0.589]$? As previously, it is 0.

What can we say about the PMF of X ? It is always zero!

So another question we might ask is:

Question

What do you think $\mathbb{P}[X \leq 5]$ is?

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}[X \leq 5] = \frac{1}{2}$$

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}[X \leq 5] = \frac{1}{2}$$

Fortunately, we have another tool in case of continuous random variables.

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}[X \leq 5] = \frac{1}{2}$$

Fortunately, we have another tool in case of continuous random variables.

Definition

If X is a continuous random variable, then there exists a function $f(x) \geq 0$ such that for any $c \in \mathbb{R}$,

$$F_X(c) = \mathbb{P}[X \leq c] = \int_{-\infty}^c f(t) dt$$

The function f is called the *probability density function* or *PDF* of X .

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}[X \leq 5] = \frac{1}{2}$$

Fortunately, we have another tool in case of continuous random variables.

Definition

If X is a continuous random variable, then there exists a function $f(x) \geq 0$ such that for any $c \in \mathbb{R}$,

$$F_X(c) = \mathbb{P}[X \leq c] = \int_{-\infty}^c f(t) dt$$

The function f is called the *probability density function* or *PDF* of X .


The PDF of a continuous random variable plays essentially the same role as the PMF of a discrete random variable.

Just like the density of an object measures the concentration of mass (per unit volume), the probability density function captures the density of *probability* at point x :

$$f_X(x) = \lim_{h \rightarrow 0} \frac{\mathbb{P}[x < X \leq x + h]}{h}$$

Just like the density of an object measures the concentration of mass (per unit volume), the probability density function captures the density of *probability* at point x :

$$f_X(x) = \lim_{h \rightarrow 0} \frac{\mathbb{P}[x < X \leq x + h]}{h}$$



Properties

- 1 $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Properties

① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

② $\int_{-\infty}^{\infty} f_X(x) dx = 1$

Properties

- ① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$
- ② $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- ③ If F_X is continuous at x , $F'_X(x) = f_X(x)$

Properties

- ① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$
- ② $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- ③ If F_X is continuous at x , $F'_X(x) = f_X(x)$
- ④ $\mathbb{P}[a < X \leq b] = F_X(b) - F_X(a) = \int_a^b f_X(x) dx$

Properties

- ① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$
- ② $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- ③ If F_X is continuous at x , $F'_X(x) = f_X(x)$
- ④ $\mathbb{P}[a < X \leq b] = F_X(b) - F_X(a) = \int_a^b f_X(x) dx$
- ⑤ $\mathbb{P}[X = a] = \mathbb{P}[X \leq a] - \mathbb{P}[X < a]$

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

For $x < a$, $F_X(x) = \mathbb{P}[X \leq x] = 0$,

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

For $x < a$, $F_X(x) = \mathbb{P}[X \leq x] = 0$,

For $x \geq b$, $F_X(x) = \mathbb{P}[X \leq x] = 1$,

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.


Let's find the CDF and PDF of X .


For $x < a$, $F_X(x) = \mathbb{P}[X \leq x] = 0$,

For $x \geq b$, $F_X(x) = \mathbb{P}[X \leq x] = 1$,

For $a \leq x \leq b$, we have:

$$F_X(x) = \mathbb{P}[X \leq x] = \mathbb{P}[X \in [a, x]] = \frac{x - a}{b - a}$$





Thus,


$$F_X(x) = \begin{cases} 0 & \text{if } x < a \\ \frac{x-a}{b-a} & \text{if } a \leq x \leq b \\ 1 & \text{if } x > b \end{cases}$$

To find $f_X(x)$, we take the derivative of $F_X(x)$:

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a < x < b \\ 0 & \text{if } x < a \text{ or } x > b \end{cases}$$

To find $f_X(x)$, we take the derivative of $F_X(x)$:

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a < x < b \\ 0 & \text{if } x < a \text{ or } x > b \end{cases}$$



Summary

To summarize, all random variables come in two types:

	PMF	PDF	CDF
discrete	✓	—	✓
continuous	—	✓	✓

Independence

Recall that two events A and B are called independent if they do not affect each other, i.e.

$$\mathbb{P}[A \cap B] = \mathbb{P}[A] \cdot \mathbb{P}[B]$$

Independence

Recall that two events A and B are called independent if they do not affect each other, i.e.

$$\mathbb{P}[A \cap B] = \mathbb{P}[A] \cdot \mathbb{P}[B]$$

Similarly, we will say that two random variables X and Y are independent if their values do not affect each other (i.e. if you know the value of X , it gives you no information about the value of Y).

Independence

Recall that two events A and B are called independent if they do not affect each other, i.e.

$$\mathbb{P}[A \cap B] = \mathbb{P}[A] \cdot \mathbb{P}[B]$$

Similarly, we will say that two random variables X and Y are independent if their values do not affect each other (i.e. if you know the value of X , it gives you no information about the value of Y).

Definition

X and Y are called *independent* if

$$\mathbb{P}[X \leq a \text{ and } Y \leq b] = \mathbb{P}[X \leq a] \cdot \mathbb{P}[Y \leq b]$$

for any $a, b \in \mathbb{R}$.

So the probability of both X and Y simultaneously being less than some numbers is just their *separate* probabilities multiplied together.