Practice 1 — Set Theory

Solutions to Selected Problems

Problems 2, 7, 11, 16, 20, 54 (3, 11), 58, 60

Discrete Mathematics — QYSU

Contents

1 Problem 2 — Membership with $\varnothing$ and $\{\varnothing\}$ **2**

2 Problem 7 — $\in$ is Not Transitive **3**

3 Problem 11 — Mixing $\in$ and $\subseteq$ **4**

4 Problem 16 — Describing Sets Under Operations **6**

5 Problem 20 — Equivalent Characterizations of $\subseteq$ **8**

6 Problem 54.3 — $(A \cup B) \setminus C = (A \setminus C) \cup (B \setminus C)$ **11**

7 Problem 54.11 — $\cap$ Distributes Over $\triangle$ **13**

8 Problem 58 — $A \times B = B \times A \implies A = B$ **15**

9 Problem 60 — $\times$ Distributes Over Set Operations **17**

1 Problem 2 — Membership with $\varnothing$ and $\{\varnothing\}$

Theory Recap: Membership ($\in$) vs. Subset ($\subseteq$)

These two relations are fundamentally different:

- $a \in S$ means "a is an **element** of S."
- $A \subseteq S$ means "every element of A is also in S."

The empty set $\varnothing$ has no elements. The set $\{\varnothing\}$ is **not** empty—it has one element, namely $\varnothing$ itself. To decide "$x \in S$?" list the elements of S and check whether x is one of them.



Each wrapping step creates a new set whose sole element is the previous set.

Problem Statement

Which of the following statements are true?

2.1. $\varnothing \in \{\varnothing,\ \{\varnothing\}\}$

2.2. $\{\varnothing\} \in \{\{\varnothing\}\}$

2.3. $\varnothing \in \{\{\varnothing\}\}$

Box Analogy: How $\in$ Works

Think of every set as a **box**. The $\in$ operator opens the box and looks at what is *directly* inside—it only looks **one box deep**. It never opens inner boxes.

- $\varnothing$ is an **empty box**—you open it and find nothing inside. Therefore $|\varnothing| = 0$.
- $\{\varnothing\}$ is a **box containing one item**: another (empty) box. You open $\{\varnothing\}$ and see one thing sitting there—the empty box $\varnothing$. Therefore $|\{\varnothing\}| = 1$.

This is why $\varnothing \neq \{\varnothing\}$: an empty box is not the same as a box that contains an empty box, just as an empty envelope is not the same as an envelope containing another empty envelope.
When you test "$x \in S$?", open the box S, look at the items directly visible (one level deep), and check whether x is one of those items. **Never unwrap items inside items.**

Method. List the elements of each outer set first: $\{\varnothing,\ \{\varnothing\}\}$, $\{\{\varnothing\}\}$. Then check whether the left-hand object matches any listed element.

Solution

The strategy is always the same: *list the elements* of the outer set, then check whether the object on the left of $\in$ matches any of them.



Left object	Elements of outer set	Match?
$\varnothing$	$\varnothing$, $\{\varnothing\}$	Yes — **TRUE**
$\{\varnothing\}$	$\{\varnothing\}$	Yes — **TRUE**
$\varnothing$	$\{\varnothing\}$	No ($\varnothing \neq \{\varnothing\}$) — **FALSE**

2.1. $\varnothing \in \{\varnothing, \{\varnothing\}\}$ — **TRUE.** The set has two elements: $\varnothing$ and $\{\varnothing\}$. The object $\varnothing$ is explicitly one of them.

2.2. $\{\varnothing\} \in \{\{\varnothing\}\}$ — **TRUE.** The set $\{\{\varnothing\}\}$ has exactly one element: $\{\varnothing\}$. The object on the left is $\{\varnothing\}$, which matches.

2.3. $\varnothing \in \{\{\varnothing\}\}$ — **FALSE.** The only element is $\{\varnothing\}$. We need $\varnothing = \{\varnothing\}$, but $|\varnothing| = 0$ while $|\{\varnothing\}| = 1$, so they are not equal.

2.1 — True 2.2 — True 2.3 — False

2 Problem 7 — $\in$ is Not Transitive

Theory Recap: $\in$ is Not Transitive

$\subseteq$ **is** transitive: $A \subseteq B \wedge B \subseteq C \implies A \subseteq C$.
$\in$ is **not**: $A \in B \wedge B \in C \not\implies A \in C$.

Why? $\in$ crosses one "level" of nesting. Two $\in$-steps cross *two* levels, so A is buried inside C but is not a direct element of C.



Problem Statement

Give examples of sets A, B, C such that $A \in B$, $B \in C$, but $A \notin C$.

Real-World Analogy: Level Jumps

A person, a committee, and a task force.
Suppose Professor A is a *member* of the Curriculum Committee B, and the Curriculum Committee B is a *member* of the University Task Force C (a body whose members are entire committees, not individual people).

- $A \in B$: the professor sits on the committee.
- $B \in C$: the committee belongs to the task force.
- $A \notin C$: the professor is **not** individually a member of the task force—only whole committees are members of the task force.

The key insight: $\in$ crosses exactly **one level** of containment. Two $\in$-steps cross *two* levels, so the individual (A) is buried two levels deep inside C but is not a *direct* member of C.

Solution

Let

$$A = 1, \qquad B = \{1, 2\}, \qquad C = \big\{\{1, 2\},\ 3\big\}.$$

Condition	Check	Result
$A \in B$	Is $1 \in \{1, 2\}$?	Yes ✓
$B \in C$	Is $\{1, 2\} \in \big\{\{1, 2\},\ 3\big\}$?	Yes ✓
$A \notin C$	Is $1 \in \big\{\{1, 2\},\ 3\big\}$? The elements are $\{1, 2\}$ and 3.	$1 \notin C$ ✓

Key insight. The number 1 is an element of B, and B is an element of C, but the elements of C are $\{1, 2\}$ (a set) and 3 (a number). The number 1 is neither of these, so $1 \notin C$.

Contrast. The conclusion $A \notin C$ is not automatic—it can fail. Let $A = 1$, $B = \{1\}$, $C = \{1, \{1\}\}$. Then $A \in B$ ($1 \in \{1\}$), $B \in C$ ($\{1\} \in \{1, \{1\}\}$), and also $A \in C$ ($1 \in \{1, \{1\}\}$). So $\in$ is *sometimes* transitive by accident, but not always.

3 Problem 11 — Mixing $\in$ and $\subseteq$

Theory Recap: Mixing $\in$ and $\subseteq$

The two relations operate at different "levels":

- $A \subseteq B$: both sit at the same level—every element of A is also an element of B.
- $B \in C$: B itself becomes an object inside C—a jump *up* one level.



Problem Statement

Give examples of sets A, B, C, D, E satisfying:

$$A \subseteq B, \quad B \in C, \quad C \subseteq D, \quad D \subseteq E.$$

Solution

Color Guide: Seeing the Nesting Levels

When sets contain other sets as elements, it helps to color-code the levels:

- Blue = plain numbers (lowest level objects).
- Red braces = set-braces of an inner set that is itself an *element* of an outer set.
- Black braces = the outer set's own boundary.

For example, $C = \big\{\{1,2\},\ 3\big\}$ has two elements: the set $\{1,2\}$ (an inner set, marked in red braces) and the number 3 (a plain number in blue).

Similarly:

$$D = \big\{\{1,2\},\ 3,\ 4\big\}, \qquad E = \big\{\{1,2\},\ 3,\ 4,\ 5\big\}$$

In each case, the red-braced item is a set-element (one level up), while the blue items are number-elements.

$$A = \{1\}, \quad B = \{1,2\}, \quad C = \big\{\{1,2\},\ 3\big\}, \quad D = \big\{\{1,2\},\ 3,\ 4\big\}, \quad E = \big\{\{1,2\},\ 3,\ 4,\ 5\big\}$$

Condition	Type	Verification
$A \subseteq B$	subset	Every element of $\{1\}$ is in $\{1,2\}$: $1 \in \{1,2\}$ ✓
$B \in C$	membership	$\{1,2\}$ is listed as an element of $\big\{\{1,2\},3\big\}$ ✓
$C \subseteq D$	subset	Elements $\{1,2\}$ and 3 of C are both in D ✓
$D \subseteq E$	subset	Elements $\{1,2\}$, 3, 4 of D are all in E ✓

Level structure.

- A, B: sets of *numbers* (elements are $1, 2$).
- C, D, E: sets whose elements *include a set* ($\{1,2\}$ is an element alongside numbers like $3, 4, 5$).

The $\in$-step from B to C is where the level jump happens.

Why this works. $\subseteq$ preserves the "level": A and B contain the same kind of objects (numbers). The $\in$-step promotes B to an object inside C, after which C, D, E all live at the higher level (their elements include sets like $\{1, 2\}$).

4 Problem 16 — Describing Sets Under Operations

Theory Recap: Set Operations

For subsets of a universal set U:

Operation	Symbol	Definition
Complement	$\overline{A}$	$\{x \in U \mid x \notin A\}$
Set difference	$A \setminus B$	$\{x \in A \mid x \notin B\}$
Symmetric diff.	$A \triangle B$	$(A \setminus B) \cup (B \setminus A)$ = elements in exactly one of A, B



Complement $\quad A \triangle B \quad A \setminus B$

Problem Statement

Let $\mathbb{Z}$ be the universal set. Define:

$$A = \{x \in \mathbb{Z} \mid \exists\, y \in \mathbb{Z}^+,\ x = 2y\} \qquad \text{(positive even integers: } 2, 4, 6, 8, \ldots)$$
$$B = \{x \in \mathbb{Z} \mid \exists\, y \in \mathbb{Z}^+,\ x = 2y - 1\} \qquad \text{(positive odd integers: } 1, 3, 5, 7, \ldots)$$
$$C = \{x \in \mathbb{Z} \mid x < 10\}$$

Describe: $\overline{A}$, $A \triangle B$, $\overline{C}$, $A \setminus C$, $C \setminus (A \triangle B)$.

Solution

Visualising the sets on the number line.



Translate each operation into words first:

- $\overline{A}$: integers that are **not** positive even.

- $A \triangle B$: integers in **exactly one** of A, B.
- $\overline{C}$: integers that are **not** less than 10.
- $A \setminus C$: positive evens that are **not** in C (i.e. ≥ 10).
- $C \setminus (A \triangle B)$: integers < 10 that are **not** positive.

(a) $\overline{A}$**:** Everything in $\mathbb{Z}$ that is *not* a positive even integer:

$$\boxed{\overline{A} = \{\ldots, -2, -1, 0\} \cup \{1, 3, 5, 7, \ldots\} = \big\{x \in \mathbb{Z} \mid x \leq 0 \text{ or } x \text{ is odd}\big\}}$$

(b) $A \triangle B$**:** Since $A \cap B = \varnothing$ (no integer is both positive-even and positive-odd):

$$A \triangle B = A \cup B = \{1, 2, 3, 4, \ldots\}$$

Since every positive integer is either even or odd, and never both, $A \cup B = \mathbb{Z}^+$ and $A \cap B = \varnothing$.

$$\boxed{A \triangle B = \mathbb{Z}^+}$$

(c) $\overline{C}$**:** All integers ≥ 10: $\boxed{\overline{C} = \{10, 11, 12, \ldots\}}$

(d) $A \setminus C$**:** Positive evens that are ≥ 10: $\boxed{A \setminus C = \{10, 12, 14, \ldots\} = \{2k \mid k \geq 5\}}$

(e) $C \setminus (A \triangle B)$**:** This is a compound operation—evaluate it step by step:

Step 1: Evaluate the inner parentheses: $A \triangle B$. From part (b) above, we already know $\boxed{A \triangle B = \mathbb{Z}^+}$ (all positive integers).

Step 2: Substitute back into the outer operation: $C \setminus (A \triangle B) = C \setminus \mathbb{Z}^+$. This means: take all elements of C (integers < 10) and remove every positive integer.

Step 3: Compute the result: The elements of C that are *not* positive are $\ldots, -2, -1, 0$.

$$\boxed{C \setminus (A \triangle B) = \{\ldots, -2, -1, 0\} = \{x \in \mathbb{Z} \mid x \leq 0\}}$$

5 Problem 20 — Equivalent Characterizations of $\subseteq$

Theory Recap: Cyclic-Implication Proof Strategy

To show three conditions (a), (b), (c) are equivalent, it suffices to prove a **cycle of implications**:

$$(a) \implies (b) \implies (c) \implies (a).$$

This is more efficient than proving all six pairwise implications directly.



Problem Statement

Prove that for any subsets A, B of universal set U, the conditions in each group are equivalent:

1. $A \subseteq B \iff A \cup B = B \iff A \cap B = A$
2. $A \cap B = \varnothing \iff A \subseteq \overline{B} \iff B \subseteq \overline{A}$
3. $A \cup B = U \iff \overline{A} \subseteq B \iff \overline{B} \subseteq A$

Visual Intuition: Why These Conditions Are Equivalent

Before diving into the formal proof, consider this Venn diagram showing $A \subseteq B$ (i.e. A sits entirely inside B):



- **Why $A \cup B = B$?** Since A is already inside B, adding A to the union contributes nothing new. The shaded region of $A \cup B$ is just B.
- **Why $A \cap B = A$?** Every element of A is also in B (because A is inside B), so intersecting with B keeps all of A. Nothing from A gets removed.

This picture makes all three conditions feel like the same statement in different words. The formal cyclic-implication proof below confirms this rigorously.

Part 1: $A \subseteq B \iff A \cup B = B \iff A \cap B = A$



Key idea: if A is already inside B, adding A to the union changes nothing, and intersecting with B keeps all of A.

$(a) \Rightarrow (b)$**:** $A \subseteq B \implies A \cup B = B$.

$(\supseteq)$ $B \subseteq A \cup B$ always.

$(\subseteq)$ Let $x \in A \cup B$. Then $x \in A$ or $x \in B$. If $x \in A$, then $A \subseteq B$ gives $x \in B$. Either way, $x \in B$. $\square$

$(b) \Rightarrow (c)$**:** $A \cup B = B \implies A \cap B = A$.

$(\subseteq)$ $A \cap B \subseteq A$ always.

$(\supseteq)$ Let $x \in A$. Then $x \in A \cup B = B$, so $x \in B$. Hence $x \in A \cap B$. $\square$

$(c) \Rightarrow (a)$**:** $A \cap B = A \implies A \subseteq B$.

Let $x \in A$. Then $x \in A = A \cap B$, so $x \in B$. $\square$

Part 2: $A \cap B = \varnothing \iff A \subseteq \overline{B} \iff B \subseteq \overline{A}$



Disjoint: $A \cap B = \varnothing$

Key idea: disjointness means every element of A lies outside B, i.e. in $\overline{B}$.

$(a) \Rightarrow (b)$**:** $A \cap B = \varnothing \implies A \subseteq \overline{B}$.

Let $x \in A$. If $x \in B$, then $x \in A \cap B = \varnothing$—contradiction. So $x \notin B$, i.e. $x \in \overline{B}$. $\square$

$(b) \Rightarrow (c)$**:** $A \subseteq \overline{B} \implies B \subseteq \overline{A}$.

Let $x \in B$. If $x \in A$, then $x \in \overline{B}$ (since $A \subseteq \overline{B}$), so $x \notin B$—contradiction. Hence $x \notin A$, i.e. $x \in \overline{A}$. $\square$

$(c) \Rightarrow (a)$**:** $B \subseteq \overline{A} \implies A \cap B = \varnothing$.

Suppose $x \in A \cap B$. Then $x \in B \subseteq \overline{A}$, so $x \notin A$—contradicting $x \in A \cap B$. $\square$

Part 3: $A \cup B = U \iff \overline{A} \subseteq B \iff \overline{B} \subseteq A$



Covering: $A \cup B = U$

Key idea: if A and B together cover U, then anything outside A must be in B.

$(a) \Rightarrow (b)$**:** $A \cup B = U \implies \overline{A} \subseteq B$.

Let $x \in \overline{A}$, so $x \notin A$. Since $x \in U = A \cup B$ and $x \notin A$, we must have $x \in B$. $\square$

$(b) \Rightarrow (c)$**:** $\overline{A} \subseteq B \implies \overline{B} \subseteq A$.

Let $x \in \overline{B}$, so $x \notin B$. If $x \notin A$, then $x \in \overline{A} \subseteq B$, giving $x \in B$—contradiction. So $x \in A$. $\square$

$(c) \Rightarrow (a)$**:** $\overline{B} \subseteq A \implies A \cup B = U$.

Let $x \in U$. Either $x \in B$ (then $x \in A \cup B$) or $x \notin B$ (then $x \in \overline{B} \subseteq A$, so $x \in A \cup B$). Since also $A \cup B \subseteq U$, we get $A \cup B = U$. $\square$

6 Problem 54.3 — $(A \cup B) \setminus C = (A \setminus C) \cup (B \setminus C)$

Theory Recap: Set Difference Distributes Over Union

Mental model: "$X \setminus C$" means "start with X, **remove** everything in C."
Removing C from a union is the same as removing C from each piece separately, then taking the union of the remainders.



Both diagrams shade exactly the same region—the parts of A and B that fall outside C.

Problem Statement

Prove: $(A \cup B) \setminus C = (A \setminus C) \cup (B \setminus C)$.

Solution

Let x be arbitrary:

$$\begin{aligned}
x \in (A \cup B) \setminus C &\iff x \in (A \cup B) \ \wedge \ x \notin C \\
&\iff (x \in A \ \vee \ x \in B) \ \wedge \ x \notin C \\
&\iff (x \in A \wedge x \notin C) \ \vee \ (x \in B \wedge x \notin C) \qquad (\wedge \text{ distributes over } \vee) \\
&\iff x \in (A \setminus C) \ \vee \ x \in (B \setminus C) \\
&\iff x \in (A \setminus C) \cup (B \setminus C).
\end{aligned}$$

Every step is $\iff$, so the sets are equal. $\square$

Set-language reading. Distributing "outside C" across "in A or in B" is the set version of the logical law $p \wedge (q \vee r) \iff (p \wedge q) \vee (p \wedge r)$.

Region Numbering Method — Empirical Verification

Technique. In a three-set Venn diagram there are $2^3 = 8$ regions. Number them by which sets they belong to:

Region	Membership
1	A only
2	B only
3	C only
4	$A \cap B$ only (not C)
5	$A \cap C$ only (not B)
6	$B \cap C$ only (not A)
7	$A \cap B \cap C$
8	None ($\overline{A} \cap \overline{B} \cap \overline{C}$)

Now express each side by its region numbers:

- $A \cup B = \{1, 2, 4, 5, 6, 7\}$
- $(A \cup B) \setminus C$: remove regions in C ($\{3, 5, 6, 7\}$) $\implies \boxed{\{1, 2, 4\}}$
- $A \setminus C = \{1, 4\}, \quad B \setminus C = \{2, 4\}$
- $(A \setminus C) \cup (B \setminus C) = \{1, 4\} \cup \{2, 4\} = \boxed{\{1, 2, 4\}}$

Both sides give exactly regions $\{1, 2, 4\}$, confirming the identity empirically.

This method does not replace a proof, but it is an excellent way to build confidence or catch errors before writing one.

Logic correspondence. The key step uses the propositional law $p \wedge (q \vee r) \iff (p \wedge q) \vee (p \wedge r)$, where $p = (x \notin C)$, $q = (x \in A)$, $r = (x \in B)$.

7 Problem 54.11 — $\cap$ Distributes Over $\triangle$

Theory Recap: Intersection Distributes Over Symmetric Difference

Recall: $X \triangle Y = \{x \mid x \text{ belongs to exactly one of } X, Y\}$.

The identity $A \cap (B \triangle C) = (A \cap B) \triangle (A \cap C)$ says:

> *"The part of A that sees exactly one of B, C"* equals *"exactly one of (A's overlap with B) and (A's overlap with C)."*



Algebraic analogy. Under the correspondence $\cap \leftrightarrow \cdot$ and $\triangle \leftrightarrow +$ (mod 2), the power set $(\mathcal{P}(U), \triangle, \cap)$ forms a commutative ring. This identity is the ring distributive law.

Problem Statement

Prove: $A \cap (B \triangle C) = (A \cap B) \triangle (A \cap C)$.

Intermediate Shading Steps — Building the Venn Diagram

To build the Venn diagram for $A \cap (B \triangle C)$ from scratch, follow these stages:

Stage 1: Shade $B \triangle C$: Shade all regions that belong to exactly one of B or C (i.e. B-only and C-only regions, but *not* the $B \cap C$ overlap). Using the region numbering from the previous problem (8 regions for three sets): B occupies regions $\{2, 4, 6, 7\}$ and C occupies regions $\{3, 5, 6, 7\}$. Therefore $B \setminus C = \{2, 4\}$ and $C \setminus B = \{3, 5\}$, so $B \triangle C = (B \setminus C) \cup (C \setminus B)$ covers regions $\{2, 3, 4, 5\}$.

Stage 2: Identify A: The regions inside A are $\{1, 4, 5, 7\}$.

Stage 3: Intersect—highlight the overlap: $A \cap (B \triangle C)$ is the intersection of stages 1 and 2: $\{2, 3, 4, 5\} \cap \{1, 4, 5, 7\} = \{4, 5\}$. These are exactly region 4 ($A \cap B \cap \overline{C}$) and region 5 ($A \cap C \cap \overline{B}$)—the two shaded regions in the diagram above.

The same two regions emerge from the RHS: $(A \cap B) \triangle (A \cap C) = \{4, 7\} \triangle \{5, 7\} = \{4, 5\}$.

Solution

Let x be arbitrary. We expand both sides to the same logical form.

Left-hand side:

$$\begin{aligned} x \in A \cap (B \triangle C) &\iff x \in A \ \wedge\ x \in B \triangle C \\ &\iff x \in A \ \wedge\ \big[(x \in B \wedge x \notin C) \ \vee\ (x \notin B \wedge x \in C)\big] \\ &\iff \underbrace{(x \in A \wedge x \in B \wedge x \notin C)}_{\text{case 1}} \vee \underbrace{(x \in A \wedge x \notin B \wedge x \in C)}_{\text{case 2}} \qquad (\star) \end{aligned}$$

Right-hand side:

$$x \in (A \cap B) \triangle (A \cap C) \iff \big[x \in A \cap B \ \wedge\ x \notin A \cap C\big] \ \vee\ \big[x \notin A \cap B \ \wedge\ x \in A \cap C\big]$$

First disjunct: $x \in A \cap B$ and $x \notin A \cap C$.

- $x \in A \cap B \implies x \in A$ and $x \in B$.
- $x \notin A \cap C$ means $\neg(x \in A \wedge x \in C)$. Since $x \in A$, this forces $x \notin C$.
- Combined: $x \in A,\ x \in B,\ x \notin C$ (= case 1).

Second disjunct: $x \notin A \cap B$ and $x \in A \cap C$.

- $x \in A \cap C \implies x \in A$ and $x \in C$.
- $x \notin A \cap B$ means $\neg(x \in A \wedge x \in B)$. Since $x \in A$, this forces $x \notin B$.
- Combined: $x \in A,\ x \notin B,\ x \in C$ (= case 2).

Both sides reduce to $(\star)$, so:

$$A \cap (B \triangle C) = (A \cap B) \triangle (A \cap C). \qquad \blacksquare$$

Verification by truth table. Every element belongs to some combination of A, B, C. The table below checks that LHS and RHS agree in all cases.

$x \in A$	$x \in B$	$x \in C$	LHS: $A \cap (B \triangle C)$	RHS: $(A \cap B) \triangle (A \cap C)$
✓	✓	✓	×	×
✓	✓	×	✓	✓
✓	×	✓	✓	✓
✓	×	×	×	×
×	✓	✓	×	×
×	✓	×	×	×
×	×	✓	×	×
×	×	×	×	×

All rows match, confirming the identity.

8 Problem 58 — $A \times B = B \times A \implies A = B$

Theory Recap: Cartesian Product

$$A \times B = \{(a, b) \mid a \in A,\ b \in B\}.$$

Order in the pair matters: $(a, b) \neq (b, a)$ when $a \neq b$.



When $A \neq B$, the grids have different shapes, so $A \times B \neq B \times A$.

Problem Statement

Prove that $A \times B = B \times A$ implies $A = B$.

Note: More precisely, $A \times B = B \times A$ implies $A = B$, or $A = \varnothing$, or $B = \varnothing$.

Concrete Mini-Example: Why $A \neq B$ Breaks Equality

Let $A = \{1, 2\}$ and $B = \{3\}$. Then:

$$A \times B = \{(1, 3),\ (2, 3)\}, \qquad B \times A = \{(3, 1),\ (3, 2)\}.$$

Visualise these as grids:



The grids have different "shapes" (2×1 vs. 1×2), and the pairs themselves are different: $(1, 3) \neq (3, 1)$. So $A \times B \neq B \times A$ whenever $A \neq B$ (and both are non-empty).

Solution

Hidden assumption. We first assume $A \neq \varnothing$ and $B \neq \varnothing$, because if either is empty, both products are $\varnothing$ and the equality holds trivially without $A = B$. (Example: $A = \varnothing$, $B = \{1\}$.)

Assume $A \times B = B \times A$ with $A \neq \varnothing$ and $B \neq \varnothing$. We show $A \subseteq B$ and $B \subseteq A$.

$A \subseteq B$: Let $a \in A$. Since $B \neq \varnothing$, pick some $b \in B$.

$$(a, b) \in A \times B = B \times A \quad \implies \quad \text{first component } a \in B.$$

Since a was arbitrary, $A \subseteq B$.

$B \subseteq A$**:** Let $b \in B$. Since $A \neq \varnothing$, pick some $a \in A$.

$$(b, a) \in B \times A = A \times B \quad \Longrightarrow \quad \text{first component } b \in A.$$

Since b was arbitrary, $B \subseteq A$.

By antisymmetry of $\subseteq$:

$$\boxed{A \subseteq B \ \wedge \ B \subseteq A \implies A = B.}$$

$\square$

Note: the non-emptiness assumption was used above to "pick $b \in B$" and "pick $a \in A$".

9 Problem 60 — $\times$ Distributes Over Set Operations

Theory Recap: $\times$ Distributes Over $\cup$, $\cap$, $\setminus$

All three proofs follow the same three-step recipe:

(1) **Unpack:** $(x, y) \in A \times B \iff x \in A \wedge y \in B$.
(2) **Apply a propositional-logic law** ($\wedge/\vee$ distribution, or negation).
(3) **Repack** into set notation.

The operation in the second component "passes through" $\times$:

$$A \times (B \diamond C) = (A \times B) \diamond (A \times C) \qquad \text{for } \diamond \in \{\cup, \cap, \setminus\}.$$

Mirror identities $(B \diamond C) \times A$ work the same way on the *first* component. We prove the three non-mirror versions; the mirrors follow by swapping $x \leftrightarrow y$.



Problem Statement

Prove the following identities:

1. $A \times (B \cup C) = (A \times B) \cup (A \times C)$
3. $A \times (B \cap C) = (A \times B) \cap (A \times C)$
5. $A \times (B \setminus C) = (A \times B) \setminus (A \times C)$

Template method. To prove any Cartesian-product identity, always start with "Let (x, y) be arbitrary" and unpack $(x, y) \in$ LHS. The ordered-pair structure ensures the proof reduces to a propositional-logic equivalence.

Vertical Truth Table — Verifying the Propositional Equivalences

Each identity reduces to a propositional equivalence. Let $P = (x \in A)$, $Q = (y \in B)$, $R = (y \in C)$. The table below stacks the cases vertically for easier reading:

P Q R	**Identity 1** $P \wedge (Q \vee R) \stackrel{?}{=} (P \wedge Q) \vee (P \wedge R)$	**Identity 3** $P \wedge Q \wedge R \stackrel{?}{=} (P \wedge Q) \wedge (P \wedge R)$	**Identity 5** $P \wedge Q \wedge \neg R \stackrel{?}{=} (P \wedge Q) \wedge \neg(P \wedge$
Row 1: P=T, Q=T, R=T	T = T ✓	T = T ✓	F = F ✓
Row 2: P=T, Q=T, R=F	T = T ✓	F = F ✓	T = T ✓
Row 3: P=T, Q=F, R=T	T = T ✓	F = F ✓	F = F ✓
Row 4: P=T, Q=F, R=F	F = F ✓	F = F ✓	F = F ✓
Row 5: P=F, Q=T, R=T	F = F ✓	F = F ✓	F = F ✓
Row 6: P=F, Q=T, R=F	F = F ✓	F = F ✓	F = F ✓
Row 7: P=F, Q=F, R=T	F = F ✓	F = F ✓	F = F ✓
Row 8: P=F, Q=F, R=F	F = F ✓	F = F ✓	F = F ✓

All 8 rows match for each identity, confirming the propositional equivalences used in the proofs below.

Identity 1: $A \times (B \cup C) = (A \times B) \cup (A \times C)$

Let (x, y) be an arbitrary ordered pair:

$$\begin{aligned}
(x,y) \in A \times (B \cup C) &\iff x \in A \ \wedge \ y \in B \cup C \\
&\iff x \in A \ \wedge \ (y \in B \ \vee \ y \in C) \\
&\iff (x \in A \wedge y \in B) \ \vee \ (x \in A \wedge y \in C) \qquad (\wedge \text{ dist. over } \vee) \\
&\iff (x,y) \in A \times B \ \vee \ (x,y) \in A \times C \\
&\iff (x,y) \in (A \times B) \cup (A \times C). \qquad \blacksquare
\end{aligned}$$

Identity 3: $A \times (B \cap C) = (A \times B) \cap (A \times C)$

Let (x, y) be an arbitrary ordered pair:

$$\begin{aligned}
(x,y) \in A \times (B \cap C) &\iff x \in A \ \wedge \ y \in B \cap C \\
&\iff x \in A \ \wedge \ y \in B \ \wedge \ y \in C \\
&\iff (x \in A \wedge y \in B) \ \wedge \ (x \in A \wedge y \in C) \quad (\text{idempotency of } \wedge) \\
&\iff (x,y) \in A \times B \ \wedge \ (x,y) \in A \times C \\
&\iff (x,y) \in (A \times B) \cap (A \times C). \qquad \blacksquare
\end{aligned}$$

Identity 5: $A \times (B \setminus C) = (A \times B) \setminus (A \times C)$

Let (x, y) be an arbitrary ordered pair:

$$\begin{aligned}
(x,y) \in A \times (B \setminus C) &\iff x \in A \ \wedge \ y \in B \ \wedge \ y \notin C \\
&\iff \underbrace{(x \in A \wedge y \in B)}_{(x,y) \in A \times B} \ \wedge \ \underbrace{\neg(x \in A \wedge y \in C)}_{(x,y) \notin A \times C} \qquad (\star) \\
&\iff (x,y) \in (A \times B) \setminus (A \times C).
\end{aligned}$$

Justification of $(\star)$*:* We need $(p \wedge q \wedge \neg r) \iff (p \wedge q) \wedge \neg(p \wedge r)$, where $p = (x \in A)$, $q = (y \in B)$, $r = (y \in C)$.

$(\Rightarrow)$ $p, q, \neg r$ all hold, so $p \wedge q$ holds. Since $\neg r$, the conjunction $p \wedge r$ is false, so $\neg(p \wedge r)$ holds.

$(\Leftarrow)$ From $p \wedge q$ we have p and q. From $\neg(p \wedge r)$ and p, we get $\neg r$. $\square$

Identities 2, 4, 6 are the mirror versions $(B \diamond C) \times A = (B \times A) \diamond (C \times A)$ for $\diamond \in \{\cup, \cap, \setminus\}$. Replace "$x \in A$" with "$y \in A$" throughout—the proofs are identical.