

MAT 185 Lecture Notes

Hei Shing Cheung

Linear Algebra, Winter 2024

MAT 185

The up-to-date version of this document can be found at <https://github.com/HaysonC/skulenotes>

1 Vector Space

1.1 Fundamental Properties

Definiton 1.1.1 (Vector Space in \mathbb{R} , Field). This course concerns with real vector spaces. A vectors space, V , over \mathbb{R} is a collection of **object** $\mathbf{v} \in V$ s.t. the follow axioms are followed

1. Addition Axioms

- (a) **Closure Under Addition:**
 $\mathbf{x}, \mathbf{y} \in V \implies \mathbf{x} + \mathbf{y} \in V$
- (b) **Associativity of Addition:**
 $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V \implies (\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$
- (c) **Existence of Additive Identity:**
 $\exists \mathbf{0} \in V$ such that $\mathbf{x} \in V \implies \mathbf{x} + \mathbf{0} = \mathbf{x}$
- (d) **Existence of Additive Inverse:**
 $\forall \mathbf{x} \in V, \exists -\mathbf{x} \in V$ such that $\mathbf{x} + -\mathbf{x} = \mathbf{0}$

2. Scalar Multiplication Axioms

- (a) **Closure Under Scalar Multiplication:**
 $\forall \mathbf{x} \in V$ and $\forall \alpha \in \mathbb{R}, \alpha \mathbf{x} \in V$
- (b) **Associativity of Scalar Multiplication:**
 $\forall \mathbf{x} \in V$ and $\forall \alpha, \beta \in \mathbb{R}, (\alpha\beta)\mathbf{x} = \alpha(\beta\mathbf{x})$
- (c) **Distributive Property of Scalar Multiplication:**
 $\forall \mathbf{x} \in V$ and $\forall \alpha, \beta \in \mathbb{R}, (\alpha + \beta)\mathbf{x} = \alpha\mathbf{x} + \beta\mathbf{x}$
- (d) **Existence of Multiplicative Identity:**
 $\forall \mathbf{x} \in V, 1\mathbf{x} = \mathbf{x}$

Note It could be shown that the axiom imply the commutativity of in addition, namely $\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}, \mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$

Example \mathbb{R}^n is a vector space over \mathbb{R} , where $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$, then $\mathbf{x} + \mathbf{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$ and $\alpha \mathbf{x} = (\alpha x_1, \alpha x_2, \dots, \alpha x_n)$

Theorem 1.1.2 (Cancellation, Part 1). Let V be a vector space and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$, then:

$$\begin{aligned}\mathbf{x} + \mathbf{z} &= \mathbf{y} + \mathbf{z} \\ \mathbf{x} &= \mathbf{y}\end{aligned}$$

Proof.

$$\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$$

From additive inverse we know that $-\mathbf{z}$ exists

$$(\mathbf{x} + \mathbf{z}) + (-\mathbf{z}) = (\mathbf{y} + \mathbf{z}) + (-\mathbf{z})$$

By order of addition we have:

$$\begin{aligned}\mathbf{x} + (\mathbf{z} + (-\mathbf{z})) &= \mathbf{y} + (\mathbf{z} + (-\mathbf{z})) \\ \mathbf{x} + \mathbf{0} &= \mathbf{y} + \mathbf{0}\end{aligned}$$

By additive identity we have:

$$\mathbf{x} = \mathbf{y}$$

□

Theorem 1.1.3 (Cancellation, Part 2). Let V be a vector space and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$, then:

$$\begin{aligned}\mathbf{z} + \mathbf{x} &= \mathbf{z} + \mathbf{y} \\ \mathbf{x} &= \mathbf{y}\end{aligned}$$

To proof that, it would require the following proposition:

Lemma 1.1.4. Let V be a vector space and $\mathbf{z} \in V$, then $-\mathbf{z} + \mathbf{z} = \mathbf{0}$

Proof.

We know:

$$\begin{aligned}-\mathbf{z} + \mathbf{z} &= (-\mathbf{z} + \mathbf{z}) + \mathbf{0} \\ &= (-\mathbf{z} + \mathbf{z}) + (-\mathbf{z} + -(-\mathbf{z}))\end{aligned}$$

By the order of addition,

$$= -\mathbf{z} + (\mathbf{z} + (-\mathbf{z} + -(-\mathbf{z})))$$

Again, by the order of addition,

$$\begin{aligned}&= -\mathbf{z} + ((\mathbf{z} + -\mathbf{z}) + -(-\mathbf{z})) \\ &= -\mathbf{z} + -(-\mathbf{z}) \\ &= \mathbf{0}\end{aligned}$$

□

1.1 Fundamental Properties

Now, to prove the part 2 of the Cancellation Theorem:

Proof.

$$\mathbf{z} + \mathbf{x} = \mathbf{z} + \mathbf{y}$$

From additive inverse we know that $-\mathbf{z}$ exists

$$-\mathbf{z} + (\mathbf{z} + \mathbf{x}) = -\mathbf{z} + (\mathbf{z} + \mathbf{y})$$

$$(-\mathbf{z} + \mathbf{z}) + \mathbf{x} = (-\mathbf{z} + \mathbf{z}) + \mathbf{y}$$

From above, we have $-\mathbf{z} + \mathbf{z} = 0$

$$0 + \mathbf{x} = 0 + \mathbf{y}$$

$$\mathbf{x} = \mathbf{y}$$

□

Lemma 1.1.5 (Inverse of an inverse). Let V be a vector space and $\mathbf{x} \in V$, then:

$$-(-\mathbf{x}) = \mathbf{x}$$

Proof. Assume $0, 0^*$ are the additive identity of V and $-\mathbf{x}, -\mathbf{x}^*$ are the additive inverse of \mathbf{x} . We have:

$$u + 0 = u + 0^*$$

By Cancellation Theorem, we have $0 = 0^*$. Thus, the additive identity is unique. Now, we have:

$$\mathbf{x} + -\mathbf{x} = 0$$

$$\mathbf{x} + -\mathbf{x}^* = 0$$

Thus, by the uniqueness of the additive inverse, we have:

$$\mathbf{x} + -\mathbf{x} = \mathbf{x} + -\mathbf{x}^*$$

By the Cancellation Theorem, we have:

$$-\mathbf{x} = -\mathbf{x}^*$$

Now, we have that the additive inverse is unique. Thus, $-(-\mathbf{x})$ must be unique and necessarily \mathbf{x} . □

Additional Results from Above

1. The additive identity is unique
2. The additive inverse is unique

Definiton 1.1.6 (Subtraction). Let V be a vector space and $\mathbf{x}, \mathbf{y} \in V$, then:

$$\mathbf{x} - \mathbf{y} \stackrel{\text{def}}{=} \mathbf{x} + -\mathbf{y}$$

Theorem 1.1.7 (Addition is Commutative). Let V be a vector space and $\mathbf{x}, \mathbf{y} \in V$, then:

$$\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$$

Proof.

$$\mathbf{x} + \mathbf{y} =$$

□

1.2 Vector Subspace

Definiton 1.2.1 (Vector Subspace). Let V be a vector space and $W \subseteq V$, then W is a vector subspace of V if W is a vector space.

Theorem 1.2.2 (Subspace Test, I). Let V be a vector space over \mathbb{R} and $W \subseteq V$ with the usual addition and scalar multiplication, then W is a subspace of V iff for all $\mathbf{x}, \mathbf{y} \in W$ and $\alpha \in \mathbb{R}$:

1. **Closure Under Addition:**

$$\mathbf{x} + \mathbf{y} \in W$$

2. **Closure Under Scalar Multiplication:**

$$\alpha \mathbf{x} \in W$$

3. **Additive Identity:**

$$\mathbf{0} \in W$$

Proof. (\Rightarrow) Assume W is a subspace of V , then W is a vector space. Thus, the axioms of vector space are satisfied.

(\Leftarrow) Assume the three conditions are satisfied, then W is a vector space. Thus, W is a subspace of V . □

Definiton 1.2.3 (Null Space). Let V be a vector space and $A \in {}^m\mathbb{R}^{n1}$, then:

$$\text{null}(A) = \{\mathbf{x} \in {}^n\mathbb{R} \mid A\mathbf{x} = \mathbf{0}\} \quad (1)$$

is the null space of A , otherwise known as the **kernel** of A or the solution space of $A\mathbf{x} = \mathbf{0}$

We can use the Subspace Test I to show that the null space of a matrix is a subspace of ${}^n\mathbb{R}$.

1. **Existence of Additive Identity:**

The zero vector is in the null space of A as the trivial solution to the equation $A\mathbf{x} = \mathbf{0}$

2. **Closure Under Addition:**

Let $\mathbf{x}, \mathbf{y} \in \text{null}(A)$, then $A(\mathbf{x} + \mathbf{y}) = A\mathbf{x} + A\mathbf{y} = \mathbf{0} + \mathbf{0} = \mathbf{0}$. Thus, $\mathbf{x} + \mathbf{y} \in \text{null}(A)$. This holds since ${}^n\mathbb{R}$ is a vector space.

¹Same as $\mathbb{R}^{m \times n}$, The set of all $m \times n$ matrices with real entries

3. Closure Under Scalar Multiplication:

Let $\mathbf{x} \in \text{null}(A)$ and $\alpha \in \mathbb{R}$, then $A(\alpha\mathbf{x}) = \alpha A\mathbf{x} = \alpha\mathbf{0} = \mathbf{0}$. Thus, $\alpha\mathbf{x} \in \text{null}(A)$. This holds since ${}^n\mathbb{R}^m$ is a vector space under usual addition and scalar multiplication.

Theorem 1.2.4 (Subspace Test, II). Let V be a vector space over \mathbb{R} and $W \subseteq V$ with the usual addition and scalar multiplication, then W is a subspace of V iff for all $\mathbf{x}, \mathbf{y} \in W$ and $\alpha, \beta \in \mathbb{R}$:

1. $\alpha\mathbf{x} + \beta\mathbf{y} \in W$

Proof. (\Rightarrow) Assume W is a subspace of V , then W is a vector space. Thus, the axioms of vector space are satisfied.

(\Leftarrow) Assume the condition is satisfied, then W is a vector space. Thus, W is a subspace of V . \square

Definiton 1.2.5 (Intersection of Sets). Let A and B be sets, then:

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\} \quad (2)$$

Definiton 1.2.6 (Union of Sets). Let A and B be sets, then:

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\} \quad (3)$$

2 Linear combinations

Definiton 2.0.1 (Linear Combination). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$ and $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{R}$, then:

$$\alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n$$

is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$

2.1 Linear Independence, Span, and Basis

Definiton 2.1.1 (Linear independence). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$ and $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{R}$. If and only if the equation:

$$\alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n = \mathbf{0}$$

has **only** the trivial solution $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$.

Then, $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are said to be linearly independent; otherwise, they are linearly dependent.

Definiton 2.1.2 (Span). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$, then:

$$\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n) = \{\alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n \mid \alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{R}\} \quad (4)$$

is the set of all linear combinations of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$

2.2 Fundamental Theorem of Linear Algebra

Theorem 2.1.3 (Unique Representation). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$ and $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{R}/0$, and that they span V . For any $\mathbf{v}_{n+1} \in V$, if $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent, then the equation:

$$\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n = \mathbf{v}_{n+1}$$

is the only way to express \mathbf{v}_{n+1} as a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$.

We can prove by contrapositive:

Proof. Assume that there exists another way to express \mathbf{v}_{n+1} as a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$, then:

$$\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n = \beta_1 \mathbf{v}_1 + \beta_2 \mathbf{v}_2 + \dots + \beta_n \mathbf{v}_n$$

Rearranging the equation, we have:

$$(\alpha_1 - \beta_1) \mathbf{v}_1 + (\alpha_2 - \beta_2) \mathbf{v}_2 + \dots + (\alpha_n - \beta_n) \mathbf{v}_n = \mathbf{0}$$

Since $(\alpha_i - \beta_i)$ for $i = 1, 2, \dots, n$ are not all zero, then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly dependent. Thus, the contrapositive is true. \square

Theorem 2.1.4 (Growing and Pruning). The following are the two theorems describe the relationship between linear independence and the span:

Growing Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$ be linearly independent. If $\mathbf{v}_{n+1} \in V$ and $\mathbf{v}_{n+1} \notin \text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n)$, then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n, \mathbf{v}_{n+1}$ are linearly independent.

Pruning Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$. If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly dependent, then there exists a \mathbf{v}_i such that $\mathbf{v}_i \in \text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{i-1}, \mathbf{v}_{i+1}, \dots, \mathbf{v}_n)$.

Theorem 2.1.5 (Span and Linear Independence). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$. For every \mathbf{v}_k with $k = 1, 2, \dots, n$, $\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{k-1}, \mathbf{v}_{k+1}, \dots, \mathbf{v}_n) \subset \text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_n)$ if and only if $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent.

Definiton 2.1.6 (Basis). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$. If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent and $\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n) = V$, then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is a basis of V .

2.2 Fundamental Theorem of Linear Algebra

Theorem 2.2.1 (Fundamental Theorem of Linear Algebra). Let V be a vector space and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$. If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent and $\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n) = V$, then:

1. Any basis of V has the same number of elements.
2. Any linearly independent set of V has at most n elements.
3. Any spanning set of V has at least n elements.

2.2 Fundamental Theorem of Linear Algebra

Example 2.2.2 (Dimension of Even 4-degree polynomial). Given $W \subseteq \mathbb{P}_4$ where $W = \{p(x) \mid p(x) = p(-x)\}$, then W is a subspace of \mathbb{P}_4 . Since $x^3, x \in P_4$ but $x^3, x \notin W$, $\text{Span}(\{W, x, x^3\}) = \dim(W) + 2$. So $\dim(W) \leq 3$; We can find 3 linearly independent vectors in W such as $1, x^2, x^4$. Thus, $\dim(W) \geq 3$. Thus, $\dim(W) = 3$.

Definiton 2.2.3 (Column Space). Let $a_1, a_2, \dots, a_n \in {}^m\mathbb{R}$, then:

$$\text{Col}(A) = \text{Span}(a_1, a_2, \dots, a_n) \quad (5)$$

is the column space of A .

Note that the span of set is a subspace of the vector space. Thus, the column space of a matrix is a subspace of ${}^m\mathbb{R}$. This is also call the set of images of the transformation. We can also note that matrix multiplication is commutative with scalar multiplication. We can conduct proofs using such properties.

Definiton 2.2.4 (Rank). Let $A \in {}^m\mathbb{R}^n$, then:

$$\text{rank}(A) = \dim(\text{Col}(A)) \quad (6)$$

is the rank of A .