

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/265135825>

Antipsychotic-induced akathisia and neuroleptic malignant syndrome in anti-NMDAR encephalitis

Article in *Annals of Clinical Psychiatry* · August 2014

Source: PubMed

CITATIONS

6

READS

338

5 authors, including:



Shekeeb S Mohammad

Children's Hospital at Westmead

51 PUBLICATIONS **508** CITATIONS

[SEE PROFILE](#)



Geoff B Wallace

Queensland Government

25 PUBLICATIONS **1,250** CITATIONS

[SEE PROFILE](#)



Sudarshini Ramanathan

Westmead Hospital, Sydney; The University of Sydney, Sydney

37 PUBLICATIONS **624** CITATIONS

[SEE PROFILE](#)

Antipsychotic-induced akathisia and neuroleptic malignant syndrome in anti-NMDAR encephalitis

ANNALS OF CLINICAL PSYCHIATRY 2014;26(3):E1-E2

Shekeeb S. Mohammad, MBBS, FRACP; Geoffrey Wallace, FRACP; Sudarshini Ramanathan, FRACP; Fabienne Brilot, PhD; Russell C. Dale, PhD

KEYWORDS: antipsychotic, NMDAR, akathisia

TO THE EDITOR:

In a recent review in *Annals of Clinical Psychiatry*, Kruse et al¹ highlighted the importance of recognizing psychiatric symptoms as a common early feature of anti-N-methyl-D-aspartate receptor (NMDAR) encephalitis and the challenges in managing these symptoms while waiting for a patient to respond to immunotherapy. We present 2 cases to highlight the risk of extrapyramidal symptoms (EPS) in patients with anti-NMDAR encephalitis who were treated with antipsychotics for psychiatric symptoms.

Case 1

A 4-year-old boy presented with a 4-week history of aggressive behavior, hallucinations, and seizures. His cerebrospinal fluid (CSF) had 30 mononuclear cells/ μ L and was positive for IgG antibodies against NMDAR. He was treated with 2 g/kg of IV immunoglobulin (IVIg) and 3 days of IV methylprednisolone, 30 mg/kg/d. Distressing episodes of aggressive behavior, mood swings, and hallucinations persisted despite receiving nitrazepam, 2.5 mg twice daily. Quetiapine was introduced, 5 mg in the morning, and when the drug was increased to 10 mg/d the boy was noted to have continuous restless leg and body movements that were worse during sleep over 2 nights. He had a constant urge to keep moving during the day. The phenomenology was suggestive of akathisia. Quetiapine was stopped, and the akathisia resolved within 2 days. His ongoing disease was treated with 3 doses of rituximab (375 mg/m²) and further IVIg 2 g/kg with complete recovery over the ensuing 12 months.

Case 2

A 13-year-old male presented with lethargy, visual hallucinations, behavioral change, dystonia, stereotypical movements, and increasing agitation. CSF showed pleocytosis (13 cells/ μ L) and autoimmune encephalitis was suspected and he was treated with IV methylprednisolone, 30 mg/kg/d for 3 days. Retrospective testing of acute serum and CSF was positive for NMDAR antibodies. Worsening visual hallucinations, agitation, dystonia, and stereotypical writhing movements were treated with droperidol, 2.5 mg twice daily (0.05 mg/kg/dose) and benzotropine, 1 mg twice daily (0.02 mg/kg/dose). After 3 days on droperidol, he developed hypertension, sweating, increased rigidity, and elevated creatine kinase. Neuroleptic malignant syndrome (NMS) was suspected, droperidol was stopped, and benzotropine continued. The patient's NMS resolved within a few days of stopping droperidol. After a 3-month admission, the patient made a complete recovery and has been well for 5 years.

DISCUSSION

Antipsychotics and other psychotropics often are administered in cases of anti-NMDAR encephalitis with associated psychiatric symptoms with poor efficacy¹ and can precipitate or exacerbate motor symptoms.^{2,3} It can be challenging to differentiate EPS of antipsychotics from the movement disorders seen in anti-NMDAR encephalitis, as highlighted by previous descriptions of NMS-like syndrome in anti-NMDAR encephalitis,^{2,4} even in the absence of antipsychotic administration. In the 2 cases we describe, onset of new motor symptoms with antipsychotic initiation, worsening of motor symptoms with increased doses of antipsychotics,² and improvement in symptoms when the antipsychotic was stopped supports the hypothesis that these symptoms were medication effects rather than part of disease evolution alone. In both of our cases, akathisia and NMS were clearly associated with starting antipsychotics, and resolved on stopping the respective medication. Interestingly, anecdotal reports have described higher prevalence of adverse reactions to antipsychotics in Sydenham chorea, another autoimmune movement disorder.⁵ These observations

LETTERS TO THE EDITOR

suggest a vulnerability to classic neuroleptic side effects in patients with anti-NMDAR encephalitis. ■

DISCLOSURES: Drs. Brilot and Dale have received funding from the Australian National Health and

Medical Research Council, and the Petre Foundation. Drs. Mohammad, Wallace, and Ramanathan report no financial relationships with any company whose products are mentioned in this article or with manufacturers of competing products.

REFERENCES

1. Kruse JL, Jeffrey JK, Davis MC, et al. Anti-N-methyl-D-aspartate receptor encephalitis: A targeted review of clinical presentation, diagnosis, and approaches to psychopharmacologic management. *Ann Clin Psychiatry*. 2014;26:111-119.
2. Chapman MR, Vause HE. Anti-NMDA receptor encephalitis: diagnosis, psychiatric presentation, and treatment. *Am J Psychiatry*. 2011;168:245-251.
3. Mann A, Machado NM, Liu N, et al. A multidisciplinary approach to the treatment of anti-NMDA-receptor antibody encephalitis: a case and review of the literature. *J Neuropsychiatry Clin Neurosci*. 2012;24:247-254.
4. Consoli A, Ronen K, An-Gourfinkel I, et al. Malignant catatonia due to anti-NMDA-receptor encephalitis in a 17-year-old girl: case report. *Child Adolesc Psychiatry Ment Health*. 2011;5:15.
5. Teixeira AL, Cardoso F, Maia DP, et al. Sydenham's chorea may be a risk factor for drug induced parkinsonism. *J Neurol Neurosurg Psychiatry*. 2003;74:1350-1351.

Shekeeb S. Mohammad, MBBS, FRACP
Neuroimmunology Group
Institute for Neuroscience and Muscle Research
The Kids Research Institute at the Children's
Hospital at Westmead
University of Sydney
Sydney, Australia

Sudarshini Ramanathan, FRACP
Neuroimmunology Group
Institute for Neuroscience and Muscle Research
The Kids Research Institute at the Children's
Hospital at Westmead
University of Sydney
Sydney, Australia

Fabienne Brilot, PhD
Neuroimmunology Group
Institute for Neuroscience and Muscle Research
The Kids Research Institute at the Children's
Hospital at Westmead
University of Sydney
Sydney, Australia

Russell C. Dale, PhD
Neuroimmunology Group
Institute for Neuroscience and Muscle Research
The Kids Research Institute at the Children's
Hospital at Westmead
University of Sydney
Sydney, Australia

Geoffrey Wallace, FRACP
Department of Neurosciences
Royal Children's Hospital, Herston
School of Medicine
University of Queensland
Brisbane, Australia

CORRESPONDENCE: Russell C. Dale, PhD, Clinical School, the Children's Hospital at Westmead, Locked Bag 4001, NSW 2145 Australia **E-MAIL:** russell.dale@health.nsw.gov.au

