NMDA receptor encephalitis causing reversible caudate changes on MRI and PET imaging

William O. Tobin, PhD Edythe A. Strand, PhD Heather M. Clark, PhD Val J. Lowe, MD Carrie E. Robertson, MD Sean J. Pittock, MD

Practical Implications

NMDA receptor (NMDAR) antibodies may cause reversible caudate MRI and PET brain abnormalities associated with a restricted phenotype of NMDAR encephalitis. ince the original description of the classic phenotype of memory deficits, psychiatric symptoms, decreased consciousness, and hypoventilation was described in association with NMDAR antibodies, the phenotype of NMDAR encephalitis has expanded to include patients with a pure psychiatric presentation, insomnia, isolated dystonia, paroxysmal limb weakness, and eye movement abnormalities.¹ Despite prominent dystonia and dyskinesias, reversible basal ganglia abnormalities on structural MRI and FDG-PET have not been reported to date, even in patients with isolated hemidystonia.²

Case report


A 21-year-old man presented with episodic anxiety, palpitations, diaphoresis, diarrhea, weight loss, and tinnitus of 1 month's duration. Over the next month, he developed bradykinesia, bilateral arm numbness, inappropriate laughter, psychosis, cognitive impairment, an exaggerated response to alcohol, and dysarthria. Spinal fluid analysis revealed 12 leukocytes per microliter (86% lymphocytes) and mildly elevated protein. Brain MRI 4 months after symptom onset demonstrated T2 fluid-attenuated inversion recovery high signal involving medial temporal lobes, caudate heads and bodies, and lentiform nuclei (figure, B). FDG-PET/CT scan demonstrated absent metabolism in the basal ganglia bilaterally, with otherwise normal brain activity (figure, A). No malignancy or tumor was found on whole-body FDG-PET/CT or testicular ultrasound.

Testing for the NMDAR antibody was positive in serum and spinal fluid using a transfected HEK293 cell-binding assay (Euroimmun, Lübeck, Germany; each sample scored positive or negative by 3 experienced readers). The patient's serum and CSF underwent comprehensive neural antibody evaluation with a standardized indirect immunofluorescence assay on a composite substrate of mouse cerebellum, midbrain, basal ganglia, thalamus, cerebral cortex, hippocampus, stomach, and kidney to detect immunoglobulin G (IgG) autoantibodies binding selectively to

Departments of Neurology (WOT, EAS, HMC, CER, SJP), Speech Pathology (EAS, HMC), Radiology (VJL), and Laboratory Medicine and Pathology (SJP), and College of Medicine, Mayo Clinic, Rochester, MN. Funding information and disclosures are provided at the end of the article. Full disclosure form information provided by the authors is available with the **full text of this article at Neurology.org/cp**.

Correspondence to: pittock.sean@mayo.edu

Figure Imaging



PET demonstrates hypometabolism in the caudate heads (dashed arrow) and lentiform nuclei (solid arrow) prior to treatment (A) and following 3 months of treatment with IV steroids and immunoglobulin (C). Brain MRI demonstrates T2 fluid-attenuated inversion recovery high signal in the caudate head (dashed arrow) and lentiform nuclei (solid arrow) prior to treatment (B) and following 3 months of treatment with IV steroids and immunoglobulin (D).

neuronal and glial nuclei (antineuronal nuclear antibodies, type 1 [anti-Hu], type 2 [anti-Ri], and type 3; antiglial/neuronal nuclear antibody, type 1 [AGNA or SOX 1 antibody]), neuronal cytoplasm (Purkinje cell antibodies [types 1 (anti Yo), 2, and -Tr], collapsin response—mediator protein 5-IgG and amphiphysin-IgG), or hippocampal and basal ganglionic synapses.

Antibodies reactive with neural cation channel complexes (neuronal voltage-gated calcium channels [P/Q type and N type], voltage-gated potassium channel complexes, nicotinic acetylcholine receptors of skeletal muscle type [α 1 subunit] and neuronal ganglionic type [α 3 subunit]) and glutamic acid decarboxylase 65 kDa isoform (GAD65) were sought using a radioimmunoprecipitation assay. Skeletal muscle striational antibodies were tested by ELISA. IgGs targeting other specific neurotransmitter receptors in hippocampal synapses (AMPA [GluA1 and GluA2] and GABA-B) were also sought by indirect immunofluorescence on HEK293 cells transfected with appropriate cDNAs (Euroimmun). The only other neural antibody detected was GAD65 antibody level at 0.03 nmol/L (normal range <0.02 nmol/L) in spinal fluid. No other antibody was detected.

Speech assessment demonstrated a mixed dysarthria, with primarily hypokinetic features as well as spastic/dystonic features (video 1 at Neurology.org/cp). The patient was treated with IV steroids, 1 g daily for 3 consecutive days, then once weekly for 5 weeks, followed by once every 2 weeks for 6 weeks. IV immunoglobulin was given concurrently at a dose of 0.4 g/kg daily for 3 days followed by 0.4 g/kg every week for 6 weeks and then every 2 weeks for 6 weeks. At the same time, he received mycophenolate 500 mg twice daily, increasing after 2 weeks to 1,000 mg twice daily.

Three months after commencing treatment, neuroimaging abnormalities had almost completely resolved (figure, C and D). Examination revealed a residual action dystonia in the patient's right hand, with associated mild bradykinesia. Neuropsychometric and speech testing were normal (video 2).

DISCUSSION

This is the first reported case of reversible caudate MRI and PET abnormalities in a patient with NMDAR antibody–associated encephalitis. A patient with isolated generalized dystonia, caudate MRI abnormalities, and NMDAR antibodies has been described previously,³ although reversibility was not demonstrated despite treatment. Our patient had some unusual features including tinnitus and an exaggerated response to alcohol. His speech abnormalities were presumably caused by basal ganglia dysfunction, as evidenced by complete resolution in tandem with resolution of MRI changes. The patient's action dystonia correlated with a residual area of reduced metabolism on FDG-PET imaging, supporting a functional correlation. Brain MRI abnormalities are typically subtle or absent in patients with NMDAR

Video

Neurology.org/cp

encephalitis.⁴ In the largest case series of patients with NMDAR encephalitis, 5 out of 100 patients had basal ganglia abnormalities on brain MRI.⁴ Other areas where MRI abnormalities have been described include the medial temporal lobe; frontal, parietal, and occipital lobe cortex; and cerebellar cortex.^{4–6} FDG-PET global hypometabolism and cerebellar hypermetabolism, which subsequently became hypometabolic, has been reported in a 15-year-old boy with NMDAR encephalitis.⁷ Advanced imaging techniques have demonstrated alterations of functional connectivity in patients with normal standard MRIs of the brain.⁸ Orofacial dyskinesias, along with "piano playing" movements of the hands, are frequent, and isolated hemidystonia has been described, in the absence of MRI abnormalities.² The current case may represent the extreme end of a spectrum of basal ganglia abnormalities in this condition, and lends support to the theory of neuronal dysfunction, in the absence of neuronal destruction in the condition.

The patient demonstrated a dramatic recovery, typical of NMDAR encephalitis, despite moderately severe deficits prior to immunomodulatory treatment. The NMDAR antibody is directed against a cell surface antigen and is associated with a reduction in cell surface and synaptic NMDAR density⁹ and is thought to be pathogenic, in contrast to antibodies directed against intracellular antigens.⁹ In contrast to antibody-mediated inflammatory diseases such as neuromyelitis optica, binding of NMDAR antibodies does not lead to complement activation.⁹ This may protect from neuronal death and allow a full recovery if treated appropriately, despite significant apparent structural changes in this case.

REFERENCES

- 1. Marques IB, Teotonio R, Cunha C, Bento C, Sales F. Anti-NMDA receptor encephalitis presenting with total insomnia: a case report. J Neurol Sci 2014;336:276–280.
- 2. Rubio-Agusti I, Dalmau J, Sevilla T, Burgal M, Beltran E, Bataller L. Isolated hemidystonia associated with NMDA receptor antibodies. Mov Disord 2011;26:351–352.
- 3. Tzoulis C, Vedeler C, Haugen M, et al. Progressive striatal necrosis associated with anti-NMDA receptor antibodies. BMC Neurol 2013;13:55.
- Dalmau J, Gleichman AJ, Hughes EG, et al. Anti-NMDA-receptor encephalitis: case series and analysis
 of the effects of antibodies. Lancet Neurol 2008;7:1091–1098.
- 5. Goldberg EM, Taub KS, Kessler SK, Abend NS. Anti-NMDA receptor encephalitis presenting with focal non-convulsive status epilepticus in a child. Neuropediatrics 2011;42:188–190.
- 6. Dalmau J, Tuzun E, Wu HY, et al. Paraneoplastic anti-N-methyl-D-aspartate receptor encephalitis associated with ovarian teratoma. Ann Neurol 2007;61:25–36.
- Maqbool M, Oleske DA, Huq AH, Salman BA, Khodabakhsh K, Chugani HT. Novel FDG-PET findings in anti-NMDA receptor encephalitis: a case based report. J Child Neurol 2011;26:1325–1328.
- 8. Finke C, Kopp UA, Scheel M, et al. Functional and structural brain changes in anti-N-methyl-D-aspartate receptor encephalitis. Ann Neurol 2013;74:284–296.
- Martinez-Hernandez E, Horvath J, Shiloh-Malawsky Y, Sangha N, Martinez-Lage M, Dalmau J. Analysis of complement and plasma cells in the brain of patients with anti-NMDAR encephalitis. Neurology 2011;77:589–593.

STUDY FUNDING

Supported by the Mayo Foundation.

DISCLOSURES

W. Tobin reports no disclosures. E. Strand receives publishing royalties for *Management of Speech and Swallowing in Degenerative Neurologic Disease* (Communication Skill Builders; 1995; 3rd ed, 2013), *Clinical Management of Motor Speech Disorders of Children* (Thieme Publishing Co., 1999), and *Clinical Management of Motor Speech Disorders* (Little, Brown & Co.; 2nd ed, 1999; 3rd ed, 2010); and receives research support from NIH/NIDCD. H. Clark receives publishing royalties for *Neurogenic Disorders of Language* (Cengage, 2005). V. Lowe serves on scientific advisory boards for Piramal Imaging and Bayer Schering Pharma and receives research support from GE Healthcare, Siemens Molecular Imaging, AVID Radiopharmaceuticals, NIH (NIA, NCI), the Minnesota Partnership for Biotechnology and Medical Genomics, the Elsie and Marvin Dekelboum Family Foundation, and the Leukemia & Lymphoma Society. C. Robertson receives royalties from *UpToDate*. S. Pittock is a named

inventor on patents (12/678,350 filed 2010 and 12/573,942 filed 2008) that relate to functional AQP4/NMO-IgG assays and NMO-IgG as a cancer marker; receives research support from Alexion Pharmaceuticals, Inc., the Guthy-Jackson Charitable Foundation, and the NIH; and has provided consultation to Alexion Pharmaceuticals, Medimmune, and Chugai Pharma USA but has received no personal fees or personal compensation for these consulting activities. All compensation for consulting activities is paid directly to Mayo Clinic. Full disclosure form information provided by the authors is available with the **full text of this article at Neurology.org/cp**.

Share Your Knowledge

Neurology® Clinical Practice encourages readers to share their insights, expertise, and experiences.

- How are you employing drugs and devices in your field?
- What ethical challenges do you face?
- Do you have a case report that is illustrative of a clinical challenge?
- What challenges have you faced or successes have you enjoyed in bringing greater efficiency to your practice?

If you are interested in delivering a high-quality, peer-reviewed message to your colleagues in practice, submit your paper **online**.