

## Letters to the Editor

**Anti-myelin oligodendrocyte glycoprotein antibody neuritis optica following anti-NMDA receptor encephalitis**

Koji Yokoyama, Mutsuki Hori and Akira Yoshida


*Department of Pediatrics, Japanese Red Cross Wakayama Medical Center, Wakayama, Japan*

A 9-year-old girl was admitted to hospital with fever, somnolence, hemiplegia, dysphagia, and memory impairment. Cell-based assay indicated that she was positive for anti-*N*-methyl-D-aspartate receptor (anti-NMDAR) antibodies in serum and cerebrospinal fluid (CSF). On brain magnetic resonance imaging (MRI) T2-weighted (T2)/fluid-attenuated inversion recovery, abnormality was noted in the left temporal and parietal lobes (Fig. 1). Electroencephalogram demonstrated slowing of the background rhythm in the left cerebral hemisphere, but no epileptiform discharge. On lumbar puncture, 44 white blood cells had mixed cellularity and protein was 22 mg/dL. Bacterial culture and tests for various viruses were negative. Oligoclonal IgG band and anti-aquaporin-4 antibody (AQP4) were not present. The patient had no ophthalmologic abnormalities or associated tumors; she was therefore diagnosed with anti-NMDAR encephalitis and treated with three courses of pulse i.v. methylprednisolone (IVMP) and i.v. gammaglobulin (IVIG). She was symptom free at discharge and was treated with oral prednisolone (gradually tapered over 5 months). Seventeen days after cessation of prednisolone, she was readmitted due to right visual impairment, pain, and headache. The right optic disc was swollen, she had a reduced light reflex and could discern hand movement but nothing else through the right eye. Visual acuity in the left eye was intact. Brain MRI indicated contrast enhancement in the right optic nerve, but no abnormality in the cerebral hemisphere or spinal cord (Fig. 2). Serum and CSF were strongly positive for anti-myelin oligodendrocyte glycoprotein (anti-MOG) antibodies, but negative for anti-AQP4 and anti-NMDAR antibodies. There was no oligoclonal IgG band. Myelin basic protein in the CSF was elevated (149 pg/mL, standard value, <102). She was diagnosed with anti-MOG antibody-positive neuromyelitis optica spectrum disorder (NMOSD) and treated with three courses of IVMP and IVIG. Visual acuity at discharge was normal. On re-testing of stocked serum, anti-MOG antibodies were positive at the same time that symptoms of anti-NMDAR encephalitis were evident. At 18 months after initial presentation, she received oral prednisolone (5 mg/day).


Correspondence: Koji Yokoyama, MD, Department of Pediatrics, Japanese Red Cross Wakayama Medical Center, Komatsubara 4-20, Wakayama City, Wakayama 640-8558, Japan. Email: [kojiy@kuhp.kyoto-u.ac.jp](mailto:kojiy@kuhp.kyoto-u.ac.jp)

Received 5 March 2016; revised 27 May 2016; accepted 7 June 2016.

doi: 10.1111/ped.13053



**Fig. 1** T2/fluid-attenuated inversion recovery-weighted magnetic resonance imaging the brain showing high signal intensity in the left temporal and parietal lobes.



**Fig. 2** T1-weighted magnetic resonance imaging of the brain showing gadolinium enhancement in the right optic nerve.

Anti-NMDAR encephalitis is an immune-mediated disorder associated with IgG antibodies against the GluR1 subunit of NMDAR.<sup>1</sup> It is very difficult to distinguish anti-NMDAR encephalitis from acute disseminated encephalomyelitis. Thus, further investigation of NMDAR antibody level is needed to facilitate diagnosis. Neuromyelitis optica (NMO) and NMOSD are autoimmune inflammatory disorders of the central nervous

system (CNS) characterized by acute or subacute onset of optic neuritis and myelitis.<sup>2</sup> Patients with NMOSD and NMO-like symptoms do not fulfill the diagnosis of NMO,<sup>3</sup> although they are expected to respond to the same treatment as patients with standard NMO.<sup>4</sup> MOG is localized to oligodendrocytes and the outer surface of the myelin sheath; thus MOG-related experimental allergic encephalomyelitis can mimic a neurological syndrome closely resembling NMO.<sup>5</sup> The finding that anti-NMDAR encephalitis and a demyelinating disorder may occur in the same patient is important because the treatment and outcome for each vary; therefore, physicians should check for each type of antibody. Anti-NMDAR encephalitis has a mortality rate of 7%. Anti-MOG antibody-positive NMO may be associated with severe and sustained visual loss and atrophy of the retinal nerve fiber layer.<sup>6</sup> Greater understanding of these types of diseases and their treatment will improve prognosis. Physicians should test for antibodies to NMDAR, MOG, and AQP4. The present case shows that immunological activation can generate anti-neuro-antibodies, resulting in various diseases. The coexistence of multiple autoimmune diseases is increasingly recognized among those with autoantibody-associated neurological syndromes,<sup>7,8</sup> but the mechanisms underlying autoimmunity and autoantibody production are unclear.<sup>8</sup> Future studies should identify factors that predispose individuals to develop autoantibodies against multiple CNS antigens.

### Acknowledgments

We wish to thank Dr Keiko Tanaka, Dr Ichiro Miya, Dr Kazuyuki Takahashi, and Dr Syu Watanabe for anti-neuro-antibody analysis, Dr Masaaki Shimizu for cytokine analysis, and Dr Yoshinori Kawamura for analysis of viral agents.

### Disclosure

The authors declare no conflicts of interest.

### Author contributions

K.Y., M.H., and A.Y. designed the study; K.Y. and M.H. collected and analyzed data; K.Y. wrote the manuscript; M.H. and A.Y. gave technical support and conceptual advice. All authors read and approved the final manuscript.

### References

- 1 Goldberg EM, Titulaer M, de Blank PM, Sievert A, Ryan N. Anti-N-methyl-D-aspartate receptor-mediated encephalitis in infants and toddlers: Case report and review of the literature. *Pediatr. Neurol.* 2014; **50**: 181–4.
- 2 Wingerchuk DM, Banwell B, Bennett JL *et al.* International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 2015; **85**: 177–89.
- 3 Masaki K, Suzuki SO, Matsushita T *et al.* Connexin 43 astrocytopathy linked to rapidly progressive multiple sclerosis and neuromyelitis optica. *PLoS One* 2013; **8**: e72919.
- 4 Fujihara K, Sato DK. AQP4 antibody serostatus: Is its luster being lost in the management and pathogenesis of NMO? *Neurology* 2013; **81**: 1186–8.
- 5 Tanaka M, Tanaka K. Anti-MOG antibodies in adult patients with demyelinating disorders of the central nervous system. *J. Neuroimmunol.* 2014; **270**: 98–9.
- 6 Ramanathan S, Dale RC, Brilot F. Anti-MOG antibody: The history, clinical phenotype, and pathogenicity of a serum biomarker for demyelination. *Autoimmun. Rev.* 2016; **15**: 307–24.
- 7 Titulaer MJ, Hofstberger R, Iizuka T *et al.* Overlapping demyelinating syndromes and anti-N-methyl-D-aspartate receptor encephalitis. *Ann. Neurol.* 2014; **75**: 411–28.
- 8 Zoccarato M, Saddi MV, Serra G *et al.* Aquaporin-4 antibody neuromyelitis optica following anti-NMDA receptor encephalitis. *J. Neurol.* 2013; **260**: 3185–7.

## Pediatrician role in public health: Domestic animal-related injury alerts for neonates

Hiroshi Yamaguchi,<sup>1</sup> Masashi Kasai,<sup>2</sup> Eru Kozuki,<sup>1</sup> Yusuke Ishida,<sup>1</sup> Hiroki Takeda<sup>1</sup> and Yoshiyuki Uetani<sup>1</sup>

Departments of <sup>1</sup>Emergency and Critical Care Medicine, and <sup>2</sup>Infectious Disease Medicine, Hyogo Prefectural Kobe Children's Hospital, Kobe, Japan

Correspondence: Hiroshi Yamaguchi, MD, Department of Emergency and Critical Care Medicine, Hyogo Prefectural Kobe Children's Hospital, 1-1-1 Takakuradai, Suma-Ku, Kobe, Hyogo 654-0081 Japan. Email: hiyamaguchi\_kch@hp.pref.hyogo.jp

Received 17 March 2016; revised 17 June 2016; accepted 21 June 2016.

doi: 10.1111/ped.13071

My colleagues and I had previously reported on a neonate with meningitis and bacteremia caused by *Pasteurella multocida* following a cat scratch.<sup>1</sup> *P. multocida* is a Gram-negative coccobacillus that is part of the usual human oral flora or is found in the nails; moreover, it exists in many animals, especially cats.<sup>2</sup> In the previous study, a 23-day-old girl presented with decreased appetite and irritability for >2 days. Eighteen