

I Definitions

Let $W(x) := \left(-\frac{x}{2} + \left(\frac{1}{27} + \frac{x^2}{4}\right)^{1/2}\right)^{1/3} - \left(\frac{x}{2} + \left(\frac{1}{27} + \frac{x^2}{4}\right)^{1/2}\right)^{1/3}$

and define $Lz := -az - b\frac{\partial z}{\partial x}$, where $a(x) := 1 + \frac{W(x)}{x} + \frac{\partial W}{\partial x}(x)$ and $b(x) := \frac{3x}{2} + W(x)$.

II Computation in Sobolev spaces

Let $\langle f, g \rangle := \langle f, g \rangle_{L^2} = \int_{\mathbb{R}} f(x)g(x) dx$ be the usual inner product of $L^2(\mathbb{R})$.

Let $w, z \in H^k(\mathbb{R})$ for k large enough. For simplicity, we will denote $\frac{\partial z}{\partial x} := z'$.

II.1 Symetric part in L^2 space

In L^2 , the symetric part is computed as follows :

$$\begin{aligned} \langle Lz, w \rangle_{L^2} &= \langle -az - bz', w \rangle = \langle z, -aw \rangle - \langle z, b'w + bw' \rangle \\ &= \langle z, (-a + b')w + bw' \rangle = \langle z, L^*w \rangle \end{aligned}$$

Thus, $\frac{1}{2}(L + L^*)z = \frac{1}{2}(-az - bz' - az + b'z + bz') = -az + \frac{b'}{2}z$ in L^2 .

II.2 Quadratic form in H^1 space

In H^1 , the quadratic form is computed as follows :

$$\begin{aligned} \langle Lz, z \rangle_{H^1} &= \langle -az - bz', z \rangle + \langle -a'z - az' - b'z' - bz'', z' \rangle \\ &= \langle -az, z \rangle + \langle -bz - a'z, z' \rangle + \langle -az' - b'z', z' \rangle + \langle -bz', z'' \rangle \\ &= \langle -az, z \rangle + \langle \frac{1}{2}(b' + a'')z, z \rangle + \langle (-a - b')z', z' \rangle + \langle \frac{1}{2}b'z', z'' \rangle \\ &= \langle (-a + \frac{b'}{2} + \frac{a''}{2})z, z \rangle + \langle (-a - \frac{b'}{2})z', z' \rangle \end{aligned}$$

REMARQUE. The operator $(Lz)'$ is not defined on H^1 as it involves second derivatives of z , but it is a classical fact that the quadratic form of an operator as a larger domain than the operator itself.

II.3 Quadratic form in H^2 space

In H^2 , the quadratic form is computed as follows :

$$\begin{aligned} \langle (Lz)'', z'' \rangle &= \langle -a''z - a'z' - a'z' - az'' - b''z' - b'z'' - b'z'' - bz^{(3)}, z'' \rangle \\ &= \langle -a''z, z'' \rangle + \langle (-2a' - b'')z', z'' \rangle + \langle (-a - 2b')z'', z'' \rangle + \langle -bz^{(3)}, z'' \rangle \\ &= \langle a^{(3)}z + a''z', z' \rangle + \langle \frac{1}{2}(2a'' + b^{(3)})z', z' \rangle + \langle (-a - 2b')z'', z'' \rangle + \langle \frac{1}{2}b'z'', z'' \rangle \\ &= \langle -\frac{1}{2}a^{(4)}z, z \rangle + \langle 2a'' + \frac{1}{2}b^{(3)}z', z' \rangle + \langle (-a - \frac{3}{2}b')z'', z'' \rangle \end{aligned}$$

Thus, we have in H^2 :

$$\langle Lz, z \rangle_{H^2} = \langle (-a + \frac{b'}{2} + \frac{a''}{2} - \frac{a^{(4)}}{2})z, z \rangle + \langle (-a - \frac{b'}{2} + 2a'' + \frac{b^{(3)}}{2})z', z' \rangle + \langle (-a - \frac{3}{2}b')z'', z'' \rangle$$

II.4 Quadratic form in H^3 space

In H^3 , the quadratic form is computed as follows :

$$\begin{aligned}
\langle (Lz)^{(3)}, z^{(3)} \rangle &= \langle -a'''z - 3a''z' - 3a'z'' - az^{(3)} - b'''z' - 3b''z'' - 3b'z^{(3)} - bz^{(4)}, z^{(3)} \rangle \\
&= \langle -a'''z, z^{(3)} \rangle + \langle (-3a'' - b''')z', z^{(3)} \rangle + \langle (-3a' - 3b'')z'', z^{(3)} \rangle + \langle (-a - 3b')z^{(3)}, z^{(3)} \rangle \\
&\quad + \langle -bz^{(4)}, z^{(3)} \rangle \\
&= \langle a^{(4)}z + a'''z', z'' \rangle + \langle (3a''' + b^{(4)})z' + (3a'' + b''')z'', z'' \rangle + \langle \frac{3}{2}(a'' + b''')z'', z'' \rangle \\
&\quad + \langle (-a - 3b')z^{(3)}, z^{(3)} \rangle + \langle \frac{1}{2}b'z^{(3)}, z^{(3)} \rangle \\
&= \langle -a^{(5)}z - a^{(4)}z', z' \rangle + \langle -\frac{1}{2}a^{(4)}z', z' \rangle + \langle \frac{1}{2}(-3a^{(4)} - b^{(5)})z', z' \rangle + \langle (3a'' + b''')z'', z'' \rangle \\
&\quad + \langle \frac{3}{2}(a'' + b''')z'', z'' \rangle + \langle (-a - 3b')z^{(3)}, z^{(3)} \rangle + \langle \frac{1}{2}b'z^{(3)}, z^{(3)} \rangle \\
&= \langle \frac{a^{(6)}}{2}z, z \rangle + \langle (-3a^{(4)} - \frac{1}{2}b^{(5)})z', z' \rangle + \langle (\frac{9}{2}a'' + \frac{5}{2}b^{(3)})z'', z'' \rangle + \langle (-a - \frac{5}{2}b')z^{(3)}, z^{(3)} \rangle
\end{aligned}$$

Thus, we have in H^3 :

$$\begin{aligned}
\langle Lz, z \rangle_{H^3} &= \langle (-a + \frac{b'}{2} + \frac{a''}{2} - \frac{a^{(4)}}{2} + \frac{a^{(6)}}{2})z, z \rangle + \langle (-a - \frac{b'}{2} + 2a'' + \frac{b^{(3)}}{2} - 3a^{(4)} - \frac{1}{2}b^{(5)})z', z' \rangle \\
&\quad + \langle (-a - \frac{3}{2}b' + \frac{9}{2}a'' + \frac{5}{2}b^{(3)})z'', z'' \rangle + \langle (-a - \frac{5}{2}b')z^{(3)}, z^{(3)} \rangle
\end{aligned}$$

III Compact part of the quadratic form

We proved in the previous section that the quadratic form associated with L in H^3 is of the form :

$$\langle Lz, z \rangle_{H^3} = \langle \varphi_0 z, z \rangle + \langle \varphi_1 z', z' \rangle + \langle \varphi_2 z'', z'' \rangle + \langle \varphi_3 z^{(3)}, z^{(3)} \rangle$$

In the next section, we will show that φ_3 has a sign and is bounded. This leaves to study the lower order terms, and we will prove that there exists a compact operator M such that

$$\langle Mz, z \rangle_{H^3} = \langle \varphi_0 z, z \rangle + \langle \varphi_1 z', z' \rangle + \langle \varphi_2 z'', z'' \rangle$$

Combining those results yield the following energy estimate :

$$\langle Lz, z \rangle_{H^3} \leq -\delta \|z\|_{H^3}^2 + \langle Mz, z \rangle_{H^3}$$

III.1 Base case

We want to find M_0 such that $\langle M_0 z, z \rangle_{H^3} = \langle \varphi_0 z, z \rangle$. We will use the Fourier transform, with the following convention :

$$\hat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i x \xi} dx$$