

Assignment 3: Duality in linear optimization

Assigned: October 03; Due: October 11.

Exercise 4.5 from *Introduction to Linear Optimization*, Bertsimas & Tsitsiklis [30 pts]

Exercise 4.9 from *Introduction to Linear Optimization*, Bertsimas & Tsitsiklis [30 pts]

Problem: simple rules for order fulfillment [40 pts]

Consider the transportation problem. Let $i = 1, \dots, m$ index facilities and $j = 1, \dots, n$ index customers. We denote by s_i the supply at facility i ; by d_j the demand from customer j ; and by c_{ij} the cost of shipping a unit from facility i to customer j . We define the following decision variable:

x_{ij} = quantity shipped from facility $i = 1, \dots, m$ to customer $j = 1, \dots, n$

The transportation problem minimizes the total shipment costs, subject to supply and demand constraints. It is formulated mathematically as follows:

$$\begin{aligned} \min \quad & \sum_{i=1}^m \sum_{j=1}^n c_{ij} x_{ij} \\ \text{s.t.} \quad & \sum_{i=1}^m x_{ij} \geq d_j, & \forall j = 1, \dots, n \\ & \sum_{j=1}^n x_{ij} \leq s_i, & \forall i = 1, \dots, m \\ & x_{ij} \geq 0, & \forall i, j \end{aligned}$$

We consider a geographical area that can be modeled as a 10×10 miles rectangle, with 100 facilities and 500 customers. We assume that the unit shipment cost c_{ij} is proportional to the Euclidean distance between facility i and customer j . If facility i is located in the point of coordinates (X_i, Y_i) and if customer j is located in the point of coordinates (X_j, Y_j) , then the distance from i to j is given by $\sqrt{(X_i - X_j)^2 + (Y_i - Y_j)^2}$.

You have access to the following data files:

- facilities.csv: A matrix of size 100×3 that indicates, for each of the 100 facilities, (i) its x-coordinate, in miles (0 to 10), (ii) its y-coordinate, in miles (0 to 10) and (iii) its supply.
 - customers.csv: A matrix of size 500×3 that indicates, for each of the 500 customers, (i) its x-coordinate, in miles (0 to 10), (ii) its y-coordinate, in miles (0 to 10) and (iii) its demand.
- a. Implement the problem computationally and report the optimal cost. Out of the 500 customers, how many are fully served from the closest facility? Comment briefly.

- b. Using duality concepts, propose an index β_{ij} that can be amenable to threshold-based implementation. In other words, propose an index β_{ij} such that each customer $j = 1, \dots, n$ is served from the facility, or facilities, that minimize β_{ij} .
- c. We now assume that products are shipped from facilities through warehouses eventually to customers. We index the warehouses by $k = 1, \dots, K$ and denote by C_k the capacity of warehouse k (i.e., the maximum quantity it can handle). We denote by c_{ik}^1 the unit shipment cost from facility i to warehouse k , and by c_{kj}^2 the unit shipment cost from warehouse k to customer j . Modify the problem's formulation to minimize the shipment costs, while adhering to warehouse capacities.
- d. Using duality concepts, propose an index γ_{ijk} that can be amenable to threshold-based implementation. In other words, propose an index γ_{ijk} such that each customer $j = 1, \dots, n$ is served from the facility, or facilities, and the warehouse, or warehouses, that minimize γ_{ijk} .