

## Exercise sheet 11

### supporting the lecture Mathematical Statistics

(Submission of Solutions: 8. February 2016, 12:00 o'clock; Discussion: 10. February 2016)

#### Exercise 1.

(4 points)

Let  $(X_i)_{i \in \mathbb{N}}$  a sequence of i.i.d. distributed random variables with unknown density  $p$ . We want to test

$$H : p = p_0 \quad \text{versus} \quad K : p = p_1,$$

where  $\{x \in \mathbb{R} | p_1(x) > 0\} \subset \{x \in \mathbb{R} | p_0(x) > 0\}$ . Using the Neyman-Pearson Lemma a UMP Test with level  $\alpha$  is then defined via the critical region  $\prod_{i=1}^n r(X_i) \geq C_n(\alpha)$  with  $r(x) := p_1(x)/p_0(x)$ , where  $C_n(\alpha)$  is a constant depending on  $n$  and  $\alpha$ .

- a) Proof that the critical region from above can also be written as

$$\frac{1}{\sqrt{n}} \left( \sum_{i=1}^n \log r(X_i) - \mathbb{E}_0 [\log r(X_i)] \right) \geq k_n(\alpha)$$

with a suitable  $k_n(\alpha)$ .

- b) Assume that it holds  $\sigma_0^2 = \text{Var}_0 [\log r(X_i)] < \infty$ . Show that

$$k_n \rightarrow \sigma_0 u_{1-\alpha},$$

where  $u_\alpha$  is the  $\alpha$  quantile of a  $\mathcal{N}(0, 1)$  distribution.

- c) Show that the sequence of tests is consistent for  $p_1 \neq p_0$ . Use Lemma 5.16.

*Comment: Comment:*

- $\mathbb{E}_0, \text{Var}_0$  denote the expected value respectively the variance under the density  $p_0$ .
- The exercise shows that each Neyman-Person test already exhibits a built-in asymptotic.

#### Exercise 2.

(4 points)

We look at the setting stated in Example 7.6 from the lecture notes.

- a) Determine the maximum likelihood estimator in the sets  $\Delta$  and  $\Theta$ .  
 b) Determine  $T = -2 \log(\lambda(Z))$  with  $Z = (X_1, \dots, X_m, Y_1, \dots, Y_n)$ .

**Exercise 3.**

(4 points)

Proof the following claim from Theorem 7.7: Let  $A \sim \chi_d^2$ ,  $B \sim \chi_c^2$  with  $d > c$ ,  $A - B$  and  $B$  be independent. Then it holds that

$$A - B \sim \chi_{d-c}^2$$

*Hint:* As a first step determine the characteristic function of a  $\chi_n^2$  distributed random variable.

**Exercise 4.**

(4 points)

Let  $Y_1, \dots, Y_n$  be independent with  $Y_i = ax_i + \epsilon_i$  where  $a$  is an unknown parameter,  $x_1, \dots, x_n$  fixed and  $\epsilon_1, \dots, \epsilon_n$  are i.i.d.  $\mathcal{N}(0, 1)$  distributed.

- a) Determine the maximum likelihood estimator  $\hat{a}$  for the parameter  $a$ .
- b) Derive the likelihood quotient test for the hypotheses

$$H : a = 0 \quad \text{versus} \quad K : a \neq 0.$$