



CS201 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: wangqi@sustech.edu.cn

Euclidean Algorithm

- The Euclidean algorithm in pseudocode

ALGORITHM 1 The Euclidean Algorithm.

```
procedure gcd(a, b: positive integers)
  x := a
  y := b
  while y ≠ 0
    r := x mod y
    x := y
    y := r
  return x{gcd(a, b) is x}
```

The number of **divisions** required to find $\text{gcd}(a, b)$ is $O(\log b)$, where $a \geq b$. (this will be proved later.)



Euclidean Algorithm

- The Euclidean algorithm in pseudocode

ALGORITHM 1 The Euclidean Algorithm.

```
procedure gcd( $a, b$ : positive integers)
 $x := a$ 
 $y := b$ 
while  $y \neq 0$ 
     $r := x \bmod y$ 
     $x := y$ 
     $y := r$ 
return  $x$  {gcd( $a, b$ ) is  $x$ }
```

The number of **divisions** required to find $\text{gcd}(a, b)$ is $O(\log b)$, where $a \geq b$. (this will be proved later.)

Why ?



Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n .$$

Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\cdot \\&\cdot \\&\cdot \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

Observation:

$$r_{i+2} = r_i \bmod r_{i+1}$$

Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n.$$

Observation:

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that $r_{i+2} < \frac{1}{2} r_i$

Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\vdots \\&\vdots \\&\vdots \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

Observation:

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that $r_{i+2} < \frac{1}{2} r_i$

Case (i): $r_{i+1} \leq \frac{1}{2} r_i$: $r_{i+2} < r_{i+1} \leq \frac{1}{2} r_i$.

Case (ii): $r_{i+1} > \frac{1}{2} r_i$: $r_{i+2} = r_i \bmod r_{i+1} = r_i - r_{i+1} < \frac{1}{2} r_i$.

Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n.$$

See [Theorem 1 p. 347].

Observation:

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that $r_{i+2} < \frac{1}{2} r_i$

Case (i): $r_{i+1} \leq \frac{1}{2} r_i$: $r_{i+2} < r_{i+1} \leq \frac{1}{2} r_i$.

Case (ii): $r_{i+1} > \frac{1}{2} r_i$: $r_{i+2} = r_i \bmod r_{i+1} = r_i - r_{i+1} < \frac{1}{2} r_i$.

Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

- ◇ **linear**: it is a linear combination of previous terms



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of a_j 's



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of a_j 's
- ◇ **degree k** : a_n is expressed by the previous k terms



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of a_j 's
- ◇ **degree k** : a_n is expressed by the previous k terms
- ◇ **constant coefficients**: coefficients are constants



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of a_j 's
- ◇ **degree k** : a_n is expressed by the previous k terms
- ◇ **constant coefficients**: coefficients are constants

By induction, such a recurrence relation is **uniquely** determined by this recurrence relation, and **k initial conditions** a_0, a_1, \dots, a_{k-1} .



Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Examples

$$P_n = (1.11)P_{n-1}$$

$$f_n = f_{n-1} + f_{n-2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Examples

$$P_n = (1.11)P_{n-1} \quad \text{linear homogeneous recurrence relation of degree 1}$$

$$f_n = f_{n-1} + f_{n-2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Examples

$$P_n = (1.11)P_{n-1} \quad \text{linear homogeneous recurrence relation of degree 1}$$

$$f_n = f_{n-1} + f_{n-2} \quad \text{linear homogeneous recurrence relation of degree 2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Examples

$P_n = (1.11)P_{n-1}$ linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$ linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$ NOT linear

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Examples

$P_n = (1.11)P_{n-1}$ linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$ linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$ NOT linear

$H_n = 2H_{n-1} + 1$ NOT homogeneous

$B_n = nB_{n-1}$



Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Examples

$P_n = (1.11)P_{n-1}$ linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$ linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$ NOT linear

$H_n = 2H_{n-1} + 1$ NOT homogeneous

$B_n = nB_{n-1}$ coefficients are not constants



Solving Linear Recurrence Relations

- **Example** Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2},$$

Which of the following are solutions?

◇ $a_n = 3n$:

◇ $a_n = 2^n$:

◇ $a_n = 5$:



Solving Linear Recurrence Relations

- **Example** Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2},$$

Which of the following are solutions?

◇ $a_n = 3n$: YES

◇ $a_n = 2^n$: NO

◇ $a_n = 5$: YES



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Basic idea: Look for solutions of the form $a_n = r^n$, where r is a constant.



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Basic idea: Look for solutions of the form $a_n = r^n$, where r is a constant.

- ◇ Bring $a_n = r^n$ back to the recurrence relation:

$$\begin{aligned} & r^n = c_1 r^{n-1} + c_2 r^{n-2} + \cdots + c_k r^{n-k}, \\ \text{i.e.,} \quad & r^{n-k} (r^k - c_1 r^{k-1} - \cdots - c_k) = 0 \end{aligned}$$



Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree k* with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where c_1, c_2, \dots, c_k are real numbers, and $c_k \neq 0$.

Basic idea: Look for solutions of the form $a_n = r^n$, where r is a constant.

- ◇ Bring $a_n = r^n$ back to the recurrence relation:

$$\begin{aligned} & r^n = c_1 r^{n-1} + c_2 r^{n-2} + \cdots + c_k r^{n-k}, \\ \text{i.e.,} \quad & r^{n-k} (r^k - c_1 r^{k-1} - \cdots - c_k) = 0 \end{aligned}$$

- ◇ The solutions to the *characteristic equation* can yield an explicit formula for the sequence.

$$(r^k - c_1 r^{k-1} - \cdots - c_k) = 0$$



Recall: Problem IV

■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$



Recall: Problem IV

■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$

◇ What is the closed-form expression of F_n ?



Recall: Problem IV

■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$

◇ What is the closed-form expression of F_n ?

Consider $x^n = x^{n-1} + x^{n-2}$, with $x \neq 0$. There are two different roots

$$\phi = \frac{1 + \sqrt{5}}{2}, \quad \psi = \frac{1 - \sqrt{5}}{2}$$

Then F_n can be the form of $a\phi^n + b\psi^n$. By $F_0 = 0$ and $F_1 = 1$, we have $a + b = 0$ and $\phi a + \psi b = 1$, leading to $a = \frac{1}{\sqrt{5}}$, $b = -a$. Therefore,

$$F_n = \frac{\phi^n - \psi^n}{\sqrt{5}}$$



Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$



Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$



Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

Theorem If this CE has 2 roots $r_1 \neq r_2$, then the sequence $\{a_n\}$ is a solution of the recurrence relation **if and only if** $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$ for $n \geq 0$ and constants α_1, α_2 .



Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

Theorem If this CE has 2 roots $r_1 \neq r_2$, then the sequence $\{a_n\}$ is a solution of the recurrence relation **if and only if** $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$ for $n \geq 0$ and constants α_1, α_2 .

Proof?



Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

Theorem If this CE has 2 roots $r_1 \neq r_2$, then the sequence $\{a_n\}$ is a solution of the recurrence relation **if and only if** $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$ for $n \geq 0$ and constants α_1, α_2 .

Proof?

See [Theorem 1 p. 515].



Solving Linear Recurrence Relations of degree 2

- **Example 1** $a_n = a_{n-1} + 2a_{n-2}$, with $a_0 = 2$, $a_1 = 7$



Solving Linear Recurrence Relations of degree 2

- **Example 1** $a_n = a_{n-1} + 2a_{n-2}$, with $a_0 = 2$, $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$



Solving Linear Recurrence Relations of degree 2

- **Example 1** $a_n = a_{n-1} + 2a_{n-2}$, with $a_0 = 2$, $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and -1 . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$



Solving Linear Recurrence Relations of degree 2

- **Example 1** $a_n = a_{n-1} + 2a_{n-2}$, with $a_0 = 2$, $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and -1 . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 - \alpha_2 = 7$$



Solving Linear Recurrence Relations of degree 2

- **Example 1** $a_n = a_{n-1} + 2a_{n-2}$, with $a_0 = 2$, $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and -1 . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 - \alpha_2 = 7$$

We get $\alpha_1 = 3$ and $\alpha_2 = -1$. Thus,

$$a_n = 3 \cdot 2^n - (-1)^n$$



Solving Linear Recurrence Relations of degree 2

- **Example 2** $a_n = 7a_{n-1} - 10a_{n-2}$, with $a_0 = 2$, $a_1 = 1$



Solving Linear Recurrence Relations of degree 2

- **Example 2** $a_n = 7a_{n-1} - 10a_{n-2}$, with $a_0 = 2$, $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$



Solving Linear Recurrence Relations of degree 2

- **Example 2** $a_n = 7a_{n-1} - 10a_{n-2}$, with $a_0 = 2$, $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$



Solving Linear Recurrence Relations of degree 2

- **Example 2** $a_n = 7a_{n-1} - 10a_{n-2}$, with $a_0 = 2$, $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 + 5\alpha_2 = 1$$



Solving Linear Recurrence Relations of degree 2

- **Example 2** $a_n = 7a_{n-1} - 10a_{n-2}$, with $a_0 = 2$, $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 + 5\alpha_2 = 1$$

We get $\alpha_1 = 3$ and $\alpha_2 = -1$. Thus,

$$a_n = 3 \cdot 2^n - 5^n$$



Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$



Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$



Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

Theorem If this CE has k distinct roots r_i , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all $n \geq 0$, where the α_i 's are constants.



Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

Theorem If this CE has k distinct roots r_i , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all $n \geq 0$, where the α_i 's are constants.

Example $a_n = 2a_{n-1} + 5a_{n-2} - 6a_{n-3}$



The Case of Degenerate Roots

- **Theorem** If the CE $r^2 - c_1 r - c_2 = 0$ has **only 1** root r_0 , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all $n \geq 0$ and two constants α_1 and α_2 .



The Case of Degenerate Roots

- **Theorem** If the CE $r^2 - c_1 r - c_2 = 0$ has **only 1** root r_0 , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all $n \geq 0$ and two constants α_1 and α_2 .

Proof?



The Case of Degenerate Roots

- **Theorem** If the CE $r^2 - c_1 r - c_2 = 0$ has **only 1** root r_0 , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all $n \geq 0$ and two constants α_1 and α_2 .

Proof?

Exercise.



The Case of Degenerate Roots

- **Example** $a_n = 4a_{n-1} - 4a_{n-2}$, with $a_0 = 1$, $a_1 = 0$



The Case of Degenerate Roots

- **Example** $a_n = 4a_{n-1} - 4a_{n-2}$, with $a_0 = 1$, $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$



The Case of Degenerate Roots

- **Example** $a_n = 4a_{n-1} - 4a_{n-2}$, with $a_0 = 1$, $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$



The Case of Degenerate Roots

- **Example** $a_n = 4a_{n-1} - 4a_{n-2}$, with $a_0 = 1$, $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 = 1$$

$$a_1 = 2\alpha_1 + 2\alpha_2 = 0$$



The Case of Degenerate Roots

- **Example** $a_n = 4a_{n-1} - 4a_{n-2}$, with $a_0 = 1$, $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 = 1$$

$$a_1 = 2\alpha_1 + 2\alpha_2 = 0$$

We get $\alpha_1 = 1$ and $\alpha_2 = -1$. Thus,

$$a_n = 2^n - n2^n$$



The Case of Degenerate Roots in General

- **Theorem** [Theorem 4, p.519] Suppose that there are t roots r_1, \dots, r_t with multiplicities m_1, \dots, m_t . Then

$$a_n = \sum_{i=1}^t \left(\sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all $n \geq 0$ and constants $\alpha_{i,j}$.



The Case of Degenerate Roots in General

- **Theorem** [Theorem 4, p.519] Suppose that there are t roots r_1, \dots, r_t with multiplicities m_1, \dots, m_t . Then

$$a_n = \sum_{i=1}^t \left(\sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all $n \geq 0$ and constants $\alpha_{i,j}$.

Example

$$a_n = -3a_{n-1} - 3a_{n-2} - a_{n-3} \text{ with } a_0 = 1, a_1 = -2, \\ a_2 = -1$$



Linear Nonhomogeneous Recurrence Relations

- **Definition** A *linear nonhomogeneous relation* with constant coefficients may contain some terms $F(n)$ that depend only on n

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n).$$

The recurrence relation

$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}$ is called the *associated homogeneous recurrence relation*.



Linear Nonhomogeneous Recurrence Relations

- **Theorem** If $a_n = p(n)$ is any particular solution to the linear nonhomogeneous relation with constant coefficients,

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n),$$

Then all its solutions are of the form

$$a_n = p(n) + h(n),$$

where $a_n = h(n)$ is any solution to the associated homogeneous recurrence relation

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}.$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- t polynomial as the particular solution $p(n)$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- t polynomial as the particular solution $p(n)$.

Let $p(n) = cn + d$, then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means} \\ (2c + 2)n + (2d - 3c) = 0.$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n).$$

We try a degree- t polynomial as the particular solution $p(n)$.

Let $p(n) = cn + d$, then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means} \\ (2c + 2)n + (2d - 3c) = 0.$$

We get $c = -1$ and $d = -3/2$. Thus,

$$p(n) = -n - 3/2$$



Generating Functions

- We may use *generating functions* to **characterize** sequences.



Generating Functions

- We may use *generating functions* to **characterize** sequences.
 - ◇ The sequence $\{a_k\}$ with $a_k = 3$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$



Generating Functions

- We may use *generating functions* to characterize sequences.

- ◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

- ◇ The sequence $\{a_k\}$ with $a_k = 2^k$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence $\{a_k\}$ with $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence $\{a_k\}$ with $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$

Definition The *generating function* for the sequence $a_0, a_1, \dots, a_k, \dots$ of real numbers is the infinite series

$$G(x) = a_0 + a_1x + \dots + a_kx^k$$



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function* $G(x)$ of this infinite sequence $\{a_n\}$ is a polynomial of degree n , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function* $G(x)$ of this infinite sequence $\{a_n\}$ is a polynomial of degree n , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence a_0, a_1, \dots, a_m , with $a_k = C(m, k)$?



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function* $G(x)$ of this infinite sequence $\{a_n\}$ is a polynomial of degree n , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence a_0, a_1, \dots, a_m , with $a_k = C(m, k)$?

$$G(x) = C(m, 0) + \dots + C(m,$$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$1, 1, 1, 1, 1, \dots$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$

$1, a, a^2, a^3, a^4, \dots$

◇ $G(x) = 1/(1 - x)^2$ for $|x| < 1$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$

◇ $G(x) = 1/(1 - x)^2$ for $|x| < 1$

$$1, 2, 3, 4, 5, \dots$$



Operations of Generating Functions

- **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$



Operations of Generating Functions

■ **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1-x)^2$ for $|x| < 1$



Operations of Generating Functions

■ **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1-x)^2$ for $|x| < 1$

$$f(x) = 1/(1-x), \quad g(x) = 1/(1-x)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k 1 \right) x^k = \sum_{k=0}^{\infty} (k+1) x^k$$



Operations of Generating Functions

■ **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1 - ax)^2$ for $|x| < 1$



Operations of Generating Functions

■ **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1 - ax)^2$ for $|x| < 1$

$$f(x) = 1/(1 - ax), \quad g(x) = 1/(1 - ax)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a^j a^{k-j} \right) x^k = \sum_{k=0}^{\infty} (k+1) a^k x^k$$



Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$



Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k)x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k)a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k)x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$



Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$

$$\frac{1}{(1-x)^2} = \sum_{k=0}^{\infty} (k+1) x^k = 1 + 2x + 3x^2 + \dots$$



Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$



Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\ln(1+x) = \sum_{k=0}^{\infty} \frac{(-1)^{k+1} x^k}{k} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$



Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers?



Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers?

This is **equivalent** to the problem of r -combinations from a set with n elements when **repetition** is allowed.



Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers?

This is **equivalent** to the problem of r -combinations from a set with n elements when **repetition** is allowed.

$$C(n + r - 1, r) = C(19, 17) = C(19, 2)$$



Counting and Generating Functions

- **Problem 2** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers with $2 \leq x_1 \leq 5$,
 $3 \leq x_2 \leq 6$, $4 \leq x_3 \leq 7$.



Counting and Generating Functions

- **Problem 2** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers with $2 \leq x_1 \leq 5$,
 $3 \leq x_2 \leq 6$, $4 \leq x_3 \leq 7$.

Using *generating functions*, the number is the **coefficient** of x^{17} in the expansion of

$$(x^2 + x^3 + x^4 + x^5)(x^3 + x^4 + x^5 + x^6)(x^4 + x^5 + x^6 + x^7)$$



Counting and Generating Functions

- **Problem 3** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?



Counting and Generating Functions

- **Problem 3** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?

The coefficient of x^8 in the expansion

$$(x^2 + x^3 + x^4)^3$$



Counting and Generating Functions

- **Problem 4** Use **generating functions** to find the number of **k -combinations of a set with n elements**, $C(n, k)$.



Counting and Generating Functions

- **Problem 4** Use **generating functions** to find the number of **k -combinations of a set with n elements**, $C(n, k)$.

Each of the n elements in the set contributes the term $(1 + x)$ to the generating function $f(x) = \sum_{k=0}^n a^k x^k$.
Hence, $f(x) = (1 + x)^n$.

Then by the **binomial theorem**, we have $a_k = \binom{n}{k}$.



Next Lecture

- relation ...

