



CS201 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: wangqi@sustech.edu.cn

Binary Relations

- Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$, the *Cartesian product* $A \times B$ is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$

Definition: Let A and B be two sets. A *binary relation from A to B* is a *subset* of a *Cartesian product* $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.



Binary Relations

- Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$, the *Cartesian product* $A \times B$ is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$

Definition: Let A and B be two sets. A *binary relation from A to B* is a *subset* of a *Cartesian product* $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Definition: A *relation on the set A* is a relation *from A to itself*.



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.



Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.



Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.


Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.

Transitive Relation: A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.



Powers of R

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$



Powers of R

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Theorem The relation R on a set A is **transitive if and only if** $R^n \subseteq R$ for $n = 1, 2, 3, \dots$



Representing Relations


- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$

Representing Relations

- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*


Representing Relations

- Some ways to represent *n*-ary relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*




Representing Relations


- Some ways to represent *n*-ary relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*




reflexive



irreflexive



symmetric



antisymmetric

Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ is not in R .



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ is not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ is not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?

How to make R reflexive by minimum number of additions?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ is not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?

How to make R reflexive by minimum number of additions?

Add $(2, 2)$ and $(3, 3)$

Then $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 2), (3, 3)\} \supseteq R$



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ is not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?

How to make R reflexive by minimum number of additions?

Add $(2, 2)$ and $(3, 3)$

Then $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 2), (3, 3)\} \supseteq R$

The minimal set $S \supseteq R$ is called the reflexive closure of R .



Reflexive Closure

- The set S is called *the reflexive closure of R* if it:



Reflexive Closure

- The set S is called *the reflexive closure of R* if it:
 - ◇ contains R
 - ◇ is reflexive
 - ◇ is minimal (is contained in every reflexive relation Q that contains R ($R \subseteq Q$), i.e., $S \subseteq Q$)



Closures on Relations

- Relations can have different properties:
 - reflexive
 - symmetric
 - transitive



Closures on Relations

- Relations can have different properties:
 - reflexive
 - symmetric
 - transitive

We define:

- reflexive closures
- symmetric closures
- transitive closures



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (symmetric closure)

$$R = \{(1, 2), (1, 3), (2, 2)\} \text{ on } A = \{1, 2, 3\}$$

What is the symmetric closure S of R ?



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (symmetric closure)

$$R = \{(1, 2), (1, 3), (2, 2)\} \text{ on } A = \{1, 2, 3\}$$

What is the symmetric closure S of R ?

$$R = \{(1, 2), (1, 3), (2, 2)\} \cup \{(2, 1), (3, 1)\}$$



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is R transitive?



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is R transitive? **No**

How to make it transitive?



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is R transitive? No

How to make it transitive?

$$S = \{(1, 2), (2, 2), (2, 3)\} \cup \{(1, 3)\}$$



Transitive Closure

- We can represent the relation on the graph.



Transitive Closure

- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path



Transitive Closure

- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Transitive Closure


- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Transitive Closure


- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Transitive Closure



- We can represent the relation on the graph.

Finding a **transitive closure** corresponds to **finding all pairs** of elements that are connected **with a directed path**

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Paths in Directed Graphs

- **Definition** A *path* from a to b in the directed graph G is a *sequence of edges* $(x_0, x_1), (x_1, x_2), \dots, (x_{n-1}, x_n)$ in G , where n is nonnegative and $x_0 = a$ and $x_n = b$. A path of length $n \geq 1$ that begins and ends at the same vertex is called a *circuit* or *cycle*.



Paths in Directed Graphs

- **Definition** A *path* from a to b in the directed graph G is a *sequence of edges* $(x_0, x_1), (x_1, x_2), \dots, (x_{n-1}, x_n)$ in G , where n is nonnegative and $x_0 = a$ and $x_n = b$. A path of length $n \geq 1$ that begins and ends at the same vertex is called a *circuit* or *cycle*.

Theorem Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.



Path Length

- **Theorem** Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.


Proof (by induction)



Path Length

- **Theorem** Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.

Proof (by induction)




Path Length

- **Theorem** Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.

Proof (by induction)



Path of length 1



Path of length $n+1$

Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$



Connectivity Relation


- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .


$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$


Example:

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:


$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$

$$R^4 = \emptyset$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:

$$A = \{1, 2, 3, 4\}$$


$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$

$$R^4 = \emptyset$$

$$R^* = \{(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)\}$$



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)

- ◇ There are at most n different elements we can visit on a path if the path does not have loops



Connectivity


- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)

- ◇ There are at most n different elements we can visit on a path if the path does not have loops



- ◇ Loops may increase the length but the same node is visited more than once



Connectivity


- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)

- ◇ There are at most n different elements we can visit on a path if the path does not have loops



- ◇ Loops may increase the length but the same node is visited more than once



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$R^* = \bigcup_{k=1}^n R^k$$



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$R^* = \bigcup_{k=1}^n R^k$$

Theorem: The transitive closure of a relation R equals the connectivity relation R^* .



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$R^* = \bigcup_{k=1}^n R^k$$

Theorem: The transitive closure of a relation R equals the connectivity relation R^* .

Recall Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof



- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
2. $R^* \subseteq S$ whenever S is a transitive relation containing R



- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
2. $R^* \subseteq S$ whenever S is a transitive relation containing R

1. If $(a, b) \in R^*$ and $(b, c) \in R^*$, then there are paths from a to b and from b to c in R . Thus, there is a path from a to c in R . This means that $(a, c) \in R^*$.



- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
 2. $R^* \subseteq S$ whenever S is a transitive relation containing R
2. Suppose that S is a transitive relation containing R .



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
 2. $R^* \subseteq S$ whenever S is a transitive relation containing R
2. Suppose that S is a transitive relation containing R .
Then S^n is also transitive and $S^n \subseteq S$. Why?



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
2. $R^* \subseteq S$ whenever S is a transitive relation containing R

2. Suppose that S is a transitive relation containing R .

Then S^n is also transitive and $S^n \subseteq S$. Why?

We have $S^* \subseteq S$. Thus, $R^* \subseteq S^* \subseteq S$



Find Transitive Closure

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.



Find Transitive Closure

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$\mathbf{M}_{R^*} = \mathbf{M}_R \vee \mathbf{M}_R^{[2]} \vee \mathbf{M}_R^{[3]} \vee \dots \vee \mathbf{M}_R^{[n]}$$



Find Transitive Closure

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$\mathbf{M}_{R^*} = \mathbf{M}_R \vee \mathbf{M}_R^{[2]} \vee \mathbf{M}_R^{[3]} \vee \dots \vee \mathbf{M}_R^{[n]}$$

Example

$$\mathbf{M}_R = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

$$\mathbf{M}_{R^*} = ?$$



Simple Transitive Closure Algorithm

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

procedure transClosure (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$A := B := \mathbf{M}_R$;

for $i := 2$ to n

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

return B

// B is the zero-one matrix for R^*



Simple Transitive Closure Algorithm

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

procedure transClosure (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$A := B := \mathbf{M}_R$;

for $i := 2$ to n

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

return B

// B is the zero-one matrix for R^*

This algorithm takes $\Theta(n^4)$ time.



Simple Transitive Closure Algorithm

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

procedure transClosure (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$A := B := \mathbf{M}_R$;

for $i := 2$ to n

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

return B

// B is the zero-one matrix for R^*

This algorithm takes $\Theta(n^4)$ time. Why?



Roy-Warshall Algorithm

```
procedure Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)  
  // computes  $R^*$  with zero-one matrices  
   $W := \mathbf{M}_R$ ;  
  for  $k := 1$  to  $n$   
    for  $i := 1$  to  $n$   
      for  $j := 1$  to  $n$   
         $w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$   
  return  $W$   
  //  $W$  is the zero-one matrix for  $R^*$ 
```



Roy-Warshall Algorithm

procedure Warshall (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$W := \mathbf{M}_R$;

for $k := 1$ to n

for $i := 1$ to n

for $j := 1$ to n

$w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$

return W

// W is the zero-one matrix for R^*

$w_{ij} = 1$ means there is a path from i to j going only through nodes $\leq k$.

$$W_{ij}^{[k]} = W_{ij}^{[k-1]} \vee \left(W_{ik}^{[k-1]} \wedge W_{kj}^{[k-1]} \right)$$



Roy-Warshall Algorithm

```
procedure Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)  
  // computes  $R^*$  with zero-one matrices  
   $W := \mathbf{M}_R$ ;  
  for  $k := 1$  to  $n$   
    for  $i := 1$  to  $n$   
      for  $j := 1$  to  $n$   
         $w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$   
  return  $W$   
  //  $W$  is the zero-one matrix for  $R^*$ 
```

$w_{ij} = 1$ means there is a path from i to j going only through nodes $\leq k$.


$$W_{ij}^{[k]} = W_{ij}^{[k-1]} \vee \left(W_{ik}^{[k-1]} \wedge W_{kj}^{[k-1]} \right)$$

This algorithm takes $\Theta(n^3)$ time.



Example

Find the matrices W_0 , W_1 , W_2 , W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .




Let $v_1 = a$, $v_2 = b$, $v_3 = c$, $v_4 = d$.



Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .




Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .




Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .



Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.


$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ \textcolor{red}{1} & 0 & 1 & \textcolor{red}{1} \end{bmatrix}$$

Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .



Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ \textcolor{red}{1} & 0 & 1 & \textcolor{red}{1} \end{bmatrix}$$

$$W_4 = \begin{bmatrix} \textcolor{red}{1} & 0 & \textcolor{red}{1} & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & \textcolor{red}{1} & 1 \\ \textcolor{red}{1} & 0 & 1 & \textcolor{red}{1} \end{bmatrix}$$

Equivalence Relation

- **Definition** A relation R on a set A is called an *equivalence relation* if it is *reflexive*, *symmetric*, and *transitive*.



Equivalence Relation

- **Definition** A relation R on a set A is called an *equivalence relation* if it is *reflexive*, *symmetric*, and *transitive*.

Example:

$$A = \{0, 1, 2, 3, 4, 5, 6\}$$

$$R = \{(a, b) : a \equiv b \pmod{3}\}$$



Equivalence Relation

- **Definition** A relation R on a set A is called an *equivalence relation* if it is *reflexive*, *symmetric*, and *transitive*.

Example:

$$A = \{0, 1, 2, 3, 4, 5, 6\}$$

$$R = \{(a, b) : a \equiv b \pmod{3}\}$$

R has the following pairs:

- $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
- $(1, 1), (1, 4), (4, 1), (4, 4)$
- $(2, 2), (2, 5), (5, 2), (5, 5)$



Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$



Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$


Is R reflexive?



Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$


Is R reflexive?



Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$

Is R reflexive? Yes




Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$

Is R reflexive? Yes

Is R symmetric?




Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$

Is R reflexive? Yes

Is R symmetric? Yes




Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$

Is R reflexive? Yes

Is R symmetric? Yes

Is R transitive?




Equivalence Relation

- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$

Is R reflexive? Yes

Is R symmetric? Yes

Is R transitive? Yes



Equivalence Relation


- Relation R on $A = \{0, 1, 2, 3, 4, 5, 6\}$ has the pairs:
 - $(0, 0), (0, 3), (3, 0), (0, 6), (6, 0), (3, 3), (3, 6), (6, 3), (6, 6)$
 - $(1, 1), (1, 4), (4, 1), (4, 4)$
 - $(2, 2), (2, 5), (5, 2), (5, 5)$

Is R reflexive? Yes

Is R symmetric? Yes

Is R transitive? Yes

R is an equivalence relation



Examples of Equivalence Relations

■ Examples

“Strings a and b have the same length.”

“Integers a and b have the same absolute value.”

“Real numbers a and b have the same fractional part (i.e., $a - b \in \mathbf{Z}$).”



Examples of Equivalence Relations

■ Examples

“Strings a and b have the same length.”

“Integers a and b have the same absolute value.”

“Real numbers a and b have the same fractional part (i.e., $a - b \in \mathbf{Z}$).”

“The relation \geq between real numbers.”

“has a common factor greater than 1 between natural numbers.”



Equivalence Class

- **Definition** Let R be an **equivalence relation** on a set A . The **set of all elements** that are related to an element a of A is called the **equivalence class** of a , denoted by $[a]_R$. When only one relation is considered, we use the notation $[a]$.



Equivalence Class

- **Definition** Let R be an **equivalence relation** on a set A . The **set of all elements** that are related to an element a of A is called the **equivalence class** of a , denoted by $[a]_R$. When only one relation is considered, we use the notation $[a]$.

$$[a]_R = \{b : (a, b) \in R\}$$



Equivalence Class

- **Definition** Let R be an **equivalence relation** on a set A . The **set of all elements** that are related to an element a of A is called the **equivalence class** of a , denoted by $[a]_R$. When only one relation is considered, we use the notation $[a]$.

$$[a]_R = \{b : (a, b) \in R\}$$

$$A = \{0, 1, 2, 3, 4, 5, 6\}$$

$$R = \{(a, b) : a \equiv b \pmod{3}\}$$



Equivalence Class

- **Definition** Let R be an **equivalence relation** on a set A . The **set of all elements** that are related to an element a of A is called the **equivalence class** of a , denoted by $[a]_R$. When only one relation is considered, we use the notation $[a]$.

$$[a]_R = \{b : (a, b) \in R\}$$

$$A = \{0, 1, 2, 3, 4, 5, 6\}$$

$$R = \{(a, b) : a \equiv b \pmod{3}\}$$

$$[0] = [3] = [6] = \{0, 3, 6\}$$

$$[1] = [4] = \{1, 4\}$$

$$[2] = [5] = \{2, 5\}$$



Examples of Equivalence Classes

■ Examples

“Strings a and b have the same length.”

“Integers a and b have the same absolute value.”

“Real numbers a and b have the same fractional part (i.e., $a - b \in \mathbf{Z}$).”



Examples of Equivalence Classes

■ Examples

“Strings a and b have the same length.”

$[a]$ = the set of all strings of the same length as a

“Integers a and b have the same absolute value.”

$[a]$ = the set $\{a, -a\}$

“Real numbers a and b have the same fractional part (i.e., $a - b \in \mathbf{Z}$).”

$[a]$ = the set $\{\dots, a - 2, a - 1, a, a + 1, a + 2, \dots\}$



Equivalence Class

- **Theorem** Let R be an **equivalence relation** on a set A . The following statements are **equivalent**:

(i) $a R b$

(ii) $[a] = [b]$

(iii) $[a] \cap [b] \neq \emptyset$



Equivalence Class

- **Theorem** Let R be an **equivalence relation** on a set A . The following statements are **equivalent**:

$$(i) \ a R b$$

$$(ii) \ [a] = [b]$$

$$(iii) \ [a] \cap [b] \neq \emptyset$$

Proof

$$(i) \rightarrow (ii):$$



Equivalence Class

- **Theorem** Let R be an **equivalence relation** on a set A . The following statements are **equivalent**:

$$(i) \ a R b$$

$$(ii) \ [a] = [b]$$

$$(iii) \ [a] \cap [b] \neq \emptyset$$

Proof

$$(i) \rightarrow (ii): \quad \text{prove } [a] \subseteq [b] \text{ and } [b] \subseteq [a]$$



Equivalence Class

- **Theorem** Let R be an **equivalence relation** on a set A . The following statements are **equivalent**:

$$(i) \ a R b$$

$$(ii) \ [a] = [b]$$

$$(iii) \ [a] \cap [b] \neq \emptyset$$

Proof

$$(i) \rightarrow (ii): \quad \text{prove } [a] \subseteq [b] \text{ and } [b] \subseteq [a]$$

$$(ii) \rightarrow (iii):$$



Equivalence Class

- **Theorem** Let R be an equivalence relation on a set A . The following statements are equivalent:

$$(i) \quad a R b$$

$$(ii) \quad [a] = [b]$$

$$(iii) \quad [a] \cap [b] \neq \emptyset$$

Proof

$$(i) \rightarrow (ii): \quad \text{prove } [a] \subseteq [b] \text{ and } [b] \subseteq [a]$$

$$(ii) \rightarrow (iii): \quad [a] \text{ is not empty (} R \text{ reflexive)}$$



Equivalence Class

- **Theorem** Let R be an **equivalence relation** on a set A . The following statements are **equivalent**:

$$(i) \quad a R b$$

$$(ii) \quad [a] = [b]$$

$$(iii) \quad [a] \cap [b] \neq \emptyset$$

Proof

$$(i) \rightarrow (ii): \quad \text{prove } [a] \subseteq [b] \text{ and } [b] \subseteq [a]$$

$$(ii) \rightarrow (iii): \quad [a] \text{ is not empty (} R \text{ reflexive)}$$

$$(iii) \rightarrow (i):$$



Equivalence Class

- **Theorem** Let R be an **equivalence relation** on a set A . The following statements are **equivalent**:

(i) $a R b$

(ii) $[a] = [b]$

(iii) $[a] \cap [b] \neq \emptyset$

Proof

(i) \rightarrow (ii): prove $[a] \subseteq [b]$ and $[b] \subseteq [a]$

(ii) \rightarrow (iii): $[a]$ is **not empty** (R **reflexive**)

(iii) \rightarrow (i): there exists a c s.t. $c \in [a]$ and $c \in [b]$



Partition of a Set S

- **Definition** Let S be a set. A collection of nonempty subsets of S A_1, A_2, \dots, A_k is called *a partition of S* if:


$$A_i \cap A_j = \emptyset, \ i \neq j \text{ and } S = \bigcup_{i=1}^k A_i$$



Partition of a Set S

- **Definition** Let S be a set. A collection of nonempty subsets of S A_1, A_2, \dots, A_k is called *a partition of S* if:

$$A_i \cap A_j = \emptyset, \quad i \neq j \quad \text{and} \quad S = \bigcup_{i=1}^k A_i$$



Partition of a Set S

- **Definition** Let S be a set. A collection of nonempty subsets of S A_1, A_2, \dots, A_k is called *a partition of S* if:

$$A_i \cap A_j = \emptyset, \quad i \neq j \quad \text{and} \quad S = \bigcup_{i=1}^k A_i$$

Example:

$$A = \{0, 1, 2, 3, 4, 5, 6\}$$

$$A_1 = \{0, 3, 6\}, \quad A_2 = \{1, 4\}, \quad A_3 = \{2, 5\}$$



Partition of a Set S

- **Definition** Let S be a set. A collection of nonempty subsets of S A_1, A_2, \dots, A_k is called *a partition of S* if:

$$A_i \cap A_j = \emptyset, \quad i \neq j \quad \text{and} \quad S = \bigcup_{i=1}^k A_i$$

Example:

$$A = \{0, 1, 2, 3, 4, 5, 6\}$$

$$A_1 = \{0, 3, 6\}, \quad A_2 = \{1, 4\}, \quad A_3 = \{2, 5\}$$

Is A_1, A_2, A_3 a partition of S ?



Equivalence Classes and Partitions

- **Theorem** Let R be an equivalence relation on a set A . Then union of all the equivalence classes of R is A :

$$A = \bigcup_{a \in A} [a]_R$$



Equivalence Classes and Partitions

- **Theorem** Let R be an equivalence relation on a set A . Then union of all the equivalence classes of R is A :

$$A = \bigcup_{a \in A} [a]_R$$

Theorem The equivalence classes form a partition of A .



Equivalence Classes and Partitions

- **Theorem** Let R be an equivalence relation on a set A . Then union of all the equivalence classes of R is A :

$$A = \bigcup_{a \in A} [a]_R$$

Theorem The equivalence classes form a partition of A .

Theorem Let $\{A_1, A_2, \dots, A_i, \dots\}$ be a partition of S . Then there is an equivalence relation R on S , that has the sets A_i as its equivalence classes.



Next Lecture

- relation, graph ...

