

Confidence intervals

Table of Contents

Introduction.....	1
Example 1 Probability distribution of the mean of normal random variables.....	1
Example 2 Plot area under normal distribution.....	4
Example 3 Computing confidence intervals using Student t distribution.....	5
Exersizes.....	6

Copyright (C) 2022 Miodrag Bolic

This program is free software: you can redistribute it and/or modify it under the terms of the GNU General Public License as published by the Free Software Foundation, either version 3 of the License, or (at your option) any later version. This program is distributed in the hope that it will be useful, but WITHOUT ANY WARRANTY; without even the implied warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the GNU General Public License for more details <<https://www.gnu.org/licenses/>>.

This code was developed by Miodrag Bolic for the book PERVASIVE CARDIOVASCULAR AND RESPIRATORY MONITORING DEVICES

```
% Changing the path from main_folder to a particular chapter
main_path=fileparts(which('Main_Content.mlx'));
if ~isempty(main_path)
    %addpath(append(main_path, '/Chapter2'))
    cd (append(main_path, '/Chapter1/CLT/'))
    addpath(append(main_path, '/Service'))
end
SAVE_FLAG=0; % saving the figures in a file
```


Introduction

Example 1 Probability distribution of the mean of normal random variables

Plot the empirical distribution of the mean of a) $n=1$, b) $n=2$ and c) $n=20$ uniform random variables.

```
close all
clear_all_but('SAVE_FLAG');

% a) Plot hystogram for n=1
n1=rand(1,2000);
figure
histogram(n1, 'Normalization','probability')
ylabel('Relative frequency of X')
xlabel('x')
title('The empirical distribution without averaging random variables')
```




```

annotation_save('a','Fig1.1a.jpg', SAVE_FLAG);

%annotation('textbox',...
%   [0.0117142857142857 0.00952380952380952 0.0615 0.0547619047619048],...
%   'String',{'a')'},...
%   'FitBoxToText','off',...
%   'EdgeColor',[1 1 1]);

% b) Plot hystogram for n=2
n2=rand(2,2000);
m=mean(n2);
figure
histogram(m, 'Normalization','probability')
ylabel('Relative frequency of X=X_1+X_2')
xlabel('x')
title('The empirical distribution after averaging 2 variables')

```




```

annonation_save('b)', "Fig1.1b.jpg", SAVE_FLAG);

% c) Plot hystogram for n=20
n2=rand(20,2000);
m=mean(n2);
figure
histogram(m, 'Normalization','probability')
ylabel('Relative frequency of  $X = X_1 + \dots + X_{20}$ ')
xlabel('x')
title('The empirical distribution after averaging 20 variables')

```



```

annnotation_save('c','Fig1.1c.jpg', SAVE_FLAG);

%histfit(m)

```

Example 2 Plot area under normal distribution

This code is from: <https://www.mathworks.com/matlabcentral/answers/128788-need-help-plotting-confidence-intervals>

```

alpha = 0.05;           % significance level - change to 0.34 for 66% confidence
mu = 0;                 % mean
sigma = 1;              % std
x = linspace(mu-5*sigma, mu+5*sigma, 500);

cutoff1 = norminv(alpha/2, mu, sigma);           % Lower 95% CI is p = 0.025
cutoff2 = norminv(1-alpha/2, mu, sigma);         % Upper 95% CI is p = 0.975
y = normpdf(x, mu, sigma);

xci = [linspace(mu-5*sigma, cutoff1); linspace(cutoff2, mu+5*sigma)];
yci = normpdf(xci, mu, sigma);


figure(1)
plot(x, y, '-k', 'LineWidth', 1.5)
patch(x, y, [0.9 0.9 0.9])
patch([xci(1,:) cutoff1], [yci(1,:) 0], [1 1 1])
patch([cutoff2 xci(2,:)], [0 yci(2,:)], [1 1 1])
xticklabels({})

```

```

text(-2.5,-0.02,{ ' \mu-2\sigma' })
text(-0.5,-0.02,{ ' \mu=0' })
text(1.5,-0.02,{ ' \mu+2\sigma' })
message = sprintf('95 %% of the area \n under the curve');
text(-1.5,0.1,message)
title('PDF of a normal distribution N(0,1)')

```



Example 3 Computing confidence intervals using Student t distribution

Generate a random sample of size 50 drawn from a normal population with mean 2V and standard deviation 0.5V. Compute confidence intervals at 95% confidence. This solution is from Mathworks example at <https://www.mathworks.com/help/stats/tinv.html>.

```

mu = 2;
sigma = 0.5;
n = 50;

x = normrnd(mu,sigma,n,1);

```

Compute the sample mean, standard error, and degrees of freedom.

```

xbar = mean(x);
se = std(x)/sqrt(n);
nu = n - 1;

```

Find the upper and lower confidence bounds for the 95% confidence interval.

```

conf = 0.95;
alpha = 1 - conf;
pLo = alpha/2;
pUp = 1 - alpha/2;

```

Compute the critical values for the confidence bounds.

```
crit = tinv([pLo pUp], nu);
```

Determine the confidence interval for the population mean.

```
ci = xbar + crit*se
```

```

ci = 1x2
    1.7935    2.0828

```

Uncertainty at 95% confidence is:

```
U_95=(ci(2)-xbar)/2
```

```
U_95 = 0.0723
```

If we replace the t-distribution with the normal distribution at 95% confidence, we will obtain similar results:

```

k=2;
ci_normal = [xbar - k*se, xbar + k*se]

```

```

ci_normal = 1x2
    1.7942    2.0821

```

Uncertainty at 95% confidence is:

```
U_95_normal=(ci_normal(2)-xbar)/2
```

```
U_95_normal = 0.0720
```

Exersizes

Excercise 1: Repeat the analysis for Example 3 and comment on the difference in confidence intervals when a) $n=5$, $n=15$, $n=200$. Replace $k=2$ with $k=1.96$.

Excercise 2: Repeat the analysis for Example 3 but compute confidence intervals at 99% confidence.