Death Rate among Surgical Inpatients with Serious Treatable Complications Technical Specifications

Patient Safety Indicators 04 (PSI 04)

AHRQ Quality Indicators™, Version 5.0 March 2015 Provider-Level Indicator

Type of Score: Rate

Description

In-hospital deaths per 1,000 surgical discharges, among patients ages 18 through 89 years or obstetric patients, with serious treatable complications (deep vein thrombosis/ pulmonary embolism, pneumonia, , sepsis, shock/cardiac arrest or gastrointestinal hemorrhage/acute ulcer). Includes metrics for the number of discharges for each type of complication. Excludes cases transferred to an acute care facility.

[NOTE: The software provides the rate per hospital discharge. However, common practice reports the measure as per 1,000 discharges. The user must multiply the rate obtained from the software by 1,000 to report in-hospital deaths per 1,000 hospital discharges.]

Numerator

Overall:

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Stratum 04A: DEEP VEIN THROMBOSIS / PULMONARY EMBOLISM (DVT/PE)

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Stratum 04B: PNEUMONIA

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Stratum 04C: SEPSIS

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Stratum 04D: SHOCK / CARDIAC ARREST

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator

Stratum 04E: GASTROINTESTINAL (GI) HEMORRHAGE / ACUTE ULCER

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Denominator

Overall:

Surgical discharges, for patients ages 18 through 89 years or MDC 14 (pregnancy, childbirth, and puerperium), with all of the following:

- any-listed ICD-9-CM procedure codes for an operating room procedure; and
- the principal procedure occurring within 2 days of admission or an admission type of elective (ATYPE=3); and
- meet the inclusion and exclusion criteria for Stratum A (deep vein thrombosis or pulmonary embolism), Stratum B (pneumonia), , Stratum C (sepsis), Stratum D (shock or cardiac arrest), or Stratum E (gastrointestinal hemorrhage or acute ulcer) defined below.

Surgical discharges are defined by specific DRG or MS-DRG codes.

See Patient Safety Indicators Appendices:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs
- Appendix E Surgical Discharge MS-DRGs

Exclude cases:

- transferred to an acute care facility (DISP = 2)
- with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), or principal diagnosis (DX1=missing)

To ensure that the strata are mutually exclusive within each indicator, the strata were prioritized in order of their relative prevalence in the reference population data. In the event that

a discharge record meets the denominator criteria for several strata, the software assigns the record to the one (and only one) candidate stratum whose outcome is most prevalent in the reference population. That is to say that if a record met the criteria to be in the denominator for strata C and D, and if the outcome in D were more prevalent than the one in C in the reference population dataset, then the software would only assign the record to stratum D, and would not assign it to C.

For PSI 04, prioritization to ensure mutual exclusivity for the strata is as follows:

STRATUM D: SHOCK/CARDIAC ARREST

STRATUM C: SEPSIS

STRATUM B: PNEUMONIA STRATUM A: DVT/PE

STRATUM E: GI HEMORRHAGE/ACUTE ULCER

Stratum 04A: DEEP VEIN THROMBOSIS / PULMONARY EMBOLISM (DVT/PE)

Surgical discharges, for patients ages 18 through 89 years or MDC 14 (pregnancy, childbirth, and puerperium), with all of the following:

- any-listed ICD-9-CM procedure codes for an operating room procedure; and
- the principal procedure occurring within 2 days of admission or an admission type of elective (ATYPE=3); and
- any secondary ICD-9-CM diagnosis codes for deep vein thrombosis or pulmonary embolism.

Surgical discharges are defined by specific DRG or MS-DRG codes.

See Patient Safety Indicators Appendices:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs
- Appendix E Surgical Discharge MS-DRGs

ICD-9-CM Pulmonary embolism or deep vein thrombosis diagnosis codes:

4151	PULMON EMBOLISM/INFARCT (end 1995)	45181	ILIAC THROMBOPHLEBITIS
41511	IATROGEN PILM EMB/INFARC `	4519	THROMBOPHLEBITIS NOS (used with data
			prior to FY 2010)
41513	SADDLE EMBOL PULMON ART	45340	AC DVT/EMBL LOW EXT NOS
41519	PULM EMBOL/INFARCT NEC	45341	AC DVT/EMB PROX LOW EXT
45111	FEMORAL VEIN PHLEBITIS	45342	AC DVT/EMB DISTL LOW EXT
45119	DEEP PHLEBITIS-LEG NEC	4538	VENOUS THROMBOSIS NEC (end 2009)
4512	THROMBOPHLEBITIS LEG NOS	4539	VENOUS THROMBOSIS NOS (used with
			data prior to FY 2010)

Exclude cases:

- with a principal ICD-9-CM diagnosis code for deep vein thrombosis or pulmonary embolism (see above)
- with a principal ICD-9-CM diagnosis code for abortion-related or postpartum obstetric pulmonary embolism
- transferred to an acute care facility (DISP = 2)

• with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), or principal diagnosis (DX1=missing)

ICD-9-CM Abortion-related or postpartum obstetric pulmonary embolism diagnosis codes:

63460	SPON ABORT W EMBOL-UNSP	63761	AB NOS W EMBOLISM-INC
63461	SPON ABORT W EMBOL-INC	63762	AB NOS W EMBOLISM-COMP
63462	SPON ABOT W EMBOL-COMP	6386	ATTEMP ABORT W EMBOLISM
63560	LEGAL ABORT W EMBOL-UNSP	6396	POSTABORTION EMBOLISM
63561	LEGAL ABORT W EMBOL-INC	67320	OB PULM EMBOL NOS-UNSPEC
63562	LEGAL AMOT W EMBOL-COMP	67321	PULM EMBOL NOS-DELIVERED
63660	ILLEG AB W EMBOLISM-UNSP	67322	PULM EMBOL NOS-DEL W P/P
63661	ILLEG AB W EMBOLISM-INC	67323	PULM EMBOL NOS-ANTEPART
63662	ILLEG AB W EMBOLISM-COMP	67324	PULM EMBOL NOS-POSTPART
63760	AB NOS W EMBOLISM-UNSP		

Stratum B: PNEUMONIA

Surgical discharges, for patients ages 18 through 89 years or MDC 14 (pregnancy, childbirth, and puerperium), with all of the following:

- any-listed ICD-9-CM procedure codes for an operating room procedure; and
- the principal procedure occurring within 2 days of admission or an admission type of elective (ATYPE=3); and
- any secondary ICD-9-CM diagnosis codes for pneumonia.

Surgical discharges are defined by specific DRG or MS-DRG codes.

See Patient Safety Indicators Appendices:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs
- Appendix E Surgical Discharge MS-DRGs

ICD-9-CM Pneumonia diagnosis codes:

4820	K. PNEUMONIAE PNEUMONIA	48249	STAPH PNEUMONIA NEC
4821	PSEUDOMONAL PNEUMONIA	4828	BACTERIAL PNEUMONIA NEC
4822	H.INFLUENZAE PNEUMONIA	48281	PNEUMONIA ANAEROBES
4823	STREPTOCOCCAL PNEUMONIA	48282	PNEUMONIA E COLI
48230	STREPTOCOCCAL PNEUMN NOS	48283	PNEUMO OTH GRM-NEG BACT
48231	PNEUMONIA STRPTOCOCCUS A	48284	LEGIONNAIRES DISEASE
48232	PNEUMONIA STRPTOCOCCUS B	48289	PNEUMONIA OTH SPCF BACT
48239	PNEUMONIA OTH STREP	4829	BACTERIAL PNEUMONIA NOS
4824	STAPHYLOCOCCAL PNEUMONIA	485	BRONCHOPNEUMONIA, ORGANISM
			UNSPECIFIED
48240	STAPHYLOCOCCAL PNEU NOS	486	PNEUMONIA, ORGANISM UNSPECIFIED
48241	METH SUS PNEUM D/T STAPH	5070	DUE TO INHALATION OF FOOD OR
			VOMITUS
48242	METH RES PNEU D/T STAPH	514	PULMONARY CONGESTION AND
			HYPOSTASIS

Exclude cases:

- with a principal ICD-9-CM diagnosis code for pneumonia (see above)
- with a principal ICD-9-CM diagnosis code for respiratory complications
- with any-listed ICD-9-CM diagnosis codes for viral pneumonia or influenza
- with any-listed ICD-9-CM diagnosis codes or any-listed ICD-9-CM procedure codes for

immunocompromised state

- with any-listed ICD-9-CM procedure codes for lung cancer
- MDC 4 (diseases/disorders of respiratory system)
- transferred to an acute care facility (DISP = 2)
- with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), or principal diagnosis (DX1=missing)

See Patient Safety Indicators Appendices:

• Appendix I – Immunocompromised State Diagnosis and Procedure Codes

ICD-9-CM Respiratory complications diagnosis codes:

9973	RESPIRATORY COMPLICATIONS (end 2008)	99732	POSTPROC ASPIRATION PNEU
99731	VENTLTR ASSOC PNEUMONIA	99739	RESPIRATORY COMP NEC

ICD-9-CM Viral pneumonia or influenza diagnosis codes:

ADENOVIRAL PNEUMONIA	4870	INFLUENZA W/ PNEUMONIA
RESP SYNCYT VIRAL PNEUM	4871	FLU W RESP MANIFEST NEC
PARAINFLUENZA VIRAL PNEUM	4878	FLU W MANIFESTATION NEC
PNEUMONIA DUE TO SARS	488	FLU D/T AVIAN FLU VIRUS (end 2009)
VIRAL PNEUMONIA NEC	4880	FLU D/T IDENTIF AVIAN FLU (end 2010)
VIRAL PNEUMONIA NOS	48801	FLU DT IDEN AVIAN W PNEU
PNEUMOCOCCAL PNEUMONIA	48802	FLU DT AVIAN W OTH RESP
PNEUMONIA ORGANISM NEC (end 1992)	48809	FLU DT AVIAN MANFEST NEC
PNEU MYCPLSM PNEUMONIAE	4881	FLU DT NOVEL H1N1 FLU (end 2010)
PNEUMONIA D/T CHLAMYDIA	48811	FLU DT 2009 H1N1 W PNEU
PNEUMON OTH SPEC ORGNSM	48812	FLU-2009 H1N1 W OTH RESP
PNEUM W CYTOMEG INCL DIS	48819	FLU-2009 H1N1 W OTH MAN
PNEUMONIA IN WHOOP COUGH	48881	FLU DT NVL A VRS W PNEU
PNEUMONIA IN ANTHRAX	48882	FLU DT NVL A W OTH RESP
PNEUMONIA IN ASPERGILLOSIS	48889	FLU DT NOVEL A W OTH MAN
PNEUMONIA IN OTH SYS MYCOSES		
PNEUMONIA IN INFECT DIS NEC		
	RESP SYNCYT VIRAL PNEUM PARAINFLUENZA VIRAL PNEUM PNEUMONIA DUE TO SARS VIRAL PNEUMONIA NEC VIRAL PNEUMONIA NOS PNEUMOCOCCAL PNEUMONIA PNEUMONIA ORGANISM NEC (end 1992) PNEU MYCPLSM PNEUMONIAE PNEUMONIA D/T CHLAMYDIA PNEUMON OTH SPEC ORGNSM PNEUM W CYTOMEG INCL DIS PNEUMONIA IN WHOOP COUGH PNEUMONIA IN ANTHRAX PNEUMONIA IN ASPERGILLOSIS PNEUMONIA IN OTH SYS MYCOSES	RESP SYNCYT VIRAL PNEUM 4871 PARAINFLUENZA VIRAL PNEUM 4878 PNEUMONIA DUE TO SARS 488 VIRAL PNEUMONIA NEC 4880 VIRAL PNEUMONIA NOS 48801 PNEUMOCOCCAL PNEUMONIA 48802 PNEUMONIA ORGANISM NEC (end 1992) 48809 PNEU MYCPLSM PNEUMONIAE 4881 PNEUMONIA D/T CHLAMYDIA 48811 PNEUMON OTH SPEC ORGNSM 48812 PNEUM W CYTOMEG INCL DIS 48819 PNEUMONIA IN WHOOP COUGH 48881 PNEUMONIA IN ANTHRAX 48882 PNEUMONIA IN ASPERGILLOSIS 48889 PNEUMONIA IN OTH SYS MYCOSES

ICD-9-CM Lung cancer procedure codes:

3230	THORACOSCOPIC SURGERY	3249	LOBECTOMY OF LUNG NEC
3239	OTH SEG LUNG RESECT NOS	3250	THORACOSCOPIC SURGERY
3241	THORACOSCOPIC SURGERY	3259	OTHER PNEUMONECTOMY NOS

Stratum 04C: SEPSIS

Surgical discharges, for patients ages 18 through 89 years or MDC 14 (pregnancy, childbirth, and puerperium), with all of the following:

- any-listed ICD-9-CM procedure codes for an operating room procedure; and
- the principal procedure occurring within 2 days of admission or an admission type of elective (ATYPE=3); and
- any secondary ICD-9-CM diagnosis codes for sepsis.

Surgical discharges are defined by specific DRG or MS-DRG codes.

See Patient Safety Indicators Appendices:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs
- Appendix E Surgical Discharge MS-DRGs

ICD-9-CM Sepsis diagnosis codes:

0380	STREP SEPTICEMIA	03842	ESCHERICHIA COLI
0381	STAPH SEPTICEMIA	03843	PSEUDOMONAS
03810	STAPH SEPTICEMIA, UNSPEC	03844	SERRATIA
03811	METH SUSC STAPH AUR SEPT	03849	SEPTICEM DT OTHER GRAM-NEG ORG
03812	MRSA SEPTICEMIA	0388	OTHER SPECIFIED SEPTICEMIAS
03819	OTHER STAPH SEPTICEMIA	0389	UNSPECIFIED SEPTICEMIA
0382	PNEUMOCL SEPTICEMIA	78552	SEPTIC SHOCK
0383	SEPTICEMIA DUE TO ANAEROBES	99591	SIRS DT INFECT WO ORGAN DYSFUN
03840	GRAM-NEGATIVE ORG, UNSPEC	99592	SIRS DT INFECT W ORGAN DYSFUN
03841	HEMOPHILUS INFLUENZAE	99802	POSTOPERATIVE SHOCK, SEPTIC

Exclude cases:

- with a principal ICD-9-CM diagnosis code for sepsis (see above)
- with a principal ICD-9-CM diagnosis code for infection or pressure ulcer
- with any-listed ICD-9-CM diagnosis codes or any-listed ICD-9-CM procedure codes for immunocompromised state
- with a length of stay of less than 4 days
- transferred to an acute care facility (DISP = 2)
- with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), or principal diagnosis (DX1=missing)

See Patient Safety Indicators Appendices:

- Appendix F Infection Diagnosis Codes
- Appendix I Immunocompromised State Diagnosis and Procedure Codes

ICD-9-CM pressure ulcer codes (for discharges prior to 2008)

7070	DECUBITUS ULCER	70704	PRESSURE ULCER, HIP
70700	PRESSURE ULCER, SITE NOS	70705	PRESSURE ULCER, BUTTOCK
70701	PRESSURE ULCER, ELBOW	70706	PRESSURE ULCER, ANKLE
70702	PRESSURE ULCER, UPR BACK	70707	PRESSURE ULCER, HEEL
70703	PRESSURE ULCER, LOW BACK	70709	PRESSURE ULCER, SITE NEC

CARDIAC ARREST

Surgical discharges, for patients ages 18 through 89 years or MDC 14 (pregnancy, childbirth, and puerperium), with all of the following:

- any-listed ICD-9-CM procedure codes for an operating room procedure; and
- the principal procedure occurring within 2 days of admission or an admission type of elective (ATYPE=3); and
- any secondary ICD-9-CM diagnosis codes or any-listed ICD-9-CM procedure codes for shock or cardiac arrest.

Surgical discharges are defined by specific DRG or MS-DRG codes.

See Patient Safety Indicators Appendices:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs
- Appendix E Surgical Discharge MS-DRGs

ICD-9-CM Shock or cardiac arrest diagnosis codes¹

4275	CARDIAC ARREST	4275	CARDIAC ARREST
63450	SPONTANEOUS ABORTION	63450	SPONTANEOUS ABORTION
	UNSPECIFIED COMPLICATED BY		UNSPECIFIED COMPLICATED BY
	SHOCK		SHOCK
63451	SPONTANEOUS ABORTION	63451	SPONTANEOUS ABORTION
	UNSPECIFIED COMPLICATED BY		UNSPECIFIED COMPLICATED BY
62452	SHOCK SPONTANEOUS ABORTION	63452	SHOCK SPONTANEOUS ABORTION
63452	UNSPECIFIED COMPLICATED BY	03432	UNSPECIFIED COMPLICATED BY
	SHOCK		SHOCK
63550	LEGALLY INDUCED ABORTION	63550	LEGALLY INDUCED ABORTION
	UNSPECIFIED COMPLICATED BY		UNSPECIFIED COMPLICATED BY
	SHOCK		SHOCK
63551	LEGALLY INDUCED ABORTION	63551	LEGALLY INDUCED ABORTION
	UNSPECIFIED COMPLICATED BY		UNSPECIFIED COMPLICATED BY
	SHOCK		SHOCK
63552	LEGALLY INDUCED ABORTION	63552	LEGALLY INDUCED ABORTION
	UNSPECIFIED COMPLICATED BY		UNSPECIFIED COMPLICATED BY
63650	SHOCK ILLEGAL ABORTION UNSPECIFIED	63650	SHOCK ILLEGAL ABORTION UNSPECIFIED
03030	COMPLICATED BY SHOCK	03030	COMPLICATED BY SHOCK
63651	ILLEGAL ABORTION UNSPECIFIED	63651	ILLEGAL ABORTION UNSPECIFIED
00001	COMPLICATED BY SHOCK	00001	COMPLICATED BY SHOCK
63652	ILLEGAL ABORTION UNSPECIFIED	63652	ILLEGAL ABORTION UNSPECIFIED
	COMPLICATED BY SHOCK		COMPLICATED BY SHOCK
63750	LEGALLY UNSPECIFIED TYPE OF	63750	LEGALLY UNSPECIFIED TYPE OF
	ABORTION UNSPECIFIED		ABORTION UNSPECIFIED
	COMPLICATED BY SHOCK		COMPLICATED BY SHOCK

ICD-9-CM Shock or cardiac arrest (resuscitation) procedure codes:

9393	NONMECHAN RESUSCITATION	9963	CLOSED CHEST CARD MASSAG
9960	CARDIOPULM RESUSCITA NOS		

Exclude cases:

- with a principal ICD-9-CM diagnosis code for shock or cardiac arrest (see above)
- with a principal ICD-9-CM diagnosis code for trauma
- with a principal ICD-9-CM diagnosis code for hemorrhage
- with a principal ICD-9-CM diagnosis code for gastrointestinal hemorrhage
- with a principal ICD-9-CM diagnosis code for abortion-related shock
- with any-listed ICD-9-CM procedure codes for lung cancer resection
- MDC 4 (diseases/disorders of respiratory system)
- MDC 5 (diseases/disorders of circulatory system)
- transferred to an acute care facility (DISP = 2)
- with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), or principal diagnosis (DX1=missing)

See Patient Safety Indicators Appendices:

• Appendix G – Trauma Diagnosis Codes

ICD-9-CM Hemorrhage diagnosis codes:

2851	AC POSTHEMORRHAG ANEMIA	9582	SECONDARY/RECUR HEMORR
4590	HEMORRHAGE NOS	99811	HEMORHAGE COMPLIC PROC
56881	HEMOPERITONEUM		

ICD-9-CM Gastrointestinal hemorrhage diagnosis codes:

ı	CD-3-C	ivi Gasti oli itestiliai nemorriage ulagnosi	is codes) i
	4560	ESOPH VARICES W BLEED	53361	PEPTIC ULC, SITE UNSPEC, CHR OR UNSPEC W HEM AND PERF W OBST
	45620	ESOPH VARICES IN DISEASES	53400	GASTROJEJ ULC, AC W HEM WO
	10020	CLASSIFIED ELSEWHERE W BLEED	00100	MENTION OF OBST
	5307	GASTROESOPH LACERATION- HEM	53401	GASTROJEJ ULC, AC W HEM W OBST
	5507	SYNDROME	33701	GACTICOLO DEC, AC W HEM W OBOT
	53021	ULC OF ESOPH W BLEED	53420	GASTROJEJ ULC, AC W HEM AND PERF
	33021	OLO OF LOOF IT W BELLED	33720	WO MENTION OF OBST
	53082	ESOPH HEM	53421	GASTROJEJ ULC, AC W HEM AND PERF
	33002	L301 TITILW	33421	W OBST
	53100	GAST ULC AC W HEM WO MENTION OF	53440	GASTROJEJ ULC, CHR OR UNSPEC W
	33100	OBST	33440	HEM WO MENTION OF OBST
	53101	GAST ULC AC W HEM W OBST	53441	
	55101	GAST ULC AC W HEIN W OBST	33441	GASTROJEJ ULC, CHR OR UNSPEC W HFM W OBST
	E2420	CACTURE AC WILLEM AND DEDE MO	E2460	020.
	53120	GAST ULC AC W HEM AND PERF WO MENTION OF OBST	53460	GASTROJEJ ULC, CHR OR UNSPEC W HEM AND PERF WO MENTION OF OBST
	E0404		E0.404	
	53121	GAST ULC, AC W HEM AND PERF W	53461	GASTROJEJ ULC, CHR OR UNSPEC W
	50440	OBST	50504	HEM AND PERF W OBST
	53140	GAST ULC CHR OR UNSPEC W HEM -	53501	GAST AND DUOD, AC GASTRITIS W
	50444	WO MENTION OF OBST	50544	HEM
	53141	GAST ULC CHR OR UNSPEC W HEM W	53511	GAST AND DUOD, ATROPHIC
	5 0400	OBST	50504	GASTRITIS W HEM
	53160	GAST ULC CHR OR UNSPEC W HEM	53521	GAST AND DUOD, GAST MUCOSAL
	50404	AND PERF WO MENTION OF OBST	50504	HYPERTROPHY, W HEM
	53161	GAST ULC CHR OR UNSPEC W HEM	53531	GAST AND DUOD, ALCOHOLIC
	=0000	AND PERF W OBST	50544	GASTRITIS, W HEM
	53200	DUODENAL ULC AC W HEM WO	53541	GAST AND DUOD, OTHER SPECIFIED
	50004	MENTION OF OBST	50554	GASTRITIS W HEM
	53201	DUODENAL ULC AC W HEM W OBST	53551	GAST AND DUOD, UNSPEC GASTRITIS
	50000	DUODENAL LILO AO WULEMAND DEDE	=0=0 <i>:</i>	AND GASTDUODENITIS W HEM
	53220	DUODENAL ULC AC W HEM AND PERF WO MENTION OF OBST	53561	GAST AND DUOD, DUODENITIS W HEM

AHRQ QI™ Research Version 5.0, Patient Safety Indicators 04, Technical Specifications, Death Rate among Surgical Inpatients with Serious Treatable Complications www.qualityindicators.ahrq.gov

53221	DUODENAL ULC AC W HEM AND PERF W OBST	53783	OTH SPEC DIS STOM AND DUOD, ANGIODYSPLASIA W HEM
53240	DUODENAL ULC CHR OR UNSPEC W HEM WO MENTION OF OBST	53784	DIEULAFOY LESION (HEMORR) OF STOMACH AND DUODENUM
53241	DUODENAL ULC CHR OR UNSPEC W HEM W OBST	56202	DIVERTICULOSIS OF SMALL INTESTINE -W HEM
53260	DUODENAL ULC CHR OR UNSPEC W HEM AND PERF WO MENTION OF OBST	56203	DIVERTICULITIS OF SMALL INTESTINE - W HEM
53261	DUODENAL ULC CHR OR UNSPEC W HEM AND PERF W OBST	56212	DIVERTICULOSIS OF COLON W HEM
53300	PEPTIC ULC, SITE UNSPEC AC W HEM WO MENTION OF OBST	56213	DIVERTICULITIS OF COLON W HEM
53301	PEPTIC ULC, SITE UNSPEC, AC W HEM W OBST	5693	HEM OF RECTUM AND ANUS
53320	PEPTIC ULC, SITE UNSPEC, AC W HEM AND PERF WO MENTION OF OBST	56985	ANGIODYSPLASIA OF INTESTINE W HEM
53321	PEPTIC ULC, SITE UNSPEC, AC W HEM AND PERF W OBST	56986	DIEULAFOY LESION (HEMORRHAGIC) OF INTESTINE
53340	PEPTIC ULC, SITE UNSPEC CHR OR UNSPEC W HEM WO MENTION OF OBST	5780	GI HEM, HEMATEMESIS
53341	PEPTIC ULC, SITE UNSPEC, CHR OR UNSPEC W HEM W OBST	5781	GI HEM, BLOOD IN STOOL
53360	PEPTIC ULC, SITE UNSPEC, CHR OR UNSPEC W HEM AND PERF -WO MENTION OF OBST	5789	GI HEM, HEM OF GI TRACT, UNSPEC

ICD-9-CM Abortion-related shock diagnosis codes:

63450	SPON ABORT W SHOCK-UNSP	63651	ILLEG ABORT W SHOCK-INC
63451	SPON ABORT W SHOCK-INC	63652	ILLEG ABORT W SHOCK-COMP
63452	SPON ABORT W SHOCK-COMP	63750	ABORT NOS W SHOCK-UNSO
63550	LEGAL ABORT W SHOCK-UNSO	63751	ABORT NOS W SHOCK-INC
63551	LEGAL ABORT W SHOCK-INC	63752	ABORT NOS W SHOCK-COMP
63552	LEGAL ABORT W SHOCK-COMP	6385	ATTEM ABORTION W SHOCK
63650	ILLEG AB W SHOCK-UNSO		

ICD-9-CM Lung cancer resection procedure codes:

3239	OTH SEG LUNG RESECT NOS	3259	OTHER PNEUMONECTOMY NOS
3249	LOBECTOMY OF LUNG NEC		

Stratum 04E: GASTROINTESTINAL (GI) HEMORRHAGE / ACUTE ULCER

Surgical discharges, for patients ages 18 through 89 years or MDC 14 (pregnancy, childbirth, and puerperium), with all of the following:

- any-listed ICD-9-CM procedure codes for an operating room procedure; and
- the principal procedure occurring within 2 days of admission or an admission type of elective (ATYPE=3); and
- any secondary ICD-9-CM diagnosis codes for gastrointestinal hemorrhage or acute ulcer. Surgical discharges are defined by specific DRG or MS-DRG codes.

See Patient Safety Indicators Appendices:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs

• Appendix E – Surgical Discharge MS-DRGs

ICD-9-CM Gastrointestinal	hemorrhage	or acute ulcer	diagnosis codes:
100 0 Oili Casti Cilitostiliai	monitorinage	or acate areer	alagilosis coacs.

4560	ESOPHAGEAL VARICES WITH	53390	
4000	BLEEDING	00000	ACUTE OR CHRONIC, WITHOUT
			MENTION OF HEMORRHAGE OR
			PERFORATION - WITHOUT MENTION OF
			OBSTRUCTION
45620	ESOPHAGEAL VARICES IN DISEASES	53391	SITE UNSPECIFIED UNSPECIFIED AS
	CLASSIFIED ELSEWHERE WITH		ACUTE OR CHRONIC, WITHOUT
	BLEEDING		MENTION OF HEMORRHAGE OR
			PERFORATION - WITH OBSTRUCTION
5307	GASTROESOPHAGEAL LACERATION-	53400	ACUTE WITH HEMORRHAGE -WITHOUT
	HEMORRHAGE SYNDROME		MENTION OF OBSTRUCTION
53082	ESOPHAGEAL HEMORRHAGE	53401	ACUTE WITH HEMORRHAGE - WITH
			OBSTRUCTION
53100	ACUTE WITH HEMORRHAGE - WITHOUT	53410	ACUTE WITH PERFORATION - WITHOUT
	MENTION OF OBSTRUCTION		MENTION OF OBSTRUCTION
53101	ACUTE WITH HEMORRHAGE - WITH	53411	ACUTE WITH PERFORATION - WITH
	OBSTRUCTION		OBSTRUCTION
53110	ACUTE WITH PERFORATION - WITHOUT	53420	ACUTE WITH HEMORRHAGE AND
	MENTION OF OBSTRUCTION		PERFORATION - WITHOUT MENTION OF
			OBSTRUCTION
53111	ACUTE WITH PERFORATION - WITH	53421	ACUTE WITH HEMORRHAGE AND
	OBSTRUCTION		PERFORATION - WITH OBSTRUCTION
53120	ACUTE WITH HEMORRHAGE AND	53430	ACUTE WITHOUT MENTION OF
	PERFORATION - WITHOUT MENTION OF		HEMORRHAGE OR PERFORATION -
	OBSTRUCTION		WITHOUT MENTION OF OBSTRUCTION
53121	ACUTE WITH HEMORRHAGE AND	53431	ACUTE WITHOUT MENTION OF
	PERFORATION - WITH OBSTRUCTION		HEMORRHAGE OR PERFORATION -
			WITH OBSTRUCTION
53130	ACUTE WITHOUT MENTION OF	53490	UNSPECIFIED AS ACUTE OR CHRONIC,
	HEMORRHAGE OR PERFORATION -		WITHOUT MENTION OF HEMORRHAGE
	WITHOUT MENTION OF OBSTRUCTION		OR PERFORATION - WITHOUT MENTION
			OF OBSTRUCTION
53131	ACUTE WITHOUT MENTION OF	53491	UNSPECIFIED AS ACUTE OR CHRONIC,
	HEMORRHAGE OR PERFORATION -		WITHOUT MENTION OF HEMORRHAGE
	WITH OBSTRUCTION		OR PERFORATION - WITH
E2400	LINEDECIFIED AS ACUTE OF CURONIC	E0E04	OBSTRUCTION WITH
53190	UNSPECIFIED AS ACUTE OR CHRONIC, WITHOUT MENTION OF HEMORRHAGE	53501	ACUTE GASTRITIS - WITH
			HEMORRHAGE
	OR PERFORATION - WITHOUT MENTION OF OBSTRUCTION		
53191	UNSPECIFIED AS ACUTE OR CHRONIC,	53511	ATROPHIC GASTRITIS - WITH
55191	WITHOUT MENTION OF HEMORRHAGE	55511	HEMORRHAGE
	OR PERFORATION - WITH		HEWORKHAGE
	OBSTRUCTION		
53200	ACUTE WITH HEMORRHAGE - WITHOUT	53521	GASTRIC MUCOSAL HYPERTROPHY -
33200	MENTION OF OBSTRUCTION	33321	WITH HEMORRHAGE
53201	ACUTE WITH HEMORRHAGE - WITH	53531	ALCOHOLIC GASTRITIS - WITH
33201	OBSTRUCTION	JJJJ 1	HEMORRHAGE
53210	ACUTE WITH PERFORATION - WITHOUT	53541	OTHER SPECIFIED GASTRITIS - WITH
00210	MENTION OF OBSTRUCTION	000-1	HEMORRHAGE
53211	ACUTE WITH PERFORATION - WITH	53551	UNSPECIFIED GASTRITIS AND
00211	OBSTRUCTION	00001	GASTRODUODENITIS - WITH
			HEMORRHAGE
53220	ACUTE WITH HEMORRHAGE AND	53561	DUODENITIS - WITH HEMORRHAGE
	PERFORATION - WITHOUT MENTION OF		
	OBSTRUCTION		
53221	ACUTE WITH HEMORRHAGE AND	53783	ANGIODYSPLASIA OF STOMACH AND
	PERFORATION - WITH OBSTRUCTION		DUODENUM WITH HEMORRHAGE
53230	ACUTE WITHOUT MENTION OF	53784	DIEULAFOY LESION (HEMORRHAGIC)
			,

	HEMORRHAGE OR PERFORATION -		OF STOMACH AND DUODENUM
53231	WTHOUT MENTION OF OBSTRUCTION ACUTE WITHOUT MENTION OF HEMORRHAGE OR PERFORATION - WITH OBSTRUCTION	56202	DIVERTICULOSIS OF SMALL INTESTINE WITH HEMORRHAGE
53290	UNSPECIFIED AS ACUTE OR CHRONIC, WITHOUT MENTION OF HEMORRHAGE OR PERFORATION - WITHOUT MENTION	56203	DIVERTICULITIS OF SMALL INTESTINE WITH HEMORRHAGE
53291	OF OBSTRUCTION UNSPECIFIED AS ACUTE OR CHRONIC, WITHOUT MENTION OF HEMORRHAGE OR PERFORATION - WITH OBSTRUCTION	56212	DIVERTICULOSIS OF COLON WITH HEMORRHAGE
53300	SITE UNSPECIFIED ACUTE WITH HEMORRHAGE - WITHOUT MENTION OF OBSTRUCTION	56213	DIVERICULITIS OF COLON WITH HEMORRHAGE
53301	SITE UNSPECIFIED ACUTE WITH HEMORRHAGE - WITH OBSTRUCTION	5693	HEMORRHAGE OF RECTUM AND ANUS
53310	SITE UNSPECIFIED ACUTE WITH PERFORATION - WITHOUT MENTION OF OBSTRUCTION	56985	ANGIODYSPLASIA OF INTESTINE WITH HEMORRHAGE
53311	SITE UNSPECIFIED ACUTE WITH - PERFORATION WITH OBSTRUCTION	56986	DIEULAFOY LESION (HEMORRHAGIC) OF INTESTINE
53320	SITE UNSPECIFIED ACTE WITH HEMORRHAGE AND PERFORATION - WITHOUT MENTION OF OBSTRUCTION	5780	HEMATEMESIS
53321	SITE UNSPECIFIED ACUTE WITH HEMORRHAGE AND PERFORATION - WITHOUT MENTION OF OBSTRUCTION	5781	BLOOD IN STOOL
53330	SITE UNSPECIFIED ACUTE WITHOUT MENTION OF HEMORRHAGE AND PERFFORATION - WITHOUT MENTION OF OBSTRUCTION	5789	HEMORRHAGE OF GASTROINTESTINAL TRACT, UNSPECIFIED
53331	SITE UNSPECIFIED ACUTE WITHOUT MENTION OF HEMORRHAGE AND PERFORATION - WITH OBSTRUCTION		

Exclude cases:

- with a principal ICD-9-CM diagnosis code for gastrointestinal hemorrhage or acute ulcer (see above)
- with a principal ICD-9-CM diagnosis code for trauma
- with a principal ICD-9-CM diagnosis code for alcoholism
- with a principal ICD-9-CM diagnosis code for anemia
- MDC 6 (diseases and disorders of the digestive system)
- MDC 7 (diseases and disorders of the hepatobiliary system and pancreas)
- transferred to an acute care facility (DISP = 2)
- with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), or principal diagnosis (DX1=missing)

See Patient Safety Indicators Appendices:

• Appendix G – Trauma Diagnosis Codes

ICD-9-CM Alcoholism diagnosis codes:

2910	DELERIUM TREMENS	30392	ALCOH DEP NEC/NOS-EPISOD
2911	ALCOHOL AMNESTIC DISORDR	30393	ALCOH DEP NEC/NOS-REMISS

AHRQ QI™ Research Version 5.0, Patient Safety Indicators 04, Technical Specifications, Death Rate among Surgical Inpatients with Serious Treatable Complications www.qualityindicators.ahrq.gov

2912	ALCOHOL PERSIST DEMENTIA	30500	ALCOHOL ABUSE-UNSPEC
2913	ALCOH PSY DIS W HALLUCIN	30501	ALCOHOL ABUSE-CONTINUOUS
2914	PATHOLOGIC ALCOHOL INTOX	30502	ALCOHOL ABUSE-EPISODIC
2915	ALCOH PSYCH DIS W DELUS	30503	ALCOHOL ABUSE-IN REMISS
29181	ALCOHOL WITHDRAWAL	4255	ALCOHOLIC CARDIOMYOPATHY
29182	ALCOH INDUCE SLEEP DISOR	53530	ALCHL GASTRTIS W/O HMRHG
29189	ALCOHOL MENTAL DISOR NEC	53531	ALCHL GSTRITIS W HMRHG
2919	ALCOHOL MENTAL DISOR NOS	5710	ALCOHOLIC FATTY LIVER
30300	AC ALCOHOL INTOX-UNSPEC	5711	AC ALCOHOLIC HEPATITIS
30301	AC ALCOHOL INTOX-CONTIN	5712	ALCOHOL CIRRHOSIS LIVER
30302	AC ALCOHOL INTOX-EPISOD	5713	ALCOHOL LIVER DAMAGE NOS
30303	AC ALCOHOL INTOX-REMISS	9800	TOXIC EFF EHTYL ALCOHOL
30390	ALCOH DEP NEX/NOS-UNSPEC	9809	TOXIC EFFECT ALCOHOL NOS

ICD-9-CM Anemia diagnosis codes: 2800 CHR BLOOD LOSS ANEMIA

2800 CHR BLOOD LOSS ANEMIA 2851 AC POSTHEMORRHAG ANEMIA