expert™ Imaging & Path Lab
Accuracy Matters....

Delhi Pollution Control Committee | An ISO Certified Lab | Government Of India Atomic Energy Regulatory Board Radiological Safety Division

Pt's Name	Mrs. ANJALI	Date of Sample	26/10/2024
Age / Sex	25 YrsF	Date of Report	26/10/2024
Referred By	SELF	Printing Time	18:29:07
Panel ID	ROHIT_PRO	Access ID	58
Barcode ID		Patient ID	2410260056

Test Description	Result	Unit	Biological Ref.Renge

TYPHIDOT (IgG & IgM)

Method : Immuno chromatography

DOT EIA FOR DECTECTION OF IgM AND IgG ANTIBODIES TO SALMONELLA TYPHI

TYPHIDOT - IgM	**POSITIVE**
TYPHIDOT - IgG	**POSITIVE**

COMMENT:- IgM Positive- Acute Typhoid fever.Typhi DOT has high sensitivity and specificity (95%) as compared to Blood Culture,Stool Culture and Widal test. It is useful for early, definite diagnosis of Typhoid (IgM levels in patient are usually detected by typhidot at the onset of fever).Detection of S. typhi specific IgM antibodies serves as a marker for early detection of recent infection.However the test should be used as a screening test in clinically suspected cases only,and its result should be confirmed by other supplemental method before taking clinical decisions.In some studies it has been reported that low titer IgM antibodies to S.typhi persist for about 4 months post infection.Typhidot is done on a dot ELISA kit that detects IgM and IgG antibodies against the outer membrane protein (OMP) of the Salmonella typhi. The Typhidot test is expected to become positive within 2-3 days of infection.The test is based on the presence of specific IgM and IgG antibodies. IgM shows recent infection whereas IgG signifies remote infection. Typhidot was 67% sensitive and 54% specific, with 85% positive and 81% NPVs.

Sony Singh.

DR. SONY SINGH
MBBS, MD , Clinical Pathologist
Microbiologist Regd No. 85520

REDEFINING QUALITY STANDARDS IN LABORATORY MEDICINE

- If the test results are unexpected please contact the Lab.
- This is only a professional opinion and not the diagnosis it should be clinically correlated.
- This report is for perusal of Doctors only not for medico-legal purpose.

Monday to Saturday : Timings : 8.00 a.m. to 9.00 p.m. (Sunday : 8.00 am to 4.00 pm)

ABCB ACCREDITED

expert™ Imaging & Path Lab
Accuracy Matters....

Delhi Pollution Control Committee | An ISO Certified Lab | Government Of India Atomic Energy Regulatory Board Radiological Safety Division

Pt's Name	Mrs. ANJALI	Date of Sample	26/10/2024
Age / Sex	25 YrsF	Date of Report	26/10/2024
Referred By	SELF	Printing Time	18:29:07
Panel ID	ROHIT_PRO	Access ID	58
Barcode ID		Patient ID	2410260056

Test Description	Result	Unit	Biological Ref.Renge
NS-1 ANTIGEN DENGUE (Method: Immuno chromatography)	NON REACTIVE		NON-REACTIVE

REMARKS :-
In tropical and subtropical regions, dengue is the most important viruses in terms of mortality and morality. An easy-to-perform, rapid and specific test is needed to confirm infection during the acute phase of the infection in order to implement appropriate treatment during early course of the disease. NS1 antigen is an non-structural protein recognized as a marker of acute phase of dengue infection, a period for which traditional serological antibodies based methods are of limited value. NS1 antigen was found circulating in sample of infected patients from the first day up to 9 days after onset of fever (1,2 & 3)Dengue NS1 Antigen Strip is an individual test for qualitative detection of Dengue virus NS1 antigen in human serum or plasma as an aid in the diagnosis of acute dengue infection.

Sony Singh.
DR. SONY SINGH
MBBS, MD , Clinical Pathologist
Microbiologist Regd No. 85520

REDEFINING QUALITY STANDARDS IN LABORATORY MEDICINE
- If the test results are unexpected please contact the Lab.
- This is only a professional opinion and not the diagnosis it should be clinically correlated.
- This report is for perusal of Doctors only not for medico-legal purpose.
Monday to Saturday : Timings : 8.00 a.m. to 9.00 p.m. (Sunday : 8.00 am to 4.00 pm)
ABCB ACCREDITED