

2022 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: Developed by the task force for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death of the European Society of Cardiology (ESC) Endorsed by the Association for European Paediatric and Congenital Cardiology (AEPC) *Free*

Katja Zeppenfeld

(Chairperson)

(Netherlands)

Corresponding authors: Katja Zeppenfeld, Department of Cardiology, Leiden University Medical Centre, Albinusdreef 2, 2333 ZA Leiden, Netherlands. Tel +31 715262020, E-mail: K.Zeppenfeld@LUMC.nl

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Jacob Tfelt-Hansen

(Chairperson)

(Denmark)

Jacob Tfelt-Hansen, The Department of Cardiology, The Heart Centre, Copenhagen University Hospital, Rigshospitalet, Blegdamsvej 9, 2100, Copenhagen, Denmark. Tel +45 61360399, E-mail: jacob.tfelt@regionh.dk



<https://orcid.org/0000-0003-3895-9316>

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Marta de Riva

(Task Force Coordinator)

(Netherlands)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Bo Gregers Winkel

(Task Force Coordinator)

(Denmark)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Elijah R Behr

(United Kingdom)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Nico A Blom

(Netherlands)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Philippe Charron

(France)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Domenico Corrado

(Italy)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Nikolaos Dagres

(Germany)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Christian de Chillou

(France)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

... [Show more](#)

[Lars Eckardt](#)

Lars Eckardt

(Germany)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Tim Friede](#)

Tim Friede

(Germany)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Kristina H Haugaa](#)

Kristina H Haugaa

(Norway)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Mélèze Hocini](#)

Mélèze Hocini

(France)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Pier D Lambiase](#)

Pier D Lambiase

(United Kingdom)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Eloi Marijon](#)

Eloi Marijon

(France)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Jose L Merino](#)

Jose L Merino

(Spain)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Petr Peichl](#)

Petr Peichl

(Czech Republic)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Silvia G Priori](#)

Silvia G Priori

(Italy)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Tobias Reichlin](#)

Tobias Reichlin

(Switzerland)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Jeanette Schulz-Menger](#)

Jeanette Schulz-Menger

(Germany)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Christian Sticherling](#)

Christian Sticherling

(Switzerland)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Stylios Tzeis](#)

Stylios Tzeis

(Greece)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Axel Verstrael](#)

Axel Verstrael

(Belgium)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[Maurizio Volterrani](#)

Maurizio Volterrani

(Italy)

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

[ESC Scientific Document Group](#)

ESC Scientific Document Group

Search for other works by this author on:

[Oxford Academic](#) [PubMed](#) [Google Scholar](#)

Katja Zeppenfeld and Jacob Tfelt-Hansen chairpersons contributed equally to the document and are joint corresponding authors.

Marta de Riva and Bo Gregers Winkel task force Coordinators contributed equally to the document.

Author/task force Member affiliations are listed in Author information.

Representing the Association for European Paediatric and Congenital Cardiology (AEPC).

ESC Clinical Practice Guidelines (CPG) Committee: listed in the Appendix.

ESC subspecialty communities having participated in the development of this document:

Associations: Association for Acute CardioVascular Care (ACVC), Association of Cardiovascular Nursing & Allied Professions (ACNAP), European Association of Cardiovascular Imaging (EACVI), European Association of Percutaneous Cardiovascular Interventions (EAPCI), European Heart Rhythm Association (EHRA), Heart Failure Association (HFA).

Working Groups: Cardiac Cellular Electrophysiology, Myocardial and Pericardial Diseases.

Patient Forum

The content of these European Society of Cardiology (ESC) Guidelines has been published for personal and educational use only. No commercial use is authorized. No part of the ESC Guidelines may be translated or reproduced in any form without written permission from the ESC. Permission can be obtained upon submission of a written request to Oxford University Press, the publisher of the *European Heart Journal* and the party authorized to handle such permissions on behalf of the ESC (Journals.Permissions@oup.com).

Disclaimer: The ESC Guidelines represent the views of the ESC and were produced after careful consideration of the scientific and medical knowledge and the evidence available at the time of their publication. The ESC is not responsible in the event of any contradiction, discrepancy, and/or ambiguity between the ESC Guidelines and any other official recommendations or guidelines issued by the relevant public health authorities, in particular in relation to good use of healthcare or therapeutic strategies. Health professionals are encouraged to take the ESC Guidelines fully into account when exercising their clinical judgment, as well as in the determination and the implementation of preventive, diagnostic, or therapeutic medical strategies; however, the ESC Guidelines do not override, in any way whatsoever, the individual responsibility of health professionals to make appropriate and accurate decisions in consideration of each patient's health condition and in consultation with that patient and, where appropriate and/or necessary, the patient's caregiver. Nor do the ESC Guidelines exempt health professionals from taking into full and careful consideration the relevant official updated recommendations or guidelines issued by the competent public health authorities, in order to manage each patient's case in light of the scientifically accepted data pursuant to their respective ethical and professional obligations. It is also the health professional's responsibility to verify the applicable rules and regulations relating to drugs and medical devices at the time of prescription.

© The European Society of Cardiology 2022. All rights reserved. For permissions please e-mail:

Journals.Permissions@oup.com

All experts involved in the development of these guidelines have submitted declarations of interest. These have been compiled in a report and published in a supplementary document simultaneously to the guidelines. The report is also available on the ESC website www.escardio.org/Guidelines

[Author Notes](#)

European Heart Journal, Volume 43, Issue 40, 21 October 2022, Pages 3997a–4126,
<https://doi.org/10.1093/eurheartj/ehac262>

Published:

26 August 2022

[Guidelines](#), [Anti-arrhythmic drugs](#), [Cardiac magnetic resonance](#), [Cardiomyopathies](#), [Catheter ablation](#), [Chronic coronary artery disease](#), [Genetic testing](#), [Implantable cardioverter defibrillator](#), [Premature ventricular complex](#), [Primary electrical disease](#), [Recommendations](#), [Risk calculator](#), [Risk stratification](#), [Sudden cardiac death](#), [Sudden death](#), [Ventricular arrhythmia](#), [Ventricular fibrillation](#), [Ventricular tachycardia](#)

 See the *European Heart Journal* online for supplementary data that includes background information and detailed discussion of the data that have provided the basis of the guidelines.



Click here to access the corresponding ESC CardioMed chapters.

Table of contents

- 1. Preamble [4003](#)
- 2. Introduction [4005](#)
- [2.1. What is new](#) [4005](#)
- 3. Definitions [4011](#)
- [3.1. Ventricular arrhythmia subtypes](#) [4011](#)
- [3.2. Sudden cardiac death](#) [4011](#)
- [3.3. Syncope](#) [4011](#)
- [3.4. Specialized centres](#) [4011](#)
- [3.5. Genetics](#) [4011](#)
- 4. Epidemiology of sudden cardiac death, public awareness, and risk stratification [4011](#)
- [4.1. Incidence of sudden cardiac death](#) [4011](#)
- [4.2. Causes of sudden cardiac death in different age groups](#) [4012](#)
- [4.3. Population vs. individual risk prediction](#) [4012](#)
- [4.4. Risk calculators for sudden cardiac death and review of the methodology](#) [4012](#)
- [4.5. Awareness and intervention: public basic life support, and access to automated external defibrillators](#) [4012](#)
- 5. Evaluation and treatment. General aspects [4014](#)
- [5.1. Diagnostic tools](#) [4014](#)
- [5.1.1. History and physical examination](#) [4014](#)
- [5.1.2. Laboratory testing](#) [4014](#)
- [5.1.3. Non-invasive and invasive tests](#) [4014](#)
- [5.1.3.1. Electrocardiogram and ambulatory electrocardiographic monitoring](#) [4014](#)
- [5.1.3.2. Signal-averaged electrocardiogram](#) [4014](#)
- [5.1.3.3. Exercise testing](#) [4014](#)
- [5.1.3.4. Imaging](#) [4014](#)
- [5.1.3.5. Provocative diagnostic tests](#) [4015](#)
- [5.1.3.6. Electrophysiological study](#) [4015](#)
- [5.1.4. Genetic testing](#) [4015](#)

- 5.2. Diagnostic evaluation at first presentation with ventricular arrhythmia in patients without known cardiac disease 4018
- 5.2.1. Scenario 1: Incidental finding of a non-sustained ventricular tachycardia 4018
- 5.2.2. Scenario 2: First presentation of sustained monomorphic ventricular tachycardia 4018
- 5.2.3. Scenario 3: Sudden cardiac arrest survivor 4020
- 5.2.4. Scenario 4: Sudden death victim 4023
- 5.2.5. Scenario 5: Relatives of sudden arrhythmic death syndrome decedents 4024
- 6. Therapies for ventricular arrhythmias. General aspects 4027
- 6.1. Acute management 4027
- 6.1.1. Treatment of reversible causes 4027
- 6.1.2. Acute management of sustained monomorphic ventricular tachycardia 4029
- 6.1.3. Management of electrical storm and incessant ventricular tachycardia 4030
- 6.2. Long-term management 4034
- 6.2.1. Pharmacotherapy 4034
- 6.2.2. Device therapy 4038
- 6.2.2.1. Implantable cardioverter defibrillator. 4038
- 6.2.2.2. Adding cardiac resynchronization therapy 4040
- 6.2.2.3. Wearable cardioverter defibrillator. 4040
- 6.2.3. Special aspects of device therapy 4041
- 6.2.3.1. Optimization of device programming 4041
- 6.2.3.2. Concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy 4041
- 6.2.3.3. Psychosocial impact of implantable cardioverter defibrillator treatment 4042
- 6.2.3.4. Patients with left ventricular assist devices 4042
- 6.2.3.5. Complications of devices 4042
- 6.2.3.6. End-of-life issues 4043
- 6.2.4. Interventional therapy 4043
- 6.2.4.1. Catheter ablation 4043
- 6.2.4.1.1. Patients with structural heart disease 4043
- 6.2.4.1.2. Patients without apparent structural heart disease 4044
- 6.2.4.2. Autonomic modulation 4044
- 7. Diagnostic evaluation, management, and risk stratification according to clinical presentation and known (likely) disease 4044
- 7.1. Specific structural heart diseases 4044
- 7.1.1. Coronary artery disease 4044
- 7.1.1.1. Acute coronary syndromes and vasospasm 4044
- 7.1.1.1.1. Acute coronary syndromes 4044

- [7.1.1.1.2. Vasospasm](#)4045
- [7.1.1.2. Early after myocardial infarction](#)4046
- [7.1.1.3. Chronic coronary artery disease](#)4046
- [7.1.1.3.1. Primary prevention of sudden cardiac death in patients with reduced ejection fraction](#)4046
- [7.1.1.3.2. Primary prevention of sudden cardiac death in patients with preserved or mildly reduced ejection fraction](#)4047
- [7.1.1.3.3. Secondary prevention of sudden cardiac death](#)4047
- [7.1.1.3.4. Management of patients with haemodynamically tolerated ventricular tachycardia and preserved and mildly reduced ejection fraction](#)4048
- [7.1.1.3.5. Management of recurrent ventricular tachycardia in implantable cardioverter defibrillator carriers](#)4049
- [7.1.1.4. Coronary anomalies](#)4049
- [7.1.2. Idiopathic premature ventricular complexes/ventricular tachycardia and premature ventricular complex-induced cardiomyopathy](#)4050
- [7.1.2.1. Idiopathic premature ventricular complexes/ventricular tachycardia](#)4050
- [7.1.2.2. Premature ventricular complex-induced/-aggravated cardiomyopathy](#)4052
- [7.1.3. Cardiomyopathies](#)4054
- [7.1.3.1. Dilated cardiomyopathy and hypokinetic nondilated cardiomyopathy](#)4054
- [7.1.3.1.1. Diagnostic evaluation and risk stratification..](#) 4054
- [7.1.3.1.2. Primary prevention of sudden cardiac death](#) 4055
- [7.1.3.1.3. Secondary prevention of sudden cardiac death and management of ventricular arrhythmias](#)4057
- [7.1.3.2. Arrhythmogenic right ventricular cardiomyopathy](#)4058
- [7.1.3.2.1. Risk stratification](#)4058
- [7.1.3.2.2. Treatment](#)4060
- [7.1.3.3. Hypertrophic cardiomyopathy](#)4060
- [7.1.3.3.1. Risk stratification and primary prevention of sudden cardiac death](#)4061
- [7.1.3.3.2. Treatment to prevent ventricular arrhythmia recurrence](#)4062
- [7.1.3.4. Left ventricular non-compaction](#)4063
- [7.1.3.5. Restrictive cardiomyopathy](#)4063
- [7.1.3.6. Neuromuscular disorders](#)4063
- [7.1.4. Inflammatory cardiac diseases](#)4066
- [7.1.4.1. Myocarditis](#)4066
- [7.1.4.2. Cardiac sarcoidosis](#)4067
- [7.1.4.3. Chagas](#) [cardiomyopathy](#)4068
- [7.1.5. Valvular heart disease](#)4070
- [7.1.6. Congenital heart disease](#)4071

- 7.2. Primary electrical disease 4073
- 7.2.1. Idiopathic ventricular fibrillation 4073
- 7.2.2. Long QT syndrome (including acquired long QT syndrome) 4074
- 7.2.3. Andersen-Tawil syndrome Type 1a 4079
- 7.2.4. Brugada syndrome 4079
- 7.2.5. Early repolarization syndromes 4081
- 7.2.6. Catecholaminergic polymorphic ventricular tachycardia 4084
- 7.2.7. Short QT syndrome 4086
- 8. Special aspects in selected populations 4088
- 8.1. Pregnant patients and peri-partum cardiomyopathy 4088
- 8.1.1. Electrocardioversion and implantable cardioverter defibrillator therapy in pregnancy 4088
- 8.1.2. Pharmacological treatment 4088
- 8.1.3. Catheter ablation 4088
- 8.2. Heart transplantation 4089
- 8.3. Sudden cardiac death in athletes 4089
- 8.4. Wolff-Parkinson-White syndrome 4090
- 8.5. Prevention of sudden cardiac death in the elderly 4090
- 9. Key messages 4090
- 9.1. General aspects 4090
- 9.2. Structural heart disease 4090
- 9.3. Primary electrical disease 4091
- 10. Gaps in evidence 4091
- 10.1. General aspects 4091
- 10.2. Structural heart disease – general 4091
- 10.3. Idiopathic premature ventricular complexes/ventricular tachycardia 4091
- 10.4. Coronary artery disease 4091
- 10.5. Cardiomyopathies 4091
- 10.6. Valvular heart disease 4091
- 10.7. Congenital heart disease 4091
- 10.8. Primary electrical disease 4091
- 11. What to do and what not to do – messages from the Guidelines 4092
- 12. Quality indicators 4099
- 13. Supplementary data 4099
- 14. Data availability statement 4099
- 15. Author information 4099
- 16. Appendix 4100

- 17. Referencesâ€¦â€¦â€¦4101

Tables of Recommendations

- Recommendation Table 1â€¦â€¦â€¦Recommendations for public basic life support and access to automated external defibrillatorsâ€¦â€¦â€¦4014
- Recommendation Table 2â€¦â€¦â€¦Recommendations for genetic testingâ€¦â€¦â€¦4018
- Recommendation Table 3â€¦â€¦â€¦Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmiaâ€¦â€¦â€¦4018
- Recommendation Table 4â€¦â€¦â€¦Recommendations for evaluation of patients presenting with a first episode of sustained monomorphic ventricular tachycardiaâ€¦â€¦â€¦4020
- Recommendation Table 5â€¦â€¦â€¦Recommendations for evaluation of sudden cardiac arrest survivorsâ€¦â€¦â€¦4022
- Recommendation Table 6â€¦â€¦â€¦Recommendations for evaluation of sudden death victimsâ€¦â€¦â€¦4023
- Recommendation Table 7â€¦â€¦â€¦Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedentsâ€¦â€¦â€¦4027
- Recommendation Table 8â€¦â€¦â€¦Recommendations for treatment of reversible conditionsâ€¦â€¦â€¦4029
- Recommendation Table 9â€¦â€¦â€¦Recommendations for the acute management of sustained ventricular tachycardia and electrical stormâ€¦â€¦â€¦4033
- Recommendation Table 10â€¦â€¦â€¦Recommendations for treatment with heart failure medicationâ€¦â€¦â€¦4034
- Recommendation Table 11â€¦â€¦â€¦Recommendations for implantable cardioverter defibrillator implantation (general aspects)â€¦â€¦â€¦4040
- Recommendation Table 12â€¦â€¦â€¦Recommendations for secondary prevention of sudden cardiac deathâ€¦â€¦â€¦4040
- Recommendation Table 13â€¦â€¦â€¦Recommendations for subcutaneous implantable cardioverter defibrillatorâ€¦â€¦â€¦4040
- Recommendation Table 14â€¦â€¦â€¦Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillatorâ€¦â€¦â€¦4040
- Recommendation Table 15â€¦â€¦â€¦Recommendations for wearable cardioverter defibrillatorâ€¦â€¦â€¦4040
- Recommendation Table 16â€¦â€¦â€¦Recommendations for optimization of device programmingâ€¦â€¦â€¦4041
- Recommendation Table 17â€¦â€¦â€¦Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapyâ€¦â€¦â€¦4042
- Recommendation Table 18â€¦â€¦â€¦Recommendations for psychosocial management after implantable cardioverter defibrillator implantationâ€¦â€¦â€¦4042
- Recommendation Table 19â€¦â€¦â€¦Recommendations for implantable cardioverter defibrillator implantation in left ventricular assist device recipientsâ€¦â€¦â€¦4042
- Recommendation Table 20â€¦â€¦â€¦Recommendations for prevention of implantable cardioverter defibrillator complicationsâ€¦â€¦â€¦4043
- Recommendation Table 21â€¦â€¦â€¦Recommendations for end-of-life issues in implantable cardioverter defibrillator carriersâ€¦â€¦â€¦4043
- Recommendation Table 22â€¦â€¦â€¦Recommendations for treatment of ventricular arrhythmias in acute coronary syndrome and vasospasmâ€¦â€¦â€¦4046
- Recommendation Table 23â€¦â€¦â€¦Recommendations for risk stratification and treatment of ventricular arrhythmias early after myocardial infarctionâ€¦â€¦â€¦4046
- Recommendation Table 24â€¦â€¦â€¦Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in chronic coronary artery diseaseâ€¦â€¦â€¦4047

- Recommendation Table 25a^{ADSCCH} Recommendations for sudden cardiac death prevention in patients with coronary anomalies^{ADNBH}4050
- Recommendation Table 26a^{ADSCCH} Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia^{ADNBH}4052
- Recommendation Table 27a^{ADSCCH} Recommendations for the management of patients with premature ventricular complex-induced or premature ventricular complex-aggravated cardiomyopathy^{ADNBH}4053
- Recommendation Table 28a^{ADSCCH} Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy^{ADNBH}4057
- Recommendation Table 29a^{ADSCCH} Recommendations for diagnostic, risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in arrhythmogenic right ventricular cardiomyopathy^{ADNBH}4060
- Recommendation Table 30a^{ADSCCH} Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in hypertrophic cardiomyopathy^{ADNBH}4062
- Recommendation Table 31a^{ADSCCH} Recommendations for implantable cardioverter defibrillator implantation in left ventricular non-compaction^{ADNBH}4063
- Recommendation Table 32a^{ADSCCH} Recommendations for implantable cardioverter defibrillator implantation in patients with cardiac amyloidosis^{ADNBH}4063
- Recommendation Table 33a^{ADSCCH} Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in neuromuscular diseases^{ADNBH}4065
- Recommendation Table 34a^{ADSCCH} Recommendations for sudden cardiac death prevention and treatment of ventricular arrhythmias in myocarditis^{ADNBH}4066
- Recommendation Table 35a^{ADSCCH} Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in cardiac sarcoidosis^{ADNBH}4067
- Recommendation Table 36a^{ADSCCH} Recommendations for the treatment of ventricular arrhythmias in Chagas^{ADNBH}4070
- Recommendation Table 37a^{ADSCCH} Recommendations for sudden cardiac death prevention and treatment of ventricular arrhythmias in valvular heart disease^{ADNBH}4070
- Recommendation Table 38a^{ADSCCH} Recommendations for risk stratification and primary prevention of sudden cardiac death in congenital heart disease^{ADNBH}4072
- Recommendation Table 39a^{ADSCCH} Recommendations for secondary prevention of sudden cardiac death and treatment of ventricular arrhythmia in congenital heart disease^{ADNBH}4072
- Recommendation Table 40a^{ADSCCH} Recommendations for the management of patients with idiopathic ventricular fibrillation^{ADNBH}4073
- Recommendation Table 41a^{ADSCCH} Recommendations for the management of patients with long QT syndrome^{ADNBH}4077
- Recommendation Table 42a^{ADSCCH} Recommendations for management of patients with Andersen^{ADSTST}Tawil syndrome^{ADNBH}4079
- Recommendation Table 43a^{ADSCCH} Recommendations for management of patients with Brugada syndrome^{ADNBH}4080
- Recommendation Table 44a^{ADSCCH} Recommendations for the management of patients with early repolarization pattern/syndrome^{ADNBH}4083
- Recommendation Table 45a^{ADSCCH} Recommendations for the management of patients with catecholaminergic polymorphic ventricular tachycardia^{ADNBH}4086
- Recommendation Table 46a^{ADSCCH} Recommendations for the management of patients with short QT syndrome^{ADNBH}4087

- Recommendation Table 47a Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy 4088
- Recommendation Table 48a Recommendations for the prevention of sudden cardiac death before and after heart transplantation 4089
- Recommendation Table 49a Recommendations for risk stratification and prevention of sudden cardiac death in athletes 4089
- Recommendation Table 50a Recommendations for implantable cardioverter defibrillator implantation in the elderly 4090

List of tables

- Table 1 Classes of recommendations 4003
- Table 2 Levels of evidence 4004
- Table 3 New sections and concepts 4005
- Table 4 New recommendations in 2022 4005
- Table 5 Changes in recommendations since 2015 4010
- Table 6 Intravenous provocative diagnostic tests 4016
- Table 7 Genetic tests and suggested work-up of probands and relatives with primary electrical diseases 4017
- Table 8 Anti-arrhythmic drugs (acute and chronic treatment) 4035
- Table 9 Summary of the recommendations for the treatment of patients with frequent idiopathic premature ventricular complexes/ventricular tachycardia or premature ventricular complex-induced cardiomyopathy 4051
- Table 10 Modified long QT syndrome diagnostic score 4076

List of figures

- Figure 1 Central figure 4013
- Figure 2 Algorithm for the evaluation of patients presenting with an incidental finding of non-sustained ventricular tachycardia 4019
- Figure 4 Algorithm for the evaluation of patients presenting with a first sustained monomorphic ventricular tachycardia episode 4021
- Figure 3 Typical idiopathic ventricular tachycardia morphologies 4020
- Figure 5 Bundle branch re-entrant ventricular tachycardia 4022
- Figure 6 Part One. Algorithm for the evaluation of sudden cardiac arrest survivors 4024
- Figure 6 Part Two. Algorithm for the evaluation of sudden cardiac arrest survivors 4025
- Figure 7 Algorithm for the evaluation of sudden death victims 4026
- Figure 8 Algorithm for the evaluation of relatives of unexplained sudden death decedents 4028
- Figure 9 Algorithm for the acute management of regular wide QRS complex tachycardia 4029
- Figure 10 Repetitive runs of ventricular tachycardia interrupted by occasional sinus beats 4030
- Figure 11 Part One. Management of patients with electrical storm or repeated implantable cardioverter defibrillator discharges 4031
- Figure 11 Part Two. Management of patients with electrical storm or repeated implantable cardioverter defibrillator discharges 4032

- Figure 11 Part Three. Management of patients with electrical storm or repeated implantable cardioverter defibrillator dischargesâ€¦4033
- Figure 12 Algorithm for evaluation before initiation and follow-up of patients requiring sodium channel blocking agentsâ€¦4038
- Figure 13 Algorithm for evaluation before initiation and follow-up of patients requiring drugs associated with QT prolongationâ€¦4039
- Figure 14 Algorithm for the prevention and management of ventricular arrhythmias in ST-elevation myocardial infarctionâ€¦4045
- Figure 15 Algorithm for risk stratification and primary prevention of sudden cardiac death in patients with chronic coronary artery disease and reduced ejection fractionâ€¦4048
- Figure 16 Algorithm for the management of sustained monomorphic ventricular tachycardia in patients with chronic coronary artery diseaseâ€¦4049
- Figure 17 Algorithm for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia and non-apparent structural heart diseaseâ€¦4051
- Figure 18 Algorithm for the management of patients with premature ventricular complex-induced/-aggravated cardiomyopathyâ€¦4053
- Figure 19 Algorithm for risk stratification and primary prevention of sudden cardiac death in patients with dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathyâ€¦4055
- Figure 20 Typical features of dilated cardiomyopathy associated with lamin A/C gene mutation with ventricular arrhythmiasâ€¦4056
- Figure 21 Typical features of arrhythmogenic right ventricular cardiomyopathy associated with ventricular arrhythmiasâ€¦4059
- Figure 22 Typical features of hypertrophic cardiomyopathy associated with sustained monomorphic ventricular tachycardiaâ€¦4061
- Figure 23 Algorithm for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in myotonic dystrophyâ€¦4064
- Figure 24 Algorithm for sudden cardiac death prevention and treatment of ventricular arrhythmia in patients with cardiac sarcoidosisâ€¦4068
- Figure 25 Typical features of cardiac sarcoidosis associated with sustained monomorphic ventricular tachycardiaâ€¦4069
- Figure 26 Algorithm for the management of sustained ventricular arrhythmia in patients with congenital heart diseaseâ€¦4071
- Figure 27 Algorithm for the management of patients with idiopathic ventricular fibrillationâ€¦4074
- Figure 28 Idiopathic ventricular fibrillation triggered by short-coupled premature ventricular complexesâ€¦4075
- Figure 29 Long QT syndrome electrocardiograms and torsade-de-pointes ventricular tachycardiaâ€¦4076
- Figure 30 Brisk standing electrocardiogram changes and T wave alternans in Long QT syndrome patientsâ€¦4077
- Figure 31 Algorithm for the management of patients with long QT syndromeâ€¦4078
- Figure 32 Typical examples of (A) Brugada type 1 electrocardiogram, and (B) early repolarization pattern electrocardiogramâ€¦4080
- Figure 33 Part One. Algorithm for the management of patients with Brugada pattern electrocardiogramâ€¦4082
- Figure 33 Part Two. Algorithm for the management of patients with Brugada pattern electrocardiogramâ€¦4083
- Figure 34 Management of patients with early repolarization pattern/syndromeâ€¦4084
- Figure 35 Management of patients with catecholaminergic polymorphic ventricular tachycardiaâ€¦4085

- Figure 36 Exercise test of a patient with catecholaminergic polymorphic ventricular tachycardia ⁴⁰⁸⁶
- Figure 37 Short QT syndrome ⁴⁰⁸⁷

Abbreviations and acronyms

[A](#)

- AAD
Anti-arrhythmic drug

[A](#)

- ACE-I
Angiotensin-converting enzyme inhibitor

[A](#)

- ACS
Acute coronary syndrome

[A](#)

- AED
Automated external defibrillator

[A](#)

- AF
Atrial fibrillation

[A](#)

- AH
Atrial ⁴⁰⁸⁸His interval

[A](#)

- ALS
Advanced life support

[A](#)

- ARB
Angiotensin receptor blocker

[A](#)

- ARNI
Angiotensin receptor neprilysin inhibitor

[A](#)

- ARVC
Arrhythmogenic right ventricular cardiomyopathy

[A](#)

- ATP
Anti-tachycardia pacing

[A](#)

- AV
Atrioventricular

[A](#)

- AVRT
AV re-entry tachycardia

[A](#)

- BBR-VT
Bundle branch re-entrant ventricular tachycardia

[A](#)

- BrS

Brugada syndrome

[Â](#)

- CA

Cardiac arrest

[Â](#)

- cAMP

Cyclic adenosine monophosphate

[Â](#)

- CAD

Coronary artery disease

[Â](#)

- CAG

Coronary angiogram

[Â](#)

- CCB

Calcium channel blocker

[Â](#)

- CHD

Congenital heart disease

[Â](#)

- CIED

Cardiac implantable electronic devices

[Â](#)

- CMR

Cardiac magnetic resonance

[Â](#)

- CPR

Cardiopulmonary resuscitation

[Â](#)

- CPVT

Catecholaminergic polymorphic ventricular tachycardia

[Â](#)

- CRT

Cardiac resynchronization therapy

[Â](#)

- CT/CTA

Computed tomography/Computed tomography angiography

[Â](#)

- DCM

Dilated cardiomyopathy

[Â](#)

- DNA

Deoxyribonucleic acid

[Â](#)

- ECG

Electrocardiogram

[Â](#)

- EF

Ejection fraction

[Â](#)

- ERP
Early repolarization pattern

[Â](#)

- ERS
Early repolarization syndrome

[Â](#)

- FBI
Fast, broad, irregular

[Â](#)

- HCM
Hypertrophic cardiomyopathy

[Â](#)

- HFrEF
Heart failure with reduced ejection fraction

[Â](#)

- HNDCM
Hypokinetic non-dilated cardiomyopathy

[Â](#)

- HTX
Heart transplantation

[Â](#)

- HV
His-Purkinje ventricular interval

[Â](#)

- ICD
Implantable cardioverter defibrillator

[Â](#)

- ILR
Implantable loop recorder

[Â](#)

- IVF
Idiopathic ventricular fibrillation

[Â](#)

- LBBB
Left bundle branch block

[Â](#)

- LCSD
Left cardiac sympathetic denervation

[Â](#)

- LGE
Late gadolinium enhancement

[Â](#)

- LMNA
Lamin A/C

[Â](#)

- LQTS
Long QT syndrome

[Â](#)

- LV
Left ventricular

[Â](#)

- LVAD
Left ventricular assist device

[Â](#)

- LVEF
Left ventricular ejection fraction

[Â](#)

- LVH
Left ventricular hypertrophy

[Â](#)

- LVNC
Left ventricular non-compaction

[Â](#)

- LVOT
Left ventricle outflow tract

[Â](#)

- MI
Myocardial infarction

[Â](#)

- MRA
Mineralocorticoid receptor antagonist

[Â](#)

- MVP
Mitral valve prolapse

[Â](#)

- MVT
Monomorphic ventricular tachycardia

[Â](#)

- NSVT
Non-sustained ventricular tachycardia

[Â](#)

- NYHA
New York Heart Association

[Â](#)

- OHCA
Out-of-hospital cardiac arrest

[Â](#)

- OMT
Optimal medical treatment

[Â](#)

- PCI
Percutaneous coronary intervention

[Â](#)

- PCR
Polymerase chain reaction

[Â](#)

- PES
Programmed electrical stimulation

[Â](#)

- PET-CT
Positron emission tomography computed tomography

[Â](#)

- PPCM
Peri-partum cardiomyopathy

[Â](#)

- PVC
Premature ventricular complex

[Â](#)

- PVT
Polymorphic ventricular tachycardia

[Â](#)

- QI
Quality indicators

[Â](#)

- RBBB
Right bundle branch block

[Â](#)

- RCM
Restrictive cardiomyopathy

[Â](#)

- RCT
Randomized control trial

[Â](#)

- RV
Right ventricular

[Â](#)

- RVOT
Right ventricle outflow tract

[Â](#)

- SADS
Sudden arrhythmic death syndrome

[Â](#)

- SaECG
Signal-averaged ECG

[Â](#)

- SCA
Sudden cardiac arrest

[Â](#)

- SCD
Sudden cardiac death

[Â](#)

- SD
Sudden death

[^](#)

- SGLT2
Sodium-glucose co-transporter 2

[^](#)

- SHD
Structural heart disease

[^](#)

- S-ICD
Subcutaneous implantable cardioverter defibrillator

[^](#)

- SMVT
Sustained monomorphic ventricular tachycardia

[^](#)

- SPTV
Sustained polymorphic ventricular tachycardia

[^](#)

- SQTS
Short QT syndrome

[^](#)

- STEMI
ST elevation myocardial infarction

[^](#)

- SVT
Supraventricular tachycardia

[^](#)

- TAVI
Transcatheter aortic valve implantation

[^](#)

- TdP
Torsades de pointes

[^](#)

- TOF
Tetralogy of Fallot

[^](#)

- VA
Ventricular arrhythmia

[^](#)

- VF
Ventricular fibrillation

[^](#)

- VT
Ventricular tachycardia

[^](#)

- WCD
Wearable cardioverter defibrillator

[^](#)

- WPW
Wolff-Parkinson-White

1. Preamble

Guidelines summarize and evaluate available evidence with the aim of assisting health professionals in proposing the best management strategies for an individual patient with a given condition. Guidelines and their recommendations should facilitate decision making of health professionals in their daily practice. However, guidelines are not a substitute for the patientâpractitioner relationship with their practitioner. The final decisions concerning an individual patient must be made by the responsible health professional(s), based on what they consider to be the most appropriate in the circumstances. These decisions are made in consultation with the patient and caregiver as appropriate.

Guidelines are intended for use by health professionals. To ensure that all users have access to the most recent recommendations, the ESC makes its guidelines freely available. The ESC warns readers that the technical language may be misinterpreted and declines any responsibility in this respect.

A great number of guidelines have been issued in recent years by the ESC. Because of their impact on clinical practice, quality criteria for the development of guidelines have been established to make all decisions transparent to the user. The recommendations for formulating and issuing ESC Guidelines can be found on the ESC website (<https://www.escardio.org/Guidelines>). The ESC Guidelines represent the official position of the ESC on a given topic and are regularly updated.

In addition to the publication of Clinical Practice Guidelines, the ESC carries out the EURObservational Research Programme of international registries of cardiovascular diseases and interventions that are essential to assess diagnostic/therapeutic processes, use of resources, and adherence to guidelines. These registries aim at providing a better understanding of medical practice in Europe and around the world, based on high-quality data collected during routine clinical practice.

Furthermore, the ESC develops sets of quality indicators (QIs), which are tools to evaluate the level of implementation of the guidelines and may be used by the ESC, hospitals, healthcare providers, and professionals to measure clinical practice, and in educational programmes, alongside the key messages from the guidelines, to improve quality of care and clinical outcomes.

The Members of this task force were selected by the ESC to represent professionals involved with the medical care of patients with this pathology. The selection procedure aimed to ensure that there is a representative mix of members predominantly from across the whole of the ESC region and from relevant ESC Subspecialty Communities. Consideration was given to diversity and inclusion, notably with respect to gender and country of origin. A critical evaluation of diagnostic and therapeutic procedures was performed, including assessment of the riskâbenefit ratio. The level of evidence and the strength of the recommendation of particular management options were weighed and scored according to predefined scales, as outlined below. The task force followed the ESC voting procedures. All recommendations subject to a vote achieved at least 75% among voting members.

The experts of the writing and reviewing panels provided declaration of interest forms for all relationships that might be perceived as real or potential sources of conflicts of interest. Their declarations of interest were reviewed according to the ESC declaration of interest rules and can be found on the ESC website (<http://www.escardio.org/Guidelines>) and have been compiled in a report and published in a supplementary document simultaneously to the guidelines.

This process ensures transparency and prevents potential biases in the development and review processes. Any changes in declarations of interest that arise during the writing period were notified to the ESC and updated. The task force received its entire financial support from the ESC without any involvement from the healthcare industry.

The ESC CPG Committee supervises and coordinates the preparation of new guidelines. The Committee is also responsible for the approval process of these guidelines. The ESC Guidelines undergo extensive review by the CPG Committee and external experts, including a mix of members from across the whole of the ESC region and from relevant ESC Subspecialty Communities and National Cardiac Societies. After appropriate revisions, the guidelines are signed off by all the experts involved in the task force. The finalized document is signed off by the CPG Committee for publication in the *European Heart Journal*. The guidelines were developed after careful consideration of the scientific and medical knowledge and the evidence available at the time of their writing.

The task of developing ESC Guidelines also includes the creation of educational tools and implementation programmes for the recommendations, including condensed pocket guideline versions, summary slides, summary cards for non-specialists, and an electronic version for digital applications (smartphones, etc.). These versions are abridged and thus, for more detailed information, the user should always access the full-text version of the guidelines, which is freely available via the ESC website and the *European Heart Journal*. The National Cardiac Societies of the ESC are encouraged to endorse, adopt, translate, and implement all ESC Guidelines. Implementation programmes are

needed because it has been shown that the outcome of disease may be favourably influenced by the thorough application of clinical recommendations.

Health professionals are encouraged to take the ESC Guidelines fully into account when exercising their clinical judgement, as well as in the determination and the implementation of preventive, diagnostic, or therapeutic medical strategies. However, the ESC Guidelines do not override in any way whatsoever the individual responsibility of health professionals to make appropriate and accurate decisions in consideration of each patient's health condition and in consultation with that patient or the patient's caregiver where appropriate and/or necessary. It is also the health professional's responsibility to verify the rules and regulations applicable in each country to drugs and devices at the time of prescription and, where appropriate, to respect the ethical rules of their profession.

Off-label use of medication may be presented in this guideline if sufficient level of evidence shows that it can be considered medically appropriate to a given condition and if patients could benefit from the recommended therapy. However, the final decisions concerning an individual patient must be made by the responsible health professional giving special consideration to:

1. the specific situation of the patient. In this respect, it is specified that, unless otherwise provided for by national regulations, off-label use of medication should be limited to situations where it is in the patient's interest to do so, with regard to the quality, safety, and efficacy of care, and only after the patient has been informed and has provided consent;
2. country-specific health regulations, indications by governmental drug regulatory agencies, and the ethical rules to which health professionals are subject, where applicable.

2. Introduction

This document presents an update of the 2015 ESC Guidelines for the management of patients with ventricular arrhythmias (VA) and the prevention of sudden cardiac death (SCD). New insights into the epidemiology of SCD, new evidence on genetics, imaging, and clinical findings for risk stratification for VA and SCD, and advances in diagnostic evaluation and therapeutic strategies made this revision necessary. The committee was composed of 25 members including 23 expert physicians, one methodologist, and one patient representative. Experts were selected to cover all areas of VA and SCD as well as subspecialties of cardiology with the assistance of related ESC working groups.

All 25 members of the task force committee approved the guideline recommendations after an anonymous voting process. Ninety-nine peer reviewers reviewed the document. A systematic literature survey was conducted, after instructions by the methodologist in the group, that led to the incorporation of 1155 references, of which 485 were selected to support the recommendations and further specified in the table of evidence ([Supplementary data](#)).

2.1. What is new

The diagnostic and management parts of the guidelines have been adapted to facilitate their use in everyday clinical decision-making.

Table 1

[Open in new tab](#)

Classes of recommendations

Classes of recommendations

Definition		Wording to use
Class I	Evidence and/or general agreement that a given treatment or procedure is beneficial, useful, effective.	Is recommended or is indicated
Class II	Conflicting evidence and/or a divergence of opinion about the usefulness/efficacy of the given treatment or procedure.	
Class IIa	Weight of evidence/opinion is in favour of usefulness/efficacy.	Should be considered
Class IIb	Usefulness/efficacy is less well established by evidence/opinion.	May be considered
Class III	Evidence or general agreement that the given treatment or procedure is not useful/effective, and in some cases may be harmful.	Is not recommended

©ESC 2022

Classes of recommendations	Definition		Wording to use
	Class I	Evidence and/or general agreement that a given treatment or procedure is beneficial, useful, effective.	Is recommended or is indicated
	Class II	Conflicting evidence and/or a divergence of opinion about the usefulness/ efficacy of the given treatment or procedure.	
	Class IIa	Weight of evidence/opinion is in favour of usefulness/efficacy.	Should be considered
	Class IIb	Usefulness/efficacy is less well established by evidence/opinion.	May be considered
	Class III	Evidence or general agreement that the given treatment or procedure is not useful/effective, and in some cases may be harmful.	Is not recommended

Table 1

[Open in new tab](#)

Classes of recommendations

Classes of recommendations	Definition		Wording to use
	Class I	Evidence and/or general agreement that a given treatment or procedure is beneficial, useful, effective.	Is recommended or is indicated
	Class II	Conflicting evidence and/or a divergence of opinion about the usefulness/ efficacy of the given treatment or procedure.	
	Class IIa	Weight of evidence/opinion is in favour of usefulness/efficacy.	Should be considered
	Class IIb	Usefulness/efficacy is less well established by evidence/opinion.	May be considered
	Class III	Evidence or general agreement that the given treatment or procedure is not useful/effective, and in some cases may be harmful.	Is not recommended

Classes of recommendations	Definition		Wording to use
	Class I	Evidence and/or general agreement that a given treatment or procedure is beneficial, useful, effective.	Is recommended or is indicated
	Class II	Conflicting evidence and/or a divergence of opinion about the usefulness/ efficacy of the given treatment or procedure.	
	Class IIa	Weight of evidence/opinion is in favour of usefulness/efficacy.	Should be considered
	Class IIb	Usefulness/efficacy is less well established by evidence/opinion.	May be considered
	Class III	Evidence or general agreement that the given treatment or procedure is not useful/effective, and in some cases may be harmful.	Is not recommended

Table 2

[Open in new tab](#)

Levels of evidence

Level of evidence A	Data derived from multiple randomized clinical trials or meta-analyses.
Level of evidence B	Data derived from a single randomized clinical trial or large non-randomized studies.
Level of evidence C	Consensus of opinion of the experts and/or small studies, retrospective studies, registries.

©ESC 2022

Level of evidence A	Data derived from multiple randomized clinical trials or meta-analyses.
Level of evidence B	Data derived from a single randomized clinical trial or large non-randomized studies.
Level of evidence C	Consensus of opinion of the experts and/or small studies, retrospective studies, registries.

©ESC 2022

TableÂ 2

[Open in new tab](#)

Levels of evidence

Level of evidence A	Data derived from multiple randomized clinical trials or meta-analyses.
Level of evidence B	Data derived from a single randomized clinical trial or large non-randomized studies.
Level of evidence C	Consensus of opinion of the experts and/or small studies, retrospective studies, registries.

©ESC 2022

Level of evidence A	Data derived from multiple randomized clinical trials or meta-analyses.
Level of evidence B	Data derived from a single randomized clinical trial or large non-randomized studies.
Level of evidence C	Consensus of opinion of the experts and/or small studies, retrospective studies, registries.

©ESC 2022

The first general part has new sections on diagnostic evaluation, including pharmacologic provocative tests, genetic testing, and a systematic work-up of probands and relatives with primary electrical diseases. Comprehensive flowcharts and recommendations for the diagnostic evaluation at first presentation with a VA of patients without a previously known cardiac disease are provided for five frequently encountered clinical scenarios. Practical recommendations for optimization of implantable cardioverter defibrillator (ICD) programming and algorithms for management of patients experiencing regular wide complex tachycardia and electrical storm are presented ([TableÂ 3](#)).

Table 3

[Open in new tab](#)

New sections and concepts

New sections and concepts	Section
Provocative diagnostic tests	5.1.3.5
Genetic testing	5.1.4
Diagnostic evaluation at first presentation with VA in patients without known cardiac disease	5.2
Management of patients with electrical storm	6.1.3
Special aspects of device therapy	6.2.3

New sections and concepts	Section
Provocative diagnostic tests	5.1.3.5
Genetic testing	5.1.4
Diagnostic evaluation at first presentation with VA in patients without known cardiac disease	5.2
Management of patients with electrical storm	6.1.3
Special aspects of device therapy	6.2.3

© ESC 2022

Table 3

[Open in new tab](#)

New sections and concepts

New sections and concepts	Section
Provocative diagnostic tests	5.1.3.5
Genetic testing	5.1.4
Diagnostic evaluation at first presentation with VA in patients without known cardiac disease	5.2
Management of patients with electrical storm	6.1.3
Special aspects of device therapy	6.2.3

New sections and concepts	Section
Provocative diagnostic tests	5.1.3.5
Genetic testing	5.1.4
Diagnostic evaluation at first presentation with VA in patients without known cardiac disease	5.2
Management of patients with electrical storm	6.1.3
Special aspects of device therapy	6.2.3

© ESC 2022

The second part of the guidelines is structured according to disease-specific management, providing a link to the updated  [ESC CardioMed](#) chapter for additional content. Risk stratification, SCD prevention, treatment of VA, and management of family members are addressed in a systematic fashion. Indications for cardiac magnetic resonance (CMR) imaging, genetic testing, and updated indications for catheter ablation of ventricular arrhythmias are presented. Flowcharts summarizing the workflow for diagnostic and treatment are provided for the disease entities. The colour-coding of the flowcharts reflects the class of recommendation according to this guideline and other ESC

Guidelines.^{1a} Another novel concept of this document is the table of evidence (see [Supplementary data](#)). The trials and studies that have been selected to support a recommendation are systematically described in the table of evidence after careful review of the available data and the applied methodology, prioritizing papers published after 2015. Recommendations with level of evidence C that are not accompanied by a reference are supported by this panel of experts. To assist physicians in their daily clinical practice, diagnostic and therapeutic procedures with promising usefulness, typically classified as class IIb, but for which evidence is limited and difficult to collect in the near future, the related recommendations are not only described in the narratives but are listed in the table of recommendation.

3. Definitions

3.1. Ventricular arrhythmia subtypes

Premature ventricular complex (PVC): Premature occurrence of an abnormal QRS complex (duration typically ≤ 120 ms, corresponding T-wave typically broad and in the opposite direction of the major QRS deflection, no preceding P-wave).

Unifocal or monomorphic PVCs: PVCs with a single QRS morphology.

Multifocal, multiform, or polymorphic PVCs: PVCs with different QRS morphologies.

Short-coupled PVC: A PVC that interrupts the T-wave of the preceding conducted beat.

Ventricular tachycardia (VT): ≥ 3 consecutive beats with a rate >100 b.p.m. originating from the ventricles, independent from atrial and atrioventricular (AV) nodal conduction.

Non-sustained ventricular tachycardia (NSVT): Run of consecutive ventricular beats persisting for 3 beats to 30 s.

Monomorphic ventricular tachycardia (MVT): Same QRS morphology from beat to beat.

Polymorphic ventricular tachycardia (PVT): Continually changing QRS morphology.

Sustained monomorphic/polymorphic ventricular tachycardia (SMVT/SPVT): Continuous VT for at least 30 s, or which requires an intervention for termination.

Bidirectional ventricular tachycardia: Beat to beat alternation of the frontal QRS axis (e.g. in catecholaminergic polymorphic ventricular tachycardia [CPVT], Andersen-Tawil, digoxin toxicity, acute myocarditis).

Torsades de pointes ventricular tachycardia (TdP): Subtype of a polymorphic VT in the context of QT prolongation with continually changing QRS complexes that appear to spiral around the baseline of the electrocardiogram (ECG) lead in a sinusoidal pattern.

Ventricular fibrillation (VF): A chaotic rhythm with undulations that are irregular in timing and morphology, without discrete QRS complexes on the surface ECG.

Electrical storm: VA that occurs 3 or more times within 24 h (separated by at least 5 min), each requiring termination by an intervention.

Incessant VT: Continuous sustained VT that recurs promptly despite repeated intervention for termination over several hours.

3.2. Sudden cardiac death

Sudden cardiac arrest (SCA): Sudden cessation of normal cardiac activity with haemodynamic collapse.

Sudden cardiac death (SCD): Sudden natural death presumed to be of cardiac cause that occurs within 1 h of onset of symptoms in witnessed cases, and within 24 h of last being seen alive when it is unwitnessed. SCD in autopsied cases is defined as the natural unexpected death of unknown or cardiac cause.

Sudden unexplained death: Unexplained sudden death occurring in an individual older than 1 year.

Sudden infant death syndrome (SIDS): Unexplained sudden death occurring in an individual younger than 1 year with negative pathological and toxicological assessment and negative forensic examination of the circumstances of death.

Sudden arrhythmic death syndrome (SADS): Unexplained sudden death occurring in an individual older than 1 year with negative pathological and toxicological assessment. Note: Synonymous with *autopsy-negative sudden unexplained death*.

3.3. Syncope

Unexplained syncope: Transient loss of consciousness due to cerebral hypoperfusion, characterized by a rapid onset, short duration, and spontaneous complete recovery, but unexplained after conventional workup. Work-up and differential diagnosis are provided in the 2018 ESC Guidelines for the diagnosis and management of syncope.¹

Arrhythmic syncope: as above, but highly suspicious for intermittent bradycardia, rapid supraventricular tachycardia (SVT), or VA.

3.4. Specialized centres

Multidisciplinary teams: A multidisciplinary team across specialties is characterized by open communication, positive management and leadership, appropriate resources, and a mix of skills. Decision-making should be shared in the team.

Specialized centre for catheter ablation of VA: Patient and procedural complexity vary widely. Some patients require a more experienced operator and a centre with more capabilities, which is more likely in patients with a non-ischaemic aetiology. A specialized centre has at least one operator with appropriate experience in interventions that may be required for a successful procedure (e.g. percutaneous epicardial access). The centre performs catheter ablation of VT in structural heart disease (SHD) on a regular basis. In addition, the centre has the required resources to manage medical conditions, comorbidities, and potential complications in patients undergoing complex VA ablation; this includes interventional cardiology expertise, acute placement of mechanical circulatory assist devices, and cardiothoracic surgical back-up. Considering the variable availability across European countries, it is preferable to treat complex patients in the most experienced centre within reasonable distance.

3.5. Genetics

Pathogenic variant and likely pathogenic variant: The American College of Medical Genetics has provided a framework for the interpretation of disease causation by genetic variants standardized into classes. The genetic variants most likely to cause an associated disease are termed V, a pathogenic, and IV, a likely pathogenic.

Mutation: This term is used in this document to mean a Class IV or V variant.

Variant of uncertain significance: A change in a gene's deoxyribonucleic acid (DNA) sequence that has an unknown effect on a person's health.

4. Epidemiology of sudden cardiac death, public awareness, and risk stratification

4.1. Incidence of sudden cardiac death

SCD accounts for approximately 50% of all cardiovascular deaths, with up to 50% being the first manifestation of cardiac disease.^{19,24} Ideally, cases suspicious of SCD should be identified from multiple sources and undergo autopsy, which is required to reliably exclude non-cardiac causes of sudden death (SD).

The incidence of SCD increases markedly with age. With a very low incidence during infancy and childhood (1 per 100 000 person-years),²⁵ the incidence is approximately 50 per 100 000 person-years in middle-aged individuals (in the fifth to sixth decades of life).^{28,30} In the eighth decade of life, it reaches an annual incidence of at least 200 per 100 000 person-years.²⁰ At any age, males have higher SCD rates compared with females, even after adjustment for risk factors of coronary artery disease (CAD).^{24,31,33} Ethnic background also seems to have large effects.^{34,35} It is estimated that 10–20% of all deaths in Europe are SCD.^{36,37} Approximately 300 000 people in Europe have out-of-hospital cardiac arrest (OHCA) treated by emergency medical systems every year.^{38,39}

In the Western world, the epidemiology of SCD is closely related to CAD, which is responsible for up to 75–80% of SCD cases.⁴⁰ While the prevalence of CAD has not decreased, there has been a significant decline in mortality due

to CAD. Reports show that the incidence of SCD is declining,^{40,42} but the risk of SCD as a proportion of the overall cardiovascular deaths may have increased.^{43,44}


Although regular physical activity benefits cardiovascular health, sport, particularly when practiced vigorously, has been shown to be associated with SCD during or shortly after exercise in selected populations.⁴⁵ Reports have suggested that the majority of sports-related SCD occurs in a recreational^{52,53} rather than competitive setting, especially among middle-aged male participants, suggesting that CAD is the most common underlying cause.^{46,54,55}

4.2. Causes of sudden cardiac death in different age groups

Cardiac diseases associated with SCD vary depending on the individual's age. In the young there is a predominance of primary electric diseases and cardiomyopathies, as well as myocarditis, and coronary anomalies.^{25,27,56} However, half of SCD cases during the fourth decade are related to CAD, especially acute coronary syndrome (ACS).^{62,63}

In older populations, chronic structural diseases predominate (CAD either through acute coronary events or chronic coronary stenoses, valvular heart diseases, and heart failure), while potentially inherited electrical diseases or structural non-ischaemic diseases may cause more than 50% of SCD in individuals under the age of 50 years.²⁷

Age distribution at presentation with ventricular arrhythmias and sudden cardiac death, dominant arrhythmia subtypes, triggers, genetic factors, and gender associated with increased risk for ventricular arrhythmias in selected primary electrical and structural heart diseases are presented in [FigureÂ 1](#).



FigureÂ 1
Central figure.

Genetic risk for VA/SCD, typical triggers for VA/SCD, age at presentation with VA/SCD, sex predominance, and typical VA (PVT/VF vs. MVT) in different diseases associated with VA/SCD. ACS, acute coronary syndrome; ARVC, arrhythmogenic right ventricular cardiomyopathy; BrS, Brugada syndrome; CAD, coronary artery disease; CPVT, catecholaminergic polymorphic ventricular tachycardia; DCM, dilated cardiomyopathy; HCM, hypertrophic cardiomyopathy; LQT, long QT syndrome; MVT, monomorphic ventricular tachycardia; PVT, polymorphic ventricular tachycardia; rTOF, repaired tetralogy of Fallot; SCD, sudden cardiac death; VA, ventricular arrhythmia; VF, ventricular fibrillation.

[Open in new tab](#)[Download slide](#)

4.3. Population vs. individual risk prediction

In the general population (individuals without known heart disease), the most effective approach for preventing SCD resides in quantification of the individual risk of developing CAD based on risk score charts.^{64,65} Several studies have provided evidence that there is a genetic predisposition to die suddenly during acute ischaemia.^{66a,67,70} The goal would be to identify the relatively small, high-risk subgroups from the general population at risk of SCD as their first cardiac event. Models for novel SCD risk stratification in the general population have recently been proposed.^{71a,72,73} There are no clear data supporting the benefit of mass screening programs in the general population for preventing SCD.^{74a,75,76}

For decades, investigators have envisioned a broad range of ^{77a,78,79}indicators^{80,81} for SCD, especially occurring in the setting of CAD. Several non-invasive markers of risk have been proposed (including late potentials, heart rate variability, periodic repolarization dynamics, and baroreflex sensitivity).⁷⁷ Despite the promising outcomes of initial studies, however, these ^{82a,83,84}predictors⁸⁵ have not yet influenced clinical practice. Left ventricular ejection fraction (LVEF) is only used, often in combination with New York Heart Association (NYHA) class, for primary prevention indication of an ICD in the setting of chronic CAD and dilated cardiomyopathy (DCM). Risk stratification schemes and calculators have been developed for inheritable arrhythmogenic diseases, such as hypertrophic cardiomyopathy (HCM), arrhythmogenic right ventricular cardiomyopathy (ARVC), and lamin A/C (LMNA) cardiomyopathy.^{78a,79,82}

What is considered low, intermediate, or high risk will depend on the type of event (e.g. fatal or not) and the risk of an event in the background population. For instance, mortality depends on age, sex, and other risk factors including comorbidities. The situation is further complicated when considering a specific type of death, e.g. SCD. Then non-SCD deaths are competing events (or competing risks) to SCD in the sense that their occurrence prevents the observation of SCD, and may make interventions to prevent SCD, such as an ICD, of limited benefit.

4.4. Risk calculators for sudden cardiac death and review of the methodology

A number of risk calculators for SCD have been proposed for adult and paediatric populations.^{80,81,83a,84,85} The field of prediction modelling has evolved over the past decades, establishing standards for the development, validation (internal and external), and reporting of prediction models in SCD.^{86,87} In addition to discrimination measures such as the c-index, measures of calibration such as the calibration slope have recently received more attention because it is not only important to distinguish patients with higher risks from those with lower risks but also to obtain a robust quantification of the risk itself from the risk calculators.⁸⁸ Typical shortcomings in the development and validation of risk calculators include, but are not limited to, the use of historic samples not representative of contemporary patient cohorts, missing values, composite outcomes with composite events of different clinical importance, lack of external validation, and missing calibration. In this document different cut-off for 5 years risk of SCD/VA is used for indication of ICD. Each cut-off was chosen by the original authors and the task force group considering the competing risk, the outcome measured (SCD vs VA) and the robustness of each risk calculator.

4.5. Awareness and intervention: public basic life support, and access to automated external defibrillators

Survival rate remains strikingly low after OHCA,^{89a,90,95} although major regional disparities have been described.⁹⁶ Early implementation of resuscitative interventions, especially prior to emergency medical system arrival, has been identified as key elements to improve survival.^{95,97} Bystander cardiopulmonary resuscitation (CPR) and use of public automated external defibrillators (AEDs) have demonstrated improvement of neurological and functional outcome as well as survival of OHCA patients. Data support the need for increased availability of public access defibrillators and community training in basic life support,^{89a,90,95} preferably initiated in childhood and repetitive.^{98a,99,100} Finally, dispatch of basic life support-trained volunteers, through specific mobile apps networks, has been shown to increase the rate of bystander-initiated CPR, and as a result, greatly reduces the resuscitation-free interval and improves the outcome of OHCA victims.^{101a,102,103} Education of public officials and community members regarding the importance of increasing rates of bystander CPR and promoting the use of early defibrillation by lay and professional rescuers is critical to increasing survival rates.

Recommendation TableÂ 1

[Open in new tab](#)

Recommendations for public basic life support and access to automated external defibrillators

Recommendations	Class ^a	Level ^b
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ^{c,90–92}	I	B
Prompt CPR by bystanders is recommended at OHCA. ^{93–95}	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use. ^{93,97,104}	I	B
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered. ^{101–103,105}	IIa	B

© ESC 2022

AED, automated external defibrillator; CPR, cardiopulmonary resuscitation; OHCA, out-of-hospital cardiac arrest.
^aClass of recommendation.
^bLevel of evidence.
^cShopping centres, stadiums, public transport stations, casinos.

Recommendations	Class ^a	Level ^b
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ^{c,90–92}	I	B
Prompt CPR by bystanders is recommended at OHCA. ^{93–95}	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use. ^{93,97,104}	I	B
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered. ^{101–103,105}	IIa	B

© ESC 2022

AED, automated external defibrillator; CPR, cardiopulmonary resuscitation; OHCA, out-of-hospital cardiac arrest.
^aClass of recommendation.
^bLevel of evidence.
^cShopping centres, stadiums, public transport stations, casinos.

Recommendation Table 1

[Open in new tab](#)

Recommendations for public basic life support and access to automated external defibrillators

Recommendations	Class ^a	Level ^b
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ^{c,90–92}	I	B
Prompt CPR by bystanders is recommended at OHCA. ^{93–95}	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use. ^{93,97,104}	I	B
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered. ^{101–103,105}	IIa	B

© ESC 2022

AED, automated external defibrillator; CPR, cardiopulmonary resuscitation; OHCA, out-of-hospital cardiac arrest.
^aClass of recommendation.
^bLevel of evidence.
^cShopping centres, stadiums, public transport stations, casinos.

Recommendations	Class ^a	Level ^b
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ^{c,90–92}	I	B
Prompt CPR by bystanders is recommended at OHCA. ^{93–95}	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use. ^{93,97,104}	I	B
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered. ^{101–103,105}	IIa	B

© ESC 2022

AED, automated external defibrillator; CPR, cardiopulmonary resuscitation; OHCA, out-of-hospital cardiac arrest.
^aClass of recommendation.
^bLevel of evidence.
^cShopping centres, stadiums, public transport stations, casinos.

5. Evaluation and treatment. General aspects

5.1. Diagnostic tools

5.1.1. History and physical examination

History should focus on a **red flags**, including features of arrhythmic syncope, e.g. absence of vagal prodrome, and family history of premature or SCD including, e.g. drowning or car accident in long QT syndrome (LQTS), and CPVT.^{1,106} Subtle features to suggest inherited causes include a family history of epilepsy, sudden infant death syndrome, deafness (LQTS), heart failure, or pacemaker implantation at <50 years old. Features of diseases related to pro-arrhythmic conditions include mid-systolic click in mitral valve prolapse (MVP) and outflow tract murmurs with Valsalva in HCM. Specific skin features may be relevant, e.g. Lupus pernio, erythema nodosum in sarcoidosis, angiokeratoma in Fabry's disease, xanthelasma/xanthoma, and palmoplantar keratosis in ARVC.

5.1.2. Laboratory testing

Natriuretic peptides (b-type-natriuretic peptide, or N-terminal pro-b-type-natriuretic peptide) may have a role in the identification of individuals at increased risk of SCD in the general population^{107,108} or in patients with CAD.¹⁰⁹ There is not sufficient evidence to use b-type-natriuretic peptide as a method to select the need for an ICD.^{110,111}

5.1.3. Non-invasive and invasive tests

5.1.3.1. Electrocardiogram and ambulatory electrocardiographic monitoring²

The 12-lead ECG is an important tool for the diagnosis of underlying disease, for risk stratification in selected populations, and for the diagnosis of the VA subtype, if captured. Documentation of arrhythmias related to symptoms is clinically pivotal but may be challenging with sporadic events. The type of ECG-monitoring device and the recording time should therefore match the frequency of clinical events. Monitoring over a period of 24 h (typically a Holter recording) is appropriate for daily arrhythmias,¹¹² while intermittent monitoring over a longer period, with patient-activated ECG recorders (or mobile-health/smartphones), should be preferred for infrequent events.¹¹³ Implantable loop recorders (ILR) can be useful in diagnosing arrhythmias in patients with potentially life-threatening symptoms, such as unexplained syncope.¹¹⁴

5.1.3.2. Signal-averaged electrocardiogram

Signal-averaged electrocardiogram (SaECG) can detect very low amplitude signals (late potentials) in the terminal QRS segment¹¹⁵ using three time-domain measurements: QRS duration, low-amplitude (<40 µV) signal duration and root mean square voltage of terminal 40 ms QRS.¹¹² Abnormalities in the SaECG can also be assessed by frequency-domain analysis.¹¹² SaECG can contribute to the diagnosis of ARVC.¹¹⁶

5.1.3.3. Exercise testing

Exercise testing is useful for the diagnosis and for evaluating response to therapy in patients with suspected/proven adrenergic-dependent rhythm disturbances, such as exercise-induced idiopathic MVT, PVT, or bidirectional VT in CPVT.^{117,118} The 4-minute recovery QTc after exercise testing can contribute to the diagnosis LQTS.¹¹⁹

5.1.3.4. Imaging

Imaging is crucial to assess cardiac function and detect cardiomyopathies (ESC CardioMed chapter 10).¹²⁰ A negative imaging study supports primary electrical disease in a patient with VA. Echocardiography is a readily available and first-line diagnostic and risk stratification tool for valve diseases, CAD and DCM, HCM, ARVC,¹²¹ and left ventricular non-compaction (LVNC). Echocardiographic strain-rate imaging allows differentiation between active and passive movement of myocardial segments and early detection of myocardial dysfunction. Wall motion abnormalities can indicate previous infarcts, cardiomyopathies, or inflammatory disease. Global longitudinal strain is a robust measure of left ventricular (LV) function and can detect subtle changes in LV function while LVEF is still preserved.¹²² Strain imaging can assess mechanical dispersion reflecting inhomogeneous contraction that could be associated with increased risk of VA.^{122,125}

CMR currently provides the most accurate and reproducible measurement of atrial, biventricular global and regional systolic function, and can detect myocardial oedema, fibrosis, infiltration, and perfusion defects (ESC CardioMed chapter 11.4).¹²⁶ CMR is more sensitive than echocardiography to diagnose ARVC,¹²⁷ is diagnostic in LVNC, and can detect apical aneurysms in HCM. Fibrosis detection by late gadolinium enhancement (LGE) contributes to VA risk stratification in HCM,¹²⁸ DCM,¹²⁹ and potentially in mitral valve prolapse arrhythmic syndrome.^{130,131} Novel myocardial mapping techniques can detect diffuse fibrosis and can suggest the aetiology of left ventricular hypertrophy (LVH) for specific therapy guiding, e.g. Fabry disease and amyloidosis. The prognostic value remains to be evaluated.

Cardiac imaging by computed tomography (CT) has the advantage of a high spatial resolution (ESC CardioMed chapter 12.1).¹³² ECG synchronization, additional breath-hold sequences, and beta-blocker to lower the heart rate improve the quality. Radiation exposure is in the range of an invasive coronary angiogram (CAG). Coronary computed tomography angiography (CTA) is the preferred method to rule out coronary artery stenosis in patients with low probability of CAD.^{133,134} The quality of almost all image modalities is influenced by the presence of frequent PVCs.

Table 4

[Open in new tab](#)

New recommendations in 2022

Recommendations	Class
Public basic life support and access to AEDs	
It is recommended that public-access defibrillation be available at sites where cardiac arrest is more likely to occur. ^a	I
Prompt CPR by bystanders is recommended at OHCA.	I
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered.	IIa
Treatment of VA. General aspects	
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good-quality survival >1 year.	I
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered.	IIa
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered.	IIb
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb
The WCD may be considered in the early phase after MI in selected patients.	IIb
Coronary artery disease	
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered.	IIa
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT.	IIa
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT and NSVT, if they are inducible for SMVT by PES.	IIa

In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^b	Ila
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blocker or sotalol treatment.	Ila
Idiopathic PVC/VT and PVC-induced cardiomyopathy	
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.	I
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram.	Ila
Beta-blockers, non-dihydropyridine CCBs or flecainide should be considered when catheter ablation is not available, not desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles.	Ila
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	Ila
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered.	Ila
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered.	Ila
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered.	Ila
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up.	IIb
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III
DCM/HNDCM	
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: - the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or - there is a family history of DCM/HNDCM, or premature unexpected SD.	I
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD.	Ila

Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age or with signs suspicious for an inherited aetiology.	IIa
ICD implantation should be considered in DCM/HNDCM patients with an LVEF <50% and ≥ 2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG, and an echocardiogram may be considered.	IIb
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation.	III
ARVC	
In patients with suspected ARVC, CMR is recommended.	I
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and either NSVT or inducibility of SMVT at PES.	IIa
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered.	IIa
Avoidance of high-intensity ^e exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype.	IIb
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification.	IIb
HCM	
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I
Genetic counselling and testing are recommended in HCM patients.	I
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to <6%) ^f , and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF <50%; or (c) abnormal blood pressure response during exercise test ^g ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation.	IIa
In children <16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^h), ICD implantation should be considered.	IIa
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa

In patients with HCM and recurrent, symptomatic VA, or recurrent symptomatic ICD therapy, AAD treatment should be considered.	IIa
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers.	IIb
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD (<4%), ^f and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF < 50%; or (c) LV apical aneurysm.	IIb
Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb
LVNC and RCM	
In patients with an LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/HNDCM recommendations.	IIa
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa
Neuromuscular diseases	
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA.	I
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration.	IIa
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias, or who are older than 40 years and have significant LGE on CMR	IIa
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered.	IIa
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered.	IIa
In patients with limb-girdle type 1B or Emery–Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered.	IIa
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR.	IIb
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ⁱ for VA and SCD.	IIb
In patients with myotonic dystrophy, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III

Inflammatory diseases	
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered.	Ila
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	Ila
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired.	Ila
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis who have an LVEF >35% but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis who have an LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	Ila
In patients with cardiac sarcoidosis, LVEF 35–50%, and inducible SMVT at PES, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	Ila
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT.	Ila
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered.	Ila
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^b	Ilb
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD recipients with recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	Ilb
Congenital heart disease	
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended.	I
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation, and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered.	Ila
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered.	Ilb

In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy.	IIb
Idiopathic VF	
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF.	IIa
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF.	IIa
Clinical testing (history, ECG, and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered.	IIb
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered.	IIb
Long QT syndrome	
In patients with clinically diagnosed LQTS, genetic testing, and genetic counselling are recommended.	I
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I
Mexiletine is indicated in LQT3 patients with a prolonged QT interval.	I
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval.	IIa
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients).	IIb
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III
Andersen–Tawil syndrome	
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome.	I
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT.	I
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.	IIa

Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA.	IIa
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT.	IIb
Brugada syndrome	
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS.	I
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of the following: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope.	IIa
BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG.	IIb
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG.	IIb
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III
Catheter ablation in asymptomatic BrS patients is not recommended.	III
Early repolarization syndrome	
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I
In a SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered.	IIa
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^j	IIa
ILR should be considered in individuals with ERP and at least one risk feature ^k or arrhythmic syncope.	IIa
Isoproterenol infusion should be considered for ERS patients with electrical storm.	IIa
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients.	IIa
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment.	IIa
Genetic testing in ERS patients may be considered.	IIb
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^k	IIb

ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^l in the presence of a family history of unexplained juvenile SD.	IIb
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III
CPVT	
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT.	I
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb
Short QT syndrome	
Genetic testing is indicated in patients diagnosed with SQTS.	I
SQTS should be considered in the presence of a QTc ≤ 320 ms.	IIa
SQTS should be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and arrhythmic syncope.	IIa
ILR should be considered in young SQTS patients.	IIa
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa
SQTS may be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and a family history of SD at age <40 years.	IIb
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD.	IIb
Isoproterenol may be considered in SQTS patients with an electrical storm.	IIb
Selected populations	
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I
Continuation of beta-blockers should be considered during pregnancy in women with ARVC.	IIa
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs.	IIa

In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered.	IIb	© ESC 2022
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered.	IIb	

AAD, anti-arrhythmic drug; ACE-Is, angiotensin-converting enzyme inhibitors; AED, automated external defibrillator; ARBs, angiotensin receptor blockers; ARNIs, angiotensin receptor neprilysin inhibitors; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HND, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; LGE, late gadolinium enhancement; LMNA, lamin A/C; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; LVNC, left ventricular non-compaction; MI, myocardial infarction; MRAs, mineralocorticoid receptor antagonists; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OMT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; RCM, restrictive cardiomyopathy; RVOT, right ventricular outflow tract; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SPVT, sustained polymorphic ventricular tachycardia; SQTS, short QT syndrome; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia; WCD, wearable cardioverter defibrillator.

^aShopping malls, stadiums, public transport stations, casinos.

^bVT non-inducibility and elimination of electrograms consistent with conduction delay.

^cIncluding but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry.

^dPresyncope or palpitations suggestive of VA.

^eThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^fBased on the HCM Risk SCD: <https://doc2do.com/hcm/webHCM.html>

^gDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^hBased on the HCM Kid Risk score: <https://hcmriskkids.org>

ⁱFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6-9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^jERP high risk features: J waves > 2 mm, dynamic changes in J point and ST morphology.

^kHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.

Recommendations	Class
Public basic life support and access to AEDs	
It is recommended that public-access defibrillation be available at sites where cardiac arrest is more likely to occur. ^a	I
Prompt CPR by bystanders is recommended at OHCA.	I
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered.	IIa
Treatment of VA. General aspects	
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good-quality survival >1 year.	I
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered.	IIa
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered.	IIb
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb
The WCD may be considered in the early phase after MI in selected patients.	IIb
Coronary artery disease	
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered.	IIa
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT.	IIa
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT and NSVT, if they are inducible for SMVT by PES.	IIa

In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^b	Ila
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blocker or sotalol treatment.	Ila
Idiopathic PVC/VT and PVC-induced cardiomyopathy	
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.	I
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram.	Ila
Beta-blockers, non-dihydropyridine CCBs or flecainide should be considered when catheter ablation is not available, not desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles.	Ila
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	Ila
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered.	Ila
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered.	Ila
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered.	Ila
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up.	IIb
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III
DCM/HNDCM	
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: - the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or - there is a family history of DCM/HNDCM, or premature unexpected SD.	I
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD.	Ila

Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age or with signs suspicious for an inherited aetiology.	IIa
ICD implantation should be considered in DCM/HNDCM patients with an LVEF <50% and ≥ 2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG, and an echocardiogram may be considered.	IIb
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation.	III
ARVC	
In patients with suspected ARVC, CMR is recommended.	I
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and either NSVT or inducibility of SMVT at PES.	IIa
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered.	IIa
Avoidance of high-intensity ^e exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype.	IIb
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification.	IIb
HCM	
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I
Genetic counselling and testing are recommended in HCM patients.	I
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to <6%) ^f , and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF <50%; or (c) abnormal blood pressure response during exercise test ^g ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation.	IIa
In children <16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^h), ICD implantation should be considered.	IIa
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa

In patients with HCM and recurrent, symptomatic VA, or recurrent symptomatic ICD therapy, AAD treatment should be considered.	Ila
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers.	Ilb
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD (<4%), ^f and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF < 50%; or (c) LV apical aneurysm.	Ilb
Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	Ilb
LVNC and RCM	
In patients with an LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/ HNDCM recommendations.	Ila
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	Ila
Neuromuscular diseases	
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA.	I
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration.	Ila
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias, or who are older than 40 years and have significant LGE on CMR	Ila
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered.	Ila
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered.	Ila
In patients with limb-girdle type 1B or Emery–Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered.	Ila
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR.	Ilb
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ⁱ for VA and SCD.	Ilb
In patients with myotonic dystrophy, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III

Inflammatory diseases	
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered.	IIa
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	IIa
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired.	IIa
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	IIa
In patients with cardiac sarcoidosis who have an LVEF >35% but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered.	IIa
In patients with cardiac sarcoidosis who have an LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	IIa
In patients with cardiac sarcoidosis, LVEF 35–50%, and inducible SMVT at PES, ICD implantation should be considered.	IIa
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	IIa
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT.	IIa
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered.	IIa
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^b	IIb
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD recipients with recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb
Congenital heart disease	
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended.	I
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation, and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered.	IIa
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered.	IIb

In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy.	IIb
Idiopathic VF	
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF.	IIa
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF.	IIa
Clinical testing (history, ECG, and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered.	IIb
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered.	IIb
Long QT syndrome	
In patients with clinically diagnosed LQTS, genetic testing, and genetic counselling are recommended.	I
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I
Mexiletine is indicated in LQT3 patients with a prolonged QT interval.	I
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval.	IIa
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients).	IIb
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III
Andersen–Tawil syndrome	
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome.	I
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT.	I
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.	IIa

Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA.	IIa
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT.	IIb
Brugada syndrome	
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS.	I
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of the following: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope.	IIa
BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG.	IIb
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG.	IIb
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III
Catheter ablation in asymptomatic BrS patients is not recommended.	III
Early repolarization syndrome	
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I
In a SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered.	IIa
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^j	IIa
ILR should be considered in individuals with ERP and at least one risk feature ^k or arrhythmic syncope.	IIa
Isoproterenol infusion should be considered for ERS patients with electrical storm.	IIa
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients.	IIa
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment.	IIa
Genetic testing in ERS patients may be considered.	IIb
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^k	IIb

ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^l in the presence of a family history of unexplained juvenile SD.	IIb
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III
CPVT	
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT.	I
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb
Short QT syndrome	
Genetic testing is indicated in patients diagnosed with SQTS.	I
SQTS should be considered in the presence of a QTc ≤ 320 ms.	IIa
SQTS should be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and arrhythmic syncope.	IIa
ILR should be considered in young SQTS patients.	IIa
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa
SQTS may be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and a family history of SD at age <40 years.	IIb
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD.	IIb
Isoproterenol may be considered in SQTS patients with an electrical storm.	IIb
Selected populations	
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I
Continuation of beta-blockers should be considered during pregnancy in women with ARVC.	IIa
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs.	IIa

In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered.	IIb	© ESC 2022
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered.	IIb	

AAD, anti-arrhythmic drug; ACE-Is, angiotensin-converting enzyme inhibitors; AED, automated external defibrillator; ARBs, angiotensin receptor blockers; ARNIs, angiotensin receptor neprilysin inhibitors; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HND, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; LGE, late gadolinium enhancement; LMNA, lamin A/C; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; LVNC, left ventricular non-compaction; MI, myocardial infarction; MRAs, mineralocorticoid receptor antagonists; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OMT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; RCM, restrictive cardiomyopathy; RVOT, right ventricular outflow tract; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SPVT, sustained polymorphic ventricular tachycardia; SQTS, short QT syndrome; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia; WCD, wearable cardioverter defibrillator.

^aShopping malls, stadiums, public transport stations, casinos.

^bVT non-inducibility and elimination of electrograms consistent with conduction delay.

^cIncluding but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry.

^dPresyncope or palpitations suggestive of VA.

^eThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^fBased on the HCM Risk SCD: <https://doc2do.com/hcm/webHCM.html>

^gDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^hBased on the HCM Kid Risk score: <https://hcmriskkids.org>

ⁱFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6-9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^jERP high risk features: J waves > 2 mm, dynamic changes in J point and ST morphology.

^kHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.

Table 4

[Open in new tab](#)

New recommendations in 2022

Recommendations	Class
Public basic life support and access to AEDs	
It is recommended that public-access defibrillation be available at sites where cardiac arrest is more likely to occur. ^a	I
Prompt CPR by bystanders is recommended at OHCA.	I
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered.	IIa
Treatment of VA. General aspects	
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good-quality survival >1 year.	I
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered.	IIa
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered.	IIb
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb
The WCD may be considered in the early phase after MI in selected patients.	IIb
Coronary artery disease	
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered.	IIa
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT.	IIa
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT and NSVT, if they are inducible for SMVT by PES.	IIa

In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^b	Ila
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blocker or sotalol treatment.	Ila
Idiopathic PVC/VT and PVC-induced cardiomyopathy	
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.	I
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram.	Ila
Beta-blockers, non-dihydropyridine CCBs or flecainide should be considered when catheter ablation is not available, not desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles.	Ila
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	Ila
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered.	Ila
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered.	Ila
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered.	Ila
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up.	IIb
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III
DCM/HNDCM	
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: - the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or - there is a family history of DCM/HNDCM, or premature unexpected SD.	I
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD.	Ila

Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age or with signs suspicious for an inherited aetiology.	IIa
ICD implantation should be considered in DCM/HNDCM patients with an LVEF <50% and ≥ 2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG, and an echocardiogram may be considered.	IIb
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation.	III
ARVC	
In patients with suspected ARVC, CMR is recommended.	I
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and either NSVT or inducibility of SMVT at PES.	IIa
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered.	IIa
Avoidance of high-intensity ^e exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype.	IIb
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification.	IIb
HCM	
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I
Genetic counselling and testing are recommended in HCM patients.	I
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to <6%) ^f , and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF <50%; or (c) abnormal blood pressure response during exercise test ^g ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation.	IIa
In children <16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^h), ICD implantation should be considered.	IIa
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa

In patients with HCM and recurrent, symptomatic VA, or recurrent symptomatic ICD therapy, AAD treatment should be considered.	IIa
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers.	IIb
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD (<4%), ^f and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF < 50%; or (c) LV apical aneurysm.	IIb
Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb
LVNC and RCM	
In patients with an LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/ HNDCM recommendations.	IIa
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa
Neuromuscular diseases	
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA.	I
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration.	IIa
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias, or who are older than 40 years and have significant LGE on CMR	IIa
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered.	IIa
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered.	IIa
In patients with limb-girdle type 1B or Emery–Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered.	IIa
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR.	IIb
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ⁱ for VA and SCD.	IIb
In patients with myotonic dystrophy, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III

Inflammatory diseases	
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered.	Ila
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	Ila
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired.	Ila
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis who have an LVEF >35% but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis who have an LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	Ila
In patients with cardiac sarcoidosis, LVEF 35–50%, and inducible SMVT at PES, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	Ila
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT.	Ila
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered.	Ila
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^b	Ilb
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD recipients with recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	Ilb
Congenital heart disease	
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended.	I
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation, and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered.	Ila
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered.	Ilb

In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy.	IIb
Idiopathic VF	
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF.	IIa
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF.	IIa
Clinical testing (history, ECG, and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered.	IIb
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered.	IIb
Long QT syndrome	
In patients with clinically diagnosed LQTS, genetic testing, and genetic counselling are recommended.	I
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I
Mexiletine is indicated in LQT3 patients with a prolonged QT interval.	I
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval.	IIa
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients).	IIb
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III
Andersen–Tawil syndrome	
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome.	I
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT.	I
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.	IIa

Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA.	IIa
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT.	IIb
Brugada syndrome	
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS.	I
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of the following: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope.	IIa
BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG.	IIb
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG.	IIb
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III
Catheter ablation in asymptomatic BrS patients is not recommended.	III
Early repolarization syndrome	
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I
In a SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered.	IIa
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^j	IIa
ILR should be considered in individuals with ERP and at least one risk feature ^k or arrhythmic syncope.	IIa
Isoproterenol infusion should be considered for ERS patients with electrical storm.	IIa
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients.	IIa
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment.	IIa
Genetic testing in ERS patients may be considered.	IIb
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^k	IIb

ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^l in the presence of a family history of unexplained juvenile SD.	IIb
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III
CPVT	
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT.	I
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb
Short QT syndrome	
Genetic testing is indicated in patients diagnosed with SQTS.	I
SQTS should be considered in the presence of a QTc ≤ 320 ms.	IIa
SQTS should be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and arrhythmic syncope.	IIa
ILR should be considered in young SQTS patients.	IIa
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa
SQTS may be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and a family history of SD at age <40 years.	IIb
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD.	IIb
Isoproterenol may be considered in SQTS patients with an electrical storm.	IIb
Selected populations	
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I
Continuation of beta-blockers should be considered during pregnancy in women with ARVC.	IIa
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs.	IIa

In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered.	IIb	© ESC 2022
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered.	IIb	

AAD, anti-arrhythmic drug; ACE-Is, angiotensin-converting enzyme inhibitors; AED, automated external defibrillator; ARBs, angiotensin receptor blockers; ARNIs, angiotensin receptor neprilysin inhibitors; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HND, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; LGE, late gadolinium enhancement; LMNA, lamin A/C; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; LVNC, left ventricular non-compaction; MI, myocardial infarction; MRAs, mineralocorticoid receptor antagonists; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OMT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; RCM, restrictive cardiomyopathy; RVOT, right ventricular outflow tract; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SPVT, sustained polymorphic ventricular tachycardia; SQTS, short QT syndrome; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia; WCD, wearable cardioverter defibrillator.

^aShopping malls, stadiums, public transport stations, casinos.

^bVT non-inducibility and elimination of electrograms consistent with conduction delay.

^cIncluding but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry.

^dPresyncope or palpitations suggestive of VA.

^eThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^fBased on the HCM Risk SCD: <https://doc2do.com/hcm/webHCM.html>

^gDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^hBased on the HCM Kid Risk score: <https://hcmriskkids.org>

ⁱFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6-9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^jERP high risk features: J waves > 2 mm, dynamic changes in J point and ST morphology.

^kHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.

Recommendations	Class
Public basic life support and access to AEDs	
It is recommended that public-access defibrillation be available at sites where cardiac arrest is more likely to occur. ^a	I
Prompt CPR by bystanders is recommended at OHCA.	I
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I
Mobile phone-based alerting of basic life support-trained bystander volunteers to assist nearby OHCA victims should be considered.	IIa
Treatment of VA. General aspects	
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good-quality survival >1 year.	I
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered.	IIa
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered.	IIb
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb
The WCD may be considered in the early phase after MI in selected patients.	IIb
Coronary artery disease	
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered.	IIa
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT.	IIa
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT and NSVT, if they are inducible for SMVT by PES.	IIa

In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^b	Ila
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blocker or sotalol treatment.	Ila
Idiopathic PVC/VT and PVC-induced cardiomyopathy	
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.	I
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram.	Ila
Beta-blockers, non-dihydropyridine CCBs or flecainide should be considered when catheter ablation is not available, not desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles.	Ila
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	Ila
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered.	Ila
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered.	Ila
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered.	Ila
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up.	IIb
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III
DCM/HNDCM	
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: - the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or - there is a family history of DCM/HNDCM, or premature unexpected SD.	I
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD.	Ila

Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age or with signs suspicious for an inherited aetiology.	IIa
ICD implantation should be considered in DCM/HNDCM patients with an LVEF <50% and ≥ 2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG, and an echocardiogram may be considered.	IIb
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation.	III
ARVC	
In patients with suspected ARVC, CMR is recommended.	I
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and either NSVT or inducibility of SMVT at PES.	IIa
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered.	IIa
Avoidance of high-intensity ^e exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype.	IIb
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification.	IIb
HCM	
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I
Genetic counselling and testing are recommended in HCM patients.	I
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to <6%) ^f , and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF <50%; or (c) abnormal blood pressure response during exercise test ^g ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation.	IIa
In children <16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^h), ICD implantation should be considered.	IIa
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa

In patients with HCM and recurrent, symptomatic VA, or recurrent symptomatic ICD therapy, AAD treatment should be considered.	IIa
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers.	IIb
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD (<4%), ^f and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF < 50%; or (c) LV apical aneurysm.	IIb
Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb
LVNC and RCM	
In patients with an LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/HNDCM recommendations.	IIa
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa
Neuromuscular diseases	
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA.	I
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration.	IIa
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias, or who are older than 40 years and have significant LGE on CMR	IIa
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered.	IIa
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered.	IIa
In patients with limb-girdle type 1B or Emery–Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered.	IIa
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR.	IIb
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ⁱ for VA and SCD.	IIb
In patients with myotonic dystrophy, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III

Inflammatory diseases	
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered.	Ila
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	Ila
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired.	Ila
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis who have an LVEF >35% but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis who have an LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	Ila
In patients with cardiac sarcoidosis, LVEF 35–50%, and inducible SMVT at PES, ICD implantation should be considered.	Ila
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	Ila
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT.	Ila
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT, or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered.	Ila
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^b	IIb
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD recipients with recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb
Congenital heart disease	
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended.	I
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation, and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered.	Ila
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered.	IIb

In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy.	IIb
Idiopathic VF	
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF.	IIa
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF.	IIa
Clinical testing (history, ECG, and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered.	IIb
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered.	IIb
Long QT syndrome	
In patients with clinically diagnosed LQTS, genetic testing, and genetic counselling are recommended.	I
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I
Mexiletine is indicated in LQT3 patients with a prolonged QT interval.	I
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval.	IIa
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients).	IIb
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III
Andersen–Tawil syndrome	
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome.	I
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT.	I
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.	IIa

Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA.	IIa
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT.	IIb
Brugada syndrome	
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS.	I
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of the following: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope.	IIa
BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG.	IIb
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG.	IIb
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III
Catheter ablation in asymptomatic BrS patients is not recommended.	III
Early repolarization syndrome	
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I
In a SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered.	IIa
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^j	IIa
ILR should be considered in individuals with ERP and at least one risk feature ^k or arrhythmic syncope.	IIa
Isoproterenol infusion should be considered for ERS patients with electrical storm.	IIa
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients.	IIa
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment.	IIa
Genetic testing in ERS patients may be considered.	IIb
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^k	IIb

ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^l in the presence of a family history of unexplained juvenile SD.	IIb
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III
CPVT	
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT.	I
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb
Short QT syndrome	
Genetic testing is indicated in patients diagnosed with SQTS.	I
SQTS should be considered in the presence of a QTc ≤ 320 ms.	IIa
SQTS should be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and arrhythmic syncope.	IIa
ILR should be considered in young SQTS patients.	IIa
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa
SQTS may be considered in the presence of a QTc ≥ 320 ms and ≤ 360 ms and a family history of SD at age <40 years.	IIb
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD.	IIb
Isoproterenol may be considered in SQTS patients with an electrical storm.	IIb
Selected populations	
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I
Continuation of beta-blockers should be considered during pregnancy in women with ARVC.	IIa
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs.	IIa

In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered.	IIb	© ESC 2022
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered.	IIb	

AAD, anti-arrhythmic drug; ACE-Is, angiotensin-converting enzyme inhibitors; AED, automated external defibrillator; ARBs, angiotensin receptor blockers; ARNIs, angiotensin receptor neprilysin inhibitors; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HND, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; LGE, late gadolinium enhancement; LMNA, lamin A/C; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; LVNC, left ventricular non-compaction; MI, myocardial infarction; MRAs, mineralocorticoid receptor antagonists; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OMT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; RCM, restrictive cardiomyopathy; RVOT, right ventricular outflow tract; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SPVT, sustained polymorphic ventricular tachycardia; SQTS, short QT syndrome; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia; WCD, wearable cardioverter defibrillator.

^aShopping malls, stadiums, public transport stations, casinos.

^bVT non-inducibility and elimination of electrograms consistent with conduction delay.

^cIncluding but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry.

^dPresyncope or palpitations suggestive of VA.

^eThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^fBased on the HCM Risk SCD: <https://doc2do.com/hcm/webHCM.html>

^gDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^hBased on the HCM Kid Risk score: <https://hcmriskkids.org>

ⁱFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6-9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^jERP high risk features: J waves > 2 mm, dynamic changes in J point and ST morphology.

^kHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.

Table 5

[Open in new tab](#)

Changes in recommendations since 2015

	Class	
	2015	2022
Coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	IIa	I
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS.	I	IIa
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks.	IIa	IIb
PVC-induced cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	IIa	I
DCM/HNDCM		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III) and LVEF $\leq 35\%$ after ≥ 3 months of OMT.	I	IIa
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb	IIa
ARVC		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope.	IIb	IIa
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction.	IIb	IIa
Inflammatory diseases		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation is recommended.	IIa	I
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF $\leq 35\%$.	IIb	I
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA.	IIb	I
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF.	IIb	IIa
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered.	IIa	IIb

CHD		
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^a electrophysiologic evaluation, including PES, may be considered.	IIa	IIb
In patients with CHD and recurrent, symptomatic SMVT, or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered.	I	IIa
Primary electrical disease and selected populations		
ICD implantation is recommended in patients with LQTS who are symptomatic ^b while receiving beta-blockers and genotype-specific therapies.	IIa	I
ICD implantation should be considered in patients with CPVT who experience arrhythmic syncope and/or documented bidirectional/PVT while on the highest tolerated beta-blocker dose and on flecainide.	I	IIa
Pre-participation cardiovascular evaluation of competitive athletes should be considered.	I	IIa
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy.	IIb	IIa
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated.	IIb	IIa

© ESC 2022

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ARVC, arrhythmogenic right ventricular cardiomyopathy; AV, atrioventricular; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CHD, congenital heart disease; CPVT, catecholaminergic polymorphic ventricular tachycardia; DCM, dilated cardiomyopathy; HNCDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; LV, left ventricle; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RV, right ventricle; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; TdP, torsade de pointes; TOF, tetralogy of Fallot; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR, QRS duration ≥ 180 ms, and severe QRS fragmentation.

^bArrhythmic syncope or haemodynamically not-tolerated VA.

	Class	
	2015	2022
Coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	Ila	I
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS.	I	Ila
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks.	Ila	Ilb
PVC-induced cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	Ila	I
DCM/HNDCM		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III) and LVEF $\leq 35\%$ after ≥ 3 months of OMT.	I	Ila
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	Ilb	Ila
ARVC		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope.	Ilb	Ila
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction.	Ilb	Ila
Inflammatory diseases		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation is recommended.	Ila	I
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF $\leq 35\%$.	Ilb	I
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA.	Ilb	I
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF.	Ilb	Ila
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered.	Ila	Ilb

CHD		
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^a electrophysiologic evaluation, including PES, may be considered.	IIa	IIb
In patients with CHD and recurrent, symptomatic SMVT, or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered.	I	IIa
Primary electrical disease and selected populations		
ICD implantation is recommended in patients with LQTS who are symptomatic ^b while receiving beta-blockers and genotype-specific therapies.	IIa	I
ICD implantation should be considered in patients with CPVT who experience arrhythmic syncope and/or documented bidirectional/PVT while on the highest tolerated beta-blocker dose and on flecainide.	I	IIa
Pre-participation cardiovascular evaluation of competitive athletes should be considered.	I	IIa
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy.	IIb	IIa
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated.	IIb	IIa

© ESC 2022

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ARVC, arrhythmogenic right ventricular cardiomyopathy; AV, atrioventricular; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CHD, congenital heart disease; CPVT, catecholaminergic polymorphic ventricular tachycardia; DCM, dilated cardiomyopathy; HNCDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; LV, left ventricle; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RV, right ventricle; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; TdP, torsade de pointes; TOF, tetralogy of Fallot; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR, QRS duration ≥ 180 ms, and severe QRS fragmentation.

^bArrhythmic syncope or haemodynamically not-tolerated VA.

Table 5

[Open in new tab](#)

Changes in recommendations since 2015

	Class	
	2015	2022
Coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	Ila	I
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS.	I	Ila
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks.	Ila	Ilb
PVC-induced cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	Ila	I
DCM/HNDCM		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III) and LVEF $\leq 35\%$ after ≥ 3 months of OMT.	I	Ila
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	Ilb	Ila
ARVC		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope.	Ilb	Ila
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction.	Ilb	Ila
Inflammatory diseases		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation is recommended.	Ila	I
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF $\leq 35\%$.	Ilb	I
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA.	Ilb	I
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF.	Ilb	Ila
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered.	Ila	Ilb

CHD		
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^a electrophysiologic evaluation, including PES, may be considered.	IIa	IIb
In patients with CHD and recurrent, symptomatic SMVT, or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered.	I	IIa
Primary electrical disease and selected populations		
ICD implantation is recommended in patients with LQTS who are symptomatic ^b while receiving beta-blockers and genotype-specific therapies.	IIa	I
ICD implantation should be considered in patients with CPVT who experience arrhythmic syncope and/or documented bidirectional/PVT while on the highest tolerated beta-blocker dose and on flecainide.	I	IIa
Pre-participation cardiovascular evaluation of competitive athletes should be considered.	I	IIa
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy.	IIb	IIa
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated.	IIb	IIa

© ESC 2022

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ARVC, arrhythmogenic right ventricular cardiomyopathy; AV, atrioventricular; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CHD, congenital heart disease; CPVT, catecholaminergic polymorphic ventricular tachycardia; DCM, dilated cardiomyopathy; HNDKM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; LV, left ventricle; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RV, right ventricle; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; TdP, torsade de pointes; TOF, tetralogy of Fallot; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR, QRS duration ≥ 180 ms, and severe QRS fragmentation.

^bArrhythmic syncope or haemodynamically not-tolerated VA.

	Class	
	2015	2022
Coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	IIa	I
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS.	I	IIa
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks.	IIa	IIb
PVC-induced cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	IIa	I
DCM/HNDCM		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III) and LVEF $\leq 35\%$ after ≥ 3 months of OMT.	I	IIa
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT, or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated.	IIb	IIa
ARVC		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope.	IIb	IIa
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction.	IIb	IIa
Inflammatory diseases		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation is recommended.	IIa	I
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF $\leq 35\%$.	IIb	I
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA.	IIb	I
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF.	IIb	IIa
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered.	IIa	IIb

CHD		
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^a electrophysiologic evaluation, including PES, may be considered.	IIa	IIb
In patients with CHD and recurrent, symptomatic SMVT, or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered.	I	IIa
Primary electrical disease and selected populations		
ICD implantation is recommended in patients with LQTS who are symptomatic ^b while receiving beta-blockers and genotype-specific therapies.	IIa	I
ICD implantation should be considered in patients with CPVT who experience arrhythmic syncope and/or documented bidirectional/PVT while on the highest tolerated beta-blocker dose and on flecainide.	I	IIa
Pre-participation cardiovascular evaluation of competitive athletes should be considered.	I	IIa
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy.	IIb	IIa
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated.	IIb	IIa

© ESC 2022

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ARVC, arrhythmogenic right ventricular cardiomyopathy; AV, atrioventricular; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CHD, congenital heart disease; CPVT, catecholaminergic polymorphic ventricular tachycardia; DCM, dilated cardiomyopathy; HNCMD, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; LV, left ventricle; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RV, right ventricle; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; TdP, torsade de pointes; TOF, tetralogy of Fallot; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR, QRS duration ≥ 180 ms, and severe QRS fragmentation.

^bArrhythmic syncope or haemodynamically not-tolerated VA.

5.1.3.5. Provocative diagnostic tests

These are summarized in [Table 6](#). Common tests performed are sodium channel blocker testing for Brugada syndrome (BrS) and adenosine test to exclude latent pre-excitation.^{135,136} Epinephrine challenge may be useful in CPVT when exercise cannot be performed. Epinephrine test is not recommended for LQTS due to the high false positive rate and utility of exercise testing.¹³⁷ Coronary vasospasm as a cause of VF in the absence of obstructive coronary diseases/cardiomyopathy can be tested with incremental intracoronary doses of acetylcholine/ergonovine.

Table 6

[Open in new tab](#)

Intravenous provocative diagnostic tests

Diagnostic test	Indication	Protocols Dose/infusion rate/duration	Positive test	Contraindications	Criteria to stop test, counselling and management	Observation times	Location
-----------------	------------	---	---------------	-------------------	---	----------------------	----------

Ajmaline	Family history of BrS or SADS. Resuscitated CA without SHD.	1 mg/kg over 5 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min. Record in standard and high precordial leads over 30 min.	BrS type 1 ECG.	Type I BrS ECG, HF. Precaution if evidence of conduction disease (consider temporary pacing wire).	VT/VF, Type 1 BrS ECG, PVCs, QRS widening >150%. If VT/VF, administer iv isoprenaline, iv sodium bicarbonate.	30 min if negative test; 4 h if positive test.	Cath lab or outpatient testing location with full resuscitation equipment
Flecainide	Same as ajmaline.	2 mg/kg over 10 min (maximum dose 150 mg). Record in standard and high precordial leads over 30 min.	Same as ajmaline.	Same as ajmaline.	Same as ajmaline.	4 h if negative test; 24 h if positive test.	Same as ajmaline.
Epinephrine	CPVT and resuscitated CA with or without SHD when exercise test not feasible. Family history of SADS.	Rest 10 min. Start at 0.025 µg/kg/min for 10 min increase sequentially to 0.05, 0.1 and 0.2 µg/kg/min in 5 min steps.	>3 beats of PVT or bidirectional VT.	QTc prolongation >480 ms.	Systolic blood pressure >200 mmHg, non-sustained VT or PVT, >10 PVCs/min, T-wave alternans, or patient intolerance. If symptoms persist after discontinuation, iv metoprolol 2.5 mg over 1 min.	30 min.	Same as ajmaline.
Acetylcholine	Suspicion of coronary vasospasm.	Intracoronary injection: RCA: 20 and 50 µg. LCA: 20, 50, and 100 µg over 20 s. >3-min intervals between injections. Maximal dose of 50 µg in the RCA and 100 µg in the LCA.	Coronary artery spasm visualized during procedure.	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/IV HF, renal failure, severe bronchial asthma.	Temporary wire for back-up pacing. Risk of cardiogenic shock.	Normal post-procedure observational time.	Cath lab.
Ergonovine	Same as acetylcholine.	Intracoronary stepwise injection: RCA (20-60 µg)	Same as acetylcholine	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA	Temporary wire should be placed for back-up pacing. Risk of	Same as acetylcholine.	Cath lab.

		mg) LCA (20â€”60â€” mg) over a period of 2â€”5â€” min.		III/V HF, renal failure.	cardiogenic shock.		
Adenosine	Exclude latent pre- excitation.	6, 12, 18â€” mg boluses up to maximum dose 24â€” mg or until AV block or pre-excitation occurs.	Identification of accessory pathway.	Asthma, sinus node disease, allergy to adenosine.	Side effects: Bronchospasm, bradycardia, asystole, AF, seizure. Antagonist: theophylline.	5â€” min.	Same as ajmaline.

Diagnostic test	Indication	Protocols Dose/infusion rate/duration	Positive test	Contraindications	Criteria to stop test, counselling and management	Observation times	Location
Ajmaline	Family history of BrS or SADS. Resuscitated CA without SHD.	1 mg/kg over 5â€”10 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min. Record in standard and high precordial leads over 30 min.	BrS type 1 ECG.	Type I BrS ECG, HF. Precaution if evidence of conduction disease (consider temporary pacing wire).	VT/VF, Type 1 BrS ECG, PVCs, QRS widening >150%. If VT/VF, administer iv isoprenaline, iv sodium bicarbonate.	30â€” min if negative test; 4 h if positive test.	Cath lab c outpatient testing location with full resuscitati equipmen
Flecainide	Same as ajmaline.	2â€” mg/kg over 10â€” min (maximum dose 150â€” mg). Record in standard and high precordial leads over 30â€” min.	Same as ajmaline.	Same as ajmaline.	Same as ajmaline.	4 h if negative test; 24 h if positive test.	Same as ajmaline.
Epinephrine	CPVT and resuscitated CA with or without SHD when exercise test not feasible. Family history of SADS.	Rest 10â€” min. Start at 0.025â€” Âµg/ kg/min for 10â€” min increase sequentially to 0.05, 0.1 and 0.2 Âµg/kg/min in 5â€” min steps.	â€” beats of PVT or bidirectional VT.	QTc prolongation â€” ms.	Systolic blood pressure â€” 200â€” mmHg, non- sustained VT or PVT, >10 PVCs /min, T-wave alternans, or patient intolerance. If symptoms persist after discontinuation, iv metoprolol 2.5â€” mg over 1â€” min.	30â€” min.	Same as ajmaline.

Acetylcholine	Suspicion of coronary vasospasm.	Intracoronary injection: RCA: 20 and 50 μ g. LCA: 20, 50, and 100 μ g over 20 s. >3-min intervals between injections. Maximal dose of 50 μ g in the RCA and 100 μ g in the LCA.	Coronary artery spasm visualized during procedure.	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/IV HF, renal failure, severe bronchial asthma.	Temporary wire for back-up pacing. Risk of cardiogenic shock.	Normal post-procedure observational time.	Cath lab.
Ergonovine	Same as acetylcholine.	Intracoronary stepwise injection: RCA (20 μ g) LCA (20 μ g) over a period of 2 μ g/min.	Same as acetylcholine	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/V HF, renal failure.	Temporary wire should be placed for back-up pacing. Risk of cardiogenic shock.	Same as acetylcholine.	Cath lab.
Adenosine	Exclude latent pre-excitation.	6, 12, 18 μ g boluses up to maximum dose 24 μ g or until AV block or pre-excitation occurs.	Identification of accessory pathway.	Asthma, sinus node disease, allergy to adenosine.	Side effects: Bronchospasm, bradycardia, asystole, AF, seizure. Antagonist: theophylline.	5 μ g/min.	Same as ajmaline.

AF, atrial fibrillation; AV, atrioventricular; BrS, Brugada syndrome; CA, cardiac arrest; CPVT, catecholaminergic polymorphic ventricular tachycardia; ECG, electrocardiogram; HF, heart failure; LCA, left coronary artery; NYHA, New York Heart Association; PVCs, premature ventricular complexes; PVT, polymorphic ventricular tachycardia; RCA, right coronary artery; SADS, sudden arrhythmic death syndrome; SHD, structural heart disease; VF, ventricular fibrillation; VT, ventricular tachycardia.

© ESC 2022

Table 6

[Open in new tab](#)

Intravenous provocative diagnostic tests

Diagnostic test	Indication	Protocols Dose/infusion rate/duration	Positive test	Contraindications	Criteria to stop test, counselling and management	Observation times	Location
Ajmaline	Family history of BrS or SADS. Resuscitated CA without SHD.	1 mg/kg over 5 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min. Record in standard and high	BrS type 1 ECG.	Type I BrS ECG, HF. Precaution if evidence of conduction disease (consider temporary pacing wire).	VT/VF, Type 1 BrS ECG, PVCs, QRS widening >150%. If VT/VF, administer iv isoprenaline, iv	30 min if negative test; 4 h if positive test.	Cath lab or outpatient testing location with full resuscitation equipment

		precordial leads over 30 min.			sodium bicarbonate.		
Flecainide	Same as ajmaline.	2â€mg/kg over 10â€min (maximum dose 150â€mg). Record in standard and high precordial leads over 30â€min.	Same as ajmaline.	Same as ajmaline.	Same as ajmaline.	4 h if negative test; 24 h if positive test.	Same as ajmaline.
Epinephrine	CPVT and resuscitated CA with or without SHD when exercise test not feasible. Family history of SADS.	Rest 10â€min. Start at 0.025â€Âµg/kg/min for 10â€min increase sequentially to 0.05, 0.1 and 0.2 Âµg/kg/min in 5â€min steps.	â€3 beats of PVT or bidirectional VT.	QTc prolongation â€480â€ms.	Systolic blood pressure â€200â€mmHg, non-sustained VT or PVT, >10 PVCs /min, T-wave alternans, or patient intolerance. If symptoms persist after discontinuation, iv metoprolol 2.5â€mg over 1â€min.	30â€min.	Same as ajmaline.
Acetylcholine	Suspicion of coronary vasospasm.	Intracoronary injection: RCA: 20 and 50â€Âµg. LCA: 20, 50, and 100â€Âµg over 20â€s. >3-min intervals between injections. Maximal dose of 50â€Âµg in the RCA and 100 Âµg in the LCA.	Coronary artery spasm visualized during procedure.	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/IV HF, renal failure, severe bronchial asthma.	Temporary wire for back-up pacing. Risk of cardiogenic shock.	Normal post-procedure observational time.	Cath lab.
Ergonovine	Same as acetylcholine.	Intracoronary stepwise injection: RCA (20â€60â€mg) LCA (20â€60â€mg) over a period of 2â€5â€min.	Same as acetylcholine	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/V HF, renal failure.	Temporary wire should be placed for back-up pacing. Risk of cardiogenic shock.	Same as acetylcholine.	Cath lab.
Adenosine	Exclude latent pre-	6, 12, 18â€mg boluses up to	Identification of accessory	Asthma, sinus node disease,	Side effects: Bronchospasm,	5â€min.	Same as ajmaline.

	excitation.	maximum dose 240 mg or until AV block or pre-excitation occurs.	pathway.	allergy to adenosine.	bradycardia, asystole, AF, seizure. Antagonist: theophylline.		
Diagnostic test	Indication	Protocols Dose/infusion rate/duration	Positive test	Contraindications	Criteria to stop test, counselling and management	Observation times	Location
Ajmaline	Family history of BrS or SADS. Resuscitated CA without SHD.	1 mg/kg over 5 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min. Record in standard and high precordial leads over 30 min.	BrS type 1 ECG.	Type I BrS ECG, HF. Precaution if evidence of conduction disease (consider temporary pacing wire).	VT/VF, Type 1 BrS ECG, PVCs, QRS widening >150%. If VT/VF, administer iv isoprenaline, iv sodium bicarbonate.	30 min if negative test; 4 h if positive test.	Cath lab or outpatient testing location with full resuscitation equipment
Flecainide	Same as ajmaline.	2 mg/kg over 10 min (maximum dose 150 mg). Record in standard and high precordial leads over 30 min.	Same as ajmaline.	Same as ajmaline.	Same as ajmaline.	4 h if negative test; 24 h if positive test.	Same as ajmaline.
Epinephrine	CPVT and resuscitated CA with or without SHD when exercise test not feasible. Family history of SADS.	Rest 10 min. Start at 0.025 µg/kg/min for 10 min increase sequentially to 0.05, 0.1 and 0.2 µg/kg/min in 5 min steps.	≥3 beats of PVT or bidirectional VT.	QTc prolongation ≥480 ms.	Systolic blood pressure ≥200 mmHg, non-sustained VT or PVT, >10 PVCs/min, T-wave alternans, or patient intolerance. If symptoms persist after discontinuation, iv metoprolol 2.5 mg over 1 min.	30 min.	Same as ajmaline.
Acetylcholine	Suspicion of coronary vasospasm.	Intracoronary injection: RCA: 20 and 50 µg. LCA: 20, 50, and 100 µg over 20 s. >3-min intervals	Coronary artery spasm visualized during procedure.	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/IV HF, renal failure, severe bronchial asthma.	Temporary wire for back-up pacing. Risk of cardiogenic shock.	Normal post-procedure observational time.	Cath lab.

		between injections. Maximal dose of 50 µg in the RCA and 100 µg in the LCA.					
Ergonovine	Same as acetylcholine.	Intracoronary stepwise injection: RCA (20 µg) LCA (20 µg) over a period of 2 min.	Same as acetylcholine	Left main stenosis >50%, 3-vessel disease, 2-vessel disease with total occlusion, NYHA III/V HF, renal failure.	Temporary wire should be placed for back-up pacing. Risk of cardiogenic shock.	Same as acetylcholine.	Cath lab.
Adenosine	Exclude latent pre-excitation.	6, 12, 18 µg boluses up to maximum dose 24 µg or until AV block or pre-excitation occurs.	Identification of accessory pathway.	Asthma, sinus node disease, allergy to adenosine.	Side effects: Bronchospasm, bradycardia, asystole, AF, seizure. Antagonist: theophylline.	5 min.	Same as ajmaline.

AF, atrial fibrillation; AV, atrioventricular; BrS, Brugada syndrome; CA, cardiac arrest; CPVT, catecholaminergic polymorphic ventricular tachycardia; ECG, electrocardiogram; HF, heart failure; LCA, left coronary artery; NYHA, New York Heart Association; PVCs, premature ventricular complexes; PVT, polymorphic ventricular tachycardia; RCA, right coronary artery; SADS, sudden arrhythmic death syndrome; SHD, structural heart disease; VF, ventricular fibrillation; VT, ventricular tachycardia.

© ESC 2022

5.1.3.6. Electrophysiological study

Electrophysiological studies including measurement of baseline intervals (e.g. atrial-His interval [AH] and His-ventricular interval [HV]), programmed electrical stimulation (PES), and electroanatomical mapping can be used for diagnostic purposes and to guide therapy.^{145,150} The yield of PES varies with the underlying cardiac condition and its severity, the presence or absence of spontaneous VT, concomitant drug therapy, stimulation protocol, and site(s) of stimulation. Typical protocols include stimulation from 2 right ventricular (RV) sites with 2 basic drive cycle lengths, introduction of 3 extra-stimuli, and isoprenaline administration.^{148,151,152}

In the current era, PES is mainly employed to confirm the diagnosis of VT and induce mappable VAs with non-inducibility being an ablation endpoint. Patients with heart failure and LVEF ≤35% generally will have an indication for an ICD; therefore, VT/VF induction before implantation is not necessary. In patients with SHD and mildly reduced or preserved LVEF who present with unexplained syncope, induction of SMVT with PES can be helpful to identify the underlying cause and to predict subsequent events.^{146,153} PVT/VF induction in SHD is in general considered as a non-specific finding.⁴⁶

In primary electrical diseases, PES is not of prognostic value, although there is some evidence to consider its use in BrS.¹²⁷ Invasive electrophysiological evaluation can have important clinical implications in patients with myotonic dystrophy.¹⁵⁷

With advances in high-density mapping, voltage mapping, conduction/repolarisation metrics, and electrogram fractionation can be employed to identify ablation targets, or to diagnose cardiomyopathic disease. Endocardial mapping may be helpful in the differentiation of ARVC from benign outflow tract VT and for targeting biopsy in suspected myocarditis, ARVC, and sarcoidosis cases.^{158,162}

5.1.4. Genetic testing

Massive parallel or next-generation sequencing has led to increasing availability of genetic testing at reduced cost. Most diagnostic cardiac genetic testing employs large gene panels determined by associations with disease generated

by prior research, i.e. candidate genes.¹⁶³ Many previous gene associations have, however, been challenged for their diagnostic utility. Therefore, it is not recommended to include questionable genes in routine diagnostic panels.^{164,165,168} Genome-wide association studies have identified that common genetic variation single nucleotide polymorphisms can cause or modify phenotypes in BrS, LQTS, HCM, and DCM. Polygenic risk scores, measures derived from the cumulative effects of these single nucleotide polymorphisms, may therefore play a role in diagnosis and prognostication in these conditions in the future.^{168,169,173}

Sequencing produces digital data that require subsequent bioinformatic analysis, permitting accurate ascertainment of most DNA alterations affecting the coding frame of each gene.¹⁷⁴ The most common findings are single nucleotide variants causing simple amino acid substitutions (missense), premature terminations, or splicing abnormalities. Insertions and/or deletions are rarer. The clinical importance of most non-coding variants is still to be determined.^{174,175}

A framework for the interpretation of disease causation by genetic variants has standardized adjudication into five classes: V a pathogenic, IV a likely pathogenic, III a variant of uncertain significance, II a likely benign, and I a benign. A combination of evidence is employed: gene-disease association; presence of a variant in healthy and/or diseased populations; *in silico* data; *in vitro* and *in vivo* functional data; and family segregation data.¹⁷⁶

A mutation (Class IV or V variant) can be used immediately either for confirmation of diagnosis in probands (the first affected family member), or for initial diagnosis of relatives and may help to guide therapy and/or prognosis. Periodic reassessment of all class IV and III variants is indicated.¹⁷⁶

Pre-implantation genetic testing is an early form of pre-natal genetic diagnosis. Genetic diagnoses of *in vitro* fertilized embryos are identified by biopsies, thereby allowing transfer of genetically normal embryos into the uterus. If the technique is available, it is important to provide information to patients with monogenic heart diseases in child-bearing age. The legislation for pre-implantation genetic testing varies in different countries and strategies differ.

Genetic and clinical testing should be undertaken only by multidisciplinary teams including professionals with skills to counsel on the implications and the uncertainty of results and experienced cardiologists able to direct testing to the correct phenotype.^{135,177,179} A negative result does not exclude a diagnosis and should not be used for this purpose. A framework for genetic and other clinical diagnostic tests for primary electrical diseases based on evidence where available has been provided in [Table 7](#).

Table 7

[Open in new tab](#)

Genetic tests and suggested work-up of probands and relatives with primary electrical diseases

			LQTS	BrS	CPVT	Idiopathic VF	ERS
Proband	Genetic test	Cornerstone for diagnosis	Class I ^a	Class I	Class I ^a	Class II ^a	Class II ^a
			ECG Exercise test Exclude acquired LQTS	ECG and high precardial lead ECG Sodium channel blockers provocative test ^b Exclude phenocopy ^c	Exercise test Exclude phenocopy/SHD	See Section 5.2.3, scenario 3	ECG
Relatives	Follow-up	Clinical screening	ECG Exercise test (when feasible) From birth	ECG and high precardial lead ECGs start at 10 years Sodium channel blockers provocative test ^b start >16 years unless clinically indicated ^{b(1)(2)}	ECG Exercise test From birth	ECG and high precardial lead ECGs Exercise test Echocardiogram ^{b(1)}	ECG Echocardiogram
			Positive phenotype and/or Class IV variant Negative phenotype and no Class IV variant	1-3 years dependent on level of risk Discharge			

BrS, Brugada syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; ECG, electrocardiogram; ERS, early repolarization syndrome; LQTS, long QT syndrome; VF, ventricular fibrillation.
^aIncluding neonatal genetic testing.
^bA phenocopy has characteristics of a genetic disease but is produced environmentally.
^cNot in case of a documented type 1 Brugada pattern.

			LQTS	BrS	CPVT	Idiopathic VF	ERS
Proband	Genetic test	Cornerstone for diagnosis	Class I ^a	Class I	Class I ^a	Class II ^a	Class II ^a
			ECG Exercise test Exclude acquired LQTS	ECG and high precardial lead ECG Sodium channel blockers provocative test ^b Exclude phenocopy ^c	Exercise test Exclude phenocopy/SHD	See Section 5.2.3, scenario 3	ECG
Relatives	Follow-up	Clinical screening	ECG Exercise test (when feasible) From birth	ECG and high precardial lead ECGs start at 10 years Sodium channel blockers provocative test ^b start >16 years unless clinically indicated ^{b(1)(2)}	ECG Exercise test From birth	ECG and high precardial lead ECGs Exercise test Echocardiogram ^{b(1)}	ECG Echocardiogram
			Positive phenotype and/or Class IV variant Negative phenotype and no Class IV variant	1-3 years dependent on level of risk Discharge			

BrS, Brugada syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; ECG, electrocardiogram; ERS, early repolarization syndrome; LQTS, long QT syndrome; VF, ventricular fibrillation.
^aIncluding neonatal genetic testing.
^bA phenocopy has characteristics of a genetic disease but is produced environmentally.
^cNot in case of a documented type 1 Brugada pattern.

Table 7

[Open in new tab](#)

Genetic tests and suggested work-up of probands and relatives with primary electrical diseases

			LQTS	BrS	CPVT	Idiopathic VF	ERS
Proband	Genetic test		Class P ^a	Class I	Class P ^a	Class B ^a	Class B ^a
	Initial clinical test	Cornerstone for diagnosis	ECG Exercise test	ECG and high precardial lead ECG Sodium channel blockers provocative test ^b	Exercise test	See Section 3.2.3, scenario 3	ECG
	Other tests/processes		Exclude acquired LQTS	Exclude phenocopy ^b	Exclude phenocopy/SHD		Holter Echocardiography
Relatives	Follow-up		ECG	ECG and high precardial lead ECG; start at 10 years Sodium channel blockers provocative test ^b ; start >16 years unless clinically indicated ^{b,c,d}	1-3 years dependent on level of risk	ECG and high precardial lead ECGs	ECG
	Clinical screening		Exercise test (when feasible) From birth	Exercise test (when feasible) From birth	ECG Exercise test From birth	ECG and high precardial lead ECGs Exercise test Echocardiogram ^{b,d}	Echocardiogram
	Follow-up	Positive phenotype and/or Class R/V variant Negative phenotype and no Class R/V variant		1-3 years dependent on level of risk			
			Discharge				

BrS, Brugada syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; ECG, electrocardiogram; ERS, early repolarisation syndrome; LQTS, long QT syndrome; VF, ventricular fibrillation.

^aIncluding neonatal genetic testing.

^bA phenocopy has characteristics of a genetic disease but is produced environmentally.

^cNot in case of a documented type 1 Brugada pattern.

© ESC 2023

			LQTS	BrS	CPVT	Idiopathic VF	ERS	
Proband	Genetic test		Class P ^a	Class I	Class P ^a	Class B ^a	Class B ^a	
	Initial clinical test	Cornerstone for diagnosis	ECG	ECG and high precardial lead ECG	Exercise test	See Section 3.2.3, scenario 3	ECG	
	Other tests/processes	Exercise test Exclude acquired LQTS	Sodium channel blockers provocative test ^b Exclude phenocopy ^c		Exclude phenocopy/SHD		Holter Echocardiography	
Relatives	Follow-up			1-3 years dependent on level of risk				
	Clinical screening		ECG	ECG and high precardial lead ECG; start at 10 years	ECG	ECG and high precardial lead ECGs	ECG	
		Exercise test (when feasible) From birth	Sodium channel blockers provocative test ^b ; start >16 years unless clinically indicated ^{b,c,d}	Exercise test From birth	Exercise test From birth	Exercise test Echocardiogram ^{b,d}	Echocardiogram	
	Follow-up	Positive phenotype and/or Class R/V variant Negative phenotype and no Class R/V variant	1-3 years dependent on level of risk					© ESC 2022
			Discharge					

BrS, Brugada syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; ECG, electrocardiogram; ERS, early repolarisation syndrome; LQTS, long QT syndrome; VF, ventricular fibrillation.
^aIncluding neonatal genetic testing.
^bA phenocopy has characteristics of a genetic disease but is produced environmentally.
^cNot in case of a documented type 1 Brugada pattern.

Recommendation Table 2

[Open in new tab](#)

Recommendations for genetic testing

Recommendations	Class ^a	Level ^b
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD. ^{56,183}	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework. ¹⁷⁶	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team. ¹⁷⁹	I	C
It is recommended that Class III (variants of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C

© ESC 2022

SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD. ^{56,183}	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework. ¹⁷⁶	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team. ¹⁷⁹	I	C
It is recommended that Class III (variants of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C

SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 2

[Open in new tab](#)

Recommendations for genetic testing

Recommendations	Class ^a	Level ^b
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD. ^{56,183}	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework. ¹⁷⁶	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team. ¹⁷⁹	I	C
It is recommended that Class III (variants of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C

SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD. ^{56,183}	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework. ¹⁷⁶	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team. ¹⁷⁹	I	C
It is recommended that Class III (variants of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C

SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.


© ESC 2022

5.2. Diagnostic evaluation at first presentation with ventricular arrhythmia in patients without known cardiac disease

VA and (aborted) SCD are common first manifestations of a previously not known cardiac condition. A comprehensive diagnostic evaluation is provided for five frequently encountered scenarios.

5.2.1. Scenario 1: Incidental finding of a non-sustained ventricular tachycardia

An algorithm for the evaluation of patients presenting with an incidental finding of NSVT is presented in [FigureÂ 2](#).



FigureÂ 2

Algorithm for the evaluation of patients presenting with an incidental finding of non-sustained ventricular tachycardia.

CAD, coronary artery disease; CMR, cardiac magnetic resonance; ECG, electrocardiogram; N, No; NSVT, non-sustained ventricular tachycardia; PVC, premature ventricular complex; SCD, sudden cardiac death; SHD, structural heart disease; Y, Yes. ^aECG morphology suggestive of RVOT or fascicular origin, negative family history, normal 12-lead ECG, and echocardiogram. ^be.g. atrioventricular conduction abnormalities, Q waves, broad QRS complex, ST/T waves deviations, abnormally high or low voltages. Ventricular dysfunction/dilatation/hypertrophy/wall thinning, wall motion abnormalities, multitopic PVCs/NSVTs/increasing ventricular arrhythmia (VA) burden with exercise. ^ce.g. Brugada pattern, long/short QT, polymorphic/bidirectional VA with exercise. ^dDiagnostic test to exclude CAD according to patient profile and symptoms. ^eConsider re-evaluation in case of new symptoms or changes in patient clinical condition.

[Open in new tab](#)[Download slide](#)

Incidental NSVT is a common finding during routine cardiological evaluation (e.g. for non-cardiac diseases, pre-initiation of oncological treatments, pre-participation in sports) and monitoring before induction of anaesthesia/sedation for non-cardiac procedures.¹⁸⁴ Patients with incidentally found NSVT require further evaluation. A recent syncope suspicious for cardiac origin is a high-risk symptom and may prompt admission to hospital.^{1,185} The morphology of NSVT (polymorphic or monomorphic) is important to assess. Typical MVT morphologies ([FigureÂ 3](#)) can suggest an idiopathic origin with favourable prognosis. In contrast, short-coupled PVC initiating non-sustained PVT or monomorphic NSVT with short cycle length (usually <300 ms, average 245 ± 28, in one series) may identify patients at higher risk of SCD.^{186,187} Resting 12-lead ECG is a first-line evaluation and may show signs of SHD or primary electrical diseases ([ESC CardioMed chapter 8.6](#)).¹⁸⁸ Echocardiography is the first-line imaging modality that provides important information about cardiac function and potential SHD ([ESC CardioMed 10.3, 10.10, 10.12](#)).^{120,189,190} Holter monitoring is useful to assess the frequency of NSVT and related PVCs ([ESC CardioMed chapter 8.9](#)).¹⁹¹ In addition, an at least 3-lead Holter (V1, two inferior leads) may give a first estimate if NSVT/PVC are unifocal or multifocal and of the NSVT site(s) of origin. The latter is important if the NSVT has not been previously documented on a 12-lead ECG.¹⁹²

An exercise test can be helpful to capture the 12-lead ECG of NSVT and to identify exercise-induced arrhythmias. Increasing arrhythmias with exercise, not suggestive of idiopathic origin, should raise suspicion of SHD and may necessitate advice to refrain from physical exercise until diagnosis and initiation of appropriate treatment. Underlying significant CAD should be ruled out according to the patient's pre-test probability.

CMR should be considered when cardiomyopathies or inflammatory diseases are suspected on initial evaluation (ESC CardioMed chapter 10.4).¹⁹³ In addition, CMR can identify areas of fibrosis as substrates of NSVT.¹²⁹

Recommendation Table 3

[Open in new tab](#)

Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmia

Recommendations	Class ^a	Level ^b
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
In patients with newly documented VA (frequent PVCs, NSVT, SMVT) and suspicion of SHD other than CAD after initial evaluation, a CMR should be considered. ^{194,195}	IIa	B
In patients with an incidental finding of a NSVT, a ≥ 24 h Holter ECG should be considered.	IIa	C

CAD, coronary artery disease; CMR, cardiac magnetic resonance; ECG, electrocardiogram; NSVT, non-sustained ventricular tachycardia; PVCs, premature ventricular contractions; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
In patients with newly documented VA (frequent PVCs, NSVT, SMVT) and suspicion of SHD other than CAD after initial evaluation, a CMR should be considered. ^{194,195}	IIa	B
In patients with an incidental finding of a NSVT, a ≥ 24 h Holter ECG should be considered.	IIa	C

CAD, coronary artery disease; CMR, cardiac magnetic resonance; ECG, electrocardiogram; NSVT, non-sustained ventricular tachycardia; PVCs, premature ventricular contractions; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 3

[Open in new tab](#)

Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmia

Recommendations	Class ^a	Level ^b
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
In patients with newly documented VA (frequent PVCs, NSVT, SMVT) and suspicion of SHD other than CAD after initial evaluation, a CMR should be considered. ^{194,195}	IIa	B
In patients with an incidental finding of a NSVT, a ≥ 24 h Holter ECG should be considered.	IIa	C

CAD, coronary artery disease; CMR, cardiac magnetic resonance; ECG, electrocardiogram; NSVT, non-sustained ventricular tachycardia; PVCs, premature ventricular contractions; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
In patients with newly documented VA (frequent PVCs, NSVT, SMVT) and suspicion of SHD other than CAD after initial evaluation, a CMR should be considered. ^{194,195}	IIa	B
In patients with an incidental finding of a NSVT, a ≥ 24 h Holter ECG should be considered.	IIa	C

CAD, coronary artery disease; CMR, cardiac magnetic resonance; ECG, electrocardiogram; NSVT, non-sustained ventricular tachycardia; PVCs, premature ventricular contractions; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

5.2.2. Scenario 2: First presentation of sustained monomorphic ventricular tachycardia

An algorithm for the evaluation of patients presenting with a first SMVT episode is presented in [FigureÂ 4](#).




Figure 3

Typical idiopathic ventricular tachycardia morphologies.

LBBB, left bundle branch block; LV, left ventricle; RBBB, right bundle branch block; RVOT, right ventricular outflow tract; VT, ventricular tachycardia.

[Open in new tab](#)[Download slide](#)





Figure 4

Algorithm for the evaluation of patients presenting with a first sustained monomorphic ventricular tachycardia episode.

ARVC, arrhythmogenic RV cardiomyopathy; CAD, coronary artery disease; CAG, coronary angiography; CMR, cardiac magnetic resonance; ECG, electrocardiogram; EP, electrophysiological; LV, left ventricular; N, No; PET-CT, positron emission tomography and computed tomography; PVC, premature ventricular complex; RV, right ventricular; SCD, sudden cardiac death; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia; Y, Yes. ^aECG morphology suggestive of RV outflow tract or fascicular origin, negative family history, normal 12-lead ECG, and echocardiogram. ^be.g. Q waves, QRS fragmentation, ST/T abnormalities, wall motion abnormalities in coronary territories. ^ce.g. atrioventricular (AV) conduction abnormalities, Q waves, broad QRS complex, T-wave inversion, abnormally high or low voltages. Ventricular dysfunction/dilatation/hypertrophy/wall thinning/wall motion abnormalities/diffuse hypokinesia. ^dDiagnostic test to exclude CAD according to patient profile and symptoms. ^eAccording to revised task force criteria.¹¹⁶ ^fe.g. AV conduction abnormalities, abnormally high or low voltages, broad QRS, ST/T wave deviations, LV dilatation and dysfunction, late gadolinium enhancement with non-ischaemic distribution. ^ge.g. AV conduction abnormalities, broad QRS, ST/T deviations, multifocal PVCs, inflammatory hyperaemia and oedema, fibrosis, LV and RV systolic dysfunction, pericardial effusion.

[Open in new tab](#)[Download slide](#)

The majority of patients presenting with SMVT have underlying SHD. SMVT in SHD is mainly due to scar-related re-entry and only occasionally due to re-entry involving a diseased conduction system or due to focal sources. The diagnosis of the underlying aetiology and identification of patients with idiopathic VT is important. Initial evaluation includes a comprehensive clinical and family history, 12-lead ECG, and echocardiography. Recording of the 12-lead VT ECG is indicated as it provides important information on the VT site of origin. Specific VT morphologies (e.g. right ventricular outflow tract [RVOT] or fascicular origin) ([FigureÂ 3](#)) in the absence of a family history for cardiomyopathies and without evidence for SHD are suggestive for idiopathic VTs.¹⁹⁶ Atypical ECG morphologies and uncommon clinical presentations should raise suspicions for underlying SHD even if baseline ECG and echocardiogram are normal. In this scenario, additional evaluation with CMR should be considered.¹⁹⁴ Bundle branch re-entrant ventricular tachycardia (BBR-VT), resembling bundle branch block configuration on the ECG, is a feature of conduction impairment, e.g. in DCM, myotonic dystrophy, and post-cardiac valve surgery ([FigureÂ 5](#)).



FigureÂ 5

Bundle branch re-entrant ventricular tachycardia.

BBR-VT, bundle branch re-entrant ventricular tachycardia; ECG, electrocardiogram.

[Open in new tab](#)[Download slide](#)

If initial evaluation raises suspicion of underlying CAD, a CAG can exclude significant CAD. If ECG and echocardiography are suggestive for a cardiomyopathy, CMR provides important diagnostic information on scar distribution and tissue characteristics ([Section 5.1.3.4](#)). When non-invasive evaluation is inconclusive, electroanatomical mapping and PES may be considered for the differential diagnosis between idiopathic VT and early ARVC.¹⁹⁷ Electroanatomical mapping-guided biopsy can be of value to provide a tissue diagnosis for ARVC and inflammatory diseases with a focal distribution (e.g. cardiac sarcoidosis).^{198,199} In cases of suspected inflammatory

diseases, positron emission tomography CT (PET-CT), autoimmune serology, and biopsies of affected tissue are part of the diagnostic evaluation.[200,201](#)

Recommendation Table 4

[Open in new tab](#)

Recommendations for evaluation of patients presenting with a first episode of sustained monomorphic ventricular tachycardia

Recommendations	Class ^a	Level ^b
In patients presenting with a first SMVT episode, electrophysiological study, electroanatomical mapping, and mapping-guided biopsies may be considered for aetiological evaluation. 197–199,202	IIb	C

SMVT, sustained monomorphic ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
In patients presenting with a first SMVT episode, electrophysiological study, electroanatomical mapping, and mapping-guided biopsies may be considered for aetiological evaluation. 197–199,202	IIb	C

SMVT, sustained monomorphic ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 4

[Open in new tab](#)

Recommendations for evaluation of patients presenting with a first episode of sustained monomorphic ventricular tachycardia

Recommendations	Class ^a	Level ^b
In patients presenting with a first SMVT episode, electrophysiological study, electroanatomical mapping, and mapping-guided biopsies may be considered for aetiological evaluation. 197–199,202	IIb	C


SMVT, sustained monomorphic ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
In patients presenting with a first SMVT episode, electrophysiological study, electroanatomical mapping, and mapping-guided biopsies may be considered for aetiological evaluation. 197–199,202	IIb	C

SMVT, sustained monomorphic ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

5.2.3. Scenario 3: Sudden cardiac arrest survivor


An algorithm for the evaluation of sudden cardiac arrest survivors is presented in [Figure 6](#).



FigureÂ 6

Part One. Algorithm for the evaluation of sudden cardiac arrest survivors.

[Open in new tab](#)[Download slide](#)



FigureÂ 6

Part Two. Algorithm for the evaluation of sudden cardiac arrest survivors.

AF, atrial fibrillation; CAD, coronary artery disease; CAG, coronary angiogram; CMR, cardiac magnetic resonance; CT, computed tomography; ECG, electrocardiogram; LGE, late gadolinium enhancement; N, No; SCA, sudden cardiac arrest; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; VF, ventricular fibrillation; Y, Yes. ^aThe 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation.³ ^bRule out SHD according to patient age and characteristics; QT duration needs to be reassessed several days after arrest. ^cConsider cardiac CT/CAG depending on patient characteristics and clinical context. ^dLeft ventricular function on echocardiogram needs to be reassessed several days after arrest to exclude stunning as cause of systolic dysfunction. ^eIn case of high clinical suspicion (typical symptoms and transient ST elevation during monitoring), it can be considered to test for coronary vasospasm earlier.

[Open in new tab](#)[Download slide](#)

Urgent CAG is recommended for patients presenting with ST elevation myocardial infarction (STEMI).^{203a,206} Despite disparate findings of pooled data analyses,^{207a,211} three randomized controlled trials (RCTs) have found no significant benefit for early CAG in cardiac arrest (CA) without ST-elevation. In case of electrical instability after CA, suspicious for ongoing ischaemia, this panel found a CAG indicated. Brain and chest CT scan may acutely identify non-cardiac causes of aborted SD,²¹² as well as blood test results for pertinent toxicological analysis.^{213a,215} Retention and storage of suitable blood samples will allow subsequent diagnostic evaluation, including DNA analysis.²¹³ Any ECG tracing from emergency services, as well as recordings from interrogation of cardiovascular implantable electronic devices (CIEDs) can also contribute to diagnosis.^{216a,219} The resting 12-lead ECG (including high precordial lead)²²⁰ is fundamental and should be repeated regularly during recovery. Continuous heart rhythm monitoring is recommended until definite treatment.^{221,222} Echocardiography may allow early diagnosis to identify any structural abnormality.^{222,223} Coronary imaging will be important to exclude CAD, dissection, or anomalies.^{62,224} Coronary optical coherence tomography and/or intravascular ultrasound may be helpful to characterize stenosis/plaque stability and underlying mechanism of stenosis.²²⁵ CMR has repeatedly been shown to provide significant incremental diagnostic value, in particular for concealed cardiomyopathy.^{131,226a,228} Primary electrical diseases may be uncovered by provocative manoeuvres such as sodium channel blocker challenge,^{136,229a,231} lying to standing ECGs,^{232,233} adenosine challenge,^{144,234} epinephrine challenge,^{141,152,235a,239} ergonovine/acetylcholine,^{222,240} mental stress,^{241,242} and exercise testing.^{116,117,119,232,243} Electrophysiological study and electroanatomic mapping may be useful to provide patient-specific insights into the mechanism of CA and to offer therapeutic options in some patients.^{244a,248} Genetic testing may identify a molecular cause of SCA by identifying pathogenic mutations in genes associated with specific phenotypes.^{213,249,250}

Recommendation Table 5

[Open in new tab](#)

Recommendations for evaluation of sudden cardiac arrest survivors

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team. ^{177,251–256}	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C

In SCA survivors, brain/chest CT scan should be considered when patient characteristics, ECG, and echocardiography are not consistent with a cardiac cause. ^{212,257}	IIa	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing. ^{56,214}	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors. ^{217,218}	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG), as well as continuous cardiac monitoring, are recommended. ^{220,222}	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause. ^{62,222,223,226}	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause. ^{117,222,258–260}	I	B
In SCA survivors, ergonovine, acetylcholine, or hyperventilation testing may be considered for the diagnosis of coronary vasospasm. ^{240,261}	IIb	B

© ESC 2022

CIEDs, cardiac insertable electronic devices; CMR, cardiac magnetic resonance; CT, computed tomography; ECG, electrocardiogram; LGE, late gadolinium enhancement; SCA, sudden cardiac arrest.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team. ^{177,251–256}	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C

In SCA survivors, brain/chest CT scan should be considered when patient characteristics, ECG, and echocardiography are not consistent with a cardiac cause. ^{212,257}	IIa	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing. ^{56,214}	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors. ^{217,218}	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG), as well as continuous cardiac monitoring, are recommended. ^{220,222}	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause. ^{62,222,223,226}	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause. ^{117,222,258–260}	I	B
In SCA survivors, ergonovine, acetylcholine, or hyperventilation testing may be considered for the diagnosis of coronary vasospasm. ^{240,261}	IIb	B

© ESC 2022

CIEDs, cardiac insertable electronic devices; CMR, cardiac magnetic resonance; CT, computed tomography; ECG, electrocardiogram; LGE, late gadolinium enhancement; SCA, sudden cardiac arrest.

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table 5

[Open in new tab](#)

Recommendations for evaluation of sudden cardiac arrest survivors

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team. ^{177,251–256}	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C

In SCA survivors, brain/chest CT scan should be considered when patient characteristics, ECG, and echocardiography are not consistent with a cardiac cause. ^{212,257}	IIa	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing. ^{56,214}	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors. ^{217,218}	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG), as well as continuous cardiac monitoring, are recommended. ^{220,222}	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause. ^{62,222,223,226}	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause. ^{117,222,258–260}	I	B
In SCA survivors, ergonovine, acetylcholine, or hyperventilation testing may be considered for the diagnosis of coronary vasospasm. ^{240,261}	IIb	B

© ESC 2022

CIEDs, cardiac insertable electronic devices; CMR, cardiac magnetic resonance; CT, computed tomography; ECG, electrocardiogram; LGE, late gadolinium enhancement; SCA, sudden cardiac arrest.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team. ^{177,251–256}	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C

In SCA survivors, brain/chest CT scan should be considered when patient characteristics, ECG, and echocardiography are not consistent with a cardiac cause. ^{212,257}	IIa	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing. ^{56,214}	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors. ^{217,218}	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG), as well as continuous cardiac monitoring, are recommended. ^{220,222}	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause. ^{62,222,223,226}	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause. ^{117,222,258–260}	I	B
In SCA survivors, ergonovine, acetylcholine, or hyperventilation testing may be considered for the diagnosis of coronary vasospasm. ^{240,261}	IIb	B

© ESC 2022

CIEDs, cardiac insertable electronic devices; CMR, cardiac magnetic resonance; CT, computed tomography; ECG, electrocardiogram; LGE, late gadolinium enhancement; SCA, sudden cardiac arrest.

^aClass of recommendation.

^bLevel of evidence.

5.2.4. Scenario 4: Sudden death victim

An algorithm for the evaluation of SD victims is presented in [Figure 7](#).




Figure 7

Algorithm for the evaluation of sudden death victims.

SADS, sudden arrhythmic death syndrome; SCD, sudden cardiac death; SD, sudden death.

^aAutopsy is recommended, ideally in all cases of unexpected SD and always in those under 50 years. Autopsy should include full macroscopic examination and histopathology of all organs. The heart should ideally be examined by an expert cardiac pathologist. Samples suitable for DNA extraction should be retained when inherited causes or unexplained death are suspected. ^bBased on all circumstances, this includes negative autopsies, autopsies with uncertain findings, non-ischaemic cardiomyopathies, coronary artery disease where familial hypercholesterolaemia is suspected and thoracic aortic dissections. ^cAfter informed consent of relatives.

[Open in new tab](#)[Download slide](#)

Potential genetic cardiac disease can be identified in 25%–49% of cases of SCD in the young (<50 years of age). This may also affect relatives of the deceased.^{25, 56, 59} To find the cause of death, it is important to collect all available data on prior symptoms, comorbidities, and family history.^{25, 56, 215, 262, 263}

The main role of autopsy in SD is to establish the cause of death. An expert cardiac pathologist alters the initial diagnosis in 41% of cases, highlighting the need for expert evaluation.^{263a, 265} Inherited cardiac diseases identified at autopsy include cardiomyopathies (HCM, DCM, ARVC) and premature CAD.^{25, 27, 56, 266} A toxicology screen can reveal drug overdose or polypharmacy in 31%–56% of young SD cases.^{267, 268} In autopsy-negative cases with negative toxicology, the term sudden arrhythmic death syndrome (SADS) may be applied and primary electrical diseases are potential causes.^{56, 183, 223, 253} Retaining tissue for DNA extraction is important for post-mortem genetic analysis, where the yield can be as high as one out of three.^{183, 269, 270}

Clinical evaluation of first-degree relatives is important if the cause of death after autopsy is unknown (Section 5.2.5, scenario 5) or suspected to be inherited, with a reported combined diagnostic yield of genetic and clinical evaluation

of 18a^{PROSIS}53%.^{252,266,271} Post-mortem genetic testing on the deceased, targeted to the cause of death, identified mutations in around one-third of cases.^{56,266,269}

Recommendation Table 6

[Open in new tab](#)

Recommendations for evaluation of sudden death victims

Recommendations	Class ^a	Level ^b
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority. ^{20,25,56}	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files. ^{25,56}	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those <50 years of age. ^{183,264,265,267,269,270}	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained. ^{264,265}	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death. ^{267,268}	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended. ^{56,266,269}	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50) and/or the circumstances and/or family history support a primary electrical disease. ^{56,183,223}	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{271,272}	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{223,253,273}	I	B
Following SADS, post-mortem genetic testing in the decedent for additional genes may be considered.	IIb	C
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended. ^{274,275}	III	B

© ESC 2022

DNA, deoxyribonucleic acid; SADS, sudden arrhythmic death syndrome; SCD, sudden cardiac death; SD, sudden death.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority. ^{20,25,56}	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files. ^{25,56}	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those <50 years of age. ^{183,264,265,267,269,270}	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained. ^{264,265}	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death. ^{267,268}	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended. ^{56,266,269}	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50) and/or the circumstances and/or family history support a primary electrical disease. ^{56,183,223}	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{271,272}	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{223,253,273}	I	B
Following SADS, post-mortem genetic testing in the decedent for additional genes may be considered.	IIb	C
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended. ^{274,275}	III	B

DNA, deoxyribonucleic acid; SADS, sudden arrhythmic death syndrome; SCD, sudden cardiac death; SD, sudden death.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendation Table 6

[Open in new tab](#)

Recommendations for evaluation of sudden death victims

Recommendations	Class ^a	Level ^b
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority. ^{20,25,56}	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files. ^{25,56}	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those <50 years of age. ^{183,264,265,267,269,270}	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained. ^{264,265}	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death. ^{267,268}	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended. ^{56,266,269}	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50) and/or the circumstances and/or family history support a primary electrical disease. ^{56,183,223}	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{271,272}	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{223,253,273}	I	B
Following SADS, post-mortem genetic testing in the decedent for additional genes may be considered.	IIb	C
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended. ^{274,275}	III	B

© ESC 2022

DNA, deoxyribonucleic acid; SADS, sudden arrhythmic death syndrome; SCD, sudden cardiac death; SD, sudden death.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority. ^{20,25,56}	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files. ^{25,56}	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those <50 years of age. ^{183,264,265,267,269,270}	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained. ^{264,265}	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death. ^{267,268}	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended. ^{56,266,269}	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50) and/or the circumstances and/or family history support a primary electrical disease. ^{56,183,223}	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{271,272}	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic. ^{223,253,273}	I	B
Following SADS, post-mortem genetic testing in the decedent for additional genes may be considered.	IIb	C
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended. ^{274,275}	III	B

DNA, deoxyribonucleic acid; SADS, sudden arrhythmic death syndrome; SCD, sudden cardiac death; SD, sudden death.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

5.2.5. Scenario 5: Relatives of sudden arrhythmic death syndrome decedents

An algorithm for the evaluation of relatives of SADS decedents is presented in [Figure 8](#).




Figure 8

Algorithm for the evaluation of relatives of unexplained sudden death decedents.

CMR, cardiac magnetic resonance; ECG, electrocardiogram; N, No; SADS, sudden arrhythmic death syndrome; SCD, sudden cardiac death; Y, Yes. ^aOver 16 years old $\hat{A}$ any suspicions for Brugada syndrome on tests or decedent circumstances of death. ^bIf exercise is not feasible. ^cRe-evaluate if change in family history or new symptoms.

[Open in new tab](#)[Download slide](#)

Studies evaluating families of SADS decedents have identified underlying genetic heart disease in relatives that is presumed to be the cause of death in the absence of other findings. The overall diagnostic yield ranged from 18 to 53%, depending on population and clinical investigative protocols.²⁷⁶ Aetiologies included LQTS, BrS, CPVT, and other disorders, such as cardiomyopathy.²⁷⁶ All study protocols relied upon a similar initial approach of evaluating the decedent's pathological reports, medical history, and circumstances of death, and then offering clinical evaluation to relatives with a minimum of personal history, family history, physical examination, ECG and exercise test, and echocardiography.^{223,252,253,277a,b,c,d,e,f,g,h,i,j,k,l,m,n,o,p,q,r,s,t,u,v,w,x,y,z} Where they diverged was the frequency of usage of additional tests such as high-lead ECGs, Holter monitoring, signal-averaged ECG, CMR, and provocative testing.¹³⁵ Sodium channel blocker drug challenge and high-lead ECGs systematically performed in SADS relatives provided a yield of 28% BrS diagnoses in one study²⁸¹; however, there are concerns about false positives.¹³⁹ Furthermore, epinephrine challenge has not been studied systematically in SADS families but, in the opinion of this panel, may be of utility in CPVT-suspected patients who are unable to exercise.¹³⁷

Recent data indicated at least a 13% genetic yield in SADS cases.^{135,178,183,276,283} Routine follow-up of families without a diagnosis yields little in new diagnoses,²⁸⁴ although children of decedents may be followed for age-penetrant disease until adulthood.¹⁸¹

If an autopsy is ambiguous, or if an autopsy was not undertaken in a young SCD case with family or personal history suspicious for inherited heart disease, then the yield of familial evaluation was similar to that in clear SADS

cases, [223,253,271](#)

Recommendation Table 7

[Open in new tab](#)

Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedents

Recommendations	Class ^a	Level ^b
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years, or if there is other circumstantial data or family history to suggest heritable disease. 223,252,253,277,281	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation. 183,253,277,281	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard and high precordial lead ECG, echocardiography, and exercise testing. 223,252,253,277,281	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood. 181,284	I	C
Pharmacological testing with a sodium channel blocker should be considered in relatives of SADS decedents who are 16 years or older when baseline testing and/or proband findings increase the suspicion of BrS. 277,281	IIa	B
Ambulatory cardiac rhythm monitoring and CMR may be considered in relatives of SADS decedents. 223,253,277,281	IIb	C
Pharmacological testing including epinephrine challenge (if exercise testing is impractical) and sodium channel blocker challenge may be considered in first-degree relatives of SADS decedents with normal baseline testing. 223,281	IIb	B
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes. 181,284	III	C

© ESC 2022

BrS, Brugada syndrome; CMR, cardiac magnetic resonance; ECG, electrocardiogram; SADS, sudden arrhythmic death syndrome.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years, or if there is other circumstantial data or family history to suggest heritable disease.^{223,252,253,277,281}	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation. ^{183,253,277,281}	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard and high precordial lead ECG, echocardiography, and exercise testing. ^{223,252,253,277,281}	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood. ^{181,284}	I	C
Pharmacological testing with a sodium channel blocker should be considered in relatives of SADS decedents who are 16 years or older when baseline testing and/or proband findings increase the suspicion of BrS. ^{277,281}	IIa	B
Ambulatory cardiac rhythm monitoring and CMR may be considered in relatives of SADS decedents. ^{223,253,277,281}	IIb	C
Pharmacological testing including epinephrine challenge (if exercise testing is impractical) and sodium channel blocker challenge may be considered in first-degree relatives of SADS decedents with normal baseline testing. ^{223,281}	IIb	B
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes. ^{181,284}	III	C

BrS, Brugada syndrome; CMR, cardiac magnetic resonance; ECG, electrocardiogram; SADS, sudden arrhythmic death syndrome.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendation Table 7

[Open in new tab](#)

Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedents

Recommendations	Class ^a	Level ^b
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years, or if there is other circumstantial data or family history to suggest heritable disease.^{223,252,253,277,281}	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation. ^{183,253,277,281}	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard and high precordial lead ECG, echocardiography, and exercise testing. ^{223,252,253,277,281}	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood. ^{181,284}	I	C
Pharmacological testing with a sodium channel blocker should be considered in relatives of SADS decedents who are 16 years or older when baseline testing and/or proband findings increase the suspicion of BrS. ^{277,281}	IIa	B
Ambulatory cardiac rhythm monitoring and CMR may be considered in relatives of SADS decedents. ^{223,253,277,281}	IIb	C
Pharmacological testing including epinephrine challenge (if exercise testing is impractical) and sodium channel blocker challenge may be considered in first-degree relatives of SADS decedents with normal baseline testing. ^{223,281}	IIb	B
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes. ^{181,284}	III	C

BrS, Brugada syndrome; CMR, cardiac magnetic resonance; ECG, electrocardiogram; SADS, sudden arrhythmic death syndrome.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years, or if there is other circumstantial data or family history to suggest heritable disease.^{223,252,253,277,281}	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation. ^{183,253,277,281}	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard and high precordial lead ECG, echocardiography, and exercise testing. ^{223,252,253,277,281}	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood. ^{181,284}	I	C
Pharmacological testing with a sodium channel blocker should be considered in relatives of SADS decedents who are 16 years or older when baseline testing and/or proband findings increase the suspicion of BrS. ^{277,281}	IIa	B
Ambulatory cardiac rhythm monitoring and CMR may be considered in relatives of SADS decedents. ^{223,253,277,281}	IIb	C
Pharmacological testing including epinephrine challenge (if exercise testing is impractical) and sodium channel blocker challenge may be considered in first-degree relatives of SADS decedents with normal baseline testing. ^{223,281}	IIb	B
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes. ^{181,284}	III	C

BrS, Brugada syndrome; CMR, cardiac magnetic resonance; ECG, electrocardiogram; SADS, sudden arrhythmic death syndrome.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

6. Therapies for ventricular arrhythmias. General aspects

6.1. Acute management

6.1.1. Treatment of reversible causes

Reversible causes may account for up to 50% of SCA.^{285,286} However, most of the time it is difficult to determine the exact underlying cause of SCA and whether it is reversible. A comprehensive evaluation of patients with SCA is mandatory if the underlying cardiac disease is unknown or disease progression is suspected ([Section 5.2.3](#), scenario 3).

Electrolyte imbalances, such as hypokalaemia, may trigger VA, and a rapid rise in extracellular potassium may lead to asystole.^{287a} Other factors such as bradycardia, ischaemia, coronary spasm, thrombosis, fever, acute starvation, and dieting may contribute to the occurrence of VA.^{290a} Acute correction of these reversible factors is recommended.

Drug-induced arrhythmias should be suspected in patients being treated with agents known to alter the electrical properties of the heart (e.g. inducing QRS and/or QT prolongation) or causing electrolyte abnormalities (e.g. thiazide and loop diuretics). When drug-induced arrhythmias are presumed, any offending drug needs to be withdrawn and substances that are known to prolong QT (e.g. sotalol) should be avoided.^{293,294} Hypomagnesaemia and/or hypokalaemia may be associated with Torsades de pointes (TdP). Intravenous magnesium is an effective therapy for TdP even in the absence of hypomagnesaemia.²⁹⁵ In refractory cases of recurrent TdP in the setting of acquired long QT, the arrhythmia can be suppressed by increasing the underlying heart rate using isoproterenol (isoprenaline) or transvenous pacing.

Patients who survive SCA in the context of a presumed reversible cause may have a high mortality rate.²⁸⁶ In a recent large observational study²⁹⁶ on survivors of SCA attributed to a reversible and correctable cause, subsequent ICD implantation was associated with lower all-cause mortality except for aborted CA occurring in the presence of acute myocardial infarction (MI). Thus, the need for prophylactic ICD implantation should be considered based on the underlying cardiac disease and an individual evaluation of the future risk of life-threatening VA.

Recommendation Table 8

[Open in new tab](#)

Recommendations for treatment of reversible conditions

Recommendations	Class ^a	Level ^b
Withdrawal of offending agents is recommended whenever drug-induced VAs are suspected. ^{293,294,297}	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ^c is recommended in patients with VA. ^{292,298}	I	C
Despite a possible correctable cause for the presenting VA, the need for ICD implantation should be considered based on an individual evaluation of the risk of subsequent VA/SCD. ^{286,296,299}	IIa	C

© ESC 2022

ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.
^cList not exhaustive.

Recommendations	Class ^a	Level ^b
Withdrawal of offending agents is recommended whenever drug-induced VAs are suspected. ^{293,294,297}	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ^c is recommended in patients with VA. ^{292,298}	I	C
Despite a possible correctable cause for the presenting VA, the need for ICD implantation should be considered based on an individual evaluation of the risk of subsequent VA/SCD. ^{286,296,299}	IIa	C

© ESC 2022

ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.
^cList not exhaustive.

Recommendation Table 8

[Open in new tab](#)

Recommendations for treatment of reversible conditions

Recommendations	Class ^a	Level ^b
Withdrawal of offending agents is recommended whenever drug-induced VAs are suspected. ^{293,294,297}	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ^c is recommended in patients with VA. ^{292,298}	I	C
Despite a possible correctable cause for the presenting VA, the need for ICD implantation should be considered based on an individual evaluation of the risk of subsequent VA/SCD. ^{286,296,299}	IIa	C

© ESC 2022

ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.
^cList not exhaustive.

Recommendations	Class ^a	Level ^b
Withdrawal of offending agents is recommended whenever drug-induced VAs are suspected. ^{293,294,297}	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ^c is recommended in patients with VA. ^{292,298}	I	C
Despite a possible correctable cause for the presenting VA, the need for ICD implantation should be considered based on an individual evaluation of the risk of subsequent VA/SCD. ^{286,296,299}	IIa	C

© ESC 2022

ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.
^cList not exhaustive.

6.1.2. Acute management of sustained monomorphic ventricular tachycardia

Patients presenting with SMVT should be treated according to symptoms and aetiology ([Figure 9](#)). Patients presenting with haemodynamic instability require immediate synchronized cardioversion. If synchronization is not possible, an unsynchronized shock should be used. Cardioversion is not indicated in patients with repetitive NSVTs ([Figure 10](#)). Documentation of any haemodynamically tolerated wide QRS tachycardia on 12-lead ECG is important. Administration of adenosine³⁰⁰ or vagal manoeuvres with continuous recording of 12-lead ECG should be considered if supraventricular tachycardia (SVT) is likely. Intravenous adenosine may also terminate specific VT subtypes. Such a response supports cyclic adenosine monophosphate (cAMP)-mediated triggered activity as an underlying VT mechanism.³⁰¹ Pre-excited atrial fibrillation (AF) can be recognized by the 'FBI' (fast, broad, irregular) ECG pattern. It may mimic VT, and intravenous administration of drugs that slow AV conduction, such as adenosine, beta-blockers, and amiodarone, should be avoided.³⁰² Prompt termination of SMVT is recommended even for tolerated SMVT, as rapid haemodynamical deterioration may occur. Termination can be achieved with electrical cardioversion, anti-arrhythmic medications, or pacing techniques. All anti-arrhythmic drugs (AADs) may lead to hypotension, but the individual risk of anaesthesia/sedation required for cardioversion also needs to be considered. For pharmacological termination of a haemodynamically tolerated VT of unknown aetiology, intravenous procainamide or amiodarone can be used. In the PROCAMIO trial,³⁰³ procainamide therapy was associated with a higher proportion of tachycardia termination and fewer major cardiac adverse events than amiodarone. Intravenous procainamide should not be used in patients with severe heart failure, acute MI, and end-stage renal disease. Administration of other AADs (ajmaline, sotalol, and flecainide)^{304,305} may be considered in patients without significant heart disease, but the risk of adverse events should be carefully weighed. Availability of AADs has to be taken into account, e.g. procainamide is not available in many European countries. In patients with an ICD, manual overdrive pacing may terminate those VTs with a cycle length under the programmed ICD detection rate. In case of a known idiopathic VT ([Figure 4](#)), treatment with beta-blockers (for RVOT VT)³⁰⁶ or verapamil (for fascicular VT)³⁰⁷ is recommended for acute conversion. Although verapamil may terminate other types of idiopathic VT,³⁰⁷ important adverse effects such as severe hypotension may occur. If the VT aetiology is uncertain, intravenous administration of verapamil is not recommended.^{308,309} A comprehensive evaluation of patients presenting with SMVT is mandatory if the underlying cardiac disease is unknown or disease progression is suspected ([Section 5.2.2, scenario 2](#)).




Figure 9

Algorithm for the acute management of regular wide QRS complex tachycardia.

ALS, advanced life support; ECG, electrocardiogram; N, No; SVT, supraventricular tachycardia; VT, ventricular tachycardia; Y, Yes. ^aBesides SVT, adenosine may also terminate idiopathic VT, which then indicates triggered activity as the mechanism underlying the arrhythmia. ^bBenefit of cardioversion should be weighed against risks related to anaesthesia/sedation. ^cConsidering limited availability of the other anti-arrhythmic drugs.




Figure 10

Repetitive runs of ventricular tachycardia interrupted by occasional sinus beats.

VT, ventricular tachycardia.

6.1.3. Management of electrical storm and incessant ventricular tachycardia

An electrical storm is common in ICD patients and has been defined as three or more episodes of sustained VA occurring within 24 h, requiring either anti-tachycardia pacing (ATP) or cardioversion/defibrillation, with each event separated by at least 5 min.^{310,312} Patients who experience an electrical storm are prone to psychological disorders, heart failure decompensation, and increased mortality.^{313,314} The severity of an electrical storm can range from recurrent asymptomatic VT episodes terminated by ATP to a life-threatening electrical instability with VA that recurs frequently after multiple shocks. Frequent ICD shocks can also be inappropriately delivered ([Figure 11](#)).




Figure 11

Part One. Management of patients with electrical storm or repeated implantable cardioverter defibrillator discharges.




Figure 11

Part Two. Management of patients with electrical storm or repeated implantable cardioverter defibrillator discharges.

[Open in new tab](#) [Download slide](#)




Figure 11

Part Three. Management of patients with electrical storm or repeated implantable cardioverter defibrillator discharges.

AAD, anti-arrhythmic drug; AF, atrial fibrillation, ALS, Advanced life support; CPVT, catecholaminergic polymorphic ventricular tachycardia; ERS, early repolarization syndrome; ICD, implantable cardioverter defibrillator; N, No; PVC, premature ventricular complex; STEMI, ST-elevation myocardial infarction; SVT, supraventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; Y, Yes. ^aSpecial aspects of device therapy section. ^bNo data on effect of quinidine on PVC-triggered polymorphic VAs in patients with cardiomyopathies. ^cHigh VA burden refers to a clinical scenario of very frequent VA episodes requiring ICD shocks, when only short periods of stable rhythm can be achieved. Low VA burden refers to a clinical scenario of repeated ATPs/ICD shocks followed by

stable rhythm. ^dIf bradycardia or post-extrasystolic pauses facilitate initiation of PVT/VF. ^eOverdrive pacing (by pacing with a slightly higher rate than the baseline rhythm) can be useful to temporarily suppress slow recurrent/incessant VT.

[Open in new tab](#)[Download slide](#)

In cases of inappropriate ICD shocks (e.g. due to SVTs or lead defects) or unnecessary ICD therapy (e.g. for NSVT or for repetitive VTs that terminate and restart spontaneously), disabling of ICD therapies is recommended. If an electrophysiology specialist or programmer is not available, the ICD can be disabled by placing a magnet over the device.

In case of haemodynamic instability at initial evaluation, institution of advanced life support (ALS) is recommended.³¹⁵ Reversible conditions contributing to initiation and perpetuation of VA need to be corrected (see [Section 6.1.1](#)). Further management depends on the type of VA and the underlying aetiology.^{312,316} A multifaceted approach is often required for management, consisting of ICD reprogramming when appropriate, AAD therapy, sedation, catheter ablation, autonomic modulation, and mechanical circulatory support.

Elevated sympathetic tone needs to be addressed. For patients with recurrent ICD shocks, sedation is indicated to alleviate psychological distress and decrease pro-arrhythmogenic sympathetic tone. Initial treatment with beta-blockers, preferably non-selective beta-blockers like propranolol, which was superior to metoprolol in one study,³¹⁷ combined with amiodarone³¹⁸ is most commonly used. In patients with recurrent haemodynamically not-tolerated VTs resistant to amiodarone, landiolol (ultra-short-acting β_2 -selective blocker) was found to be effective for arrhythmia suppression in two smaller studies.^{319,320} Administration of other AADs such as procainamide,³²¹ lidocaine,³²² or quinidine^{296,297} depends on the specific situation, type of VA, and underlying aetiology. When electrical storm remains intractable, with multiple shocks in a few hours, despite available anti-arrhythmic therapies, deep sedation/intubation should be considered, along with mechanical ventilation.³²⁵ If beta-blocker treatment is insufficient or not tolerated to decrease sympathetic tone, selected patients may benefit from autonomic modulation, i.e. percutaneous ganglionic stellate blockade,³²⁶ thoracic epidural anaesthesia,³²⁷ or left cardiac sympathetic denervation.³²⁸

The most common arrhythmia underlying electrical storm is SMVT associated with SHD that is amenable to catheter ablation.^{313,329} Successful ablation was associated with a significant reduction in VT and electrical storm recurrence and improved long-term survival in retrospective analyses.^{330,331} In patients with incessant slow MVT, catheter ablation is preferred over AAD therapy, which may only further slow VT. Catheter ablation should also be considered in patients with recurrent symptomatic episodes of PVT or VF triggered by a similar PVC.^{221,332,333,334} Institution of mechanical circulatory support may be considered for haemodynamic stabilization, when conventional therapy fails, and to provide circulatory support during ablation.³³⁵ In a recent meta-analysis³³⁶ including 2465 patients, a substantially lower mortality was observed with prophylactic mechanical circulatory support treatment among patients suffering from electrical storm or high-risk PAINESD score.³³⁷ In contrast, rescue use of mechanical circulatory support during ablation was associated with a high mortality rate.³³⁸ In patients with electrical storm due to recurrent PVT/VF, the underlying aetiology determines further management ([Figure 11](#)).

Recommendation Table 9

[Open in new tab](#)

Recommendations for the acute management of sustained ventricular tachycardia and electrical storm

Recommendations	Class ^a	Level ^b
Acute management of sustained VT		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT. ^{303,339}	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C

In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended. ^{306,307}	I	C
In patients presenting with a regular haemodynamically tolerated wide QRS complex tachycardia suspected for SVT, administration of adenosine or vagal manoeuvres should be considered. ³⁰⁰	IIa	C
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered. ³⁰³	IIa	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered. ³⁰³	IIb	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of significant SHD, flecainide, ajmaline, or sotalol may be considered. ^{304,305}	IIb	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism. ^{308,309}	III	B
Management of electrical storm		
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated. ^{317,318}	I	B
Intravenous magnesium with supplementation of potassium is recommended in patients with TdP. ²⁹⁵	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs. ^{330,331}	I	B
Deep sedation/intubation should be considered in patients with an intractable electrical storm refractory to drug treatment. ³²⁵	IIa	C
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC, non-responsive to medical treatment or coronary revascularization. ^{221,332,333}	IIa	C
Quinidine may be considered in patients with CAD and electrical storm due to recurrent PVT when other AAD therapy fails. ^{323,324}	IIb	C

Autonomic modulation may be considered in patients with electrical storm refractory to drug treatment and in whom catheter ablation is ineffective or not possible. ^{326,328,340}	IIb	C
Institution of mechanical circulatory support may be considered in the management of drug-refractory electrical storm and cardiogenic shock. ³³⁵	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; DC, direct current; LQT, long QT; PVC, premature ventricular complex; PVT, polymorphic VT; RVOT, right ventricular outflow tract; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TdP, Torsades de pointes; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Acute management of sustained VT		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT. ^{303,339}	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C

In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended. ^{306,307}	I	C
In patients presenting with a regular haemodynamically tolerated wide QRS complex tachycardia suspected for SVT, administration of adenosine or vagal manoeuvres should be considered. ³⁰⁰	IIa	C
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered. ³⁰³	IIa	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered. ³⁰³	IIb	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of significant SHD, flecainide, ajmaline, or sotalol may be considered. ^{304,305}	IIb	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism. ^{308,309}	III	B
Management of electrical storm		
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated. ^{317,318}	I	B
Intravenous magnesium with supplementation of potassium is recommended in patients with TdP. ²⁹⁵	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs. ^{330,331}	I	B
Deep sedation/intubation should be considered in patients with an intractable electrical storm refractory to drug treatment. ³²⁵	IIa	C
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC, non-responsive to medical treatment or coronary revascularization. ^{221,332,333}	IIa	C
Quinidine may be considered in patients with CAD and electrical storm due to recurrent PVT when other AAD therapy fails. ^{323,324}	IIb	C

Autonomic modulation may be considered in patients with electrical storm refractory to drug treatment and in whom catheter ablation is ineffective or not possible. ^{326,328,340}	IIb	C
Institution of mechanical circulatory support may be considered in the management of drug-refractory electrical storm and cardiogenic shock. ³³⁵	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; DC, direct current; LQT, long QT; PVC, premature ventricular complex; PVT, polymorphic VT; RVOT, right ventricular outflow tract; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TdP, Torsades de pointes; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 9

[Open in new tab](#)

Recommendations for the acute management of sustained ventricular tachycardia and electrical storm

Recommendations	Class ^a	Level ^b
Acute management of sustained VT		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT. ^{303,339}	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C

In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended. ^{306,307}	I	C
In patients presenting with a regular haemodynamically tolerated wide QRS complex tachycardia suspected for SVT, administration of adenosine or vagal manoeuvres should be considered. ³⁰⁰	IIa	C
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered. ³⁰³	IIa	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered. ³⁰³	IIb	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of significant SHD, flecainide, ajmaline, or sotalol may be considered. ^{304,305}	IIb	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism. ^{308,309}	III	B
Management of electrical storm		
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated. ^{317,318}	I	B
Intravenous magnesium with supplementation of potassium is recommended in patients with TdP. ²⁹⁵	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs. ^{330,331}	I	B
Deep sedation/intubation should be considered in patients with an intractable electrical storm refractory to drug treatment. ³²⁵	IIa	C
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC, non-responsive to medical treatment or coronary revascularization. ^{221,332,333}	IIa	C
Quinidine may be considered in patients with CAD and electrical storm due to recurrent PVT when other AAD therapy fails. ^{323,324}	IIb	C

Autonomic modulation may be considered in patients with electrical storm refractory to drug treatment and in whom catheter ablation is ineffective or not possible. ^{326,328,340}	IIb	C
Institution of mechanical circulatory support may be considered in the management of drug-refractory electrical storm and cardiogenic shock. ³³⁵	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; DC, direct current; LQT, long QT; PVC, premature ventricular complex; PVT, polymorphic VT; RVOT, right ventricular outflow tract; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TdP, Torsades de pointes; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Acute management of sustained VT		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT. ^{303,339}	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C

In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended. ^{306,307}	I	C
In patients presenting with a regular haemodynamically tolerated wide QRS complex tachycardia suspected for SVT, administration of adenosine or vagal manoeuvres should be considered. ³⁰⁰	IIa	C
In patients presenting with a haemodynamically tolerated SMVT and known or suspected SHD, intravenous procainamide should be considered. ³⁰³	IIa	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of an established diagnosis, intravenous amiodarone may be considered. ³⁰³	IIb	B
In patients presenting with a haemodynamically tolerated SMVT in the absence of significant SHD, flecainide, ajmaline, or sotalol may be considered. ^{304,305}	IIb	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism. ^{308,309}	III	B
Management of electrical storm		
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated. ^{317,318}	I	B
Intravenous magnesium with supplementation of potassium is recommended in patients with TdP. ²⁹⁵	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs. ^{330,331}	I	B
Deep sedation/intubation should be considered in patients with an intractable electrical storm refractory to drug treatment. ³²⁵	IIa	C
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC, non-responsive to medical treatment or coronary revascularization. ^{221,332,333}	IIa	C
Quinidine may be considered in patients with CAD and electrical storm due to recurrent PVT when other AAD therapy fails. ^{323,324}	IIb	C

Autonomic modulation may be considered in patients with electrical storm refractory to drug treatment and in whom catheter ablation is ineffective or not possible. ^{326,328,340}	IIb	C	© ESC 2022
Institution of mechanical circulatory support may be considered in the management of drug-refractory electrical storm and cardiogenic shock. ³³⁵	IIb	C	

AAD, anti-arrhythmic drug; CAD, coronary artery disease; DC, direct current; LQT, long QT; PVC, premature ventricular complex; PVT, polymorphic VT; RVOT, right ventricular outflow tract; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TdP, Torsades de pointes; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

6.2. Long-term management

6.2.1. Pharmacotherapy

Optimal medical treatment of the underlying cardiac disease, including the maximal tolerated doses of heart failure medication, is mandatory.³⁴¹

In patients with heart failure with reduced ejection fraction (HfrEF), the 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure recommend angiotensin-converting enzyme inhibitor (ACE-I)/angiotensin receptor blocker (ARB)/angiotensin receptor neprilysin inhibitors (ARNIs), mineralocorticoid receptor antagonists (MRAs), beta-blockers, and sodium-glucose co-transporter 2 (SGLT2) inhibitors to reduce mortality due to heart failure and SCD.³⁴²

Recommendation Table 10

[Open in new tab](#)

Recommendations for treatment with heart failure medication

Recommendations	Class ^a	Level ^b	© ESC 2022
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF. ^{343–347}	I	A	

ACE-I, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNIs, angiotensin receptor neprilysin inhibitor; EF, ejection fraction; MRA, mineralocorticoid receptor antagonists; SGLT2, sodium-glucose co-transporter 2.

^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b	© ESC 2022
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF. ^{343–347}	I	A	

ACE-I, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNIs, angiotensin receptor neprilysin inhibitor; EF, ejection fraction; MRA, mineralocorticoid receptor antagonists; SGLT2, sodium-glucose co-transporter 2.

^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 10

[Open in new tab](#)

Recommendations for treatment with heart failure medication

Recommendations	Class ^a	Level ^b
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF. ^{343–347}	I	A

ACE-I, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNIs, angiotensin receptor neprilysin inhibitor; EF, ejection fraction; MRA, mineralocorticoid receptor antagonists; SGLT2, sodium–glucose co-transporter 2.

^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF. ^{343–347}	I	A

ACE-I, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNIs, angiotensin receptor neprilysin inhibitor; EF, ejection fraction; MRA, mineralocorticoid receptor antagonists; SGLT2, sodium–glucose co-transporter 2.

^aClass of recommendation.
^bLevel of evidence.

AADs have an important role as adjunctive therapy in the management of VA, especially in symptomatic patients (Table 8). Until now, no AAD except for beta-blockers has demonstrated reduction in all-cause mortality. Each drug has a significant potential for causing adverse events, including pro-arrhythmia. For example, numerous AADs, as well as a large number of drugs with other therapeutic indications, can prolong the QT interval (<http://www.crediblemeds.org>) and provoke TdP, have negative chronotropic effects, may deteriorate heart failure, and cause bradycardia. Several drugs increase the risk of VA in patients with BrS (<http://www.brugadadrugs.org>). Modification of risk factors, when possible, is important for prevention of pro-arrhythmia. In patients who require a potentially arrhythmia-inducing drug, regular ECG and other tests according to the patient's profile and AAD characteristics are recommended (Figures 12 and 13).




Figure 12

Algorithm for evaluation before initiation and follow-up of patients requiring sodium channel blocking agents.

AV, atrioventricular; BBB, bundle branch block; ECG, electrocardiogram; LV, left ventricular; MI, myocardial infarction; N, No; Y, Yes. ^a<http://www.brugadadrugs.org>. ^bFlecainide, encainide. ^cCo-administration of drugs with AV

nodal blocking effect in patients with atrial fibrillation or atrial flutter. ^dIn implantable cardioverter defibrillator carriers, a higher risk of drug-induced pro-arrhythmia might be accepted. ^eAccording to the 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation, ³⁴⁸ A Δ $\hat{f}A \hat{f}_{ECG}$ QRS >25% is not an absolute cut-off value but dependent on QRS width before drug initiation and individualized patient risk $\hat{P}ADSTS$ benefit considerations.

[Open in new tab](#)[Download slide](#)




Figure 13

Algorithm for evaluation before initiation and follow-up of patients requiring drugs associated with QT prolongation.

ECG, electrocardiogram; LV, left ventricular; N, No; Y, Yes. ^a<http://www.crediblemeds.org>. ^bIf strong indication and no alternative treatment, consult a specialist. ^cAccording to the 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation. ³⁴⁸

[Open in new tab](#)[Download slide](#)

Table 8

[Open in new tab](#)

Anti-arrhythmic drugs (acute and chronic treatment)

Anti-arrhythmic drug	Effects on ECG	Indications (specific indication)	Oral dose per day (i.v. dose)	Side effects	Contraindications, precautions, other considerations
Amiodarone	Decreases sinus node frequency, prolongs QT interval ^a	PVC, VT, VF	200 $\hat{P}ADSTS$ 400 $\hat{P}ADNIEL$ mg Loading dose: 600 $\hat{P}ADSTS$ 1200 $\hat{P}ADNIEL$ mg/24 $\hat{P}ADNIEL$ h 8 $\hat{P}ADSTS$ 10 days. (Loading dose: 5 $\hat{P}ADNIEL$ mg/kg in 20 $\hat{P}ADNIEL$ min $\hat{P}ADSTS$ 2 $\hat{P}ADNIEL$ h, 2 $\hat{P}ADSTS$ 3 times in 24 $\hat{P}ADNIEL$ h, then 600 $\hat{P}ADSTS$ 1200 $\hat{P}ADNIEL$ mg/24 $\hat{P}ADNIEL$ h 8 $\hat{P}ADSTS$ 10 days)	<i>Cardiac:</i> Bradycardia, TdP (infrequent) <i>Extracardiac:</i> Photosensitivity, corneal deposits, hypothyroidism, hyperthyroidism, pulmonary toxicity, hepatotoxicity, polyneuropathy, skin discoloration	<i>Precautions:</i> Sinus node dysfunction, severe AV conduction disturbances, hyperthyroidism <i>Other considerations:</i> Can be used in patients with heart failure. Increases the risk of myopathy when used with statins

Adenosine	Transitory AV block	Regular wide complex tachycardia of unknown origin (outflow tract VT)	No oral use (6â€¦18â€¦mg bolus)	Chest pain, flushing, bronchoconstriction	<i>Contraindications:</i> Severe asthma, pre-excited AF <i>Other considerations:</i> Antagonist: theophylline
Ajmaline	Prolongs QRS duration and QT interval ^a	VT (unmasking BrS ECG)	No oral use (1 mg/kg over 5â€¦10 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min)	<i>Cardiac:</i> VF (rare in suspected BrS), occasional TdP, negative inotrope <i>Extracardiac:</i> Cholestatic jaundice, headache, nausea, thrombocytopenia	<i>Contraindications:</i> BrS ECG type I, QT prolongation
Beta-blocker	Decreases sinus node frequency, prolongs PR interval, shortens QT interval	PVC, VT (LQTS, CPVT)	Various (various)	<i>Cardiac:</i> Bradycardia, AV block, hypotension, negative inotrope <i>Extracardiac:</i> Fatigue, bronchospasm, sexual disturbances, depression, cold extremities	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, decompensated heart failure, coronary vasospasm, severe asthma, BrS
Landiolol (Ultra-short acting Î² ₁ -selective blocker)	See beta-blocker	VT, electrical storm	No oral use 100â€¦Âµg/kg bolus in 1â€¦min, infusion 10â€¦40â€¦Âµg/kg/min (max 80â€¦Âµg/kg/min; max 24â€¦h total dose 57.6â€¦mg/kg/day)	See beta-blocker	<i>Contraindications:</i> See beta-blocker. Bradycardia, hypotension <i>Other considerations:</i> Limited experience with its use beyond 24â€¦h
Nadolol (Non-selective Î² ₁ Î² ₂ blocker)	See beta-blocker	PVC, VT (LQTS, CPVT)	40â€¦120â€¦mg	See beta-blocker	<i>Contraindications:</i> See beta-blocker <i>Other considerations:</i> Plasma half-life 20â€¦24â€¦h
Propranolol (Non-selective Î² ₁ Î² ₂ blocker)	See beta-blocker	PVC, VT (electrical storm, LQTS, CPVT)	80â€¦320â€¦mg (160â€¦mg/24 h)	See beta-blocker	<i>Contraindications:</i> See beta-blocker
Disopyramide	Increases sinus node frequency and prolongs	PVC, VT	250â€¦750â€¦mg	<i>Cardiac:</i> Negative inotrope, AV block, pro-arrhythmia (MVT, occasional TdP)	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe

	PR interval, QRS duration, and QT interval ^a			<i>Extracardiac:</i> Anticholinergic effects	intraventricular conduction disturbances, prior MI, significant SHD, hypotension <i>Other considerations:</i> Reduces LV outflow tract obstruction and symptoms in HCM
Flecainide	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT (unmasking BrS ECG ^b)	200â€”400â€” mg (1â€”2â€” mg/kg over 10â€” min)	<i>Cardiac:</i> Pro-arrhythmia (MVT, occasional TdP), negative inotrope, sinus bradycardia, AV block, 1:1 AV conduction during flutter <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, diplopia, headache)	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, inherited LQTS (other than LQTS3), severe kidney disease (CrCl <35â€” mL/min/1.73â€” m ²) <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Isoproterenol	Increases sinus node frequency, shortens QT interval	(Electrical storm in BrS, idiopathic VF, and ERS, TdP, beta-blocker overdose; acquired LQTS)	No oral use (0.5â€”10â€” Âµg/min)	<i>Cardiac:</i> Sinus tachycardia, vasodilation <i>Extracardiac:</i> Headache, sweating, tremor	<i>Contraindications:</i> ACS, LQTS <i>Other considerations:</i> Short plasma half time (2â€”min)
Lidocaine	No significant effects	(VT/VF associated with ACS)	No oral use (50â€”200â€” mg bolus, then 2â€”4â€” mg/min)	<i>Cardiac:</i> Sinoatrial arrest <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, dizziness)	<i>Precautions:</i> Reduced dose with reduced liver blood flow (e.g. shock, Î²-blockade, severe heart failure) <i>Other considerations:</i> More effective with high potassium level. Few haemodynamic side effects
Mexiletine	No significant	PVC, VT (LQT3)	600â€”1200â€” mg	<i>Cardiac:</i> Sinus bradycardia in	<i>Contraindications:</i> Sinus node

	effects		Loading dose: 400aFADNELmg initially followed by 600aFADNELmg in the first 24aFADNELh	sinus node dysfunction, hypotension <i>Extracardiac:</i> Central nervous system effects (e.g. tremor, dysarthria, dizziness), gastrointestinal complaints	dysfunction, severe AV conduction disturbances, severe heart failure
Procainamide	Prolongs PR interval, QRS duration, and QT interval ^a	VT	(100aFADNELmg bolus, can be repeated after 5aFADNELmin if no effect, max 500aFADNELmg [max 50aFADNELmg/min]. Then, 2aFADNEL6aFADNELmg/min)	<i>Cardiac:</i> Sinus bradycardia, hypotension, TdP <i>Extracardiac:</i> Rash, myalgia, vasculitis, systemic lupus, agranulocytosis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, severe LV dysfunction hypotension, BrS
Propafenone	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	450aFADNEL900aFADNELmg	<i>Cardiac:</i> Sinus bradycardia, AV block, negative inotrope, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Gastrointestinal disturbances, headache, dry mouth	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, LQTS, significant renal or liver disease <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Quinidine	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	(VF ^c , BrS, SQTS)	600aFADNEL1600aFADNELmg Loading dose: Start 200aFADNELmg every 3 h until effect, max. 3aFADNELg in first 24aFADNELh	<i>Cardiac:</i> Hypotension, TdP^d <i>Extracardiac:</i> Gastrointestinal disturbances, auditory and visual disturbances, confusion, leukopenia, haemolytic anaemia, thrombocytopenia, anaphylaxis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, previous MI, significant SHD, hypotension, LQTS
Ranolazine	Decreases sinus node frequency, prolongs QT interval ^a	VT (LQTS3)	750aFADNEL2000aFADNELmg	<i>Cardiac:</i> Sinus bradycardia, hypotension <i>Extracardiac:</i> Dizziness, nausea, constipation, gastrointestinal	<i>Contraindications:</i> Severe sinus node dysfunction, severe heart failure, LQTS (other than LQTS3) <i>Precautions:</i> Concomitant

				disturbance, headache, rash	treatments associated with QT interval prolongation
Sotalolol	Decreases sinus node frequency, prolongs QT interval ^a	VT	160â640â mg (0.5â1.5â mg/kg in 10 min. If necessary, can be repeated after 6â h)	See beta-blockers, TdP ^d (>2% of patients, close monitoring of QT interval and CrCl)	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe heart failure with reduced LVEF, significant LVH, CrCl <30â ml/min, coronary vasospasm, LQTS <i>Precautions:</i> Concomitant treatments associated with QT interval prolongation, hypokalaemia <i>Other considerations:</i> Potassium channel blocker effects require higher dose than beta-blocker effects
Verapamil	Prolongs PR interval	(LV fascicular tachycardia)	120â480â mg (5â10â mg in slow bolus. If necessary, can be repeated in 30â min)	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, AV block, negative inotrope, hypotension <i>Extracardiac:</i> Gastrointestinal disturbances, peripheral oedema, flushing	<i>Contraindications:</i> Heart failure with reduced LVEF, severe sinus node dysfunction, and severe AV conduction disturbances, VT of unknown origin, ACS, WPW syndrome <i>Other considerations:</i> Increase the risk of myopathy when used with statins

Anti-arrhythmic drug	Effects on ECG	Indications (specific indication)	Oral dose per day (i.v. dose)	Side effects	Contraindications, precautions, other considerations
Amiodarone	Decreases sinus node frequency, prolongs QT interval ^a	PVC, VT, VF	200â400â mg Loading dose: 600â1200â mg/24â h 8â10 days. (Loading dose: 5â20â mg/kg in 20â	<i>Cardiac:</i> Bradycardia, TdP (infrequent) <i>Extracardiac:</i> Photosensitivity, corneal deposits, hypothyroidism, hyperthyroidism, pulmonary toxicity,	<i>Precautions:</i> Sinus node dysfunction, severe AV conduction disturbances, hyperthyroidism <i>Other considerations:</i> Can be used in

			minâ2âh, 2â3 times in 24âh, then 600â1200âmg/24âh 8â10 days)	hepatotoxicity, polyneuropathy, skin discoloration	patients with heart failure. Increases the risk of myopathy when used with statins
Adenosine	Transitory AV block	Regular wide complex tachycardia of unknown origin (outflow tract VT)	No oral use (6â18âmg bolus)	Chest pain, flushing, bronchoconstriction	<i>Contraindications:</i> Severe asthma, pre-excited AF <i>Other considerations:</i> Antagonist: theophylline
Ajmaline	Prolongs QRS duration and QT interval ^a	VT (unmasking BrS ECG)	No oral use (1 mg/kg over 5â10 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min)	<i>Cardiac:</i> VF (rare in suspected BrS), occasional TdP, negative inotrope <i>Extracardiac:</i> Cholestatic jaundice, headache, nausea, thrombocytopenia	<i>Contraindications:</i> BrS ECG type I, QT prolongation
Beta-blocker	Decreases sinus node frequency, prolongs PR interval, shortens QT interval	PVC, VT (LQTS, CPVT)	Various (various)	<i>Cardiac:</i> Bradycardia, AV block, hypotension, negative inotrope <i>Extracardiac:</i> Fatigue, bronchospasm, sexual disturbances, depression, cold extremities	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, decompensated heart failure, coronary vasospasm, severe asthma, BrS
Landiolol (Ultra-short acting Î² ₁ -selective blocker)	See beta-blocker	VT, electrical storm	No oral use 100âÂµg/kg bolus in 1âmin, infusion 10â40âÂµg/kg/min (max 80âÂµg/kg/min; max 24âh total dose 57.6âmg/kg/day)	See beta-blocker	<i>Contraindications:</i> See beta-blocker. Bradycardia, hypotension <i>Other considerations:</i> Limited experience with its use beyond 24âh
Nadolol (Non-selective Î² ₁ /Î² ₂ blocker)	See beta-blocker	PVC, VT (LQTS, CPVT)	40â120âmg	See beta-blocker	<i>Contraindications:</i> See beta-blocker <i>Other considerations:</i> Plasma half-life 20â24âh
Propranolol (Non-selective	See beta-blocker	PVC, VT (electrical storm,	80â320âmg (160âmg/24 h)	See beta-blocker	<i>Contraindications:</i> See beta-blocker

β ₁ /β ₂ blocker)		LQTS, CPVT)			
Disopyramide	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	250â€”750â€”mg	<i>Cardiac:</i> Negative inotrope, AV block, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Anticholinergic effects	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, prior MI, significant SHD, hypotension <i>Other considerations:</i> Reduces LV outflow tract obstruction and symptoms in HCM
Flecainide	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT (unmasking BrS ECG ^b)	200â€”400â€”mg (1â€”2â€”mg/kg over 10â€”min)	<i>Cardiac:</i> Pro-arrhythmia (MVT, occasional TdP), negative inotrope, sinus bradycardia, AV block, 1:1 AV conduction during flutter <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, diplopia, headache)	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, inherited LQTS (other than LQTS3), severe kidney disease (CrCl <35â€”mL/min/1.73â€”m²) <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Isoproterenol	Increases sinus node frequency, shortens QT interval	(Electrical storm in BrS, idiopathic VF, and ERS, TdP, beta-blocker overdose; acquired LQTS)	No oral use (0.5â€”10â€”Âµg/min)	<i>Cardiac:</i> Sinus tachycardia, vasodilation <i>Extracardiac:</i> Headache, sweating, tremor	<i>Contraindications:</i> ACS, LQTS <i>Other considerations:</i> Short plasma half time (2â€”min)
Lidocaine	No significant effects	(VT/VF associated with ACS)	No oral use (50â€”200â€”mg bolus, then 2â€”4â€”mg/min)	<i>Cardiac:</i> Sinoatrial arrest <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, dizziness)	<i>Precautions:</i> Reduced dose with reduced liver blood flow (e.g. shock, β-blockade, severe heart failure) <i>Other</i>

					<i>considerations:</i> More effective with high potassium level. Few haemodynamic side effects
Mexiletine	No significant effects	PVC, VT (LQT3)	600aPADSTSL1200aPADNEL mg Loading dose: 400aPADNELmg initially followed by 600aPADNELmg in the first 24aPADNELh	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, hypotension <i>Extracardiac:</i> Central nervous system effects (e.g. tremor, dysarthria, dizziness), gastrointestinal complaints	<i>Contraindications:</i> Sinus node dysfunction, severe AV conduction disturbances, severe heart failure
Procainamide	Prolongs PR interval, QRS duration, and QT interval ^a	VT	(100aPADNELmg bolus, can be repeated after 5aPADNELmin if no effect, max 500aPADSTSL750aPADNELmg [max 50aPADNELmg/min]. Then, 2aPADSTSL6aPADNELmg/min)	<i>Cardiac:</i> Sinus bradycardia, hypotension, TdP <i>Extracardiac:</i> Rash, myalgia, vasculitis, systemic lupus, agranulocytosis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, severe LV dysfunction hypotension, BrS
Propafenone	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	450aPADSTSL900aPADNELmg	<i>Cardiac:</i> Sinus bradycardia, AV block, negative inotrope, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Gastrointestinal disturbances, headache, dry mouth	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, LQTS, significant renal or liver disease <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Quinidine	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	(VF ^c , BrS, SQTS)	600aPADSTSL1600aPADNELmg Loading dose: Start 200aPADNELmg every 3 h until effect, max. 3aPADNELg in first 24aPADNELh	<i>Cardiac:</i> Hypotension, TdP ^d <i>Extracardiac:</i> Gastrointestinal disturbances, auditory and visual disturbances, confusion, leukopenia, haemolytic anaemia, thrombocytopenia, anaphylaxis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, previous MI, significant SHD, hypotension, LQTS

Ranolazine	Decreases sinus node frequency, prolongs QT interval ^a	VT (LQTS3)	750aPADSTS2000aPADNIEL mg	<i>Cardiac:</i> Sinus bradycardia, hypotension <i>Extracardiac:</i> Dizziness, nausea, constipation, gastrointestinal disturbance, headache, rash	<i>Contraindications:</i> Severe sinus node dysfunction, severe heart failure, LQTS (other than LQTS3) <i>Precautions:</i> Concomitant treatments associated with QT interval prolongation
Sotalol	Decreases sinus node frequency, prolongs QT interval ^a	VT	160aPADSTS640aPADNIEL mg (0.5aPADSTS1.5aPADNIEL mg/kg in 10 min. If necessary, can be repeated after 6aPADNIEL h)	See beta-blockers, Tdp ^d (>2% of patients, close monitoring of QT interval and CrCl)	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe heart failure with reduced LVEF, significant LVH, CrCl <30aPADNIEL ml/min, coronary vasospasm, LQTS <i>Precautions:</i> Concomitant treatments associated with QT interval prolongation, hypokalaemia <i>Other considerations:</i> Potassium channel blocker effects require higher dose than beta-blocker effects
Verapamil	Prolongs PR interval	(LV fascicular tachycardia)	120aPADSTS480aPADNIEL mg (5aPADSTS10aPADNIEL mg in slow bolus. If necessary, can be repeated in 30aPADNIEL min)	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, AV block, negative inotrope, hypotension <i>Extracardiac:</i> Gastrointestinal disturbances, peripheral oedema, flushing	<i>Contraindications:</i> Heart failure with reduced LVEF, severe sinus node dysfunction, and severe AV conduction disturbances, VT of unknown origin, ACS, WPW syndrome <i>Other considerations:</i> Increase the risk of myopathy when used with statins

ACS, acute coronary syndrome; AF, atrial fibrillation; AV, atrioventricular; BrS, Brugada syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; CrCl, creatinine clearance; ECG, electrocardiogram; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; LVH, left ventricular hypertrophy; MI, myocardial infarction; MVT, monomorphic ventricular tachycardia; PVC, premature ventricular complex; SHD, structural heart disease; SQTS,

short QT syndrome; TdP, torsades de pointes; VF, ventricular fibrillation; VT, ventricular tachycardia; WPW, Wolff-Parkinson-White.

a

Precaution with concomitant conditions or drugs that prolong QT interval. Should be discontinued if QTc >500 ms. See [Figure 13](#) Algorithm for evaluation before initiation and follow-up of patients requiring drugs associated with QT prolongation.

b

If ajmaline not available.

c

Subacute MI, multifocal ectopic Purkinje-related premature contractions, ERS, idiopathic VF.

d

Proarrhythmic side effects require strong indication in patients without ICD.

© ESC 2022

Table 8

[Open in new tab](#)

Anti-arrhythmic drugs (acute and chronic treatment)

Anti-arrhythmic drug	Effects on ECG	Indications (specific indication)	Oral dose per day (i.v. dose)	Side effects	Contraindications, precautions, other considerations
Amiodarone	Decreases sinus node frequency, prolongs QT interval ^a	PVC, VT, VF	200–400 mg Loading dose: 600–1200 mg/24 h 8–10 days. (Loading dose: 5–20 mg/kg in 20 min, 2–3 times in 24 h, then 600–1200 mg/24 h 8–10 days)	<i>Cardiac:</i> Bradycardia, TdP (infrequent) <i>Extracardiac:</i> Photosensitivity, corneal deposits, hypothyroidism, hyperthyroidism, pulmonary toxicity, hepatotoxicity, polyneuropathy, skin discoloration	<i>Precautions:</i> Sinus node dysfunction, severe AV conduction disturbances, hyperthyroidism <i>Other considerations:</i> Can be used in patients with heart failure. Increases the risk of myopathy when used with statins
Adenosine	Transitory AV block	Regular wide complex tachycardia of unknown origin (outflow tract VT)	No oral use (6–18 mg bolus)	Chest pain, flushing, bronchoconstriction	<i>Contraindications:</i> Severe asthma, pre-excited AF <i>Other considerations:</i> Antagonist: theophylline
Ajmaline	Prolongs QRS duration and QT interval ^a	VT (unmasking BrS ECG)	No oral use (1 mg/kg over 5–10 min (maximum dose 100 mg) or 1 mg/kg at 10 mg/min)	<i>Cardiac:</i> VF (rare in suspected BrS), occasional TdP, negative inotrope <i>Extracardiac:</i> Cholestatic jaundice, headache, nausea, thrombocytopenia	<i>Contraindications:</i> BrS ECG type I, QT prolongation

Beta-blocker	Decreases sinus node frequency, prolongs PR interval, shortens QT interval	PVC, VT (LQTS, CPVT)	Various (various)	<i>Cardiac:</i> Bradycardia, AV block, hypotension, negative inotrope <i>Extracardiac:</i> Fatigue, bronchospasm, sexual disturbances, depression, cold extremities	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, decompensated heart failure, coronary vasospasm, severe asthma, BrS
Landiolol (Ultra-short acting β_1 -selective blocker)	See beta-blocker	VT, electrical storm	No oral use 100 $\mu\text{g/kg}$ bolus in 1 min, infusion 10 $\mu\text{g/kg/min}$ (max 80 $\mu\text{g/kg/min}$; max 24 h total dose 57.6 mg/kg/day)	See beta-blocker	<i>Contraindications:</i> See beta-blocker. Bradycardia, hypotension <i>Other considerations:</i> Limited experience with its use beyond 24 h
Nadolol (Non-selective $\beta_1\beta_2$ blocker)	See beta-blocker	PVC, VT (LQTS, CPVT)	40–120 mg	See beta-blocker	<i>Contraindications:</i> See beta-blocker <i>Other considerations:</i> Plasma half-life 20–24 h
Propranolol (Non-selective $\beta_1\beta_2$ blocker)	See beta-blocker	PVC, VT (electrical storm, LQTS, CPVT)	80–320 mg (160 mg/24 h)	See beta-blocker	<i>Contraindications:</i> See beta-blocker
Disopyramide	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	250–750 mg	<i>Cardiac:</i> Negative inotrope, AV block, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Anticholinergic effects	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, prior MI, significant SHD, hypotension <i>Other considerations:</i> Reduces LV outflow tract obstruction and symptoms in HCM
Flecainide	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT (unmasking BrS ECG ^b)	200–400 mg (1–2 mg/kg over 10 min)	<i>Cardiac:</i> Pro-arrhythmia (MVT, occasional TdP), negative inotrope, sinus bradycardia, AV block, 1:1 AV conduction during flutter	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances,

				<i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, diplopia, headache)	inherited LQTS (other than LQTS3), severe kidney disease (CrCl <35 mL/min/1.73 m ²) <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Isoproterenol	Increases sinus node frequency, shortens QT interval	(Electrical storm in BrS, idiopathic VF, and ERS, TdP, beta-blocker overdose; acquired LQTS)	No oral use (0.5-10 µg/min)	<i>Cardiac:</i> Sinus tachycardia, vasodilation <i>Extracardiac:</i> Headache, sweating, tremor	<i>Contraindications:</i> ACS, LQTS <i>Other considerations:</i> Short plasma half time (2 min)
Lidocaine	No significant effects	(VT/VF associated with ACS)	No oral use (50-200 mg bolus, then 2-4 mg/min)	<i>Cardiac:</i> Sinoatrial arrest <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, dizziness)	<i>Precautions:</i> Reduced dose with reduced liver blood flow (e.g. shock, 1 st -blockade, severe heart failure) <i>Other considerations:</i> More effective with high potassium level. Few haemodynamic side effects
Mexiletine	No significant effects	PVC, VT (LQT3)	600-1200 mg Loading dose: 400 mg initially followed by 600 mg in the first 24 h	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, hypotension <i>Extracardiac:</i> Central nervous system effects (e.g. tremor, dysarthria, dizziness), gastrointestinal complaints	<i>Contraindications:</i> Sinus node dysfunction, severe AV conduction disturbances, severe heart failure
Procainamide	Prolongs PR interval, QRS duration, and QT interval ^a	VT	(100 mg bolus, can be repeated after 5 min if no effect, max 500-750 mg [max 50 mg/min]. Then, 2-6 mg/min)	<i>Cardiac:</i> Sinus bradycardia, hypotension, TdP <i>Extracardiac:</i> Rash, myalgia, vasculitis, systemic lupus, agranulocytosis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, severe

					LV dysfunction hypotension, BrS
Propafenone	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	450aPACSTST;900aPACNEL mg	<i>Cardiac:</i> Sinus bradycardia, AV block, negative inotrope, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Gastrointestinal disturbances, headache, dry mouth	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, LQTS, significant renal or liver disease <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Quinidine	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	(VF ^c , BrS, SQTs)	600aPACSTST;1600aPACNEL mg Loading dose: Start 200aPACNELmg every 3 h until effect, max. 3aPACNELg in first 24aPACNELh	<i>Cardiac:</i> Hypotension, TdP^d <i>Extracardiac:</i> Gastrointestinal disturbances, auditory and visual disturbances, confusion, leukopenia, haemolytic anaemia, thrombocytopenia, anaphylaxis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, previous MI, significant SHD, hypotension, LQTS
Ranolazine	Decreases sinus node frequency, prolongs QT interval ^a	VT (LQTS3)	750aPACSTST;2000aPACNEL mg	<i>Cardiac:</i> Sinus bradycardia, hypotension <i>Extracardiac:</i> Dizziness, nausea, constipation, gastrointestinal disturbance, headache, rash	<i>Contraindications:</i> Severe sinus node dysfunction, severe heart failure, LQTS (other than LQTS3) <i>Precautions:</i> Concomitant treatments associated with QT interval prolongation
Sotalol	Decreases sinus node frequency, prolongs QT interval ^a	VT	160aPACSTST;640aPACNEL mg (0.5aPACSTST;1.5aPACNEL mg/kg in 10 min. If necessary, can be repeated after 6aPACNELh)	See beta-blockers, TdP ^d (>2% of patients, close monitoring of QT interval and CrCl)	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe heart failure with reduced LVEF, significant LVH, CrCl <30aPACNELml/min, coronary vasospasm, LQTS <i>Precautions:</i> Concomitant treatments associated with QT interval

					prolongation, hypokalaemia <i>Other considerations:</i> Potassium channel blocker effects require higher dose than beta-blocker effects
Verapamil	Prolongs PR interval	(LV fascicular tachycardia)	120â480âmg (5â10âmg in slow bolus. If necessary, can be repeated in 30âmin)	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, AV block, negative inotrope, hypotension <i>Extracardiac:</i> Gastrointestinal disturbances, peripheral oedema, flushing	<i>Contraindications:</i> Heart failure with reduced LVEF, severe sinus node dysfunction, and severe AV conduction disturbances, VT of unknown origin, ACS, WPW syndrome <i>Other considerations:</i> Increase the risk of myopathy when used with statins

Anti-arrhythmic drug	Effects on ECG	Indications (specific indication)	Oral dose per day (i.v. dose)	Side effects	Contraindications, precautions, other considerations
Amiodarone	Decreases sinus node frequency, prolongs QT interval ^a	PVC, VT, VF	200â400âmg Loading dose: 600â1200âmg/24âh 8â10 days. (Loading dose: 5â20âmg/kg in 2â3 times in 24âh, then 600â1200âmg/24âh 8â10 days)	<i>Cardiac:</i> Bradycardia, TdP (infrequent) <i>Extracardiac:</i> Photosensitivity, corneal deposits, hypothyroidism, hyperthyroidism, pulmonary toxicity, hepatotoxicity, polyneuropathy, skin discoloration	<i>Precautions:</i> Sinus node dysfunction, severe AV conduction disturbances, hyperthyroidism <i>Other considerations:</i> Can be used in patients with heart failure. Increases the risk of myopathy when used with statins
Adenosine	Transitory AV block	Regular wide complex tachycardia of unknown origin (outflow tract VT)	No oral use (6â18âmg bolus)	Chest pain, flushing, bronchoconstriction	<i>Contraindications:</i> Severe asthma, pre-excited AF <i>Other considerations:</i> Antagonist: theophylline
Ajmaline	Prolongs QRS duration	VT (unmasking BrS ECG)	No oral use (1 mg/kg over 5â10 min)	<i>Cardiac:</i> VF (rare in suspected BrS),	<i>Contraindications:</i> BrS ECG type I, QT prolongation

	and QT interval ^a		(maximum dose 100 mg) or 1 mg/kg at 10 mg/min)	occasional TdP, negative inotrope <i>Extracardiac:</i> Cholestatic jaundice, headache, nausea, thrombocytopenia	
Beta-blocker	Decreases sinus node frequency, prolongs PR interval, shortens QT interval	PVC, VT (LQTS, CPVT)	Various (various)	<i>Cardiac:</i> Bradycardia, AV block, hypotension, negative inotrope <i>Extracardiac:</i> Fatigue, bronchospasm, sexual disturbances, depression, cold extremities	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, decompensated heart failure, coronary vasospasm, severe asthma, BrS
Landiolol (Ultra-short acting β_1 -selective blocker)	See beta-blocker	VT, electrical storm	No oral use 100 μ g/kg bolus in 1 min, infusion 10 μ g/kg/min (max 80 μ g/kg/min; max 24 h total dose 57.6 mg/kg/day)	See beta-blocker	<i>Contraindications:</i> See beta-blocker. Bradycardia, hypotension <i>Other considerations:</i> Limited experience with its use beyond 24 h
Nadolol (Non-selective β_1/β_2 blocker)	See beta-blocker	PVC, VT (LQTS, CPVT)	40–120 mg	See beta-blocker	<i>Contraindications:</i> See beta-blocker <i>Other considerations:</i> Plasma half-life 20–24 h
Propranolol (Non-selective β_1/β_2 blocker)	See beta-blocker	PVC, VT (electrical storm, LQTS, CPVT)	80–320 mg (160 mg/24 h)	See beta-blocker	<i>Contraindications:</i> See beta-blocker
Disopyramide	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	250–750 mg	<i>Cardiac:</i> Negative inotrope, AV block, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Anticholinergic effects	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, prior MI, significant SHD, hypotension <i>Other considerations:</i> Reduces LV outflow tract obstruction and symptoms in HCM

Flecainide	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT (unmasking BrS ECG ^b)	200â€”400â€” mg (1â€”2â€” mg/kg over 10â€” min)	<i>Cardiac:</i> Pro-arrhythmia (MVT, occasional TdP), negative inotrope, sinus bradycardia, AV block, 1:1 AV conduction during flutter <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, diplopia, headache)	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, inherited LQTS (other than LQTS3), severe kidney disease (CrCl <35â€” mL/min/1.73â€” m²) <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Isoproterenol	Increases sinus node frequency, shortens QT interval	(Electrical storm in BrS, idiopathic VF, and ERS, TdP, beta-blocker overdose; acquired LQTS)	No oral use (0.5â€”10â€” Âµg/min)	<i>Cardiac:</i> Sinus tachycardia, vasodilation <i>Extracardiac:</i> Headache, sweating, tremor	<i>Contraindications:</i> ACS, LQTS <i>Other considerations:</i> Short plasma half time (2â€”min)
Lidocaine	No significant effects	(VT/VF associated with ACS)	No oral use (50â€”200â€” mg bolus, then 2â€”4â€” mg/min)	<i>Cardiac:</i> Sinoatrial arrest <i>Extracardiac:</i> Central nervous system effects (e.g. drowsiness, dizziness)	<i>Precautions:</i> Reduced dose with reduced liver blood flow (e.g. shock, I²-blockade, severe heart failure) <i>Other considerations:</i> More effective with high potassium level. Few haemodynamic side effects
Mexiletine	No significant effects	PVC, VT (LQT3)	600â€”1200â€” mg Loading dose: 400â€”mg initially followed by 600â€”mg in the first 24â€”h	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, hypotension <i>Extracardiac:</i> Central nervous system effects (e.g. tremor, dysarthria, dizziness), gastrointestinal complaints	<i>Contraindications:</i> Sinus node dysfunction, severe AV conduction disturbances, severe heart failure

Procainamide	Prolongs PR interval, QRS duration, and QT interval ^a	VT	(100â€”150 mg bolus, can be repeated after 5â€”15 min if no effect, max 500â€”750 mg [max 50 mg/min]. Then, 2â€”6 mg/min)	<i>Cardiac:</i> Sinus bradycardia, hypotension, TdP <i>Extracardiac:</i> Rash, myalgia, vasculitis, systemic lupus, agranulocytosis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe intraventricular conduction disturbances, severe LV dysfunction hypotension, BrS
Propafenone	Prolongs PR interval, QRS duration, and QT interval ^a	PVC, VT	450â€”900 mg	<i>Cardiac:</i> Sinus bradycardia, AV block, negative inotrope, pro-arrhythmia (MVT, occasional TdP) <i>Extracardiac:</i> Gastrointestinal disturbances, headache, dry mouth	<i>Contraindications:</i> Prior MI, significant SHD, BrS, severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, LQTS, significant renal or liver disease <i>Other considerations:</i> Discontinue if QRS widening >25% or bundle branch block
Quinidine	Increases sinus node frequency and prolongs PR interval, QRS duration, and QT interval ^a	(VF ^c , BrS, SQTS)	600â€”1600 mg Loading dose: Start 200â€”300 mg every 3 h until effect, max. 3â€”24 g in first 24 h	<i>Cardiac:</i> Hypotension, TdP^d <i>Extracardiac:</i> Gastrointestinal disturbances, auditory and visual disturbances, confusion, leukopenia, haemolytic anaemia, thrombocytopenia, anaphylaxis	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV or intraventricular conduction disturbances, previous MI, significant SHD, hypotension, LQTS
Ranolazine	Decreases sinus node frequency, prolongs QT interval ^a	VT (LQTS3)	750â€”2000 mg	<i>Cardiac:</i> Sinus bradycardia, hypotension <i>Extracardiac:</i> Dizziness, nausea, constipation, gastrointestinal disturbance, headache, rash	<i>Contraindications:</i> Severe sinus node dysfunction, severe heart failure, LQTS (other than LQTS3) <i>Precautions:</i> Concomitant treatments associated with QT interval prolongation
Sotalol	Decreases sinus node frequency, prolongs QT interval ^a	VT	160â€”640 mg (0.5â€”1.5 mg/kg in 10 min. If necessary, can be repeated after 6 h)	See beta-blockers, TdP^d (>2% of patients, close monitoring of QT interval and CrCl)	<i>Contraindications:</i> Severe sinus node dysfunction, severe AV conduction disturbances, severe heart failure with reduced LVEF,

					significant LVH, CrCl <30 ml/min, coronary vasospasm, LQTS <i>Precautions:</i> Concomitant treatments associated with QT interval prolongation, hypokalaemia <i>Other considerations:</i> Potassium channel blocker effects require higher dose than beta-blocker effects
Verapamil	Prolongs PR interval	(LV fascicular tachycardia)	120–480 mg (50–100 mg in slow bolus. If necessary, can be repeated in 30 min)	<i>Cardiac:</i> Sinus bradycardia in sinus node dysfunction, AV block, negative inotrope, hypotension <i>Extracardiac:</i> Gastrointestinal disturbances, peripheral oedema, flushing	<i>Contraindications:</i> Heart failure with reduced LVEF, severe sinus node dysfunction, and severe AV conduction disturbances, VT of unknown origin, ACS, WPW syndrome <i>Other considerations:</i> Increase the risk of myopathy when used with statins

ACS, acute coronary syndrome; AF, atrial fibrillation; AV, atrioventricular; BrS, Brugada syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; CrCl, creatinine clearance; ECG, electrocardiogram; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; LVH, left ventricular hypertrophy; MI, myocardial infarction; MVT, monomorphic ventricular tachycardia; PVC, premature ventricular complex; SHD, structural heart disease; SQTS, short QT syndrome; TdP, torsades de pointes; VF, ventricular fibrillation; VT, ventricular tachycardia; WPW, Wolff-Parkinson-White.

a

Precaution with concomitant conditions or drugs that prolong QT interval. Should be discontinued if QTc >500 ms. See [Figure 13](#) Algorithm for evaluation before initiation and follow-up of patients requiring drugs associated with QT prolongation.

b

If ajmaline not available.

c

Subacute MI, multifocal ectopic Purkinje-related premature contractions, ERS, idiopathic VF.

d

Proarrhythmic side effects require strong indication in patients without ICD.

© ESC 2022

6.2.2. Device therapy

6.2.2.1. Implantable cardioverter defibrillator

ICD is an integral part of treating patients surviving a CA due to a VA or those deemed to be at high risk thereof. Drawbacks are the high up-front device costs, device-associated complications, and the relatively high number-to-treat needed to prevent one SCD in primary prevention.

A patient-level meta-analysis of the three early ICD trials^{349,351} comparing ICD to medical therapy for secondary prevention of SCD demonstrated a 28% mortality reduction (HR 0.72; 95% CI 0.63–0.87; $P = 0.0006$) almost entirely due to reduction of arrhythmic death (HR 0.5; 95% CI 0.37–0.67; $P < 0.0001$) in the ICD group.³⁵² Therefore, the use of an ICD for secondary prevention of SCD that occurred in the absence of reversible causes is widely accepted.

Several randomized controlled trials^{353–356} have established the role of the ICD for primary prevention of SCD in heart failure patients with an LVEF $\leq 35\%$. The reported mortality reduction has recently been supported by two large contemporary prospective registries enrolling more than 5000 patients.^{357,358} In the EU-CERT-ICD trial, primary prevention ICD implantation was associated with a 27% lower mortality with similar results in CAD and DCM.³⁵⁷ Results of the DANISH trial, however, indicate that the mortality benefit may be less clear in contemporary patients with non-ischaemic heart failure (see [Section 7.1.3.1](#)).³⁵⁹

In the work-up for ICD therapy, it is of paramount importance to consider the patient's life expectancy, quality of life, and comorbidities, and to reassess and discuss these issues with the patient at the time of generator change. There is evidence that patients with end-stage renal disease, with diabetes, and elderly patients benefit less or not at all from a primary prevention ICD.^{357,360,362} Women have been underrepresented in all primary prevention trials and data suggest that they may benefit less.³⁶¹ In general, the SCD risk needs to be weighed against the individual's competing risk of a non-arrhythmic death.^{363,364}

In the context of patient-centred medicine, physicians and healthcare professionals should engage ICD candidates and recipients in a joint decision process. Communication of relevant information, ensuring a good understanding of the benefits, risks, and potential consequences of different options, to enable the patient to partake in decision-making is mandatory. This shared decision-making process should discuss the different scenarios including primary prevention ICD implantation, consideration of ICD generator replacement, and end-of-life care. Importantly, the perception of a good quality of life depends on numerous factors that are weighed differently by people with different cultural, religious, and socioeconomic backgrounds. Although clinical prediction scores, such as the MADIT-ICD benefit score,³⁶⁵ may provide helpful additional information, clinical decision-making should not only rely on such scores.

Complications of ICD therapy include inappropriate therapies, lead fractures, and device-related infections. A subcutaneous implantable cardioverter defibrillator (S-ICD) has been introduced to address problems related to transvenous leads. S-ICD has no intravascular lead and therefore cannot deliver ATP. In the PRAETORIAN trial, 849 patients with an ICD but no pacing indication were randomized to an S-ICD or a transvenous ICD.³⁶⁶ Over a mean follow-up of 49 months, non-inferiority was shown for the primary endpoint of device-related complications and inappropriate shocks. The rate of inappropriate shocks was 9.7% in the S-ICD group and 7.3% in the ICD group (HR 1.43; 95% CI 0.89–2.30) and the rate of device related complications was 5.9% in the S-ICD and 9.8% in the ICD group (HR 0.69; 95% CI 0.44–1.09). There was also no difference in the secondary endpoint of death and appropriate shocks, but the trial had not been powered to show non-inferiority for this secondary endpoint. Of note, more than 80% of the included patients were in NYHA class I and II and the included patients were younger than in prior ICD trials.

Recommendation Table 11

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation (general aspects)

Recommendations	Class ^a	Level ^b
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C

© ESC 2022

ICD, implantable cardioverter defibrillator; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C

© ESC 2022

ICD, implantable cardioverter defibrillator; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 11

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation (general aspects)

Recommendations	Class ^a	Level ^b
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C

© ESC 2022

ICD, implantable cardioverter defibrillator; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C

© ESC 2022

ICD, implantable cardioverter defibrillator; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 12

[Open in new tab](#)

Recommendations for secondary prevention of sudden cardiac death*

Recommendations	Class ^a	Level ^b
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes. ^{349–352}	I	A
In patients with VT/VF, an indication for ICD, and no contraindication for amiodarone, amiodarone may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SMVT, sustained monomorphic VT; SPVT, sustained polymorphic VT; VF, ventricular fibrillation; VT, ventricular tachycardia.

*For primary prevention and specific aspects of secondary prevention, go to [Section 7](#).

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes. ^{349–352}	I	A
In patients with VT/VF, an indication for ICD, and no contraindication for amiodarone, amiodarone may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SMVT, sustained monomorphic VT; SPVT, sustained polymorphic VT; VF, ventricular fibrillation; VT, ventricular tachycardia.

*For primary prevention and specific aspects of secondary prevention, go to [Section 7](#).

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table 12

[Open in new tab](#)

Recommendations for secondary prevention of sudden cardiac death*

Recommendations	Class ^a	Level ^b
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes. ^{349–352}	I	A
In patients with VT/VF, an indication for ICD, and no contraindication for amiodarone, amiodarone may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SMVT, sustained monomorphic VT; SPVT, sustained polymorphic VT; VF, ventricular fibrillation; VT, ventricular tachycardia.

*For primary prevention and specific aspects of secondary prevention, go to [Section 7](#).

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes. ^{349–352}	I	A
In patients with VT/VF, an indication for ICD, and no contraindication for amiodarone, amiodarone may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C
In patients with SMVT or SPVT/VF triggered by a PVC with similar morphology and an indication for ICD, catheter ablation may be considered when an ICD is not available, contraindicated for concurrent medical reasons, or declined by the patient.	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SMVT, sustained monomorphic VT; SPVT, sustained polymorphic VT; VF, ventricular fibrillation; VT, ventricular tachycardia.

*For primary prevention and specific aspects of secondary prevention, go to [Section 7](#).

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table 13

[Open in new tab](#)

Recommendations for subcutaneous implantable cardioverter defibrillator

Recommendations	Class ^a	Level ^b
Subcutaneous defibrillator should be considered as an alternative to transvenous defibrillator in patients with an indication for an ICD when pacing therapy for bradycardia, cardiac resynchronization, or ATP is not needed. ³⁶⁶	Ila	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Subcutaneous defibrillator should be considered as an alternative to transvenous defibrillator in patients with an indication for an ICD when pacing therapy for bradycardia, cardiac resynchronization, or ATP is not needed. ³⁶⁶	Ila	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 13

[Open in new tab](#)

Recommendations for subcutaneous implantable cardioverter defibrillator

Recommendations	Class ^a	Level ^b
Subcutaneous defibrillator should be considered as an alternative to transvenous defibrillator in patients with an indication for an ICD when pacing therapy for bradycardia, cardiac resynchronization, or ATP is not needed. ³⁶⁶	Ila	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Subcutaneous defibrillator should be considered as an alternative to transvenous defibrillator in patients with an indication for an ICD when pacing therapy for bradycardia, cardiac resynchronization, or ATP is not needed. ³⁶⁶	Ila	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

6.2.2.2. Adding cardiac resynchronization therapy²

Cardiac resynchronization therapy (CRT) does reduce mortality in heart failure,³⁶⁷ and careful evaluation of the potential benefit of CRT in patients with ICD indication before implantation is mandatory.² The role of the addition of a defibrillator is less well established.^{368,369} The ongoing randomized controlled RESET-CRT trial aims to determine the impact of CRT-defibrillator on overall mortality and SCD in heart failure patients with a CRT indication.

Recommendation Table 14

[Open in new tab](#)

Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillator

Recommendations	Class ^a	Level ^b
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator. ³⁶⁷	I	C

CRT, cardiac resynchronization therapy; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator. ³⁶⁷	I	C

CRT, cardiac resynchronization therapy; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 14

[Open in new tab](#)

Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillator

Recommendations	Class ^a	Level ^b
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator. ³⁶⁷	I	C

CRT, cardiac resynchronization therapy; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator. ³⁶⁷	I	C

CRT, cardiac resynchronization therapy; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

6.2.2.3. Wearable cardioverter defibrillator

The wearable cardioverter defibrillator (WCD) is an external defibrillator that has been shown to successfully detect and treat VT and VF.³⁷⁰ It is therefore suitable for patients who are at risk but temporarily not candidates for an ICD owing, for example, to extraction of an infected device and subsequent antibiotic treatment.³⁷¹ An unsolved problem is the protection of patients in the early phase (40 days) after an MI. The VEST trial enrolled 2302 patients with acute MI and an LVEF $\leq 35\%$ and randomized them early in a 2:1 fashion to receive a WCD or not under guideline-directed optimal medical treatment (OMT).³⁷² After a follow-up of 90 days there was no difference in the primary endpoint of arrhythmic death (1.6 vs. 2.4%; RR 0.67; 95% CI 0.37â€“1.21; $P = 0.18$). Concern was raised regarding the low median wear time of 18 h (IQR 3.8â€“22.7). The median wear time was higher (23.4â–22.2â–23.8) in a recent multicentre registry after structured patient education.³⁷¹ However, based on the available data, the task force does not recommend routine use of the WCD in the early post-MI phase. Nevertheless, the use of the device may be considered in selected post-MI patients deemed to be at high risk for SCD.

Data on the benefit of the WCD for primary prevention of SCD in other clinical situations (e.g. acute myocarditis, primary prevention indication during pregnancy) are sparse and no recommendations can be currently made.

Recommendation Table 15

[Open in new tab](#)

Recommendations for wearable cardioverter defibrillator

Recommendations	Class ^a	Level ^b
The WCD should be considered for adult patients with a secondary prevention ICD indication, who are temporarily not candidates for ICD implantation.	IIa	C
The WCD may be considered in the early phase after MI in selected patients. ^{371,372}	IIb	B

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
The WCD should be considered for adult patients with a secondary prevention ICD indication, who are temporarily not candidates for ICD implantation.	IIa	C
The WCD may be considered in the early phase after MI in selected patients. ^{371,372}	IIb	B

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 15

[Open in new tab](#)

Recommendations for wearable cardioverter defibrillator

Recommendations	Class ^a	Level ^b
The WCD should be considered for adult patients with a secondary prevention ICD indication, who are temporarily not candidates for ICD implantation.	IIa	C
The WCD may be considered in the early phase after MI in selected patients. ^{371,372}	IIb	B

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
The WCD should be considered for adult patients with a secondary prevention ICD indication, who are temporarily not candidates for ICD implantation.	IIa	C

The WCD may be considered in the early phase after MI in selected patients.^{371,372}

IIb

B

© ESC 2022

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.

^aClass of recommendation.

^bLevel of evidence.

6.2.3. Special aspects of device therapy

6.2.3.1. Optimization of device programming

Optimization of ICD programming is essential to minimize the burden of ICD therapy and to improve patient outcome.³⁷³⁻³⁷⁵ Detailed recommendations are available in expert consensus papers.^{376,377} Bradycardia mode should be customized to prevent unnecessary RV pacing, thus reducing heart failure hospitalizations and all-cause mortality.³⁷⁸⁻³⁸⁰ (ESC CardioMed chapter 43.21).³⁸¹ A tachycardia detection programming strategy incorporating prolonged settings and high rate thresholds (e.g. 188 b.p.m. Advance III trial,³⁸² 200 MADIT-RIT³⁸³) reduces ICD therapies and all-cause mortality without increasing the risk of syncope.^{373,382,383} The evidence is stronger for primary prevention compared to secondary prevention indications.³⁸² (ESC CardioMed chapter 43.21).³⁸¹ Consistent use of SVT/VT discriminators even in rates up to 230 b.p.m. is recommended in ICD recipients without complete heart block to reduce inappropriate therapies.³⁸³⁻³⁸⁵ Proper atrial sensing is a prerequisite for activation of dual-chamber discriminators.³⁷⁶ (ESC CardioMed chapter 43.21).³⁸¹ In general, a multi-zone detection programming is favoured to enable tailored use of detection and therapy settings at different tachycardia rates.^{383,386} A single-zone programming with a high cut-off rate may be considered in patients with a high likelihood of VF only (e.g. primary electrical diseases).³⁸⁷ In subcutaneous ICDs, a dual-zone configuration should be adopted. Standardized programming including a lower conditional shock zone with activated discrimination algorithms and a higher shock zone based on rate criteria only has been shown to reduce inappropriate shock rates without compromising patient safety.³⁸⁸⁻³⁹⁰ Systematic use of ATP before shock delivery, also for very fast ventricular tachyarrhythmias, has been shown to reduce shock therapy without increase in arrhythmic syncope.^{375,384,391} Burst ATP is preferred over ramp due to increased efficacy in first tachycardia termination.³⁹² Incorporation of remote monitoring into follow-up practice of ICD recipients should be promoted to optimize surveillance of device integrity, to enable prompt detection and management of actionable events, and to prevent occurrence of inappropriate shocks.³⁹³⁻³⁹⁶ (ESC CardioMed chapter 43.21).³⁸¹ The proposed recommendations for optimal device programming are applicable to the majority of ICD recipients. Customization may be needed in individual patients.

Recommendation Table 16

[Open in new tab](#)

Recommendations for optimization of device programming

Recommendations	Class ^a	Level ^b
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality. ³⁷³⁻³⁷⁵	I	A

In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing. ^{378–380}	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals). ^{373,382,383}	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients. ^{382,383}	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones. ^{375,384,391}	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m. ^{383–385}	I	B
It is recommended to activate lead failure alerts. ^{397–399}	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks. ³⁹⁵	I	B
Programming of burst ATP as first attempt is recommended over ramp ATP. ³⁹²	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone. ^{388–390}	I	B
For routine ICD programming, activation of more than one tachycardia detection zone should be considered. ^{383,386}	IIa	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator; SHD, structural heart disease; S-ICD, subcutaneous ICD; SVT, supraventricular tachycardia; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality. ^{373–375}	I	A

In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing. ^{378–380}	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals). ^{373,382,383}	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients. ^{382,383}	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones. ^{375,384,391}	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m. ^{383–385}	I	B
It is recommended to activate lead failure alerts. ^{397–399}	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks. ³⁹⁵	I	B
Programming of burst ATP as first attempt is recommended over ramp ATP. ³⁹²	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone. ^{388–390}	I	B
For routine ICD programming, activation of more than one tachycardia detection zone should be considered. ^{383,386}	IIa	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator; SHD, structural heart disease; S-ICD, subcutaneous ICD; SVT, supraventricular tachycardia; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 16

[Open in new tab](#)

Recommendations for optimization of device programming

Recommendations	Class ^a	Level ^b
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality. ^{373–375}	I	A

In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing. ^{378–380}	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals). ^{373,382,383}	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients. ^{382,383}	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones. ^{375,384,391}	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m. ^{383–385}	I	B
It is recommended to activate lead failure alerts. ^{397–399}	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks. ³⁹⁵	I	B
Programming of burst ATP as first attempt is recommended over ramp ATP. ³⁹²	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone. ^{388–390}	I	B
For routine ICD programming, activation of more than one tachycardia detection zone should be considered. ^{383,386}	IIa	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator; SHD, structural heart disease; S-ICD, subcutaneous ICD; SVT, supraventricular tachycardia; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality. ^{373–375}	I	A

In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing. ^{378–380}	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals). ^{373,382,383}	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients. ^{382,383}	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones. ^{375,384,391}	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m. ^{383–385}	I	B
It is recommended to activate lead failure alerts. ^{397–399}	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks. ³⁹⁵	I	B
Programming of burst ATP as first attempt is recommended over ramp ATP. ³⁹²	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone. ^{388–390}	I	B
For routine ICD programming, activation of more than one tachycardia detection zone should be considered. ^{383,386}	IIa	B

ATP, anti-tachycardia pacing; ICD, implantable cardioverter defibrillator; SHD, structural heart disease; S-ICD, subcutaneous ICD; SVT, supraventricular tachycardia; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

6.2.3.2. Concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy

Apart from optimization of device programming, pharmacological and/or invasive management may prevent inappropriate ICD therapy. Beta-blockers (carvedilol superior to metoprolol in MADIT-CRT) should be titrated to the maximal tolerated dose in heart failure patients to reduce the risk of inappropriate therapy.⁴⁰⁰ In patients with inappropriate therapies due to recurrent SVT, catheter ablation should be first-line treatment, given its high success and low complication rate.^{302,401a,402,403} In case of AF-related inappropriate therapies, unresponsive to optimized pharmacological rate control treatment, individualized treatment strategy (rate vs. rhythm) dependent on patient characteristics is suggested.³⁴⁸ In patients with early AF, adoption of a rhythm control strategy improved patient outcome in the EAST-AFNET 4 trial.⁴⁰⁴ In CRT-defibrillator patients, AV node ablation has been associated with reduced inappropriate ICD shocks and lower hospitalization rates compared to drug treatment⁴⁰⁵ (ESC CardioMed chapter 41.14).⁴⁰⁶

Recommendation Table 17

[Open in new tab](#)

Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy

Recommendations	Class ^a	Level ^b
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies. ^{401,402}	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming. ^{401,405,407}	I	C

AF, atrial fibrillation; ICD, implantable cardioverter defibrillator; SVT, supraventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies. ^{401,402}	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming. ^{401,405,407}	I	C

AF, atrial fibrillation; ICD, implantable cardioverter defibrillator; SVT, supraventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 17

[Open in new tab](#)

Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy

Recommendations	Class ^a	Level ^b
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies. ^{401,402}	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming. ^{401,405,407}	I	C

AF, atrial fibrillation; ICD, implantable cardioverter defibrillator; SVT, supraventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies. ^{401,402}	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming. ^{401,405,407}	I	C

AF, atrial fibrillation; ICD, implantable cardioverter defibrillator; SVT, supraventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

6.2.3.3. Psychosocial impact of implantable cardioverter defibrillator treatment

Almost 20% of ICD recipients suffer from anxiety and depression that are associated with increased mortality.^{408,410,411} In ICD patients, psychological distress is mainly caused by the concern of potentially receiving ICD shocks rather than by having experienced an ICD shock.^{412,413} Thus, assessment of ICD concerns is recommended in all ICD patients before ICD shocks occur. Systematic screening of ICD patients for psychological distress is feasible with the use of specific questionnaires.^{413a,416}

A significant proportion of ICD patients with clinically relevant symptoms of anxiety and depression remains undertreated.⁴¹⁷ Communication with all ICD recipients is needed to clarify misconceptions about the device function, to discuss concerns about sexual functioning, driving restrictions, and to recommend an action plan in case of shock therapy.^{418,419} Referral to mental health professionals may be needed for specific interventions.^{418,420} Cognitive behavioural therapy intervention may also be provided by trained cardiac nurses to alleviate anxiety.⁴²¹ Web-based interventions may also prove useful in improving psychosocial well-being in ICD patients with increased psychosocial distress.⁴²²

Recommendation Table 18

[Open in new tab](#)

Recommendations for psychosocial management after implantable cardioverter defibrillator implantation

Recommendations	Class ^a	Level ^b
Assessment of psychological status and treatment of distress is recommended in ICD patients. ^{421–423}	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression. ^{412,424}	I	C

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Assessment of psychological status and treatment of distress is recommended in ICD patients. ^{421–423}	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression. ^{412,424}	I	C

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 18

[Open in new tab](#)

Recommendations for psychosocial management after implantable cardioverter defibrillator implantation

Recommendations	Class ^a	Level ^b
Assessment of psychological status and treatment of distress is recommended in ICD patients. ^{421–423}	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression. ^{412,424}	I	C

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Assessment of psychological status and treatment of distress is recommended in ICD patients. ^{421–423}	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression. ^{412,424}	I	C

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

6.2.3.4. Patients with left ventricular assist devices

VAs are common among left ventricular assist device (LVAD) carriers.^{425a,426,428} VAs are usually well-tolerated, as LVADs maintain adequate cardiac output and prevent circulatory collapse.⁴²⁹ However, sustained untreated VAs may lead to circulatory collapse even in the presence of an LVAD, especially early after device implantation and in patients with higher pulmonary vascular resistance.⁴³⁰ VAs pre- and post-LVAD implantation are associated with an increased risk of cardiovascular and all-cause mortality, while ICD can significantly reduce sustained VA.^{425,426,431a,432,434} These data support ICD implantation for secondary prevention in LVAD recipients with symptomatic VA.

Observational studies in patients with previous-generation, pulsatile LVADs reported a longer survival with ICD.^{435a,436,437} (📖 [ESC CardioMed chapter 37.32](#)).⁴³⁸

Recent registries enrolling continuous-flow LVAD carriers have questioned ICD mortality reduction, but available data are inconsistent.^{425,426,428,439,440} An analysis of the INTERMACS registry, enrolling the largest continuous-flow LVAD cohort, showed that ICD presence was not associated with prolonged survival.⁴²⁸ The lack of documented survival benefit among continuous-flow LVAD carriers, in conjunction with the likelihood of VA tolerance and the associated risks of ICD placement in these patients (risk of infection, device interaction), favours an individualized approach.

Recommendation Table 19

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in left ventricular assist device recipients

Recommendations	Class ^a	Level ^b
ICD implantation should be considered in LVAD recipients with symptomatic sustained VAs. ^{425,431}	IIa	B

ICD, implantable cardioverter defibrillator; LVAD, left ventricular assist devices; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b	© ESC 2022
ICD implantation should be considered in LVAD recipients with symptomatic sustained VAs. ^{425,431}	IIa	B	

ICD, implantable cardioverter defibrillator; LVAD, left ventricular assist devices; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table 19

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in left ventricular assist device recipients

Recommendations	Class ^a	Level ^b	© ESC 2022
ICD implantation should be considered in LVAD recipients with symptomatic sustained VAs. ^{425,431}	IIa	B	

ICD, implantable cardioverter defibrillator; LVAD, left ventricular assist devices; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b	© ESC 2022
ICD implantation should be considered in LVAD recipients with symptomatic sustained VAs. ^{425,431}	IIa	B	

ICD, implantable cardioverter defibrillator; LVAD, left ventricular assist devices; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

6.2.3.5. Complications of devices

Prevention of ICD complications is important to reduce associated morbidity, mortality, and financial burden. During implantation of devices, antibiotic prophylaxis, peri-procedural patient preparation, and appropriate surgical technique should be implemented to prevent device infections and formation of pocket haematoma.^{441a,442,443} Cephalic or axillary vein access is preferred over the subclavian vein route to reduce the risk of pneumothorax and lead failure.^{444a,445,446} Proper selection of ICD systems is important. Single-chamber ICDs are recommended in primary prevention patients without atrial or AV sequential pacing indications. This reduces peri-procedural complications and generator replacements as compared to dual-chamber ICDs. This approach does not increase the risk of inappropriate shocks if optimal device programming is used.^{447a,448,450} Routine use of single-coil defibrillator leads is favoured due to reduced risk of complications during lead removal without associated differences in shock efficacy.^{451,452} Use of dual-coil leads can be considered in clinical settings where a higher defibrillator threshold is suspected, e.g. HCM, right-sided implantations.^{453,454}

Recommendation Table 20

[Open in new tab](#)

Recommendations for prevention of implantable cardioverter defibrillator complications

Recommendations	Class ^a	Level ^b
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications. ^{447,448,450}	I	A
The use of single-coil leads over dual-coil ICD leads should be considered due to lower complication rate during transvenous lead extraction. ⁴⁵¹	IIa	C

AV, atrioventricular; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications. ^{447,448,450}	I	A
The use of single-coil leads over dual-coil ICD leads should be considered due to lower complication rate during transvenous lead extraction. ⁴⁵¹	IIa	C

AV, atrioventricular; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 20

[Open in new tab](#)

Recommendations for prevention of implantable cardioverter defibrillator complications

Recommendations	Class ^a	Level ^b
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications. ^{447,448,450}	I	A
The use of single-coil leads over dual-coil ICD leads should be considered due to lower complication rate during transvenous lead extraction. ⁴⁵¹	IIa	C

AV, atrioventricular; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications. ^{447,448,450}	I	A
The use of single-coil leads over dual-coil ICD leads should be considered due to lower complication rate during transvenous lead extraction. ⁴⁵¹	IIa	C

AV, atrioventricular; ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

6.2.3.6. End-of-life issues

Patients with active ICDs experience a considerable rate of shocks in the last phase of life.⁴⁵⁵ In terminally ill patients and at the end of life, healthcare professionals can facilitate a decision by the patient and the patient's family by explaining in a sensitive and understandable manner the benefits and burdens of continued ICD therapy.⁴⁵⁶ Patients should be informed about the options of ICD deactivation. In general, deactivation of anti-bradycardia therapy is discouraged to avoid quality-of-life impairment, and in some countries deactivation in pacemaker-dependent patients may be prohibited by law.⁴⁵⁷ Deactivation should be discussed before device implantation and when there is significant deterioration of the patient's health status. Despite an increasing trend to address and perform device deactivation after careful consideration, current rates are still low and improved patient care is needed.^{458,459}

Recommendation Table 21

[Open in new tab](#)

Recommendations for end-of-life issues in implantable cardioverter defibrillator carriers

Recommendations	Class ^a	Level ^b
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration. ⁴⁵⁸	I	C

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration. ⁴⁵⁸	I	C

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 21

[Open in new tab](#)

Recommendations for end-of-life issues in implantable cardioverter defibrillator carriers

Recommendations	Class ^a	Level ^b
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration. ⁴⁵⁸	I	C

© ESC 2022

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration. ⁴⁵⁸	I	C

© ESC 2022

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

6.2.4. Interventional therapy

6.2.4.1. Catheter ablation

6.2.4.1.1. Patients with structural heart disease

In patients with SHD, SMVTs are primarily due to a scar-related re-entrant mechanism.^{460a,461b,462c,463d,464e,465f}

Because of a higher risk of SCD, an ICD implantation is usually recommended in patients with sustained VAs associated with SHD.^{466a,467b,468c} However, ICDs do not prevent VA, and many of these patients will experience symptomatic VT/VF recurrences resulting in syncope or ICD shocks and may require additional treatment.^{330,383,419,469a,470b,471c}

In patients with SHD, the choice of anti-arrhythmic agents is mostly limited to beta-blockers, sotalol, and amiodarone to help control VT/VF recurrences, and this treatment is frequently hampered by side effects.^{318,472}

Owing to advances made over the last three decades, catheter ablation has become increasingly important for the management of scar-related VTs.⁴⁷³ Since the early 1990s, catheter ablation of BBR-VT has been very successful^{474a,475b,476c,477d} and is considered as first-line therapy^{466,467} (Figure A5). Subsequently, catheter ablation has been shown to be very effective in controlling incessant VTs or electrical storms^{330,331} and in reducing subsequent VT burden. Many observational studies have shown a positive effect of VT ablation on clinical outcome in terms of VT recurrences.^{478a,479b,480c,481d,482e,483f} In patients with CAD, three randomized trials^{471,484,485} have reported that catheter ablation, compared to conventional treatment, decreases the likelihood of subsequent ICD shocks and prevents recurrent VT episodes.

The critical part of re-entrant VT circuits, referred to as the *protected VT isthmus*,^{486a,487b,488c} is the primary target for ablation,^{460,486} but it is very challenging to unmask in haemodynamically not-tolerated VT.^{487,488} Due to the high likelihood of multiple re-entry circuits in a scar and difficulties in identifying critical isthmuses, the ablation strategy has gradually evolved over the years towards more extended ablation of the arrhythmogenic substrate.^{489a,490b,491c,492d} Special attention should be paid to cases of PVT/VF initiated by similar PVCs, where the triggering PVC (often related to the Purkinje network) should be targeted for ablation.^{221,332,333,493}

The electrophysiological characteristics of VT circuits depend on the underlying SHD. Thus, post-infarct VTs are mainly related to an endocardial VT circuit (amenable to endocardial ablation), while the location of re-entrant VT circuits is more variable in patients with cardiomyopathies.^{494a,495b,496c,497d,498e,499f} Here, intramural and/or epicardial involvement are more common. This significantly contributes to the difference in the clinical outcomes of VT ablation in relation to the underlying heart disease with a better outcome in CAD as compared to non-ischaemic aetiologies.^{497a,498b,499c}

Effective ablation requires durable ablation lesions to arrhythmogenic tissue. In some cases, such as intramural VTs, difficulties remain in achieving this objective with current catheters, regardless of the approach (endocardial/epicardial).⁵⁰⁰ To improve the formation of myocardial lesions, new catheter-based techniques are being

evaluated (e.g. bipolar/needle ablation, transcatheter alcohol ablation),^{501a,506} as well as radiotherapy ablation^{507,508} or surgical ablation,⁵⁰⁹ which are currently bailout treatments.

When planning VT ablation, it is important to collect all available information about the arrhythmogenic substrate, especially to identify scars (using CMR or CT scan),^{510a,514} and to help determine the exit site of VAs with the 12-lead ECG documentation of clinical VTs or PVCs that induce PVT/VF.

The mean long-term success rate of VT ablation varies from 30% to 70%, depending on the underlying SHD.^{481,515a,518} Peri-procedural complications, in particular stroke, cardiac tamponade, or death, may occur.^{519a,522}

6.2.4.1.2. Patients without apparent structural heart disease

Idiopathic VTs⁵²³ is the term for VTs that are not associated with SHD or a genetic arrhythmic syndrome. Most idiopathic VTs are mediated by triggered activity, but a re-entrant mechanism (involving the LV Purkinje network) explains verapamil-sensitive fascicular VTs.⁵²³ Three important key features distinguish idiopathic VTs from VTs associated with SHD. First, idiopathic VTs mostly originate from a single site and specific region of the heart (namely the right or left ventricular outflow tracts,^{524,525} along the valve annuli,^{526a,528} papillary muscle,⁵²⁹ or the LV Purkinje network).⁵²³ Second, no detectable scar is found in idiopathic VTs.⁵³⁰ Finally, idiopathic VTs have a benign prognosis so that ICD implantation is generally not recommended.⁴⁶⁶

The earliest site of activation during VT is the ablation target for focal sources, while abnormal Purkinje tissue (with diastolic activity during VT) is the ablation target of left fascicular VTs.^{531,532}

Catheter ablation is curative in most idiopathic VT patients and peri-procedural complications are rare.^{533a,537}

6.2.4.2. Autonomic modulation

The role of the autonomic nervous system in promoting arrhythmias has long been recognized, leading to the concept of *Coumel's triangle of arrhythmogenesis*.⁵³⁸

Sympathetic activation has been demonstrated as playing a key role in inducing VAs in some entities, such as congenital LQTS and CPVT,^{539,540} and left cardiac sympathetic denervation was shown to be associated with a decrease in the frequency of arrhythmogenic syncope in congenital LQTS.^{541,542} The efficacy of cardiac sympathetic blockade by different approaches (thoracic epidural anaesthesia, percutaneous stellate ganglion anaesthesia, or surgical stellate ganglion resection) in reducing the arrhythmia burden in refractory VT/VF has been recognized in several small observational studies.^{326,328,340} Further studies are needed to assess which patients may benefit from a modulation of the autonomic nervous system to better control VT/VF.

7. Diagnostic evaluation, management, and risk stratification according to clinical presentation and known (likely) disease

7.1. Specific structural heart diseases

7.1.1. Coronary artery disease

7.1.1.1. Acute coronary syndromes and vasospasm

7.1.1.1.1. Acute coronary syndromes

SCD is a major cause of mortality in ACS, mostly caused by sustained VAs, in particular VF. The majority of studies have reported on patients with STEMI. Of patients with STEMI, 4a⁵⁴³12% develop VA within the first 48 h after the onset of symptoms.^{69,543,544} Pre-reperfusion VAs are more common than reperfusion-induced or post-reperfusion arrhythmias in STEMI.⁵⁴⁵ Haemodynamic instability, cardiogenic shock, LVEF < 40%, and the sum of ST-segment deviations in all leads are independent predictors of VA both in STEMI and non-STEMI.^{69,546} In addition, an early repolarization pattern has been associated with increased risk of VAs and SCD in ACS.⁵⁴⁷

Prevention of ventricular arrhythmias in ST elevation myocardial infarction

Urgent reperfusion is the most important therapy,^{292,548} as acute ischaemia triggers arrhythmias. Beta-blocker treatment is also recommended to prevent VA.^{3,549} In a recent randomized trial of patients with STEMI, early intravenous metoprolol before percutaneous coronary intervention reduced the incidence of arrhythmias in the acute phase and was not associated with an increase in adverse events.⁵⁵⁰ Prophylactic treatment with AADs has not proven beneficial, and may even be harmful.³²² Correction of electrolyte imbalances is strongly recommended.²⁸⁹

Electrical cardioversion or defibrillation is the intervention of choice to acutely terminate VAs in ACS patients (Figure 14).^{205,339} Recurrent sustained VT, especially when polymorphic, or recurrent VF can indicate incomplete reperfusion or recurrence of acute ischaemia. In this case, immediate coronary angiography is indicated.³ For recurrent PVT degenerating into VF, beta-blockers are recommended.^{551,552} In addition, deep sedation may be helpful to reduce episodes of VT or VF.⁵⁵³ Intravenous amiodarone should be considered to acutely suppress recurrent haemodynamically relevant VAs, although there is a paucity of controlled studies for amiodarone during STEMI⁵⁵⁴ and the evidence in this setting is mainly extrapolated from studies of OHCA.⁵⁵⁵ If treatment with beta-blockers and amiodarone is not effective, lidocaine may be considered.³²² The use of other AADs in ACS is not recommended.^{549,556} In haemodynamically unstable patients with refractory VA, mechanical circulatory support may be considered.^{336,557} For VA in the context of relative bradycardia or pause-related, pacing may be effective to prevent re-initiation.




Figure 14

Algorithm for the prevention and management of ventricular arrhythmias in ST-elevation myocardial infarction.

CAG, coronary angiogram; PCI, percutaneous coronary intervention; STEMI, ST-elevation myocardial infarction; VA, ventricular arrhythmias. ^aThe 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation.³ ^bIntravenous beta-blockers must be avoided in patients with hypotension, acute heart failure, AV block or severe bradycardia. ^cFlowchart for the management of electrical storm. ^dIf similar PVC triggers recurrent polymorphic VA.

[Open in new tab](#)[Download slide](#)

The prognostic significance of early ventricular arrhythmias

Early VAs are defined as VT/VF occurring within 48 h after a STEMI. In the contemporary percutaneous coronary intervention (PCI)-based revascularization era, almost all VA occur within the first 24 h.⁵⁵⁸ Early VA have been associated with an up to six-fold increase in in-hospital mortality, whereas long-term prognosis seems not to be significantly affected.^{543,559,560} In a prospective cohort study, patients with VF during the acute STEMI phase had low and very similar incidence of late SCD as compared with VF-free patients during 5-year observation.⁵⁶⁰ Of note,

early *monomorphic* VT was associated with significantly higher incidence of adequate ICD interventions compared with early VF, and was an independent predictor of death during long-term follow-up.⁵⁶¹ Thus, the prognostic significance of VT and VF occurring in the acute phase of MI may be different. The impact of VA occurring late after reperfusion (>48 h) on late SCD is less clear.

Podolecki *et al.*⁵⁵⁹ recently demonstrated that long-term all-cause mortality after STEMI was predicted by VA occurring late after reperfusion (>48 h after reperfusion), while early reperfusion VAs did not affect 5-year outcome. Further studies are required to clarify the impact of VA occurring >48 h after STEMI on late SCD in contemporary patients undergoing acute PCI.

7.1.1.1.2. Vasospasm

Coronary artery spasms might have an important role in the pathogenesis of VA. The long-term prognosis of patients with variant angina who survive a CA is worse than in other patients with variant angina.^{562,563} In a recent multicentre European survey,⁵⁶⁴ patients with life-threatening VAs secondary to coronary vasospasm were at high risk for recurrence, especially when insufficient medical therapy was administered. While calcium channel blockers (CCB) are capable of suppressing episodes, beta-blockers may trigger VAs. As medical intervention and multiple vasodilator drugs may not be sufficiently protective, placement of an ICD should still be considered in SCA survivors with variant angina.

Recommendation Table 22

[Open in new tab](#)

Recommendations for treatment of ventricular arrhythmias in acute coronary syndrome and vasospasm

Recommendations	Class ^a	Level ^b
Treatment of VAs in ACS		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated. ^{551,552}	I	B
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS. ^{552,554,555}	IIa	C
Intravenous lidocaine may be considered for the treatment of recurrent PVT/VF not responding to beta-blockers or amiodarone, or if amiodarone is contraindicated during the acute phase of ACS. ⁵⁵⁴	IIb	C
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS. ³²²	III	B
Vasospasm		
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered. ^{562–564}	IIa	C

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ICD, implantable cardioverter defibrillator; PVT, polymorphic ventricular tachycardia; SCA, sudden cardiac arrest; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Treatment of VAs in ACS		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated. ^{551,552}	I	B
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS. ^{552,554,555}	IIa	C
Intravenous lidocaine may be considered for the treatment of recurrent PVT/VF not responding to beta-blockers or amiodarone, or if amiodarone is contraindicated during the acute phase of ACS. ⁵⁵⁴	IIb	C
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS. ³²²	III	B
Vasospasm		
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered. ^{562–564}	IIa	C

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ICD, implantable cardioverter defibrillator; PVT, polymorphic ventricular tachycardia; SCA, sudden cardiac arrest; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendation Table 22

[Open in new tab](#)

Recommendations for treatment of ventricular arrhythmias in acute coronary syndrome and vasospasm

Recommendations	Class ^a	Level ^b
Treatment of VAs in ACS		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated. ^{551,552}	I	B
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS. ^{552,554,555}	IIa	C
Intravenous lidocaine may be considered for the treatment of recurrent PVT/VF not responding to beta-blockers or amiodarone, or if amiodarone is contraindicated during the acute phase of ACS. ⁵⁵⁴	IIb	C
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS. ³²²	III	B
Vasospasm		
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered. ^{562–564}	IIa	C

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ICD, implantable cardioverter defibrillator; PVT, polymorphic ventricular tachycardia; SCA, sudden cardiac arrest; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Treatment of VAs in ACS		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated. ^{551,552}	I	B
Intravenous amiodarone treatment should be considered for patients with recurrent PVT/VF during the acute phase of ACS. ^{552,554,555}	IIa	C
Intravenous lidocaine may be considered for the treatment of recurrent PVT/VF not responding to beta-blockers or amiodarone, or if amiodarone is contraindicated during the acute phase of ACS. ⁵⁵⁴	IIb	C
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS. ³²²	III	B
Vasospasm		
In SCA survivors with coronary artery spasm, implantation of an ICD should be considered. ^{562–564}	IIa	C

AAD, anti-arrhythmic drug; ACS, acute coronary syndrome; ICD, implantable cardioverter defibrillator; PVT, polymorphic ventricular tachycardia; SCA, sudden cardiac arrest; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

7.1.1.2. Early after myocardial infarction

The first weeks after a STEMI carry the highest risk both for all-cause death and for SCD, particularly in patients with reduced LVEF.^{565,566} For this reason, early assessment of LVEF, i.e. before discharge, is recommended.^{567,568} Early routine prophylactic ICD implantation in the first 40 days after MI did not reduce mortality in post-MI patients with reduced LVEF in two randomized trials (DINAMIT and IRIS),^{569,570} and is therefore not recommended. Early assessment by further non-invasive tests, apart from LVEF measurement, has also not proven to be useful for risk stratification with regard to SCD.⁵⁷¹ Limited evidence suggests that invasive risk stratification by PES in the early post-MI phase may be helpful for identification of high-risk patients with reduced LVEF.⁵⁷² However, the utility of this approach has not been confirmed so far in randomized studies. The PROTECT-ICD randomized trial (NCT03588286) is currently examining whether PES may guide the decision on ICD implantation in patients with reduced EF in the early phase after STEMI.

Reverse remodelling following an MI is associated with significantly lower rates of death, SCA, and other adverse clinical outcomes.^{573,574} Therefore, assessment of the indication for prophylactic ICD implantation, usually performed by repeat echocardiography, should take place in the post-remodelling phase of the MI after the first 6 weeks in patients with a pre-discharge LVEF $\geq$ 40%. Reassessment of the LVEF before 6 weeks following MI may not discriminate between myocardial stunning and remodelling.

In the early post-MI phase, electrical storm and/or recurrent episodes of PVT or VF are immediately life-threatening conditions. In this setting, it is important to exclude ischaemia as the triggering factor for the arrhythmias. If medical treatment is not sufficient to suppress the arrhythmic episodes, catheter ablation is potentially effective particularly if the episodes are focally triggered by similar PVCs.^{332,575} If PVT recurs despite beta-blocker and amiodarone treatment, suppression by quinidine therapy has been reported.³²³

Recommendation Table 23

[Open in new tab](#)

Recommendations for risk stratification and treatment of ventricular arrhythmias early after myocardial infarction

Recommendations	Class ^a	Level ^b
Risk stratification		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI. ^{567,568}	I	B
In patients with pre-discharge LVEF ≤40%, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation. ^{568,573,574}	I	C
Treatment of VAs		
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC non-responsive to medical treatment or coronary revascularization in the subacute phase of MI. ³³²	IIa	C

ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Risk stratification		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI. ^{567,568}	I	B
In patients with pre-discharge LVEF ≤40%, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation. ^{568,573,574}	I	C
Treatment of VAs		
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC non-responsive to medical treatment or coronary revascularization in the subacute phase of MI. ³³²	IIa	C

ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendation Table 23

[Open in new tab](#)

Recommendations for risk stratification and treatment of ventricular arrhythmias early after myocardial infarction

Recommendations	Class ^a	Level ^b
Risk stratification		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI. ^{567,568}	I	B
In patients with pre-discharge LVEF ≤40%, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation. ^{568,573,574}	I	C
Treatment of VAs		
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC non-responsive to medical treatment or coronary revascularization in the subacute phase of MI. ³³²	IIa	C

ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Risk stratification		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI. ^{567,568}	I	B
In patients with pre-discharge LVEF ≤40%, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation. ^{568,573,574}	I	C
Treatment of VAs		
Catheter ablation should be considered in patients with recurrent episodes of PVT/VF triggered by a similar PVC non-responsive to medical treatment or coronary revascularization in the subacute phase of MI. ³³²	IIa	C

ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

7.1.1.3. Chronic coronary artery disease

7.1.1.3.1. Primary prevention of sudden cardiac death in patients with reduced ejection fraction

Forty days after STEMI, approximately 5% of patients will have an LVEF $\hat{a}_{\text{HT2}} \leq 35\%$.⁵⁷⁶ These patients are at risk for SCD. Therefore, in patients with LVEF $\hat{a}_{\text{HT2}} \leq 35\%$ and heart failure symptoms NYHA class II and III, primary preventive ICD implantation is recommended.³⁵⁶ ICD implantation should also be considered for asymptomatic patients with an EF $\hat{a}_{\text{HT2}} \leq 30\%$.³⁵⁴ In this population, mortality reduction by an ICD has been demonstrated in four RCTs.^{353a,354,356} In patients with CAD, a reduced LVEF ($\hat{a}_{\text{HT2}} \leq 40\%$), and asymptomatic NSVT, inducibility by PES identifies patients who benefit from an ICD, independent from NYHA class.³⁵⁵

Since the publication of the aforementioned trials, early revascularization strategies and modern heart failure medication have reduced the overall risk of SCD in heart failure patients.⁵⁷⁷ Although total mortality has been reduced, the relative reduction by the ICD is a consistent 27%, which has been corroborated in two recent large prospective registry studies enrolling 2327 European patients between 2014 and 2018 (EU-CERT-ICD)³⁵⁷ and 2610 Swedish patients enrolled between 2000 and 2016 (SwedeHF registry).³⁵⁸

7.1.1.3.2. Primary prevention of sudden cardiac death in patients with preserved or mildly reduced ejection fraction

There are no data supporting primary prophylactic ICD implantation in post-infarct patients with preserved or mildly reduced LVEF. These patients are heterogeneous with regard to their potential arrhythmic substrate, and efforts are under way to identify those with the highest risk for SCD. PES is recommended in post-infarct patients in whom syncope remains unexplained after non-invasive evaluation to guide patient management ([Figure 15](#)).¹⁴⁶




Figure 15

Algorithm for risk stratification and primary prevention of sudden cardiac death in patients with chronic coronary artery disease and reduced ejection fraction.

CAD, coronary artery disease; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; LVEF, left ventricular ejection fraction; MI, myocardial infarction; N, No; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; PES, programmed electrical stimulation; SMVT, sustained monomorphic ventricular tachycardia; Y, Yes. ^aThe 2018 ESC Guidelines for the diagnosis and management of syncope.¹

[Open in new tab](#)[Download slide](#)

In the PRESERVE-EF study, 41 of 575 post-infarct patients with an LVEF $\geq 40\%$ and one non-invasive ECG risk factor more than 40 days post-MI were inducible for VT/VF during PES and received an ICD.¹⁵¹ During the 32-month follow-up period, no SCD occurred and 9 of 37 ICD patients received an appropriate ICD therapy. However, the role of appropriate ICD treatment as surrogate for SCD in patients with preserved LVEF is unknown, and randomized trials are needed. Prophylactic treatment with AADs other than beta-blockers is not indicated regardless of the LVEF.^{556,578,579}

7.1.1.3.3. Secondary prevention of sudden cardiac death

The three pivotal secondary prevention ICD trials enrolled 1866 patients between 1990 and 1997.^{349,350,351} A patient-level meta-analysis demonstrated a 28% mortality reduction (HR 0.72; 95% CI 0.60–0.87; $P = 0.0006$) almost entirely due to reduction of arrhythmic death (HR 0.50; 95% CI 0.37–0.67; $P < 0.0001$) in the ICD group.³⁵² This translates to an extension of survival of 4.4 months over a mean follow-up of 6 years by the ICD. Around 80% of the study population suffered from CAD. Patients with well-tolerated SMVT were excluded from secondary prevention trials ([Figure 16](#)).




Figure 16

Algorithm for the management of sustained monomorphic ventricular tachycardia in patients with chronic coronary artery disease.

CAD, coronary artery disease; ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; N, No; SMVT, sustained monomorphic ventricular tachycardia; Y, Yes. ^aIncessant ventricular tachycardia in monitor zone: consider catheter ablation. ^bIf catheter ablation is not available, not successful or not desired by the patient. ^cTo reduce ICD shocks.

[Open in new tab](#)[Download slide](#)

7.1.1.3.4. Management of patients with haemodynamically tolerated ventricular tachycardia and preserved and mildly reduced ejection fraction

With the better understanding of the mechanisms of VT after MI, as well as improved ablation and imaging technologies, catheter ablation has become an option for the treatment of haemodynamically well-tolerated VT in selected post-MI patients with preserved or mildly reduced EF, even without ICD back-up. One small monocentric retrospective trial studied patients with CAD, LVEF >40%, and haemodynamically tolerated VT who underwent catheter ablation as first-line therapy.⁵⁸⁰ The investigators could abolish 90% of the clinical and 58% of all inducible VTs. Subsequently, 42% of the patients received an ICD. After a mean follow-up of 3.8 years, 42% of the patients died irrespective of the presence or absence of an ICD ($P = 0.47$).

A larger multicentre retrospective study looked at 166 patients with an LVEF >30% presenting with well-tolerated SMVT, which were treated by catheter ablation only, and compared the outcome to a control group of 378 patients implanted with an ICD.⁴⁸⁰ Of the 166 patients undergoing ablation as first therapy, 55% suffered from CAD. The mean LVEF was 50%, and after a mean follow-up of 32 months, overall mortality did not differ between the groups (12%).

These data suggest that either ICD implantation or ablation in experienced centres should be considered in patients with a preserved or mildly reduced EF presenting with haemodynamically tolerated SMVT. Of note, although ICDs have been commonly implanted in this population, the secondary preventive ICD trials failed to show a survival benefit in patients with an LVEF $\geq 35\%$.³⁵² Although SMVT is rarely caused by ischaemia and revascularization alone does not prevent VT recurrence, exclusion or treatment of significant CAD before catheter ablation is reasonable.

7.1.1.3.5. Management of recurrent ventricular tachycardia in implantable cardioverter defibrillator carriers

Frequent, symptomatic VT in ICD recipients should be treated medically with either amiodarone or sotalol.^{318,581} In patients with CAD in whom SMVT recurs while on amiodarone treatment, catheter ablation is recommended over escalation of AAD therapy. In the VANISH trial, the composite endpoint of death, VT storm, and appropriate ICD therapy was reached significantly less often in the ablation group as compared to the escalated amiodarone-therapy group over a mean follow-up of 28 months (59% vs. 68.5%; HR 0.72; 95% CI 0.53–0.98; $P = 0.04$).⁴⁷¹ The

ongoing VANISH2 trial (ClinicalTrials.gov Identifier: NCT02830360) addresses the question of whether catheter ablation as first-line treatment is superior to AAD therapy in post-MI patients with SMVT.

Preventive VT ablation after a first documented SMVT, which was followed by ICD implantation, reduced neither mortality nor hospitalizations for arrhythmia or worsening heart failure when compared to a deferred ablation strategy only after the third ICD shock.⁵⁸² However, in patients who present with a first VT episode and in whom an ICD is indicated, catheter ablation performed immediately before or shortly after ICD implantation may be considered to decrease subsequent VT and ICD shocks.^{484,485,583}

Recommendation Table 24

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in chronic coronary artery disease

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation. ^{146,584}	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF ≤35% despite ≥3 months of OMT. ^{354,356}	I	A
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF ≤30% despite ≥3 months of OMT. ³⁵⁴	IIa	B
ICD implantation should be considered in patients with CAD, LVEF ≤40% despite ≥3 months of OMT, and NSVT, if they are inducible for SMVT by PES. ³⁵⁵	IIa	B
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended. ^{556,578,579}	III	A
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI. ^{349–351}	I	A
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy. ⁴⁷¹	I	B

The addition of oral amiodarone or beta-blocker replacement by sotalol should be considered in patients with CAD with recurrent, symptomatic SMVT, or ICD shocks for SMVT while on beta-blocker treatment. ^{318,581}	IIa	B
In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^{6,480,580}	IIa	C
ICD implantation should be considered in patients with a haemodynamically tolerated SMVT and an LVEF $\geq 40\%$ if VT ablation fails, is not available, or is not desired.	IIa	C
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blockers or sotalol treatment. ⁴⁷¹	IIa	C
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks. ^{484,485,582,583}	IIb	B

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation. ^{146,584}	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT. ^{354,356}	I	A
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT. ³⁵⁴	IIa	B
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT, and NSVT, if they are inducible for SMVT by PES. ³⁵⁵	IIa	B
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended. ^{556,578,579}	III	A
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI. ^{349–351}	I	A
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy. ⁴⁷¹	I	B

The addition of oral amiodarone or beta-blocker replacement by sotalol should be considered in patients with CAD with recurrent, symptomatic SMVT, or ICD shocks for SMVT while on beta-blocker treatment. ^{318,581}	IIa	B
In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^{c,480,580}	IIa	C
ICD implantation should be considered in patients with a haemodynamically tolerated SMVT and an LVEF $\geq 40\%$ if VT ablation fails, is not available, or is not desired.	IIa	C
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blockers or sotalol treatment. ⁴⁷¹	IIa	C
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks. ^{484,485,582,583}	IIb	B

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

Recommendation Table 24

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in chronic coronary artery disease

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation. ^{146,584}	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT. ^{354,356}	I	A
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT. ³⁵⁴	IIa	B
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT, and NSVT, if they are inducible for SMVT by PES. ³⁵⁵	IIa	B
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended. ^{556,578,579}	III	A
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI. ^{349–351}	I	A
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy. ⁴⁷¹	I	B

The addition of oral amiodarone or beta-blocker replacement by sotalol should be considered in patients with CAD with recurrent, symptomatic SMVT, or ICD shocks for SMVT while on beta-blocker treatment. ^{318,581}	IIa	B
In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^{6,480,580}	IIa	C
ICD implantation should be considered in patients with a haemodynamically tolerated SMVT and an LVEF $\geq 40\%$ if VT ablation fails, is not available, or is not desired.	IIa	C
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blockers or sotalol treatment. ⁴⁷¹	IIa	C
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks. ^{484,485,582,583}	IIb	B

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation. ^{146,584}	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT. ^{354,356}	I	A
ICD therapy should be considered in patients with CAD, NYHA class I, and LVEF $\leq 30\%$ despite ≥ 3 months of OMT. ³⁵⁴	IIa	B
ICD implantation should be considered in patients with CAD, LVEF $\leq 40\%$ despite ≥ 3 months of OMT, and NSVT, if they are inducible for SMVT by PES. ³⁵⁵	IIa	B
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended. ^{556,578,579}	III	A
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI. ^{349–351}	I	A
In patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy. ⁴⁷¹	I	B

The addition of oral amiodarone or beta-blocker replacement by sotalol should be considered in patients with CAD with recurrent, symptomatic SMVT, or ICD shocks for SMVT while on beta-blocker treatment. ^{318,581}	IIa	B
In patients with CAD and haemodynamically well-tolerated SMVT and LVEF $\geq 40\%$, catheter ablation in experienced centres should be considered as an alternative to ICD therapy, provided that established endpoints have been reached. ^{480,580}	IIa	C
ICD implantation should be considered in patients with a haemodynamically tolerated SMVT and an LVEF $\geq 40\%$ if VT ablation fails, is not available, or is not desired.	IIa	C
Catheter ablation should be considered in patients with CAD and recurrent, symptomatic SMVT, or ICD shocks for SMVT despite beta-blockers or sotalol treatment. ⁴⁷¹	IIa	C
In patients with CAD eligible for ICD implantation, catheter ablation may be considered just before (or immediately after) ICD implantation to decrease subsequent VT burden and ICD shocks. ^{484,485,582,583}	IIb	B

© ESC 2022

AAD, anti-arrhythmic drug; CAD, coronary artery disease; ICD, implantable cardioverter defibrillator; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; STEMI, ST elevation myocardial infarction; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

7.1.1.4. Coronary anomalies

Anomalous aortic origin of a coronary artery, either the left or the right, arising from the opposite sinus of Valsalva, is associated with an increased risk of SCD, especially in individuals <35 years during or following vigorous exercise.⁵⁸⁵ Anomalous aortic origin of the left coronary artery is less common but more malignant than anomalous aortic origin of the right coronary artery. Other risk factors for SCD are interarterial course between the aorta and pulmonary artery, slit-like shaped orifice, high orifice, acute-angle take-off, and intramural course and its length.^{585,586} Indications for surgical intervention, especially in asymptomatic patients, are based on the evaluation of high-risk anatomy by CTA and the assessment of exercise-induced ischaemia using advanced imaging modalities.^{586a,587,588} Cardiac stress imaging is also indicated to evaluate exercise-induced ischaemia after surgical intervention, especially in patients with an aborted CA.⁵⁸⁸

Recommendation Table 25

[Open in new tab](#)

Recommendations for sudden cardiac death prevention in patients with coronary anomalies

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia. ⁵⁸⁷	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Treatment		
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs, or angina when other causes have been excluded. ^{585,586,588}	I	C
Surgery should be considered in asymptomatic patients with anomalous aortic origin of a coronary artery and evidence of myocardial ischaemia or abnormal aortic origin of the left coronary artery with high-risk anatomy. ^{c,585,586,588}	IIa	C

© ESC 2022

CA, cardiac arrest; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

^cHigh-risk anatomy is defined as interarterial course, slit-like shaped orifice, high orifice, acute-angle take-off, and intramural course and its length.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia. ⁵⁸⁷	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Treatment		
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs, or angina when other causes have been excluded. ^{585,586,588}	I	C
Surgery should be considered in asymptomatic patients with anomalous aortic origin of a coronary artery and evidence of myocardial ischaemia or abnormal aortic origin of the left coronary artery with high-risk anatomy. ^{c,585,586,588}	IIa	C

CA, cardiac arrest; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.
^cHigh-risk anatomy is defined as interarterial course, slit-like shaped orifice, high orifice, acute-angle take-off, and intramural course and its length.

© ESC 2022

Recommendation Table 25

[Open in new tab](#)

Recommendations for sudden cardiac death prevention in patients with coronary anomalies

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia. ⁵⁸⁷	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Treatment		
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs, or angina when other causes have been excluded. ^{585,586,588}	I	C
Surgery should be considered in asymptomatic patients with anomalous aortic origin of a coronary artery and evidence of myocardial ischaemia or abnormal aortic origin of the left coronary artery with high-risk anatomy. ^{c,585,586,588}	IIa	C

© ESC 2022

CA, cardiac arrest; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

^cHigh-risk anatomy is defined as interarterial course, slit-like shaped orifice, high orifice, acute-angle take-off, and intramural course and its length.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia. ⁵⁸⁷	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Treatment		
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs, or angina when other causes have been excluded. ^{585,586,588}	I	C
Surgery should be considered in asymptomatic patients with anomalous aortic origin of a coronary artery and evidence of myocardial ischaemia or abnormal aortic origin of the left coronary artery with high-risk anatomy. ^{c,585,586,588}	IIa	C

CA, cardiac arrest; VA, ventricular arrhythmia.
^aClass of recommendation.
^bLevel of evidence.
^cHigh-risk anatomy is defined as interarterial course, slit-like shaped orifice, high orifice, acute-angle take-off, and intramural course and its length.

© ESC 2022

7.1.2. Idiopathic premature ventricular complexes/ventricular tachycardia and premature ventricular complex-induced cardiomyopathy

7.1.2.1. Idiopathic premature ventricular complexes/ventricular tachycardia

PVCs/VT in patients without SHD are defined as idiopathic ([Figure 17](#)). In patients with presumed idiopathic PVCs/VT based on a negative history and normal physical examination, 12-lead ECG and transthoracic echocardiography are important first diagnostic steps to exclude underlying SHD. Twenty-four-hour ECG Holter monitoring is usually performed to determine the PVC burden. Multiform PVCs at prolonged ECG monitoring and subtle changes on ECG or echocardiography need to be recognized.⁵⁸⁹ CMR should be performed whenever ECG and echocardiography are inconclusive to rule out SHD, or the clinical presentation raises suspicion for SHD.^{590,591}




Figure 17

Algorithm for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia and non-apparent structural heart disease.

CCB, calcium channel blocker; CMR, cardiac magnetic resonance; ECG, electrocardiogram; LVEF, left ventricular ejection fraction; N, No; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; SHD, structural heart disease; VT, ventricular tachycardia; Y, Yes. ^aNon-apparent SHD is defined by lack of significant abnormalities in physical examination, basal ECG, and echocardiogram. ^bAtypical presentation: e.g. older age, right bundle branch block morphology, sustained monomorphic VT consistent with re-entry. ^cSymptoms should be relevant and related to PVC/VT. ^dOrigin suspected by ECG or confirmed during electrophysiological evaluation. ^eConsider re-evaluation in case of new symptoms or changes in patient clinical condition.

[Open in new tab](#)[Download slide](#)

Patients need to be treated when PVCs/VT are symptomatic or associated with deterioration of cardiac function. The clinical course and responses to different treatments have been mostly studied in those originating from the RVOT or the left fascicles.

Several drugs have been used to treat idiopathic PVCs/VT. Recommendations are based on small or non-controlled series. Beta-blockers and CCBs are the most-studied drugs and both have been shown to be effective at arrhythmia suppression.³⁰⁴ The evidence for flecainide is scarce.⁵⁹² In case of a higher burden of PVCs with higher heart rate or during exercise, beta-blockers should be preferred.⁵⁹³ If there is not such a correlation, the use of flecainide or CCB drugs have been associated with more effective PVC suppression. Beta-blockers should also be selected when a focal triggered activity mechanism is suspected. CCB should be the drug of choice for fascicular PVC/VT. Although data are lacking, beta-blockers or CCBs are considered first choice for PVCs with an origin outside the RVOT or the left fascicles because flecainide may have proarrhythmic side effects. Amiodarone is associated with severe systemic toxicity and should be used only if ablation or other drugs fail or cannot be used.⁵⁹⁴ A summary of the recommendations for the treatment of idiopathic PVCs/VT and PVC-induced or aggravated cardiomyopathy with AADs is provided in [Table 9](#).

Table 9

[Open in new tab](#)

Summary of the recommendations for the treatment of patients with frequent idiopathic premature ventricular complexes/ventricular tachycardia or premature ventricular complex-induced cardiomyopathy

	Ablation	Beta-blocker	CCB	Flecainide	Amiodarone
RVOT/fascicular PVC/VT: Symptomatic, normal LV function	Class I	Class IIa	Class IIa	Class IIa	Class III
PVC/VT other than RVOT/fascicular: Symptomatic, normal LV function	Class IIa	Class I	Class I	Class IIa	Class III
RVOT/fascicular PVC/VT: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC/VT other than RVOT/fascicular: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC: Burden >20%, asymptomatic, normal LV function	Class IIb				Class III

© ESC 2022

CCB, calcium channel blocker; LV, left ventricular; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; VT, ventricular tachycardia.
^aIntravenous calcium channel blockers.
^bIn selected patients (only moderate LV dysfunction).

	Ablation	Beta-blocker	CCB	Flecainide	Amiodarone
RVOT/fascicular PVC/VT: Symptomatic, normal LV function	Class I	Class IIa	Class IIa	Class IIa	Class III
PVC/VT other than RVOT/fascicular: Symptomatic, normal LV function	Class IIa	Class I	Class I	Class IIa	Class III
RVOT/fascicular PVC/VT: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC/VT other than RVOT/fascicular: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC: Burden >20%, asymptomatic, normal LV function	Class IIb				Class III

© ESC 2022

CCB, calcium channel blocker; LV, left ventricular; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; VT, ventricular tachycardia.
^aIntravenous calcium channel blockers.
^bIn selected patients (only moderate LV dysfunction).

Table 9

[Open in new tab](#)

Summary of the recommendations for the treatment of patients with frequent idiopathic premature ventricular complexes/ventricular tachycardia or premature ventricular complex-induced cardiomyopathy

	Ablation	Beta-blocker	CCB	Flecainide	Amiodarone
RVOT/fascicular PVC/VT: Symptomatic, normal LV function	Class I	Class IIa	Class IIa	Class IIa	Class III
PVC/VT other than RVOT/fascicular: Symptomatic, normal LV function	Class IIa	Class I	Class I	Class IIa	Class III
RVOT/fascicular PVC/VT: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC/VT other than RVOT/fascicular: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC: Burden >20%, asymptomatic, normal LV function	Class IIb				Class III

© ESC 2022

CCB, calcium channel blocker; LV, left ventricular; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; VT, ventricular tachycardia.
^aIntravenous calcium channel blockers.
^bIn selected patients (only moderate LV dysfunction).

	Ablation	Beta-blocker	CCB	Flecainide	Amiodarone
RVOT/fascicular PVC/VT: Symptomatic, normal LV function	Class I	Class IIa	Class IIa	Class IIa	Class III
PVC/VT other than RVOT/fascicular: Symptomatic, normal LV function	Class IIa	Class I	Class I	Class IIa	Class III
RVOT/fascicular PVC/VT: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC/VT other than RVOT/fascicular: LV dysfunction	Class I	Class IIa	Class III*	Class IIa ^b	Class IIa
PVC: Burden >20%, asymptomatic, normal LV function	Class IIb				Class III

© ESC 2022

CCB, calcium channel blocker; LV, left ventricular; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; VT, ventricular tachycardia.
^aIntravenous calcium channel blockers.
^bIn selected patients (only moderate LV dysfunction).

A high success rate of catheter ablation of idiopathic PVCs/VT has been reported with rare complications, particularly for the RVOT and fascicular types.⁵³⁵ In a randomized study including patients with RVOT PVCs, ablation was superior to AAD therapy for arrhythmia suppression with no differences in complications.⁵⁹⁵ Ablation is therefore recommended as the first-line therapy for RVOT and fascicular PVCs/VT. The available information for other forms of idiopathic PVCs/VT is limited and mostly restricted to the acute success of ablation, which, in general, is lower and associated with more recurrences than for RVOT and fascicular PVCs/VT.⁵⁴⁰ In addition, access to and ablation at specific locations (e.g. sinus of Valsalva, LV summit) may increase the risk for procedural complications. Therefore, when the 12-lead ECG is highly suspicious of a PVC/VT source outside the RVOT or left fascicles, the level of recommendation for ablation is lower.

In general, treatment of children should be similar to that of adults. However, ablation should be deferred in young and small children due to the risk of complications and the relatively larger size of the ablation lesion as compared to the child's heart.^{596,597} Verapamil is not recommended as the first-line therapy in children <1 year old because it has been associated with hypotension in some case reports.⁵⁴² Of note, all reported patients had either heart failure, overdosing of verapamil, or other concurrent AADs at the time verapamil was given.⁵⁹⁸

Patients may present with asymptomatic frequent PVCs/VTs. Only a minority of patients with >1000 PVCs per day will develop ventricular dysfunction after 5 years follow-up.⁵⁹⁹ A PVC burden of 10% seems to be the minimal threshold for development of LV dysfunction, with higher risk when the PVC burden is >20%.^{535,600,601} Regular

assessment of LVEF in this setting is therefore indicated. To date, there are no data supporting the benefit of arrhythmia treatment for asymptomatic patients with preserved ventricular function. In addition, PVC burden often decreases spontaneously over time, particularly in children.^{599,602} In selected patients, e.g. patients who do not want to be followed, catheter ablation may be considered. In patients with PVC burden < 10%, re-evaluation may be appropriate in case of development of new symptoms or change in patient condition.

Recommendation Table 26

[Open in new tab](#)

Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia

Recommendations	Class ^a	Level ^b
General recommendations		
Regular assessment of ventricular function of patients with PVC burden >10% and normal ventricular function is indicated. ^{602,603}	I	C
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram. ¹⁹⁵	IIa	C
Treatment		
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles. ^{d,535,595,596,604}	I	B
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{304,593}	I	C
Beta-blockers, non-dihydropyridine CCBs, or flecainide should be considered when catheter ablation is not available, desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles. ^{304,592,593}	IIa	B
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{535,604,605}	IIa	C
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up. ^{535,600,601}	IIb	B
Catheter ablation of idiopathic VT/PVCs is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated. ⁵⁹⁷	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs. ⁵⁹⁴	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs. ⁶⁰⁶	III	C

© ESC 2022

AAD, anti-arrhythmic drug; CCB, calcium channel blocker; CMR, cardiac magnetic resonance; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cIncluding but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry.

^dLevel of evidence C for VT/PVCs from left fascicles.

Recommendations	Class ^a	Level ^b
General recommendations		
Regular assessment of ventricular function of patients with PVC burden >10% and normal ventricular function is indicated. ^{602,603}	I	C
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram. ¹⁹⁵	IIa	C
Treatment		
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles. ^{d,535,595,596,604}	I	B
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{304,593}	I	C
Beta-blockers, non-dihydropyridine CCBs, or flecainide should be considered when catheter ablation is not available, desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles. ^{304,592,593}	IIa	B
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{535,604,605}	IIa	C
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up. ^{535,600,601}	IIb	B
Catheter ablation of idiopathic VT/PVCs is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated. ⁵⁹⁷	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs. ⁵⁹⁴	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs. ⁶⁰⁶	III	C
AAD, anti-arrhythmic drug; CCB, calcium channel blocker; CMR, cardiac magnetic resonance; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia. ^a Class of recommendation. ^b Level of evidence. ^c Including but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry. ^d Level of evidence C for VT/PVCs from left fascicles.		

© ESC 2022

Recommendation TableÂ 26

[Open in new tab](#)

Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia

Recommendations	Class ^a	Level ^b
General recommendations		
Regular assessment of ventricular function of patients with PVC burden >10% and normal ventricular function is indicated. ^{602,603}	I	C
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram. ¹⁹⁵	IIa	C
Treatment		
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles. ^{d,535,595,596,604}	I	B
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{304,593}	I	C
Beta-blockers, non-dihydropyridine CCBs, or flecainide should be considered when catheter ablation is not available, desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles. ^{304,592,593}	IIa	B
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{535,604,605}	IIa	C
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up. ^{535,600,601}	IIb	B
Catheter ablation of idiopathic VT/PVCs is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated. ⁵⁹⁷	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs. ⁵⁹⁴	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs. ⁶⁰⁶	III	C

© ESC 2022

AAD, anti-arrhythmic drug; CCB, calcium channel blocker; CMR, cardiac magnetic resonance; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cIncluding but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry.

^dLevel of evidence C for VT/PVCs from left fascicles.

Recommendations	Class ^a	Level ^b
General recommendations		
Regular assessment of ventricular function of patients with PVC burden >10% and normal ventricular function is indicated. ^{602,603}	I	C
In patients with PVCs/VT and a presentation not typical for an idiopathic origin, ^c CMR should be considered, despite a normal echocardiogram. ¹⁹⁵	IIa	C
Treatment		
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles. ^{d,535,595,596,604}	I	B
Beta-blockers or non-dihydropyridine CCBs are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{304,593}	I	C
Beta-blockers, non-dihydropyridine CCBs, or flecainide should be considered when catheter ablation is not available, desired, or is particularly risky in symptomatic patients with idiopathic VT/PVCs from the RVOT or the left fascicles. ^{304,592,593}	IIa	B
Catheter ablation or flecainide should be considered in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles. ^{535,604,605}	IIa	C
Catheter ablation may be considered for idiopathic VT/PVCs in asymptomatic patients with repeatedly more than 20% of PVCs per day at follow-up. ^{535,600,601}	IIb	B
Catheter ablation of idiopathic VT/PVCs is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated. ⁵⁹⁷	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs. ⁵⁹⁴	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs. ⁶⁰⁶	III	C
AAD, anti-arrhythmic drug; CCB, calcium channel blocker; CMR, cardiac magnetic resonance; PVC, premature ventricular complex; RVOT, right ventricular outflow tract; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia. ^a Class of recommendation. ^b Level of evidence. ^c Including but not limited to older age, right bundle branch block (RBBB) morphology, SMVT consistent with re-entry. ^d Level of evidence C for VT/PVCs from left fascicles.		

© ESC 2022

7.1.2.2. Premature ventricular complex-induced/-aggravated cardiomyopathy

The importance of PVC-induced cardiomyopathy as a secondary and reversible cause of LV dysfunction in patients without SHD has been recognized.^{607,608} The patient's medical and family history, 12-lead ECG, Holter-ECG, and echocardiography form the cornerstones of the evaluation of patients with suspected PVC-induced cardiomyopathy (Figure 18). PVC burden has been shown to be the strongest independent predictor of PVC-induced cardiomyopathy in several studies.^{600,609,610,611} Day-by-day fluctuations of PVC burden have been reported in

patients undergoing 14-day monitoring, but most data are based on 24-h registrations.⁶¹¹ A PVC burden of at least 10% appears to be the minimal threshold for development of PVC-induced cardiomyopathy, and the risk increases further with a PVC burden >20%.^{600,611} In patients with a PVC burden <10%, other cardiomyopathy aetiologies should be suspected and further diagnostic work-up undertaken. In such patients, Holter-ECG should be repeated to assess fluctuations in PVC burden. Factors predicting adverse LV remodelling in patients with frequent PVCs include superior PVC axis, epicardial origin, NSVT, shorter coupling interval, and male gender.^{535,611a,613}




Figure 18

Algorithm for the management of patients with premature ventricular complex-induced/-aggravated cardiomyopathy.

AAD, anti-arrhythmic drug; CAD, coronary artery disease; CMR, cardiac magnetic resonance; LGE, late gadolinium enhancement; LV, left ventricular; LVEF, left ventricular ejection fraction; N, No; PVC, premature ventricular complex; SHD, structural heart disease; Y, Yes.

[Open in new tab](#)[Download slide](#)

Frequent PVCs can also aggravate LV dysfunction in patients with SHD. In such cases, LV dysfunction can either be a direct consequence of PVCs as in PVC-induced cardiomyopathy, or due to the limiting effect of PVCs on optimal biventricular pacing in CRT patients. Parameters such as smaller LV end diastolic diameter and shorter intrinsic QRS duration might help to distinguish PVC-induced cardiomyopathy from PVC-aggravated cardiomyopathy.⁶¹⁴ CMR should be considered for patients suspected to have PVC-induced cardiomyopathy to exclude subtle forms of SHD.^{590,615} In a patient with frequent PVCs, the presence of LGE suggests SHD with frequent PVCs rather than PVC-induced cardiomyopathy, in which LGE is mostly absent. Given that PVCs with a RBBB morphology have been reported to show a stronger association with LGE,⁶¹⁶ those patients should be particularly considered for CMR.

The diagnosis of PVC-induced cardiomyopathy vs. PVC-aggravated cardiomyopathy can be confirmed only after LVEF improvement/normalization (reverse remodelling) following suppression of the PVCs.

Catheter ablation of the PVCs is very efficient, with reported success rates of 75a⁶¹⁷90%, and is considered first-line treatment for PVC-induced cardiomyopathy.^{535,600,609,610,612,617a⁶¹⁸620} Factors affecting acute ablation success and clinical outcome include the site of origin of PVCs (highest for outflow tract PVCs), the number of PVC morphologies, and the absence of LGE on CMR.^{535,610,614} In patients with SHD, catheter ablation of frequent monomorphic PVCs has also been shown to improve LVEF in patients with both CAD and cardiomyopathies with and without CRT.^{609,617,621a⁶¹⁹623} Similarly, the use of anti-arrhythmic medications for PVC suppression has been shown to improve LVEF. In one RCT, amiodarone resulted in a better PVC suppression and a higher LVEF improvement compared to placebo.⁶²⁴ Sodium channel blockers can also effectively suppress PVCs.⁶²⁵ In one study, flecainide reduced the PVC burden from 36% to 10% and resulted in LVEF increase from 37% to 49%.⁶²⁶ While flecainide have a more favourable adverse effect profile regarding organ toxicity, increased mortality has previously

been shown in patients with MI.⁵⁵⁶ For selected patients with suspected PVC-induced cardiomyopathy and PVC-aggravated cardiomyopathy, flecainide can still be considered, particularly if an ICD is in place (Table 9).

A rare monogenetic cause of PVC-induced cardiomyopathy, referred to as multifocal ectopic Purkinje-related premature contractions, has been reported. It is characterized by a DCM phenotype and the presence of numerous PVCs with RBBB and/or left bundle branch block (LBBB) morphologies and an increased risk of SCD.⁶²⁷ Pathogenic mutations in the *SCN5A* gene lead to a gain of function of the sodium channel responsible for hyperexcitability of the fascicular Purkinje system.^{627,628} Limited data suggest that patients with multifocal ectopic Purkinje-related premature contractions do not respond to beta-blockers but might benefit from therapy with flecainide, quinidine, or amiodarone.^{627,628,631}

Recommendation Table 27

[Open in new tab](#)

Recommendations for the management of patients with premature ventricular complex-induced or premature ventricular complex-aggravated cardiomyopathy

Recommendation	Class ^a	Level ^b
Diagnostic evaluation		
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered. ^{600,609,610}	IIa	C
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered. ^{590,615}	IIa	B
Treatment		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended. ^{535,600,609,612,617,618,620}	I	C
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, treatment with AADs ^c should be considered if catheter ablation is not desired, suspected to be high-risk, or unsuccessful. ^{624,626}	IIa	C
In patients with SHD in whom predominately monomorphic frequent PVCs are suspected to be contributing to the cardiomyopathy, AAD (amiodarone) treatment or catheter ablation should be considered. ^{617,621,622,624}	IIa	B
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered. ⁶²³	IIa	C

© ESC 2022

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; CRT, cardiac resynchronization therapy; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricular; PVC, premature ventricular complex; SHD, structural heart disease.

^aClass of recommendation.

^bLevel of evidence.

^cFlecainide only in selected patients (ICD recipients, only moderate LV dysfunction).

Recommendation	Class ^a	Level ^b
Diagnostic evaluation		
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered. ^{600,609,610}	IIa	C
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered. ^{590,615}	IIa	B
Treatment		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended. ^{535,600,609,612,617,618,620}	I	C
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, treatment with AADs ^c should be considered if catheter ablation is not desired, suspected to be high-risk, or unsuccessful. ^{624,626}	IIa	C
In patients with SHD in whom predominately monomorphic frequent PVCs are suspected to be contributing to the cardiomyopathy, AAD (amiodarone) treatment or catheter ablation should be considered. ^{617,621,622,624}	IIa	B
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered. ⁶²³	IIa	C

© ESC 2022

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; CRT, cardiac resynchronization therapy; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricular; PVC, premature ventricular complex; SHD, structural heart disease.

^aClass of recommendation.

^bLevel of evidence.

^cFlecainide only in selected patients (ICD recipients, only moderate LV dysfunction).

Recommendation Table 27

[Open in new tab](#)

Recommendations for the management of patients with premature ventricular complex-induced or premature ventricular complex-aggravated cardiomyopathy

Recommendation	Class ^a	Level ^b
Diagnostic evaluation		
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered. ^{600,609,610}	IIa	C
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered. ^{590,615}	IIa	B
Treatment		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended. ^{535,600,609,612,617,618,620}	I	C

In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, treatment with AADs ^c should be considered if catheter ablation is not desired, suspected to be high-risk, or unsuccessful. ^{624,626}	IIa	C
In patients with SHD in whom predominately monomorphic frequent PVCs are suspected to be contributing to the cardiomyopathy, AAD (amiodarone) treatment or catheter ablation should be considered. ^{617,621,622,624}	IIa	B
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered. ⁶²³	IIa	C

© ESC 2022

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; CRT, cardiac resynchronization therapy; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricular; PVC, premature ventricular complex; SHD, structural heart disease.

^aClass of recommendation.

^bLevel of evidence.

^cFlecainide only in selected patients (ICD recipients, only moderate LV dysfunction).

Recommendation	Class ^a	Level ^b
Diagnostic evaluation		
In patients with an unexplained reduced EF and a PVC burden of at least 10%, PVC-induced cardiomyopathy should be considered. ^{600,609,610}	IIa	C
In patients with suspected PVC-induced cardiomyopathy, CMR should be considered. ^{590,615}	IIa	B
Treatment		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended. ^{535,600,609,612,617,618,620}	I	C

In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, treatment with AADs ^c should be considered if catheter ablation is not desired, suspected to be high-risk, or unsuccessful. ^{624,626}	IIa	C
In patients with SHD in whom predominately monomorphic frequent PVCs are suspected to be contributing to the cardiomyopathy, AAD (amiodarone) treatment or catheter ablation should be considered. ^{617,621,622,624}	IIa	B
In non-responders to CRT with frequent, predominately monomorphic PVCs limiting optimal biventricular pacing despite pharmacological therapy, catheter ablation or AADs should be considered. ⁶²³	IIa	C

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; CRT, cardiac resynchronization therapy; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricular; PVC, premature ventricular complex; SHD, structural heart disease.

^aClass of recommendation.

^bLevel of evidence.

^cFlecainide only in selected patients (ICD recipients, only moderate LV dysfunction).

© ESC 2022

7.1.3. Cardiomyopathies

7.1.3.1. Dilated cardiomyopathy and hypokinetic non-dilated cardiomyopathy

DCM is characterized by LV dilatation and systolic dysfunction unexplained by CAD or abnormal loading conditions.⁶³² The true prevalence is difficult to estimate. Older studies reported a prevalence of 1 in 2700 individuals.⁶³³

Historic data in adults with DCM showed a 1-year mortality of 20%^{PROBTS}25% and a 50% survival at 5 years.⁶³⁴ Recent trials of DCM patients with systolic heart failure and OMT report 5-year mortality rates in the range of 21%^{PROBTS}28%.^{356,359} SCD occurs in up to 12% of patients with DCM and still accounts for 25%^{PROBTS}35% of all deaths.^{359,634,635}

In children, the annual incidence of DCM is 0.57 cases per 100 000.⁶³⁶ The prognosis in children is poor, with a 5-year incidence of heart transplantation or heart failure-related death of 40%. However, in contrast to adults, the incidence of SCD in paediatric DCM patients is much lower, with a 5-year incidence of 2.4%^{PROBTS}3% as reported in two large paediatric cardiomyopathy registries from the United States and Australia.^{637,638}

Causes of DCM can be genetic or acquired.⁶³⁹ Genetic predisposition may also interact with extrinsic factors such as in peri-partum, alcoholic, or chemotherapy-related cardiomyopathies.⁶³⁹ The phenotype, particularly of genetic aetiologies, can change over time and/or may not meet standard disease criteria at the time of disease manifestation. As a consequence, a new category of a hypokinetic non-dilated cardiomyopathy (HNDCM) has been proposed.⁶³⁹

7.1.3.1.1. Diagnostic evaluation and risk stratification

An algorithm for risk stratification and primary prevention of SCD in patients with DCM/HNDCM is presented in [FigureÂ 19](#).




Figure 19

Algorithm for risk stratification and primary prevention of sudden cardiac death in patients with dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy.

AV, atrioventricular; CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; LMNA: lamin A/C gene; LVEF, left ventricular ejection fraction; N, No; NSVT, non-sustained ventricular tachycardia; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmias; Y, Yes. ^aRisk factors: unexplained syncope, pathogenic variants in PLN, FLNC, or RBM20, LGE on CMR, inducible SMVT at PES. ^bThe 2018 ESC Guidelines for the diagnosis and management of syncope.¹

[Open in new tab](#)[Download slide](#)

Careful diagnostic work-up, including genetic testing and CMR, should be considered to identify the underlying cause for aetiology-oriented risk stratification and treatment.^{341,639}

Pathogenic mutations are identified in 253⁶³⁴55% of DCM patients,^{634,639,640} most often with autosomal dominant inheritance. Truncated mutations in the titin gene (*TTN*) are most frequently found, followed by *LMNA*, sarcomeric, and desmosomal mutations. Mutations in genes such as *LMNA*, *PLN*, *RBM20*, and *FLNC* are associated with a higher risk of VA and SCD.^{641⁶⁴⁰645} Carriers of desmosomal and *LMNA* variants experienced the highest rate of VA/SCD, which was independent of the LVEF in one series.⁶⁴⁵

The yield of genetic testing is particularly high in DCM patients with familial forms of DCM or SCD in a first-degree relative at younger age. An inherited DCM is also more likely in patients who present at young age or with signs suspicious for a specific aetiology (e.g. prolonged AV conduction for *LMNA*) (Figure 20).^{634,639,640,646} First-degree relatives of DCM/HNDCM patients should undergo clinical diagnostic evaluation, especially if an inherited cause is suspected.




Figure 20

Typical features of dilated cardiomyopathy associated with lamin A/C gene mutation with ventricular arrhythmias.

AV, atrioventricular; CMR, cardiac magnetic resonance; ECG, electrocardiogram; LGE, late gadolinium enhancement; LV, left ventricle; RBBB, right bundle branch block; VT, ventricular tachycardia.

[Open in new tab](#)[Download slide](#)

Discrimination between high- and low-risk patients for SCD remains challenging. Beyond LVEF and NYHA class,^{357,359,635,647,650} recent data suggest that both genetic and CMR findings can contribute to risk stratification. A meta-analysis of 29 studies combining 2948 patients studied the role of CMR in patients with DCM.¹²⁹ LGE was significantly associated with the arrhythmic endpoint, even when including only studies that performed multivariate analysis (HR 6.7; $P < 0.001$). Interestingly, the association between LGE and the arrhythmic endpoint remained significant among patients with mean LVEF $>35\%$. Similarly, a recent study of 1020 consecutive patients with DCM and CMR observed that LGE and LVEF were both risk markers for all-cause mortality and cardiac death, but only LGE was significantly associated with SCD risk.⁶⁵¹

SCD risk stratification has been refined in the subgroup of patients with *LMNA* mutations, which represent 5%⁶⁵² of all DCM patients. *LMNA* mutations are associated with early atrial and ventricular arrhythmias, premature conduction disease, a high risk of SCD, and progression to end-stage heart failure.^{80,642,652,653} In a multicentre registry of 269 *LMNA* mutation carriers, NSVT, LVEF $<45\%$ at first evaluation, male sex, and non-missense mutations were identified as independent risk factors for VA.⁶⁵² VA occurred only in persons with at least two of these risk factors. The risk stratification was subsequently externally validated.⁶⁵³ Another study, which involved 589 *LMNA* mutation carriers, identified AV block as an additional predictor. Recently a risk calculator has been developed (<https://lmna-risk-vta.fr/>) to predict the risk of life-threatening VA (C-index of 0.776 [95% CI: 0.711–0.842]).⁸⁰ In patients with a 5-year estimated risk $\geq 10\%$ and a manifest cardiac phenotype (NSVT, LVEF $< 50\%$, or AV conduction delay), a primary prevention ICD implantation should be considered to avoid potential over-implantation in mutation carriers without a cardiac phenotype. In *LMNA* mutation carriers with a DCM phenotype, high-intensity exercise has been associated with a high risk of SCD and impaired LVEF and is therefore not recommended.^{654,655}

A risk score for VA prediction has been recently proposed for patients with DCM or ARVC related to the p.Arg14del mutation in the *PLN* gene.⁶⁵⁶ Validation studies are needed before it can be recommended for clinical use.

Beyond genetics and CMR, additional SCD predictors have been suggested, often derived from small cohorts with no or few replication studies. An unexplained syncope requires further evaluation and PES may determine the underlying cause. The risk for arrhythmic events in DCM patients with a low LVEF and an unexplained syncope was similar to those with prior CA⁶⁵⁷ and high, independent from the outcome of PES.¹⁴⁸ However, data in DCM patients with a

mildly reduced EF suggest an incremental value of PES. Among DCM patients with an EF $\geq 40\%$ and an unexplained syncope, who underwent ICD implantation after a positive PES, 80% had appropriate ICD therapy during follow-up. No SCD or symptomatic VA occurred in non-inducible patients.¹⁴⁶

7.1.3.1.2. Primary prevention of sudden cardiac death

In patients with DCM/HNDCM, OMT according to current 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure is mandatory.³⁴² Re-evaluation of cardiac function and clinical status after 3 months³⁴³ OMT is required before primary prevention ICD implantation. LV function is more likely to improve in DCM caused by myocarditis or *TTN* mutations.

The efficacy of primary prevention ICDs in DCM patients with HFrEF has been addressed in six RCTs.⁶³⁵

Five studies were published between 2002 and 2005 (CAT, AMIOVIRT, DEFINITE, COMPANION, and SCD-HeFT). The first three were smaller studies enrolling only DCM patients, whereas SCD-HeFT and COMPANION were larger and included patients with CAD and DCM. The COMPANION study differed from the others as it compared CRT-defibrillator, CRT-pacemaker, and OMT. A meta-analysis of the five trials (1854 patients with DCM) demonstrated a 31% reduction in all-cause mortality with ICD relative to medical therapy (RR 0.69 [95% CI 0.55, 0.87], $P < 0.002$).⁶⁵⁸ This effect persisted when COMPANION was excluded.

More recently, the DANISH trial enrolled 1116 symptomatic patients (NYHA class II or III) with non-ischaemic systolic heart failure (LVEF $\leq 35\%$). Patients were randomized to ICD or no ICD after OMT.³⁵⁹ No reduction in the primary endpoint of all-cause death for patients randomized to ICD therapy (HR 0.87; 95% CI 0.68–1.12; $P = 0.28$) was observed, despite a significant reduction in SCD in the ICD group (HR 0.50; 95% CI 0.31–0.82; $P = 0.005$). Potential explanations are the low SCD rate in the trial (4.3% in the ICD group and 8.2% in the control group), the excellent medical treatment (>90% of patients on ACE inhibitors and beta-blockers, >50% on MRA), and the high prevalence (58%) of CRT. A meta-analysis of all six primary prevention trials still demonstrated a reduction, although lower, in overall mortality with ICD therapy (RR 0.76; 95% CI 0.65–0.91; $P = 0.002$).⁶³⁵ In a sensitivity analysis, the benefit of ICD therapy was maintained after the removal of any single study from the pooled analysis.

The results of the DANISH trial emphasize the need for further refining the indication for primary prevention ICDs in contemporary DCM patients. Age and comorbidity need to be considered.⁶³⁵ In the DANISH trial, ICD implantation was associated with a significantly lower rate of death in younger patients.³⁵⁹ A further analysis of the DANISH data⁶⁴⁷ showed an association between reduced all-cause mortality and ICDs in patients ≤ 70 years of age (HR, 0.70; 95% CI 0.51–0.96; $P = 0.03$), but not in patients >70 years of age.

Prospective studies evaluating the benefit of ICDs in DCM/HNDCM patients with intermediate LV dysfunction, but with risk factors that have been associated with VA and SCD (including LGE on CMR, pathogenic mutations in *PLN*, *FLNC*, *RBM20*, unexplained syncope, and inducibility of SMVT), are lacking.^{129,641,643,644,659,660,662} Given the limitation of LVEF as the only risk marker in DCM/HNDCM, this panel of experts shares the opinion that in the presence of a combination of risk markers an ICD should be considered.

7.1.3.1.3. Secondary prevention of sudden cardiac death and management of ventricular arrhythmias

Three randomized trials (AVID, CASH, and CIDS) have compared ICD therapy and medical treatment for secondary prevention in patients after aborted CA or not-tolerated VT.^{349,351,352} The three trials enrolled a total of 1963 patients, of whom only 292 (14.8%) had non-ischaemic aetiologies. A significant reduction in death from any cause in patients with ICDs was demonstrated, which was almost entirely due to a 50% reduction in arrhythmic death.³⁵² In the subgroup of patients with cardiomyopathies there was a similar but non-significant trend in the reduction of death.³⁵² RCTs have excluded patients with haemodynamically tolerated VT. In DCM patients the VT substrate is less well defined, and disease progression needs to be considered. Therefore, despite the lack of data, this panel of experts shares the opinion that an ICD should also be considered in DCM patients with haemodynamically tolerated VT.

Optimization of ICD programming (see [Section 6.2.3.1](#)) can reduce ICD shocks delivered in response to VT, but additional therapy to reduce symptomatic VA episodes is often required. In the OPTIC trial, 412 patients with ICD implantation within 21 days of VT/VF were randomized to amiodarone plus beta-blockers, sotalol alone, or beta-blocker alone. ICD shock rates after one year were 10.3%, 24.3%, and 38.5%, respectively. The higher efficacy of amiodarone plus beta-blockers compared to sotalol needs to be weighed against the higher rate of medication-related adverse events on an individual basis.³¹⁸ Although only limited data are available, sodium channel blockers may control VT in SHD and may be beneficial in ICD recipients without advanced heart failure. The majority of SMVT is due to scar-related re-entry, which can be targeted by catheter ablation. Acute ablation outcome has been reported to be similar for CAD and DCM.^{497,663} However, VT recurrence rates are usually higher in DCM patients (VT-free

survival 40.5% in DCM vs. 57% in CAD at 1 year).⁴⁹⁷ After multiple procedures in 36% of the patients, long-term freedom of VT could be achieved in 69% of cases in a retrospective, single-centre cohort of 282 patients treated between 1999 and 2014.⁶⁶⁴

Epicardial ablation is needed in 27%–30% of the procedures.^{497,664} Outcome is particularly poor, and bailout strategies (transcoronary ethanol ablation, bipolar ablation, surgical ablation) may be required in patients with pathogenic mutations (*LMNA* gene) and in those with deep intramural, antero-septal substrate locations.^{499,665,666} Considering the complexity of the ablation procedure, DCM patients should therefore be treated only in specialized centres.

Recommendation Table 28

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years). ^{641–645}	I	B
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD. ^{129,651,667}	IIa	B
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age, or with signs suspicious for an inherited aetiology. ^{641–645}	IIa	C
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation. ⁶⁵⁵	III	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III), and LVEF ≤35% after ≥3 months of OMT. ^{357,359,635,650}	IIa	A
ICD implantation should be considered in DCM/HNDCM patients with a pathogenic mutation in <i>LMNA</i> gene, if the estimated 5-year risk of life-threatening VA is ≥10% ^c and in the presence of NSVT or LVEF < 50% or AV conduction delay. ^{80,652,653}	IIa	B
ICD implantation should be considered in DCM/HNDCM patients with a LVEF <50% and ≥2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , ^d <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa	C
In DCM/HNDM patients, electrophysiological evaluation should be considered when syncope remains unexplained after non-invasive evaluation. ^{661,668}	IIa	C

Secondary prevention of SCD and treatment of VAs

ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/VF or experience haemodynamically not-tolerated SMVT. ^{349,351,352}	I	B
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{481,497,664,669}	IIa	C
The addition of oral amiodarone or replacement of beta-blockers by sotalol should be considered in patients with DCM/HNDCM and an ICD who experience recurrent, symptomatic VA despite optimal device programming and beta-blocker treatment. ³¹⁸	IIa	B
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa	C

Management of relatives of a patient or SCD victim with DCM/HNDCM

In a first-degree relative of a DCM/HNDCM patient, an ECG and an echocardiogram are recommended if: • the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or • there is family history of DCM/HNDCM, or premature unexpected SD.⁶⁴⁶	I	C
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG and an echocardiogram may be considered. ⁶⁴⁶	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrioventricular; CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; ECG, electrocardiogram; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the risk calculator <https://lmna-risk-vta.fr/>

^dSee specific recommendations for LMNA.

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years). ^{641–645}	I	B
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD. ^{129,651,667}	IIa	B
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age, or with signs suspicious for an inherited aetiology. ^{641–645}	IIa	C
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation. ⁶⁵⁵	III	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III), and LVEF ≤35% after ≥3 months of OMT. ^{357,359,635,650}	IIa	A
ICD implantation should be considered in DCM/HNDCM patients with a pathogenic mutation in <i>LMNA</i> gene, if the estimated 5-year risk of life-threatening VA is ≥10% ^c and in the presence of NSVT or LVEF < 50% or AV conduction delay. ^{80,652,653}	IIa	B
ICD implantation should be considered in DCM/HNDCM patients with a LVEF <50% and ≥2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , ^d <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa	C
In DCM/HNDCM patients, electrophysiological evaluation should be considered when syncope remains unexplained after non-invasive evaluation. ^{661,668}	IIa	C

Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/VF or experience haemodynamically not-tolerated SMVT. ^{349,351,352}	I	B
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{481,497,664,669}	IIa	C
The addition of oral amiodarone or replacement of beta-blockers by sotalol should be considered in patients with DCM/HNDCM and an ICD who experience recurrent, symptomatic VA despite optimal device programming and beta-blocker treatment. ³¹⁸	IIa	B
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa	C
Management of relatives of a patient or SCD victim with DCM/HNDCM		
In a first-degree relative of a DCM/HNDCM patient, an ECG and an echocardiogram are recommended if: • the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or • there is family history of DCM/HNDCM, or premature unexpected SD.⁶⁴⁶	I	C
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG and an echocardiogram may be considered. ⁶⁴⁶	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrioventricular; CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; ECG, electrocardiogram; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the risk calculator <https://lmna-risk-vta.fr/>

^dSee specific recommendations for LMNA.

Recommendation TableÂ 28

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years). ^{641–645}	I	B
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD. ^{129,651,667}	IIa	B
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age, or with signs suspicious for an inherited aetiology. ^{641–645}	IIa	C
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation. ⁶⁵⁵	III	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III), and LVEF ≤35% after ≥3 months of OMT. ^{357,359,635,650}	IIa	A
ICD implantation should be considered in DCM/HNDCM patients with a pathogenic mutation in <i>LMNA</i> gene, if the estimated 5-year risk of life-threatening VA is ≥10% ^c and in the presence of NSVT or LVEF < 50% or AV conduction delay. ^{80,652,653}	IIa	B
ICD implantation should be considered in DCM/HNDCM patients with a LVEF <50% and ≥2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , ^d <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa	C
In DCM/HNDCM patients, electrophysiological evaluation should be considered when syncope remains unexplained after non-invasive evaluation. ^{661,668}	IIa	C

Secondary prevention of SCD and treatment of VAs

ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/VF or experience haemodynamically not-tolerated SMVT. ^{349,351,352}	I	B
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{481,497,664,669}	IIa	C
The addition of oral amiodarone or replacement of beta-blockers by sotalol should be considered in patients with DCM/HNDCM and an ICD who experience recurrent, symptomatic VA despite optimal device programming and beta-blocker treatment. ³¹⁸	IIa	B
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa	C

Management of relatives of a patient or SCD victim with DCM/HNDCM

In a first-degree relative of a DCM/HNDCM patient, an ECG and an echocardiogram are recommended if: • the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or • there is family history of DCM/HNDCM, or premature unexpected SD.⁶⁴⁶	I	C
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG and an echocardiogram may be considered. ⁶⁴⁶	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrioventricular; CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; ECG, electrocardiogram; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the risk calculator <https://lmna-risk-vta.fr/>

^dSee specific recommendations for LMNA.

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years). ^{641–645}	I	B
CMR with LGE should be considered in DCM/HNDCM patients for assessing the aetiology and the risk of VA/SCD. ^{129,651,667}	IIa	B
Genetic testing (including at least <i>LMNA</i> , <i>PLN</i> , <i>RBM20</i> , and <i>FLNC</i> genes) should be considered for risk stratification in patients with apparently sporadic DCM/HNDCM, who present at young age, or with signs suspicious for an inherited aetiology. ^{641–645}	IIa	C
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a <i>LMNA</i> mutation. ⁶⁵⁵	III	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with DCM/HNDCM, symptomatic heart failure (NYHA class II–III), and LVEF ≤35% after ≥3 months of OMT. ^{357,359,635,650}	IIa	A
ICD implantation should be considered in DCM/HNDCM patients with a pathogenic mutation in <i>LMNA</i> gene, if the estimated 5-year risk of life-threatening VA is ≥10% ^c and in the presence of NSVT or LVEF < 50% or AV conduction delay. ^{80,652,653}	IIa	B
ICD implantation should be considered in DCM/HNDCM patients with a LVEF <50% and ≥2 risk factors (syncope, LGE on CMR, inducible SMVT at PES, pathogenic mutations in <i>LMNA</i> , ^d <i>PLN</i> , <i>FLNC</i> , and <i>RBM20</i> genes).	IIa	C
In DCM/HNDM patients, electrophysiological evaluation should be considered when syncope remains unexplained after non-invasive evaluation. ^{661,668}	IIa	C

Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/VF or experience haemodynamically not-tolerated SMVT. ^{349,351,352}	I	B
Catheter ablation in specialized centres should be considered in patients with DCM/HNDCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{481,497,664,669}	IIa	C
The addition of oral amiodarone or replacement of beta-blockers by sotalol should be considered in patients with DCM/HNDCM and an ICD who experience recurrent, symptomatic VA despite optimal device programming and beta-blocker treatment. ³¹⁸	IIa	B
ICD implantation should be considered in patients with DCM/HNDCM and haemodynamically tolerated SMVT.	IIa	C
Management of relatives of a patient or SCD victim with DCM/HNDCM		
In a first-degree relative of a DCM/HNDCM patient, an ECG and an echocardiogram are recommended if: • the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or • there is family history of DCM/HNDCM, or premature unexpected SD.⁶⁴⁶	I	C
In a first-degree relative of a patient with apparently sporadic DCM/HNDCM, an ECG and an echocardiogram may be considered. ⁶⁴⁶	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrioventricular; CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; ECG, electrocardiogram; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical therapy; PES, programmed electrical stimulation; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the risk calculator <https://lmna-risk-vta.fr/>

^dSee specific recommendations for LMNA.

7.1.3.2. Arrhythmogenic right ventricular cardiomyopathy

ARVC is an inherited disease characterized by fibrofatty myocardial replacement.⁶⁷⁰ The prevalence ranges from 1:1000 to 1:5000 individuals, with male predominance among probands.⁶⁷¹ Patients with SCD/VF at first presentation are typically younger (median age 23 [range 13^a–57]^b) compared to those who present with a SMVT (median age 36 years [range 14^a–78]).⁶⁷²

ARVC is caused by pathogenic mutations in desmosomal genes and less commonly in non-desmosomal genes. Genetic testing is indicated and identification of a mutation (in up to 73% of probands) is a major criterion for the diagnosis.^{116,671} In 4^a^b–16% compound/digenic mutations are found, which have been associated with an increased risk of VA at younger age.^{672,673} The disease penetrance in first-degree relatives is 28^a–58%,^{674,675} supporting regular clinical evaluation of relatives.

ARVC is characterized by a predominant RV involvement. The 2010 revised international diagnostic task force criteria are based on a multiparametric strategy¹¹⁶ (Figure A 21). Tissue characterization by CMR was not included. However, RV fatty infiltration and LV LGE are frequently observed (in 29^a–53% and 35.5^a–45% of probands, respectively) and may be present before patients meet major task force imaging criteria.^{676^a–678} Both wall motion

alterations and pre-/post-contrast signal abnormalities have been suggested to enhance the diagnostic accuracy of CMR for ARVC.⁶⁷⁹ The identification of biventricular and left-dominant involvement in ARVC patients^{680,681} has recently led to the proposed term arrhythmogenic cardiomyopathy.⁶⁸²




Figure 21

Typical features of arrhythmogenic right ventricular cardiomyopathy associated with ventricular arrhythmias.

CMR, cardiac magnetic resonance; ECG, electrocardiogram; LBBB, left bundle branch block; RV, right ventricle; VT, ventricular tachycardia.

[Open in new tab](#)[Download slide](#)

Restriction from high-intensity exercise is regarded as a preventive tool in clinically affected ARVC patients to reduce the risk of VAs and disease progression.^{683a,685} Whether sport restriction is beneficial in all mutation carriers without disease expression has not been evaluated in prospective cohorts, but avoidance of competitive, high-intensity exercise seems to be reasonable.^{686,687} SCD and VA occur disproportionately during exercise in affected patients, and high-dose isoproterenol infusion may provoke PVT in >90% of ARVC patients, supporting the role of sympathetic stimulation for arrhythmogeneity.^{152,688,689} Whether beta-blockers can prevent spontaneous VA is, however, unclear. Limited data suggest a potential beneficial role of atenolol.⁶⁹⁰

7.1.3.2.1. Risk stratification

In ARVC patients not implanted with ICDs, CA occurs in 4.6a^{683a,685}6.1%, while 23% of patients experience a non-fatal SMVT during an average follow-up of 8a^{683a,685}11 years.^{691a,694} Whether a SMVT is acutely life-threatening depends on the VT cycle length, cardiac function, and the circumstances under which VT occurs. Among definite/probable ARVC patients considered at high risk for VA, 23a^{683a,685}48% will experience appropriate ICD intervention during a mean follow-up of 4.7 years. In 16a^{683a,685}19% of cases, ICD intervention is triggered by fast VT a^{683a,685}250a^{683a,685}300 b.p.m. or VF, which is considered as surrogate for a life-threatening event.^{695a,698} In a large cohort of 864 ARVC patients (38.8% with a prior VA), 43% had VT/VF during a median follow-up of 5.75 years, but only 10.8% a potentially life-threatening event. Thus, in 3 out of 4 ARVC patients, ICD therapy is appropriate but may not be considered acutely life-saving. This is relevant because risk prediction algorithms and models are based on combined arrhythmic endpoints, which are often equated with ICD indications to prevent SCD. This is particularly important for the young ARVC patients with a lifetime risk of ICD-related complications.^{695,699}

Randomized studies of ICD therapy for secondary prevention are lacking, but the high rates of appropriate ICD therapy triggered by fast VT/VF episodes in patients implanted after CA or haemodynamically poorly tolerated VT strongly suggest a survival benefit from ICD.⁷⁰⁰ In patients who present with a haemodynamically tolerated VT, the survival benefit from ICD is less clear.^{691,700a,702} Identification of ARVC patients at risk for SCD is difficult, and the evidence supporting risk factors for life-threatening VAs is limited. Arrhythmic syncope was a predictor for

subsequent events in most series of patients with definitive ARVC (pooled HR 3.67; CI 2.75â4.9)^{81,696,701,703} and, in these patients, an ICD should be considered. RV and LV dysfunction have been associated with a higher arrhythmic risk.^{675,704} Cut-off values are difficult to determine but, in patients with *severe RV dysfunction* (RV fractional area change $\leq$ 17% or RV ejection fraction $\leq$ 35%), an ICD should be considered. Likewise, ARVC patients with significant LV involvement (LVEF $\leq$ 35%) should be treated according to the current DCM recommendation regarding ICD implantation. There are conflicting data on the *independent* predictive value of NSVT for subsequent sustained arrhythmic events.^{695,696,701} Likewise, the role of PES in risk stratification, particularly in asymptomatic patients, is not well defined.^{695,696,705} In one series including mainly symptomatic patients, both NSVT and a positive PES independently predicted arrhythmic events, while others report a low diagnostic accuracy.⁶⁹⁶ Risk prediction based on a single parameter does not consider the potential combined effect and interactions between factors. Therefore, although supporting data are lacking, this panel of experts supports that an ICD should be considered⁷⁰⁶ in symptomatic ARVC patients with moderate RV (<40%) and/or LV dysfunction (<45%) and who have *either* NSVT or have SMVT inducible at PES. An internally validated risk model has recently been developed from 528 patients with definite ARVC and no prior VA, including age, sex, arrhythmic syncope, NSVT, PVC burden, leads with T-wave inversion and RV ejection fraction, to predict *any sustained* VA (c-index 0.77).⁸¹ In addition, a prediction model for specifically life-threatening VA has been promoted based on 864 ARVC patients, with predictors including male sex, age, 24âh PVC count, and leads with T-wave inversion (c-index 0.74).⁷⁰² However, validation studies are needed before these risk models can be recommended for clinical use.

7.1.3.2.2. Treatment

Up to 97.4% of all VA episodes in ARVC ICD recipients are SMVT. The very high termination rate by ATP (92% of all episodes) independent from VT cycle length strongly supports devices with the capability for ATP.⁶⁹⁸ S-ICDs reduce lead-related complications and have been proven to effectively shock terminate VA in small cohorts of ARVC patients during short-term follow-up.⁷⁰⁷

Additional treatment to suppress VA is often required. Although not confirmed by clinical data, beta-blockers are recommended as first-line therapy in symptomatic patients.

Data on AADs to prevent VT recurrence are limited to small observational studies and registries. In general, AAD therapy has limited efficacy. Although sotalol was effective to prevent inducibility of VT,⁷⁰⁸ it did not suppress clinically relevant arrhythmias.^{690,708} Treatment with amiodarone or class 1 drugs was associated with a trend to lower VT recurrence as compared with sotalol.⁷⁰⁹ The addition of flecainide to beta-blockers/sotalol was beneficial in a small cohort.⁷¹⁰ Catheter ablation provides an alternative. Endocardial and adjuvant epicardial substrate ablation, if needed, has been associated with promising VT-free survival. Potential procedural risks, drug side effects, and the patient's preference need to be taken into consideration.⁴⁸²

Recommendation TableÂ 29

[Open in new tab](#)

Recommendations for diagnostic, risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in arrhythmogenic right ventricular cardiomyopathy

Recommendations	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
In patients with suspected ARVC, CMR is recommended. ^{676–678}	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended. ^{672,673}	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC. ^{683–685}	I	B
Avoidance of high-intensity ^c exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype. ^{683,687}	IIb	C
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope. ^{696,701,711–713}	IIa	B
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction. ^{675,691}	IIa	C
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and	IIa	C

either NSVT or inducibility of SMVT at PES. ^{695,696,701,703,705}		
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification. ^{695,705}	IIb	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF. ⁷⁰⁰	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In patients with ARVC and recurrent, symptomatic SMVT or ICD shocks for SMVT despite beta-blockers, catheter ablation in specialized centres should be considered. ^{482,709,714}	IIa	C
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered. ⁶⁹⁸	IIa	B
ICD implantation should be considered in ARVC patients with a haemodynamically tolerated SMVT. ⁶⁹²	IIa	C
In patients with ARVC and recurrent, symptomatic VT despite beta-blockers, AAD treatment should be considered. ^{709,710}	IIa	C
Management of relatives of a patient with ARVC		
In a first-degree relative of a patient with ARVC, ECG and echocardiogram are recommended. ⁶⁷⁵	I	C

© ESC 2022

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained ventricular tachycardia; PES, programmed electrical stimulation; RV, right ventricle; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^dPresyncope or palpitations suggestive of VA.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
In patients with suspected ARVC, CMR is recommended. ^{676–678}	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended. ^{672,673}	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC. ^{683–685}	I	B
Avoidance of high-intensity ^c exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype. ^{683,687}	IIb	C
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope. ^{696,701,711–713}	IIa	B
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction. ^{675,691}	IIa	C
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and	IIa	C

either NSVT or inducibility of SMVT at PES. ^{695,696,701,703,705}		
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification. ^{695,705}	IIb	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF. ⁷⁰⁰	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In patients with ARVC and recurrent, symptomatic SMVT or ICD shocks for SMVT despite beta-blockers, catheter ablation in specialized centres should be considered. ^{482,709,714}	IIa	C
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered. ⁶⁹⁸	IIa	B
ICD implantation should be considered in ARVC patients with a haemodynamically tolerated SMVT. ⁶⁹²	IIa	C
In patients with ARVC and recurrent, symptomatic VT despite beta-blockers, AAD treatment should be considered. ^{709,710}	IIa	C
Management of relatives of a patient with ARVC		
In a first-degree relative of a patient with ARVC, ECG and echocardiogram are recommended. ⁶⁷⁵	I	C

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained ventricular tachycardia; PES, programmed electrical stimulation; RV, right ventricle; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^dPresyncope or palpitations suggestive of VA.

© ESC 2022

Recommendation Table 29

[Open in new tab](#)

Recommendations for diagnostic, risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in arrhythmogenic right ventricular cardiomyopathy

Recommendations	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
In patients with suspected ARVC, CMR is recommended. ^{676–678}	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended. ^{672,673}	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC. ^{683–685}	I	B
Avoidance of high-intensity ^c exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype. ^{683,687}	IIb	C
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope. ^{696,701,711–713}	IIa	B
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction. ^{675,691}	IIa	C
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and	IIa	C

either NSVT or inducibility of SMVT at PES. ^{695,696,701,703,705}		
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification. ^{695,705}	IIb	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF. ⁷⁰⁰	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In patients with ARVC and recurrent, symptomatic SMVT or ICD shocks for SMVT despite beta-blockers, catheter ablation in specialized centres should be considered. ^{482,709,714}	IIa	C
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered. ⁶⁹⁸	IIa	B
ICD implantation should be considered in ARVC patients with a haemodynamically tolerated SMVT. ⁶⁹²	IIa	C
In patients with ARVC and recurrent, symptomatic VT despite beta-blockers, AAD treatment should be considered. ^{709,710}	IIa	C
Management of relatives of a patient with ARVC		
In a first-degree relative of a patient with ARVC, ECG and echocardiogram are recommended. ⁶⁷⁵	I	C

© ESC 2022

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained ventricular tachycardia; PES, programmed electrical stimulation; RV, right ventricle; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴

^dPresyncope or palpitations suggestive of VA.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
In patients with suspected ARVC, CMR is recommended. ^{676–678}	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended. ^{672,673}	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC. ^{683–685}	I	B
Avoidance of high-intensity ^c exercise may be considered in carriers of ARVC-related pathogenic mutations and no phenotype. ^{683,687}	IIb	C
Beta-blocker therapy may be considered in all patients with a definite diagnosis of ARVC.	IIb	C
Risk stratification and primary prevention of SCD		
ICD implantation should be considered in patients with definite ARVC and an arrhythmic syncope. ^{696,701,711–713}	IIa	B
ICD implantation should be considered in patients with definite ARVC and severe RV or LV systolic dysfunction. ^{675,691}	IIa	C
ICD implantation should be considered in symptomatic ^d patients with definite ARVC, moderate right or left ventricular dysfunction, and	IIa	C

either NSVT or inducibility of SMVT at PES. ^{695,696,701,703,705}		
In patients with ARVC and symptoms highly suspicious for VA, PES may be considered for risk stratification. ^{695,705}	IIb	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF. ⁷⁰⁰	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In patients with ARVC and recurrent, symptomatic SMVT or ICD shocks for SMVT despite beta-blockers, catheter ablation in specialized centres should be considered. ^{482,709,714}	IIa	C
In ARVC patients with indication for ICDs, a device with the capability of ATP programming for SMVT up to high rates should be considered. ⁶⁹⁸	IIa	B
ICD implantation should be considered in ARVC patients with a haemodynamically tolerated SMVT. ⁶⁹²	IIa	C
In patients with ARVC and recurrent, symptomatic VT despite beta-blockers, AAD treatment should be considered. ^{709,710}	IIa	C
Management of relatives of a patient with ARVC		
In a first-degree relative of a patient with ARVC, ECG and echocardiogram are recommended. ⁶⁷⁵	I	C

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained ventricular tachycardia; PES, programmed electrical stimulation; RV, right ventricle; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.
^cThe 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease.⁴
^dPresyncope or palpitations suggestive of VA.

© ESC 2022

7.1.3.3. Hypertrophic cardiomyopathy

HCM is characterized by increased LV wall thickness not explained by abnormal loading conditions (such as hypertension or valvular disease).^{128,715} Because the natural history and management differs according to the underlying aetiology of LVH, diagnostic work-up is of paramount importance (see [Section 7.1.3.5](#)) and includes CMR and genetic testing.^{716,725} HCM is usually caused by a mutation with autosomal dominant inheritance, supporting cardiac screening in first-degree relatives, in parallel with genetic testing in the index patient. A sarcomeric gene mutation is identified in 30%–60%, most frequently in patients who are younger at diagnosis or with a family history of HCM.^{646,722}

The estimated prevalence of HCM in adults is 1 in 500.⁷²⁶ In children, the prevalence is much lower.


Annual mortality related to HCM is 1%–2% in most studies with contemporary management strategies,^{128,715} and may be as low as 0.5%.⁷²⁷ The annual rate of SCD or appropriate ICD therapy is about 0.8% but is largely dependent on age and risk profile.^{85,728,730} HCM patients are at risk for heart failure, AF, and stroke.^{128,715} Most HCM-related deaths at age <60 years occur suddenly, while older patients die more often of stroke or heart failure.⁷³¹ SCD is often related to VA, which can be a consequence of ischaemia, left ventricle outflow tract (LVOT) obstruction, or supraventricular arrhythmias. SCD may also be triggered by exercise, and participation in competitive sport has

been discouraged.⁷³² However, recent data suggest that even vigorous exercise in HCM patients without risk markers may not be associated with VA.^{733,734}

7.1.3.3.1. Risk stratification and primary prevention of sudden cardiac death

In primary prevention, the challenge remains to identify the relatively small group of patients with the highest risk of SCD. NSVT is identified on continuous (24/48 h) ambulatory ECG monitoring in 20%⁷³⁵ of patients.^{128,715} Its prevalence increases with age and correlates with LV wall thickness and LGE on CMR.⁷¹⁶⁷³⁶^{721,735} The prognostic value of NSVT for SCD is more important in young patients (<30 years).⁷³⁵ The relation between the duration, frequency, or rate of NSVT and HCM prognosis remains unclear.⁷³⁵ Documented NSVT during exercise test is very rare, but was associated with a higher risk of SCD.⁷³² Additional risk factors have been identified and a 5-year SCD risk stratification score based on seven factors (age, LV wall thickness, LA size, LVOT gradient, NSVT, unexplained syncope, and family history of SCD) has been developed⁸⁵ (HCM Risk-SCD: <https://doc2do.com/hcm/webHCM.html>) and externally validated.^{85,728,729} The calculator is not intended for use in elite athletes, or in individuals with metabolic or infiltrative diseases, or after myectomy or alcohol septal ablation.⁸⁵

Additional factors not captured by the Risk-SCD model should also be considered in patients with intermediate or low calculated risk, including LV systolic dysfunction, apical aneurysm, extensive LGE on CMR, and presence of single or multiple sarcomeric mutations (*FigureÂ 22*).^{716,717,722,736}⁷³⁹ Extensive LGE on CMR defined as a $\geq 15\%$ of LV mass has been suggested as good predictor of SCD in adults. However, thresholds may be difficult to use because LGE quantification is dependent on CMR acquisition, type, and amount of contrast.



FigureÂ 22

Typical features of hypertrophic cardiomyopathy associated with sustained monomorphic ventricular tachycardia.

CMR, cardiac magnetic resonance; ECG, electrocardiogram; LGE, late gadolinium enhancement; LV, left ventricular; RBBB, right bundle branch block; VT, ventricular tachycardia.

[Open in new tab](#)[Download slide](#)

Periodic reassessment is mandatory as part of the longitudinal evaluation of patients. VA induced by PES is considered non-specific, although conflicting results have been reported.^{740,741}

In children few data on primary prevention were available until the recent development of scores and related risk calculators.^{83,84} The HCM Risk-Kids score⁸⁴ has been developed and externally validated⁷⁴² specifically in children with HCM (12-16 years of age) and includes unexplained syncope, maximal LV wall thickness, large left atrial diameter, low LVOT gradient, and NSVT (<https://hcmriskkids.org>). In contrast to adults, including age and family history of SCD did not improve the performance of the paediatric model. Patients with prior VF or sustained VT, known inborn errors of metabolism, and syndromic causes of HCM were excluded. The predictive value of LGE in children is less defined, although preliminary data were reported.⁷⁴³

7.1.3.3.2. Treatment to prevent ventricular arrhythmia recurrence

Patients surviving a CA due to VT/VF or haemodynamically not-tolerated VT remain at high risk for future life-threatening VA, and will benefit from ICD therapy.^{128,744,746}

There are no RCTs in HCM or cohort studies supporting a significant role of drugs to prevent SCD.^{128,715,747} Amiodarone may reduce VA but with conflicting results regarding SCD prevention.^{747,748} Disopyramide and beta-blockers are efficient to control symptoms and LVOT obstruction, but there is no evidence that they reduce the risk of SCD.^{128,715} Similarly, surgical myectomy or alcohol ablation are not recommended with the aim to reduce risk of SCD in patients with LVOT obstruction.^{128,715} Following ICD implantation for primary or secondary prevention, the most common documented VA subtype is SMVT, and ATP is successful in 69%⁷⁴⁹ 76.5% of all episodes.^{749,751} Although data on drug efficacy are lacking, AADs (beta-blockers, amiodarone, sotalol, sodium channel blockers) should be considered in HCM patients with symptomatic VA. Catheter ablation may also be considered in highly selected HCM patients with SMVT, in whom AADs are ineffective, contraindicated or not tolerated, as outcome after ablation is inferior compared to other non-ischaemic aetiologies.^{752,754}

Recommendation Table 30

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in hypertrophic cardiomyopathy

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
CMR with LGE is recommended in HCM patients for diagnostic work-up. ^{716–718}	I	B
Genetic counselling and testing are recommended in HCM patients. ^{721–725}	I	B
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers. ⁷³³	IIb	C
Risk stratification and primary prevention of SCD		
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals, or when there is a change in clinical status.	I	C
ICD implantation should be considered in patients aged 16 years or more with an estimated 5-year risk of SD $\geq 6\%$. ^{c,85,728,729}	IIa	B
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to $< 6\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) abnormal blood pressure response during exercise test ^d ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation. ^{716,717,722,736–739}	IIa	B
In children less than 16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^e), ICD implantation should be considered. ^{84,742}	IIa	B
ICD implantation may be considered in HCM patients aged 16 years or more with an estimated 5-year risk of SCD of ≥ 4 to $< 6\%$. ^{c,85,728,729}	IIb	B
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD ($< 4\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) LV apical aneurysm. ^{716,717,722,736–739}	IIb	B
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF. ^{744–746}	I	B
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa	C
In patients with HCM and recurrent, symptomatic VA, or recurrent ICD therapy, AAD treatment should be considered.	IIa	C

Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AAD are ineffective, contraindicated, or not tolerated.^{753,754}

IIb

C

Management of relatives of a patient with HCM

In a first-degree relative of a patient with HCM, ECG and echocardiogram are recommended.

I

C

© ESC 2022

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; ECG, electrocardiogram; HCM, hypertrophic cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LV, left ventricle; LVEF, left ventricular ejection fraction; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the HCM Risk-SCD: <https://doc2do.com/hcm/webHCM.html>

^dDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^eBased on the HCM kid risk score: <https://hcmriskkids.org>

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
CMR with LGE is recommended in HCM patients for diagnostic work-up. ^{716–718}	I	B
Genetic counselling and testing are recommended in HCM patients. ^{721–725}	I	B
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers. ⁷³³	IIb	C
Risk stratification and primary prevention of SCD		
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals, or when there is a change in clinical status.	I	C
ICD implantation should be considered in patients aged 16 years or more with an estimated 5-year risk of SD $\geq 6\%$. ^{c,85,728,729}	IIa	B
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to $< 6\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) abnormal blood pressure response during exercise test ^d ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation. ^{716,717,722,736–739}	IIa	B
In children less than 16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^e), ICD implantation should be considered. ^{84,742}	IIa	B
ICD implantation may be considered in HCM patients aged 16 years or more with an estimated 5-year risk of SCD of ≥ 4 to $< 6\%$. ^{c,85,728,729}	IIb	B
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD ($< 4\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) LV apical aneurysm. ^{716,717,722,736–739}	IIb	B
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF. ^{744–746}	I	B
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa	C
In patients with HCM and recurrent, symptomatic VA, or recurrent ICD therapy, AAD treatment should be considered.	IIa	C

Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AAD are ineffective, contraindicated, or not tolerated. ^{753,754}	IIb	C
Management of relatives of a patient with HCM		
In a first-degree relative of a patient with HCM, ECG and echocardiogram are recommended.	I	C

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; ECG, electrocardiogram; HCM, hypertrophic cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LV, left ventricle; LVEF, left ventricular ejection fraction; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the HCM Risk-SCD: <https://doc2do.com/hcm/webHCM.html>

^dDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^eBased on the HCM kid risk score: <https://hcmriskkids.org>

© ESC 2022

Recommendation TableÂ 30

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in hypertrophic cardiomyopathy

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
CMR with LGE is recommended in HCM patients for diagnostic work-up. ^{716–718}	I	B
Genetic counselling and testing are recommended in HCM patients. ^{721–725}	I	B
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers. ⁷³³	IIb	C
Risk stratification and primary prevention of SCD		
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals, or when there is a change in clinical status.	I	C
ICD implantation should be considered in patients aged 16 years or more with an estimated 5-year risk of SD $\geq 6\%$. ^{c,85,728,729}	IIa	B
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to $< 6\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) abnormal blood pressure response during exercise test ^d ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation. ^{716,717,722,736–739}	IIa	B
In children less than 16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^e), ICD implantation should be considered. ^{84,742}	IIa	B
ICD implantation may be considered in HCM patients aged 16 years or more with an estimated 5-year risk of SCD of ≥ 4 to $< 6\%$. ^{c,85,728,729}	IIb	B
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD ($< 4\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) LV apical aneurysm. ^{716,717,722,736–739}	IIb	B
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF. ^{744–746}	I	B
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa	C
In patients with HCM and recurrent, symptomatic VA, or recurrent ICD therapy, AAD treatment should be considered.	IIa	C

Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AAD are ineffective, contraindicated, or not tolerated.^{753,754}

IIb

C

Management of relatives of a patient with HCM

In a first-degree relative of a patient with HCM, ECG and echocardiogram are recommended.

I

C

© ESC 2022

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; ECG, electrocardiogram; HCM, hypertrophic cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LV, left ventricle; LVEF, left ventricular ejection fraction; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cBased on the HCM Risk-SCD: <https://doc2do.com/hcm/webHCM.html>

^dDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.

^eBased on the HCM kid risk score: <https://hcmriskkids.org>

Recommendation	Class ^a	Level ^b
Diagnostic evaluation and general recommendations		
CMR with LGE is recommended in HCM patients for diagnostic work-up. ^{716–718}	I	B
Genetic counselling and testing are recommended in HCM patients. ^{721–725}	I	B
Participation in high-intensity exercise may be considered for asymptomatic adult HCM patients without risk markers. ⁷³³	IIb	C
Risk stratification and primary prevention of SCD		
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals, or when there is a change in clinical status.	I	C
ICD implantation should be considered in patients aged 16 years or more with an estimated 5-year risk of SD $\geq 6\%$. ^{c,85,728,729}	IIa	B
ICD implantation should be considered in HCM patients aged 16 years or more with an intermediate 5-year risk of SCD (≥ 4 to $< 6\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) abnormal blood pressure response during exercise test ^d ; or (d) LV apical aneurysm; or (e) presence of sarcomeric pathogenic mutation. ^{716,717,722,736–739}	IIa	B
In children less than 16 years of age with HCM and an estimated 5-year risk of SD $\geq 6\%$ (based on HCM Risk-Kids score ^e), ICD implantation should be considered. ^{84,742}	IIa	B
ICD implantation may be considered in HCM patients aged 16 years or more with an estimated 5-year risk of SCD of ≥ 4 to $< 6\%$. ^{c,85,728,729}	IIb	B
ICD implantation may be considered in HCM patients aged 16 years or more with a low estimated 5-year risk of SCD ($< 4\%$) ^c and with (a) significant LGE at CMR (usually $\geq 15\%$ of LV mass); or (b) LVEF $< 50\%$; or (c) LV apical aneurysm. ^{716,717,722,736–739}	IIb	B
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF. ^{744–746}	I	B
In patients with HCM presenting with haemodynamically tolerated SMVT, ICD implantation should be considered.	IIa	C
In patients with HCM and recurrent, symptomatic VA, or recurrent ICD therapy, AAD treatment should be considered.	IIa	C

Catheter ablation in specialized centres may be considered in selected patients with HCM and recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AAD are ineffective, contraindicated, or not tolerated. ^{753,754}	IIb	C
Management of relatives of a patient with HCM		
In a first-degree relative of a patient with HCM, ECG and echocardiogram are recommended.	I	C

AAD, anti-arrhythmic drug; CMR, cardiac magnetic resonance; ECG, electrocardiogram; HCM, hypertrophic cardiomyopathy; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LV, left ventricle; LVEF, left ventricular ejection fraction; SCD, sudden cardiac death; SD, sudden death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.
^cBased on the HCM Risk-SCD: <https://doc2do.com/hcm/webHCM.html>
^dDefined as a failure to increase systolic pressure by at least 20 mmHg from rest to peak exercise, or a fall of >20 mmHg from peak pressure.
^eBased on the HCM kid risk score: <https://hcmriskkids.org>

© ESC 2022

7.1.3.4. Left ventricular non-compaction

LVNC comprises a heterogeneous group of diseases. The diagnosis is challenging, and various diagnostic criteria have been proposed. The yield of genetic tests in index patients is low.⁷⁵⁵

The morphological phenotype of non-compaction based on imaging parameters may also appear in a healthy population.⁷⁵⁶

A meta-analysis including 2501 LVNC patients revealed a risk of cardiovascular mortality similar to that of DCM patients, with no relation to the extent of trabeculation.⁷⁵⁷ CMR-based detection of focal fibrosis using LGE in LVNC with preserved ejection was associated with serious cardiac events (aborted death, ICD therapy, heart transplantation [HTX]/LVAD) in another meta-analysis including 574 patients (OR 6.1; CI 2.1–17.5; $P < 0.001$).⁷⁵⁸ A combination of CMR criteria with systematic genotyping may overcome current uncertainties regarding risk stratification.⁷⁵⁹

Recommendation Table 31

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in left ventricular non-compaction

Recommendations	Class ^a	Level ^b
In patients with a LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/HNDCM recommendations.	IIa	C

CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LVNC, left ventricular non-compaction; SCD, sudden cardiac death.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
In patients with a LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/HNDCM recommendations.	IIa	C

CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LVNC, left ventricular non-compaction; SCD, sudden cardiac death.

^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 31

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in left ventricular non-compaction

Recommendations	Class ^a	Level ^b
In patients with a LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/HNDCM recommendations.	IIa	C

CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LVNC, left ventricular non-compaction; SCD, sudden cardiac death.

^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
In patients with a LVNC cardiomyopathy phenotype based on CMR or echocardiography, implantation of an ICD for primary prevention of SCD should be considered to follow DCM/HNDCM recommendations.	IIa	C

CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LVNC, left ventricular non-compaction; SCD, sudden cardiac death.

^aClass of recommendation.
^bLevel of evidence.

7.1.3.5. Restrictive cardiomyopathy

The phenotype of restrictive cardiomyopathy (RCM) is rare and can be the consequence of different aetiologies, including infiltrative disorders (e.g. amyloidosis) and storage disease (e.g. Andersen⁷⁶⁰Fabry disease), the identification of which is crucial for guiding therapy. Heart failure is one of the leading symptoms in primary and secondary RCM.⁷⁶⁰

In Fabry disease the majority of the reported cardiovascular deaths have been categorized as SCD in a recent systematic review of 13 studies including 4185 patients.⁷⁶¹ Higher age, male gender, LVH, LGE, and NSVT have been identified as potential risk factors associated with SCD events. However, SCD rates were reported in only 11 of the studies, including 623 patients. The retrospective, observational nature of most of these small, single-centre studies and the low absolute number of cardiovascular deaths (36/623) and SCD (30/623) do not currently allow guidance for primary prevention ICD implantation.⁷⁶¹

Amyloidosis can be caused by different misfolded precursor proteins leading to deposits in tissue and organs. Cardiac amyloidosis is mainly related to light-chain amyloid or transthyretin amyloid. The latter can be subdivided into wild-type, also called senile amyloid, more often complicated by AV conduction delay and atrial arrhythmias, and amyloid due to pathogenic mutations in the *TTR* gene. Disease manifestation depends on the mutation. Despite advances in the treatment of amyloid light-chain amyloidosis, outcome is still poor in patients with manifest cardiac involvement.⁷⁶²

Causes of death are progressive heart failure, autonomic dysfunction, and electromechanical dissociation.^{762,763} The benefit of primary prevention ICD implantation in patients with cardiac amyloidosis is uncertain. Currently, an ICD should be considered in patients with haemodynamically not-tolerated VT after careful discussion of the competing risks of non-arrhythmic death and non-cardiac death.

Recommendation Table 32

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in patients with cardiac amyloidosis

Recommendations	Class ^a	Level ^b
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa	C

ICD, implantable cardioverter defibrillator; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa	C

ICD, implantable cardioverter defibrillator; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 32

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in patients with cardiac amyloidosis

Recommendations	Class ^a	Level ^b
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa	C

ICD, implantable cardioverter defibrillator; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
An ICD should be considered in patients with light-chain amyloidosis or transthyretin-associated cardiac amyloidosis and haemodynamically not-tolerated VT.	IIa	C

ICD, implantable cardioverter defibrillator; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

7.1.3.6. Neuromuscular disorders

An algorithm for risk stratification, SCD prevention, and treatment of VAs in myotonic dystrophy is presented in [Figure 23](#).




Figure 23

Algorithm for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in myotonic dystrophy.

AV, atrioventricular; BBR-VT, bundle branch re-entrant ventricular tachycardia; CMR, cardiac magnetic resonance; EP, invasive electrophysiological evaluation; HV, His-to-ventricle interval; ICD, implantable cardioverter defibrillation; LGE, late gadolinium enhancement; N, No; VT, ventricular tachycardia; Y, Yes. ^aSyncope or palpitations highly suspicious of arrhythmic origin. ^bSpontaneous AV block: third or advanced second-degree AV block. ^cFactors favouring ICD implantation: age, [5,6,11](#) CTG expansion, [6,13,16](#) sudden death (SD) or family history of SD, [5](#) ECG conduction abnormalities, [16](#) PR prolongation, [13](#) left bundle branch block, [5](#) atrial arrhythmias, [6,16](#) non-sustained VT, [5](#) LV dysfunction, [17](#) significant LGE in CMR, [14,15,18](#) ^dFurther management according to outcome of EP evaluation.

[Open in new tab](#)[Download slide](#)

Arrhythmias are common and often the first manifestation of neuromuscular disorders.⁷⁶⁴ Myotonic dystrophy is the most common muscular dystrophy in the adult population (prevalence 1 in 8000). Myotonic dystrophy is the consequence of trinucleotide repeat expansion in the end of *DMPK* gene, which results in mis-splicing of *SCN5A* and cardiac conduction system delays and arrhythmia. Duchenne dystrophy also has a high incidence (1 in 3500 male births). As most patients die before the age of 20, it is rarely seen in adulthood. Other neuromuscular disorders, such as Becker (1 in 100 000 male births) and facioscapulohumeral (1 in 100 000) dystrophies, are less common. Most of these disorders are associated with conduction and rhythm disturbances, some life-threatening, and with mechanisms amenable to specific anti-arrhythmic therapy.¹⁷ Anti-arrhythmic therapy should be considered, as muscular function and life expectancy are often only moderately limited.⁷⁶⁵ It is generally recommended to treat patients with neuromuscular diseases who have survived a CA or have VAs or ventricular dysfunction in the same way as patients without extracardiac manifestations. However, the benefit of ICD implantation should be balanced with the overall prognosis in some subtypes, such as Duchenne dystrophy.

Symptoms may be the result of bradycardia or tachyarrhythmias and invasive electrophysiological evaluation is warranted unless a cause, such as AV block, has been clinically documented.¹⁵³ Sudden increase of PR interval and QRS duration have been associated with AV block and SCD in myotonic dystrophy despite the absence of symptoms, and electrophysiological evaluation should be considered.^{766,767} An HV interval ≥70 ms should prompt pacemaker implantation regardless of symptoms. Induction of BBR-VT in a symptomatic patient strongly supports BBR-VT ([ESC CardioMed chapter 42.5](#))⁷⁶⁸ as an underlying cause of symptoms, and ablation of the right

bundle branch is recommended.¹⁵³ Following bundle branch ablation, the risk of AV block at follow-up is considered particularly high due to progressive His-Purkinje disease, and a pacemaker should be implanted. Prolonged PR or QRS intervals,^{13,16,80} concurrent atrial arrhythmias,^{6,16,80} and LGE at CMR^{14,15,18} in patients older than 40 years have all been associated with AV block and SCD at follow-up in myotonic dystrophy. Accordingly, pacemaker implantation may be considered even in the absence of symptoms. Although data are lacking, implantation of an ICD rather than a pacemaker may be preferred in myotonic dystrophy patients, with additional risk factors for VA and SCD (Figure 23). Permanent pacemaker implantation may also be considered in patients with Kearns-Sayre syndrome, Emery-Dreifuss, or limb-girdle muscular dystrophy with any degree of AV block because of the substantial risk of rapid progression to total block. In patients with limb-girdle type 1B or Emery-Dreifuss muscular dystrophies who have an indication for pacing or significant LGE on CMR, ICD implantation should be considered.⁷⁶⁹

Implantation of an ICD rather than a pacemaker may be also considered in patients with Duchenne/Becker, when there is significant LGE on CMR.^{770,771} However, the general prognosis of these diseases should be taken into consideration. In patients who have survived a CA due to a SMVT, implantation of an ICD is recommended when other than BBR-VTs (e.g. scar-related VTs) are inducible or if VT is not inducible. Myotonic dystrophy patients with a syncope likely due to a VA, even if not inducible, and those with palpitations and induction of a non-BBR-VT are considered at risk for arrhythmic SCD, and ICDs should be considered.

Due to the progressive nature, annual follow-up with an ECG is recommended, even in the concealed phase of the disease when patients are asymptomatic and the ECG is normal.^{6,13} Due to slow progression and the fact that significant changes are often reflected on the surface ECG, serial electrophysiological evaluation to assess AV conduction and arrhythmia induction is not recommended in patients without arrhythmia suspicion or progression of ECG conduction disorders.^{767,772}

Recommendation Table 33

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in neuromuscular diseases

Recommendations	Class ^a	Level ^b
General recommendations		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease. ^{6,13}	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders. ^{17,765}	I	C
Risk stratification, primary and secondary prevention of SCD		
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA. ¹⁵³	I	C

ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT. ⁷⁶⁶	I	C
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration. ^{766,767}	IIa	B
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias ^c or who are older than 40 years and have significant LGE on CMR. ^{c,5,14,16,766}	IIa	B
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In patients with limb-girdle type 1B or Emery-Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered. ⁷⁶⁹	IIa	C
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR. ^{770,771}	IIb	C
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ^d for VAs and SCD.	IIb	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders. ⁷⁷²	III	C
Management of VA		
In symptomatic patients with BBR-VT, catheter ablation is recommended. ^{e,153,474,475,477}	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended. ¹⁵³	I	C

© ESC 2022

AV, atrioventricular; BBR-VT, bundle branch re-entrant ventricular tachycardia; CA, cardiac arrest; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LBBB, left bundle branch block; LGE, late gadolinium enhancement; LV, left ventricular; SCD, sudden cardiac death; SD, sudden death; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cLevel of evidence C.

^dFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6–9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^eIn other cardiac conditions (e.g. aortic valve replacement) with BBR-VT, catheter ablation is also recommended.

Recommendations	Class ^a	Level ^b
General recommendations		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease. ^{6,13}	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders. ^{17,765}	I	C
Risk stratification, primary and secondary prevention of SCD		
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA. ¹⁵³	I	C

ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT. ⁷⁶⁶	I	C
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration. ^{766,767}	IIa	B
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias ^c or who are older than 40 years and have significant LGE on CMR. ^{c,5,14,16,766}	IIa	B
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In patients with limb-girdle type 1B or Emery-Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered. ⁷⁶⁹	IIa	C
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR. ^{770,771}	IIb	C
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ^d for VAs and SCD.	IIb	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders. ⁷⁷²	III	C
Management of VA		
In symptomatic patients with BBR-VT, catheter ablation is recommended. ^{e,153,474,475,477}	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended. ¹⁵³	I	C

© ESC 2022

AV, atrioventricular; BBR-VT, bundle branch re-entrant ventricular tachycardia; CA, cardiac arrest; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LBBB, left bundle branch block; LGE, late gadolinium enhancement; LV, left ventricular; SCD, sudden cardiac death; SD, sudden death; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cLevel of evidence C.

^dFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6-9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^eIn other cardiac conditions (e.g. aortic valve replacement) with BBR-VT, catheter ablation is also recommended.

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in neuromuscular diseases

Recommendations	Class ^a	Level ^b
General recommendations		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease. ^{6,13}	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders. ^{17,765}	I	C
Risk stratification, primary and secondary prevention of SCD		
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA. ¹⁵³	I	C

ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT. ⁷⁶⁶	I	C
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration. ^{766,767}	IIa	B
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias ^c or who are older than 40 years and have significant LGE on CMR. ^{c,5,14,16,766}	IIa	B
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In patients with limb-girdle type 1B or Emery-Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered. ⁷⁶⁹	IIa	C
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR. ^{770,771}	IIb	C
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ^d for VAs and SCD.	IIb	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders. ⁷⁷²	III	C
Management of VA		
In symptomatic patients with BBR-VT, catheter ablation is recommended. ^{e,153,474,475,477}	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended. ¹⁵³	I	C

© ESC 2022

AV, atrioventricular; BBR-VT, bundle branch re-entrant ventricular tachycardia; CA, cardiac arrest; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LBBB, left bundle branch block; LGE, late gadolinium enhancement; LV, left ventricular; SCD, sudden cardiac death; SD, sudden death; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cLevel of evidence C.

^dFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6–9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^eIn other cardiac conditions (e.g. aortic valve replacement) with BBR-VT, catheter ablation is also recommended.

Recommendations	Class ^a	Level ^b
General recommendations		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease. ^{6,13}	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders. ^{17,765}	I	C
Risk stratification, primary and secondary prevention of SCD		
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA or surviving a CA. ¹⁵³	I	C

ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT. ⁷⁶⁶	I	C
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a sudden increase in the PR interval or QRS duration. ^{766,767}	IIa	B
Invasive electrophysiological evaluation should be considered in patients with myotonic dystrophy and a PR interval ≥ 240 ms or QRS duration ≥ 120 ms or who are older than 40 years and have supraventricular arrhythmias ^c or who are older than 40 years and have significant LGE on CMR. ^{c,5,14,16,766}	IIa	B
In myotonic dystrophy patients without AV conduction delay and a syncope highly suspicious for VA, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In myotonic dystrophy patients with palpitations highly suspicious for VA and induction of a non-BBR-VT, ICD implantation should be considered. ⁷⁶⁶	IIa	C
In patients with limb-girdle type 1B or Emery-Dreifuss muscular dystrophies and indication for pacing, ICD implantation should be considered. ⁷⁶⁹	IIa	C
Implantation of an ICD may be considered in patients with Duchenne/Becker muscular dystrophy and significant LGE at CMR. ^{770,771}	IIb	C
Implantation of an ICD over a permanent pacemaker may be considered in myotonic dystrophy patients with additional risk factors ^d for VAs and SCD.	IIb	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders. ⁷⁷²	III	C
Management of VA		
In symptomatic patients with BBR-VT, catheter ablation is recommended. ^{e,153,474,475,477}	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended. ¹⁵³	I	C

© ESC 2022

AV, atrioventricular; BBR-VT, bundle branch re-entrant ventricular tachycardia; CA, cardiac arrest; CMR, cardiac magnetic resonance; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LBBB, left bundle branch block; LGE, late gadolinium enhancement; LV, left ventricular; SCD, sudden cardiac death; SD, sudden death; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cLevel of evidence C.

^dFactors favouring ICD implantation: Age,^{5,6,11} CTG expansion,^{6–9,13,16} SD or family history of SD,⁵ ECG conduction abnormalities,¹⁶ PR prolongation,¹³ LBBB,⁵ atrial arrhythmias,^{6,16} non-sustained VT,⁵ LV dysfunction,¹⁷ structural abnormalities in CMR.^{14,15,18}

^eIn other cardiac conditions (e.g. aortic valve replacement) with BBR-VT, catheter ablation is also recommended.

7.1.4. Inflammatory cardiac diseases

Cardiac inflammation has been associated with altered myocardial electrical properties and arrhythmias, and has been reported in various cardiomyopathies such as, obviously, myocarditis, but also inherited cardiomyopathies or after MI.^{773,774}

Inflammatory cardiomyopathies are characterized by myocardial inflammation (i.e. myocarditis) as the *primary* cause of cardiac damage, whereas *secondary* myocardial inflammations are caused by an *initial myocardial pathology*.^{775,776}

Myocarditis, sarcoidosis, and Chagas⁷⁷⁷ disease are among the major entities within inflammatory cardiomyopathies.

7.1.4.1. Myocarditis

The diagnosis of myocarditis can be challenging. There is no pathognomonic clinical presentation of this disease,⁷⁷⁷ with cardiac manifestations ranging from subtle symptoms to severe heart failure, complete AV block, and SCD.⁷⁷⁸ In young people, it is estimated that 2⁷⁷⁹12% of SCD are related to myocarditis.^{779,780,781}

Myocarditis is diagnosed on endomyocardial biopsy by established histological, immunological, and immunohistochemical criteria, as well as polymerase chain reaction (PCR) to detect viral genomes.^{782,783} This implies that endomyocardial biopsy, although not widely used, is the diagnostic gold standard for myocarditis.^{778,784} In current practice, the diagnosis of myocarditis relies on clinical presentation, elevated troponin level, ECG changes, evidence of LV dysfunction, absence of significant CAD or valvular heart disease, and suggestive findings on CMR or PET-CT.

In the context of suspected or confirmed acute myocarditis, patients with a life-threatening presentation (fulminant myocarditis, sustained VAs, or complete AV block) are to be referred to a specialized centre.^{785,786} A specialized centre must have the capacity to perform cardiac catheterization, endomyocardial biopsies, use mechanical circulatory support devices, and manage complex VAs.

Sustained VAs may occur in acute myocarditis. In a large series of patients,⁷⁸⁶ in-hospital VF or CA was reported in 2.5% of cases. In another series of acute myocarditis in children,⁷⁸⁷ sustained tachyarrhythmias were reported in 11.5% of patients, with VAs accounting for most cases (79.5%). Tachyarrhythmias were associated with a 2.3-fold increased risk of death.⁷⁸⁷ Giant cell myocarditis, although rare, has a higher risk of life-threatening VAs, which are seen in 14% of giant cell myocarditis patients on initial presentation, and then develop as refractory arrhythmias in more than half of the patients during the course of the disease.^{788,789}

The management of acute myocarditis depends on clinical presentation, viral PCR from myocardial biopsies, and histological findings. In patients with suspected myocarditis who present with mild heart failure, it is recommended to avoid exercise and use beta-blockers and ACE inhibitors.⁷⁹⁰ Patients with VAs or AV block must be admitted to the hospital and continuously monitored.

Treatment of arrhythmias in patients with inflammatory heart disease does not differ from generally accepted clinical principles. Symptomatic VAs may require AADs such as amiodarone and/or beta-blockers.^{791,792,793} Of note, in a retrospective observational study, patients with sustained VAs during the acute phase of myocarditis (LVEF 53 $\pm$ 10%) had a high risk (45% at 3 years) of VT/VF recurrences during follow-up.⁷⁹⁴

Myocarditis may resolve without sequelae, recur, or become chronic. Thus, myocarditis is often considered a precursor of DCM.⁷⁹⁵ Development of DCM has been observed in 21% of patients with acute myocarditis over a mean follow-up period of 3 years.⁷⁹⁶ The exact proportion of chronic DCM consecutive to the progression of acute myocarditis remains unknown, but myocarditis has been identified as the cause of DCM in up to 12% of cases in a large retrospective study.⁷⁹⁷ In patients with SMVT of unclear aetiology, myocarditis should be suspected especially when CMR reveals subepicardial and/or intramural abnormal fibrotic myocardial tissue. The presence of LGE at CMR has also been associated with the late occurrence of VAs in patients with endomyocardial biopsy-proven viral myocarditis.^{798,799,800}

The management of documented VAs in chronic myocarditis is similar to that of patients with DCM, including ICD implantation and the use of AADs. Catheter ablation can be effective to treat SMVT, the arrhythmogenic substrate of which is often found on the epicardium of the lateral/basal left ventricle.^{752,801,802,803} In one publication,⁸⁰² non-inducibility of VT after catheter ablation led to avoidance of ICD implantation in one-third of patients, with subsequently no VT recurrence during follow-up.

Recommendations	Class ^a	Level ^b
General recommendations		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VAs are referred to a specialized centre. ^{786,804}	I	C
Secondary prevention of SCD and treatment of VA		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended. ^{794,805}	I	C
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered. ^{788,794,806}	IIa	C
AADs should be considered (preferably amiodarone and beta-blockers) in patients with symptomatic non-sustained or sustained VAs during the acute phase of myocarditis.	IIa	C
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	IIa	C
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired. ^{752,801,802}	IIa	C
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	IIa	C
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^c	IIb	C

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; LV, left ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

© ESC 2022

Recommendations	Class ^a	Level ^b
General recommendations		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VAs are referred to a specialized centre. ^{786,804}	I	C
Secondary prevention of SCD and treatment of VA		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended. ^{794,805}	I	C
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered. ^{788,794,806}	IIa	C
AADs should be considered (preferably amiodarone and beta-blockers) in patients with symptomatic non-sustained or sustained VAs during the acute phase of myocarditis.	IIa	C
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	IIa	C
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired. ^{752,801,802}	IIa	C
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	IIa	C
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^c	IIb	C

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; LV, left ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

Recommendation Table 34

[Open in new tab](#)

Recommendations for sudden cardiac death prevention and treatment of ventricular arrhythmias in myocarditis

Recommendations	Class ^a	Level ^b
General recommendations		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VAs are referred to a specialized centre. ^{786,804}	I	C
Secondary prevention of SCD and treatment of VA		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended. ^{794,805}	I	C
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered. ^{788,794,806}	IIa	C
AADs should be considered (preferably amiodarone and beta-blockers) in patients with symptomatic non-sustained or sustained VAs during the acute phase of myocarditis.	IIa	C
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	IIa	C
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired. ^{752,801,802}	IIa	C
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	IIa	C
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^c	IIb	C

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; LV, left ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

Recommendations	Class ^a	Level ^b
General recommendations		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VAs are referred to a specialized centre. ^{786,804}	I	C
Secondary prevention of SCD and treatment of VA		
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended. ^{794,805}	I	C
In patients with haemodynamically not-tolerated sustained VT or VF during the acute phase of myocarditis, ICD implantation before hospital discharge should be considered. ^{788,794,806}	IIa	C
AADs should be considered (preferably amiodarone and beta-blockers) in patients with symptomatic non-sustained or sustained VAs during the acute phase of myocarditis.	IIa	C
In post-myocarditis patients with recurrent, symptomatic VT, AAD treatment should be considered.	IIa	C
Catheter ablation, performed in specialized centres, should be considered in post-myocarditis patients with recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, not tolerated, or not desired. ^{752,801,802}	IIa	C
In patients with haemodynamically tolerated SMVT occurring in the chronic phase of myocarditis, ICD implantation should be considered.	IIa	C
In patients with haemodynamically well-tolerated SMVT occurring in the chronic phase of myocarditis, preserved LV function and a limited scar amenable to ablation, catheter ablation may be considered as an alternative to ICD therapy, after discussion with the patient and provided that established endpoints have been reached. ^c	IIb	C

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; LV, left ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cVT non-inducibility and elimination of electrograms consistent with conduction delay.

7.1.4.2. Cardiac sarcoidosis

Sarcoidosis is a multisystem inflammatory disease of unknown cause, with a genetic predisposition,⁸⁰⁷ characterized by the accumulation of T lymphocytes, mononuclear phagocytes, and non-caseating granulomas leading to tissue scarring.⁸⁰⁸ Although lung involvement is most frequent, any organ can be affected. It is estimated that 5% of patients with sarcoidosis have symptoms indicative of cardiac involvement. However, using advanced cardiac imaging modalities (CMR/PET-CT), subclinical cardiac sarcoidosis is increasingly diagnosed.^{808a,809,810,811}

The diagnosis of cardiac sarcoidosis is challenging. It can mimic the ARVC phenotype in the event of a predominant RV involvement⁸¹⁰ or the heart is the only organ affected by sarcoidosis (so-called **AFACS** isolated cardiac sarcoidosis⁸¹²). Cardiac electroanatomical voltage mapping may help in the differential diagnosis between

isolated cardiac sarcoidosis and ARVC.⁸¹³ The three usual cardiac manifestations of cardiac sarcoidosis are LV dysfunction, AV conduction abnormalities, and VAs. Complete AV block develops primarily during the acute inflammatory phase as opposed to sustained VT, which frequently develops in the advanced stage of the disease (Figure 24).⁸¹⁴




Figure 24

Algorithm for sudden cardiac death prevention and treatment of ventricular arrhythmia in patients with cardiac sarcoidosis.

ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; N, No; PES, programmed electrical stimulation; SMVT, sustained monomorphic ventricular tachycardia; VA, ventricular arrhythmia; VT, ventricular tachycardia; Y, Yes. ^aLGE affecting ≥9/22 segments or ≥22% of the LV mass has been associated with arrhythmic endpoints.

[Open in new tab](#)[Download slide](#)

In addition to *isolated* cardiac sarcoidosis,⁸¹² three independent factors have been found associated with an adverse outcome and, particularly, the risk of VAs: (i) an LVEF <35%⁸¹²; (ii) documentation of high-degree AV block^{815,816}; and (iii) presence of RV or LV scarring at CMR (Figure 25).⁸¹⁷ However, VAs and SCD may occur in patients with a mildly reduced or normal LVEF. PES,⁸²² PET-CT,⁸²⁵ and LGE-CMR help stratify the risk of VAs in such patients.⁸²⁷ In a series of 120 patients with biopsy-proven extracardiac sarcoidosis and preserved LV/RV systolic function, VT inducibility at PES was low (6%) but associated with the composite endpoint of VAs and SCD, which occurred in 3 patients, all with abnormal PET or CMR ($P = 0.009$).⁸²⁵ In a cohort of 66 asymptomatic patients with biopsy-proven extracardiac sarcoidosis and abnormal PET-CT or CMR, 8 (11%) were inducible at PES and 6 of 8 had VA or died during follow-up, compared to 1 of 68 non-inducible patients ($P < 0.0001$). Of note, inducible patients had a lower LVEF at PES, which further deteriorated at 2-year follow-up.⁸²³




Figure 25

Typical features of cardiac sarcoidosis associated with sustained monomorphic ventricular tachycardia.

CMR, cardiac magnetic resonance; ECG, electrocardiogram; LBBB, left bundle branch block; LGE, late gadolinium enhancement; LV, left ventricle; PET-CT, positron emission tomography CT; RV, right ventricle; RVOT, right ventricular outflow tract; VT, ventricular tachycardia. ^aNote the prominent R' wave in V1 to V3, a pattern frequently found in cardiac sarcoidosis with RV involvement.⁸⁴⁴

[Open in new tab](#)[Download slide](#)

The benefit of ICDs, both for primary and secondary prevention, has been reported.^{828,831,832} Data support the use of ICDs for primary prevention of SCD in patients with LVEF $\geq 35\%$.^{812,830} In addition, whatever the LVEF, ICD should also be considered in patients with an indication for permanent cardiac pacing or the presence of significant scar at CMR.^{833,834} Indeed, as shown in two recent meta-analyses,^{821,832} the occurrence of VA is higher in patients with complete AV block (OR = 2.19; $P < 0.01$), and the presence of scar at CMR is associated with a higher risk of the composite endpoint of ventricular arrhythmic events and mortality (OR = 10.74; $P < 0.00001$). A widely accepted definition of *significant* LGE is not available. LGE in $\geq 9/29$ segments (17 LV and 12 RV segments) and LGE affecting $\geq 22\%$ of the LV mass have been associated with arrhythmic endpoints.^{820,827,835}

Corticosteroid therapy is a mainstay of cardiac sarcoidosis treatment, but the effect of this treatment on the occurrence of VAs has only been ascertained from observational studies.^{836,837,838} Obviously, based upon pathophysiology of the disease, inflammation plays a role in the development of ventricular scar in patients with cardiac sarcoidosis. However, so far there is no evidence that VAs are directly triggered by active inflammation itself. Indeed, a scar-related intramural or epicardial substrate (with a predilection in peri-valvular regions) explains most VTs in patients with cardiac sarcoidosis.⁸³⁹ The VT substrate was more likely located in segments with scar transmural at CMR and a lower degree of inflammation on PET-scan.⁸⁴⁰ Catheter ablation may help to prevent VT,^{839,841,842} in particular if AAD treatment fails,⁸³⁷ but VT recurrences are still frequent,^{752,839} in particular if catheter ablation is performed during active inflammation, with AADs continued in many patients.^{839,843}

Recommendation Table 35

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in cardiac sarcoidosis

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
ICD implantation is recommended in patients with cardiac sarcoidosis who have a LVEF $\leq 35\%$. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF. ⁸¹⁶	IIa	C
In patients with cardiac sarcoidosis who have a LVEF $>35\%$ but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered. ^{817–819,821,833,834}	IIa	B
In patients with cardiac sarcoidosis who have a LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	IIa	C
In patients with cardiac sarcoidosis, LVEF 35–50% and inducible SMVT at PES, ICD implantation should be considered. ^{823–825}	IIa	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	IIa	C
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD-recipients with recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{839,841,842}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrio-ventricular; CA, cardiac arrest; CMR, cardiac magnetic resonance; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic VT; AV, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
ICD implantation is recommended in patients with cardiac sarcoidosis who have a LVEF $\leq 35\%$. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF. ⁸¹⁶	IIa	C
In patients with cardiac sarcoidosis who have a LVEF $>35\%$ but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered. ^{817–819,821,833,834}	IIa	B
In patients with cardiac sarcoidosis who have a LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	IIa	C
In patients with cardiac sarcoidosis, LVEF 35–50% and inducible SMVT at PES, ICD implantation should be considered. ^{823–825}	IIa	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	IIa	C
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD-recipients with recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{839,841,842}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrio-ventricular; CA, cardiac arrest; CMR, cardiac magnetic resonance; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic VT; AV, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table 35

[Open in new tab](#)

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmias in cardiac sarcoidosis

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
ICD implantation is recommended in patients with cardiac sarcoidosis who have a LVEF $\leq 35\%$. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF. ⁸¹⁶	IIa	C
In patients with cardiac sarcoidosis who have a LVEF $>35\%$ but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered. ^{817–819,821,833,834}	IIa	B
In patients with cardiac sarcoidosis who have a LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	IIa	C
In patients with cardiac sarcoidosis, LVEF 35–50% and inducible SMVT at PES, ICD implantation should be considered. ^{823–825}	IIa	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	IIa	C
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD-recipients with recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{839,841,842}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; AV, atrio-ventricular; CA, cardiac arrest; CMR, cardiac magnetic resonance; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic VT; AV, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
ICD implantation is recommended in patients with cardiac sarcoidosis who have a LVEF $\leq 35\%$. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis who have an indication for permanent cardiac pacing related to high-degree AV block, ICD implantation should be considered, regardless of LVEF. ⁸¹⁶	IIa	C
In patients with cardiac sarcoidosis who have a LVEF $>35\%$ but significant LGE at CMR after resolution of acute inflammation, ICD implantation should be considered. ^{817–819,821,833,834}	IIa	B
In patients with cardiac sarcoidosis who have a LVEF 35–50% and minor LGE at CMR, after resolution of acute inflammation, PES for risk stratification should be considered.	IIa	C
In patients with cardiac sarcoidosis, LVEF 35–50% and inducible SMVT at PES, ICD implantation should be considered. ^{823–825}	IIa	C
Secondary prevention of SCD and treatment of VAs		
ICD implantation is recommended in patients with cardiac sarcoidosis who (1) have documented sustained VT, or (2) aborted CA. ^{812,828–830,832}	I	B
In patients with cardiac sarcoidosis and recurrent, symptomatic VA, AAD treatment should be considered.	IIa	C
Catheter ablation, in specialized centres, may be considered in cardiac sarcoidosis ICD-recipients with recurrent, symptomatic SMVT or ICD shocks for SMVT, in whom AADs are ineffective, contraindicated, or not tolerated. ^{839,841,842}	IIb	C

AAD, anti-arrhythmic drug; AV, atrio-ventricular; CA, cardiac arrest; CMR, cardiac magnetic resonance; ICD, implantable cardioverter defibrillator; LGE, late gadolinium enhancement; LVEF, left ventricular ejection fraction; PES, programmed electrical stimulation; SCD, sudden cardiac death; SMVT, sustained monomorphic VT; AV, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

7.1.4.3. Chagasâ€¦ cardiomyopathy

Chagasâ€¦ disease, a myocardial disease caused by the parasite *Trypanosoma cruzi*, is the most common cause of non-ischæmic cardiomyopathy in Latin America. After the incubation period, the vast majority of infected persons have minor or no symptoms, and very few ($<1\%$) develop acute myocarditis.⁸⁴⁵ However, people are infected for life, and years to decades after the initial infection 20â€¦30% of them will develop cardiomyopathy,⁸⁴⁶ which can lead to heart failure, heart block and atrial and ventricular arrhythmias. SCD, particularly due to VF, is the most common cause of death in patients with Chagasâ€¦ cardiomyopathy.^{847,848} A risk score developed by Rassi *et al.*⁸⁴⁹ as well as the presence of myocardial fibrosis at LGE-CMR⁸⁵⁰ are useful in assessing the risk of death in Chagasâ€¦ disease patients. However, whether such risk stratification can translate into clinical benefit through ICD treatment needs to be assessed in prospective trials, also considering the high annual mortality in Chagas patients with ICDs.⁸⁵¹

Although ICD implantation may seem reasonable for the secondary prevention of SCD in patients with Chagasâ€¦ cardiomyopathy, there are controversial studies^{851aâ€¦856} about the benefits and risks of ICD implantation in these patients. Indeed, a meta-analysis of observational studies found no benefit of ICD ($n = 483$) vs. amiodarone ($n = 115$)

for the secondary prevention of SCD in patients with Chagas' disease as all-cause annual mortality rates were comparable in both groups (9.7% and 9.6%, respectively).⁸⁵⁶ Another meta-analysis found a 4.7% annual incidence of inappropriate shocks and a high 9.0% annual mortality rate in 1041 ICD patients with Chagas' disease.⁸⁵¹

Amiodarone⁸⁵⁷ and catheter ablation^{858a,859,860} have been successfully used to control recurrent VAs in some patients. As with various forms of myocarditis, the arrhythmogenic substrate is often found at the epicardium in Chagas' cardiomyopathy.

Recommendation Table 36

[Open in new tab](#)

Recommendations for the treatment of ventricular arrhythmias in Chagas' cardiomyopathy

Recommendations	Class ^a	Level ^b
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT. ⁸⁵⁷	IIa	C
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered. ^{859,860}	IIa	C
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered. ^{851,854–856}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; PVCs, premature ventricular complexes; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia.

Recommendations	Class ^a	Level ^b
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT. ⁸⁵⁷	IIa	C
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered. ^{859,860}	IIa	C
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered. ^{851,854–856}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; PVCs, premature ventricular complexes; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia.

Recommendation Table 36

[Open in new tab](#)

Recommendations for the treatment of ventricular arrhythmias in Chagas' cardiomyopathy

Recommendations	Class ^a	Level ^b
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT. ⁸⁵⁷	IIa	C
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered. ^{859,860}	IIa	C
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered. ^{851,854–856}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; PVCs, premature ventricular complexes; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia.

Recommendations	Class ^a	Level ^b
Amiodarone should be considered to reduce arrhythmia burden in patients with Chagas' cardiomyopathy who present with symptomatic PVCs or VT. ⁸⁵⁷	IIa	C
In patients with Chagas' cardiomyopathy and recurrent, symptomatic SMVT or ICD shocks for SMVT in whom AADs are ineffective, contraindicated, or not tolerated, catheter ablation in specialized centres should be considered. ^{859,860}	IIa	C
In patients with Chagas' cardiomyopathy and symptomatic VT in whom AADs (amiodarone and beta-blockers) are ineffective or not tolerated, ICD implantation may be considered. ^{851,854–856}	IIb	C

© ESC 2022

AAD, anti-arrhythmic drug; ICD, implantable cardioverter defibrillator; PVCs, premature ventricular complexes; SMVT, sustained monomorphic ventricular tachycardia; VT, ventricular tachycardia.

7.1.5. Valvular heart disease

Valvular heart diseases predispose to SCD both in the pre-operative period and after valvular surgery. In old cohorts of symptomatic, non-operated patients with severe aortic stenosis or severe mitral regurgitation, the incidence of SCD was 50%⁸⁶¹ to 60%. In these studies, the rate of SCD secondary to VA was not specified.^{861,862}

Rates of SCD in patients with prosthetic valves range from 15% to 30%, with an estimated annual risk of 0.2%⁸⁶³ to 0.9%.⁸⁶³ In a large series of 1533 patients who underwent aortic or mitral valve replacement, 6% of deaths were caused by arrhythmias.⁸⁶⁴ A recent report of 3726 patients undergoing transcatheter aortic valve implantation (TAVI) showed 5.6% SCD during 22 months of follow-up, emphasizing the remaining SCD risk also after valve replacement.⁸⁶⁵

Small observational studies have shown that patients with residual LV dysfunction after valvular surgery who undergo ICD implantation have appropriate therapy and mortality rates similar to patients with ischaemic or dilated cardiomyopathy.^{866a}^{866b}⁸⁶⁸ ICD implantation in these patients should therefore follow DCM/HNDCM recommendations.

In patients after TAVI, SCD has been associated with the presence of new-onset conduction disturbances (LBBB, QRS > 160 ms) and reduced LVEF < 40%.⁸⁶⁵ SCD in these patients may be the consequence of advanced AV block, and permanent pacing indication should follow the 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy.² The role of ICD implantation for prevention of SCD after TAVI is less clear. Two recent retrospective analyses including a total of 203 patients in whom the LVEF remained < 35% did not show survival benefit from ICD implantation, probably related to the high competing risk of death from heart failure and relevant comorbidities in this population.^{869,870}

In patients presenting with SMVT after aortic valve replacement, PES is indicated because of the high probability of BBR-VT as the underlying mechanism⁸⁷¹ and an arrhythmia that is potentially curable with catheter ablation.⁸⁷²

SCD has been reported to occur in 0.2–0.4% of patients with MVP per year.^{873,874} However, the total number of patients dying suddenly with this condition is most likely underestimated, considering that it is a common valve variant with an estimated prevalence of 2–3% in the general population.⁸⁷⁵

Using rigorous criteria for morphologic diagnosis (i.e. myxomatous mitral valve) at post-mortem, MVP has been associated with 7% of all cases of juvenile SCD in the Registry of the Veneto region of Italy.²²⁸ Recent pathological and clinical studies on SCD victims with MVP or patients with MVP and major VAs have provided insights into the potential VA substrate.^{130,228,876–880} In particular, myocardial fibrosis affecting both the infero-basal LV free wall and the papillary muscles has been recognized in pathological and LGE-CMR studies.²²⁸ The scarring process has been related to mechanical stretch secondary to the excessive mobility of the mitral valve apparatus due to mitral valvular disjunction and posterior systolic curling.⁸⁷⁶ This indicates a promising role of CMR for arrhythmic risk stratification beyond traditional electrophysiological markers.

Proposed criteria for an arrhythmic MVP syndrome⁸⁸⁰ that have been associated with an increased risk for SCD include: young adults (most often women), QTc prolongation and/or negative T-wave in inferior ECG leads, mitral annular disjunction, bi-leaflet involvement on echocardiography, frequent PVCs, or non-sustained PVT on Holter monitoring and/or exercise testing.^{130,228,876–880} Reduced LVEF and/or myocardial fibrosis on CMR may add to the risk profile. These data are mainly derived from small retrospective series of SCD survivors and post-mortem studies. Therefore, identification of the small subgroup of patients at risk of SCD remains challenging.^{228,874} Robust data to support recommendations for risk stratification of SCD in patients with MVP are not available.

Recommendation Table 37

[Open in new tab](#)

Recommendations for sudden cardiac death prevention and treatment of ventricular arrhythmias in valvular heart disease

Recommendations	Class ^a	Level ^b
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention. ^{871,872,881}	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction, (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations. ⁸⁶⁸	I	C

© ESC 2022

BBR-VT, bundle branch re-entrant ventricular tachycardia; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LV, left ventricular; PES, programmed electrical stimulation; SMVT, sustained monomorphic ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention. ^{871,872,881}	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction, (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations. ⁸⁶⁸	I	C

BBR-VT, bundle branch re-entrant ventricular tachycardia; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LV, left ventricular; PES, programmed electrical stimulation; SMVT, sustained monomorphic ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table A 37

[Open in new tab](#)

Recommendations for sudden cardiac death prevention and treatment of ventricular arrhythmias in valvular heart disease

Recommendations	Class ^a	Level ^b
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention. ^{871,872,881}	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction, (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations. ⁸⁶⁸	I	C

BBR-VT, bundle branch re-entrant ventricular tachycardia; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LV, left ventricular; PES, programmed electrical stimulation; SMVT, sustained monomorphic ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention. ^{871,872,881}	I	C

In patients with valvular heart disease and persistent LV dysfunction after surgical correction, (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations.⁸⁶⁸

I

C

© ESC 2022


BBR-VT, bundle branch re-entrant ventricular tachycardia; DCM, dilated cardiomyopathy; HNDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; LV, left ventricular; PES, programmed electrical stimulation; SMVT, sustained monomorphic ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

7.1.6. Congenital heart disease

Advances in surgical repair and medical treatment have improved the long-term prognosis of children born with congenital heart disease (CHD). More than 90% now survive to adulthood,⁸⁸² and consequently the prevalence of adult CHD has steadily increased.⁸⁸³ With fewer patients dying from peri-operative events and early heart failure, SCD has become a leading cause of death in adults with repaired CHD.⁸⁸⁴ The combination of surgical incisions, myocardial scar and residual or new anatomical abnormalities form the substrate for VAs ([FigureÂ 26](#)).



FigureÂ 26

Algorithm for the management of sustained ventricular arrhythmia in patients with congenital heart disease.

CHD, congenital heart disease; ICD, implantable cardioverter defibrillator; N, No; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VT, ventricular tachycardia; Y, Yes. ^aData derived from patients with TOF and related lesions.

^bIn TOF, anti-arrhythmic drug failure is not required.

[Open in new tab](#)[Download slide](#)

Risk stratification for SCD in CHD patients and no documented sustained VA remains difficult due to a mixed patient population, the absence of RCTs, and relatively small observational studies. In patients with biventricular physiology and a systemic LV, the standard criterion of an LVEF $\geq 35\%$ is used for patient selection for primary prevention ICD implantation.^{885,886} For patients with unexplained syncope, severe arrhythmia symptoms such as palpitations or pre-syncope and NSVT, PES should be considered.⁸⁸⁷ In asymptomatic patients with tetralogy of Fallot (TOF) but other indicators for a VA substrate, electrophysiological evaluation may refine risk stratification.⁸⁸⁸ The surgical era and the techniques used affect the VA substrate and occurrence of VAs and should be considered. The use of a transannular patch repair in TOF patients, for example, was found to be associated with a lower risk for VA.⁸⁸⁹ The benefit of primary prevention ICD therapy in patients with single or systemic RVs is less well established and requires consideration of disease- and patient-specific factors.^{890,891}

In CHD patients with sustained VA or CA survivors, a comprehensive evaluation of inciting factors, including cardiac imaging (particularly CMR) and haemodynamic assessment, is important ([Figure 26](#)).⁸⁹² If pre-existing or new anatomical abnormalities requiring intervention are detected, the treatment plan should consider pre- or intra-operative mapping and transection of the VT substrate, because accessibility might be impaired post-operatively.^{888,893,894} Pre-operative evaluation and intervention are especially important in TOF patients undergoing surgical pulmonary re-valving. In selected patients with CHD (including those with atrial baffle repair for transposition of the great arteries, Fontan operation and Ebstein anomaly), evaluation and treatment of SVT (such as intra-atrial re-entrant tachycardias or AV re-entrant tachycardias) with rapid ventricular conduction should also be considered.^{890,895,896}

In the absence of identifiable reversible factors, ICD implantation for secondary prevention of SCD is recommended.^{349,350} While transvenous ICD systems have been most frequently used and have the advantage of anti-tachycardia and anti-bradycardia pacing, subcutaneous ICDs may be an alternative in selected patients with limited venous access to the ventricle or with intracardiac shunts.

In CHD patients, MVT most often arises from re-entrant tachycardias using anatomical isthmuses bounded by valves, patch material, and surgical incisions. Early mapping and ablation studies, particularly in TOF patients, identified critical anatomical isthmuses of reproducible anatomical location.^{897,898} Following this concept, those anatomical isthmuses can be reconstructed and transected in sinus rhythm during catheter ablation procedures using electroanatomical mapping systems, achieving acute success rates of 80%.^{888,899,900,901} The use of conduction block across anatomical isthmuses as a procedural endpoint, in addition to non-inducibility of VT, has further improved long-term ablation outcomes.^{888,899} Accordingly, catheter ablation is recommended in particular in TOF patients with recurrent SMVT. Due to the complexity of CHD patients as well as the VT substrate, those procedures should be performed in centres with expertise in catheter ablation of CHD patients. Catheter ablation may be considered in lieu of ICD implantation in selected TOF patients with SMVT and preserved biventricular function, provided the combined procedural endpoint of non-inducibility and conduction block across the anatomical isthmus can be reached.^{888,899}

Recommendation Table 38

[Open in new tab](#)

Recommendations for risk stratification and primary prevention of sudden cardiac death in congenital heart disease

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
All CHD patients		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF ≤35% despite ≥3 months of OMT, ICD implantation is indicated. ^{885,886}	I	C

In patients with CHD with presumed arrhythmic syncope and with either at least moderate ventricular dysfunction or inducible SMVT on PES, ICD implantation should be considered. ^{887,889,902}	IIa	C
In patients with advanced single ventricle or systemic RV dysfunction with additional risk factors, ^c ICD implantation may be considered. ^{890,891}	IIb	C
Tetralogy of Fallot		
In patients after repair of TOF with arrhythmia symptoms and NSVT, electrophysiologic evaluation including PES should be considered. ^{889,903–905}	IIa	C
In patients after repair of TOF with arrhythmia symptoms and a positive PES, or a combination of other risk factors ^d and a positive PES, ICD implantation should be considered.	IIa	C
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^d electrophysiologic evaluation, including PES, may be considered.	IIb	C
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered. ⁸⁹⁴	IIb	C

AV, atrioventricular; CHD, congenital heart disease; CMR, cardiac magnetic resonance; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained monomorphic ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; RV, right ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; TOF, tetralogy of Fallot; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cData are sparse and risk factors may be lesion-specific, including non-sustained VT, NYHA II/III, severe AV valve regurgitation, and wide QRS ≥ 140 ms (transposition of the great arteries).

^dOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR,^{906,907} QRS duration ≥ 180 ms^{886,908} and severe QRS fragmentation.^{909,910}

© ESC 2022

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
All CHD patients		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF $\leq 35\%$ despite ≥ 3 months of OMT, ICD implantation is indicated. ^{885,886}	I	C

In patients with CHD with presumed arrhythmic syncope and with either at least moderate ventricular dysfunction or inducible SMVT on PES, ICD implantation should be considered. ^{887,889,902}	IIa	C
In patients with advanced single ventricle or systemic RV dysfunction with additional risk factors, ^c ICD implantation may be considered. ^{890,891}	IIb	C
Tetralogy of Fallot		
In patients after repair of TOF with arrhythmia symptoms and NSVT, electrophysiologic evaluation including PES should be considered. ^{889,903–905}	IIa	C
In patients after repair of TOF with arrhythmia symptoms and a positive PES, or a combination of other risk factors ^d and a positive PES, ICD implantation should be considered.	IIa	C
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^d electrophysiologic evaluation, including PES, may be considered.	IIb	C
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered. ⁸⁹⁴	IIb	C

AV, atrioventricular; CHD, congenital heart disease; CMR, cardiac magnetic resonance; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained monomorphic ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; RV, right ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; TOF, tetralogy of Fallot; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cData are sparse and risk factors may be lesion-specific, including non-sustained VT, NYHA II/III, severe AV valve regurgitation, and wide QRS ≥ 140 ms (transposition of the great arteries).

^dOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR,^{906,907} QRS duration ≥ 180 ms^{886,908} and severe QRS fragmentation.^{909,910}

© ESC 2022

Recommendation Table 38

[Open in new tab](#)

Recommendations for risk stratification and primary prevention of sudden cardiac death in congenital heart disease

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
All CHD patients		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF $\leq 35\%$ despite ≥ 3 months of OMT, ICD implantation is indicated. ^{885,886}	I	C

In patients with CHD with presumed arrhythmic syncope and with either at least moderate ventricular dysfunction or inducible SMVT on PES, ICD implantation should be considered. ^{887,889,902}	IIa	C
In patients with advanced single ventricle or systemic RV dysfunction with additional risk factors, ^c ICD implantation may be considered. ^{890,891}	IIb	C
Tetralogy of Fallot		
In patients after repair of TOF with arrhythmia symptoms and NSVT, electrophysiologic evaluation including PES should be considered. ^{889,903–905}	IIa	C
In patients after repair of TOF with arrhythmia symptoms and a positive PES, or a combination of other risk factors ^d and a positive PES, ICD implantation should be considered.	IIa	C
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^d electrophysiologic evaluation, including PES, may be considered.	IIb	C
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered. ⁸⁹⁴	IIb	C

AV, atrioventricular; CHD, congenital heart disease; CMR, cardiac magnetic resonance; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained monomorphic ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; RV, right ventricular; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; TOF, tetralogy of Fallot; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cData are sparse and risk factors may be lesion-specific, including non-sustained VT, NYHA II/III, severe AV valve regurgitation, and wide QRS ≥ 140 ms (transposition of the great arteries).

^dOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR,^{906,907} QRS duration ≥ 180 ms^{886,908} and severe QRS fragmentation.^{909,910}

© ESC 2022

Recommendations	Class ^a	Level ^b
Risk stratification and primary prevention of SCD		
All CHD patients		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF $\leq 35\%$ despite ≥ 3 months of OMT, ICD implantation is indicated. ^{885,886}	I	C

In patients with CHD with presumed arrhythmic syncope and with either at least moderate ventricular dysfunction or inducible SMVT on PES, ICD implantation should be considered. ^{887,889,902}	IIa	C
In patients with advanced single ventricle or systemic RV dysfunction with additional risk factors, ^c ICD implantation may be considered. ^{890,891}	IIb	C
Tetralogy of Fallot		
In patients after repair of TOF with arrhythmia symptoms and NSVT, electrophysiologic evaluation including PES should be considered. ^{889,903–905}	IIa	C
In patients after repair of TOF with arrhythmia symptoms and a positive PES, or a combination of other risk factors ^d and a positive PES, ICD implantation should be considered.	IIa	C
In patients after repair of TOF without arrhythmia symptoms, but with a combination of other risk factors, ^d electrophysiologic evaluation, including PES, may be considered.	IIb	C
In patients with repaired TOF undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention may be considered. ⁸⁹⁴	IIb	C

AV, atrioventricular; CHD, congenital heart disease; CMR, cardiac magnetic resonance; EF, ejection fraction; ICD, implantable cardioverter defibrillator; LV, left ventricle; NSVT, non-sustained monomorphic ventricular tachycardia; NYHA, New York Heart Association; OMT, optimal medical treatment; PES, programmed electrical stimulation; RV, right ventricle; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; TOF, tetralogy of Fallot; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

^cData are sparse and risk factors may be lesion-specific, including non-sustained VT, NYHA II/III, severe AV valve regurgitation, and wide QRS ≥ 140 ms (transposition of the great arteries).

^dOther risk factors include moderate RV or LV dysfunction, extensive RV scarring on CMR,^{906,907} QRS duration ≥ 180 ms^{886,908} and severe QRS fragmentation.^{909,910}

© ESC 2022

Recommendation Table 39

[Open in new tab](#)

Recommendations for secondary prevention of sudden cardiac death and treatment of ventricular arrhythmia in congenital heart disease

Recommendations	Class ^a	Level ^b
Secondary prevention of SCD and treatment of VA		
All CHD patients		
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended. ^{892,893}	I	B
In patients with CHD with not tolerated VT/aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes. ^{349,350}	I	C

In patients with CHD and recurrent, symptomatic SMVT or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered. ^{c 899–901}	IIa	C
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered. ^{890,895}	IIa	C
Tetralogy of Fallot		
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended. ^{899–901}	I	C
In patients with repaired TOF with SMVT who are undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention should be considered. ^{888,893,894}	IIa	C
In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy. ^{899,901}	IIb	C

CA, cardiac arrest; CHD, congenital heart disease; ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.
^cSpecific recommendations for TOF (Class I-B).

© ESC 2022

Recommendations	Class ^a	Level ^b
Secondary prevention of SCD and treatment of VA		
All CHD patients		
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended. ^{892,893}	I	B
In patients with CHD with not tolerated VT/ aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes. ^{349,350}	I	C

In patients with CHD and recurrent, symptomatic SMVT or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered. ^{c 899-901}	IIa	C
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered. ^{890,895}	IIa	C
Tetralogy of Fallot		
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended. ⁸⁹⁹⁻⁹⁰¹	I	C
In patients with repaired TOF with SMVT who are undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention should be considered. ^{888,893,894}	IIa	C
In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy. ^{899,901}	IIb	C

CA, cardiac arrest; CHD, congenital heart disease; ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.
^cSpecific recommendations for TOF (Class I-B).

© ESC 2022

Recommendation Table 39

[Open in new tab](#)

Recommendations for secondary prevention of sudden cardiac death and treatment of ventricular arrhythmia in congenital heart disease

Recommendations	Class ^a	Level ^b
Secondary prevention of SCD and treatment of VA		
All CHD patients		
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended. ^{892,893}	I	B
In patients with CHD with not tolerated VT/ aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes. ^{349,350}	I	C

In patients with CHD and recurrent, symptomatic SMVT or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered. ^{c 899–901}	IIa	C
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered. ^{890,895}	IIa	C
Tetralogy of Fallot		
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended. ^{899–901}	I	C
In patients with repaired TOF with SMVT who are undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention should be considered. ^{888,893,894}	IIa	C
In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy. ^{899,901}	IIb	C

CA, cardiac arrest; CHD, congenital heart disease; ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.
^cSpecific recommendations for TOF (Class I-B).

© ESC 2022

Recommendations	Class ^a	Level ^b
Secondary prevention of SCD and treatment of VA		
All CHD patients		
In patients with CHD presenting with sustained VAs, evaluation for residual lesions or new structural abnormalities is recommended. ^{892,893}	I	B
In patients with CHD with not tolerated VT/ aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes. ^{349,350}	I	C

In patients with CHD and recurrent, symptomatic SMVT or ICD shocks for SMVT not manageable by medical therapy or ICD reprogramming, catheter ablation performed in specialized centres should be considered. ^{c 899–901}	IIa	C
In selected patients with CHD (including atrial baffle repair for transposition of the great arteries, Fontan operation and Ebstein anomaly) presenting with CA, evaluation and treatment of SVT with rapid ventricular conduction should be considered. ^{890,895}	IIa	C
Tetralogy of Fallot		
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended. ^{899–901}	I	C
In patients with repaired TOF with SMVT who are undergoing surgical or transcatheter pulmonary valve replacement, pre-operative catheter mapping and transection of VT-related anatomical isthmuses before or during the intervention should be considered. ^{888,893,894}	IIa	C
In patients with repaired TOF with a preserved biventricular function and symptomatic SMVT, catheter ablation or concomitant surgical ablation performed in specialized centres may be considered as an alternative to ICD therapy. ^{899,901}	IIb	C

CA, cardiac arrest; CHD, congenital heart disease; ICD, implantable cardioverter defibrillator; SCD, sudden cardiac death; SMVT, sustained monomorphic ventricular tachycardia; SVT, supraventricular tachycardia; TOF, tetralogy of Fallot; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.
^cSpecific recommendations for TOF (Class I-B).

© ESC 2022

7.2. Primary electrical disease

7.2.1. Idiopathic ventricular fibrillation

The diagnosis of idiopathic ventricular fibrillation (IVF) is made in SCA survivors, preferably with documented VF, after exclusion of structural, channelopathic, metabolic, or toxicological aetiologies.^{135,222,911a,913} Diagnostic tests include blood chemistry, ECG (including high lead), cardiac CT/coronary angiography, telemetry/Holter, exercise stress test, echocardiogram, sodium-channel blocker testing and CMR (see [Section 5.2.3](#), scenario 3).^{135,222,911} Genetic testing for channelopathy and cardiomyopathy genes may be considered with a mutation yield of 3a⁹¹³17%.^{249,914,915} Clinical evaluation of first-degree family members may be considered, but the diagnostic yield is low. In particular the significance of detected early repolarization pattern (ERP) in asymptomatic relatives is uncertain.^{182,916}

In IVF patients, ICD implantation reduces the risk of arrhythmic death by up to 68% compared to amiodarone.^{352,917a,921} ([Figure 27](#)). In observational studies with a mean follow-up of 5a⁹²¹6 years, 21.0a⁹²¹29.6% of IVF patients experienced an arrhythmic recurrence, corresponding to an annual ICD shock rate of 3.6a⁹²¹5.7%, whereas 4.5a⁹²¹17.5% experienced inappropriate shocks.^{919a,921} Isoproterenol, verapamil, or quinidine have been employed for acute treatment of recurrent ICD discharges or electrical storm.^{912,913,918,922a,926} In several small studies, quinidine was highly effective in reducing or even preventing arrhythmia inducibility during programmed stimulation.^{923,924,926} Moreover, a retrospective study in 46 patients showed a reduction of mean ICD shocks from 7.5 per patient over 2.9 years to 0.9 shocks per patient over 3.7 years, with a decrease in ventricular storms from 36 to 3 after quinidine initiation.⁹²² In patients with recurrent episodes of VF triggered by a similar PVC unresponsive to medical treatment, catheter ablation has shown success

(Figure 28).^{186,221,333,493,927a,930} PVCs most commonly originate from the Purkinje system and can be eliminated with a high acute success rate of 87a^{186,221,333,493,927a,930}100%.^{186,221,333,493,927a,930} Detailed electroanatomic mapping may also reveal localized structural alterations (62.5% in one study).²⁴⁸




Figure 27

Algorithm for the management of patients with idiopathic ventricular fibrillation.

ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex.

[Open in new tab](#)[Download slide](#)




Figure 28

Idiopathic ventricular fibrillation triggered by short-coupled premature ventricular complexes.

ECG, electrocardiogram; PVC, premature ventricular complex; VF, ventricular fibrillation.

[Open in new tab](#)[Download slide](#)

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology. ^{222,911}	I	B
Clinical testing (history, ECG and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered. ^{222,278,916}	IIb	B
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered. ^{249,278,914,915}	IIb	B
Secondary prevention of SCD and treatment of VA		
ICD implantation is recommended in idiopathic VF. ^{352,917–919}	I	B
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF. ^{912,913,918,922–924,926}	IIa	C
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF. ^{923,924}	IIa	B
Catheter ablation by experienced electrophysiologists should be considered in idiopathic VF patients with recurrent episodes of VF triggered by a similar PVC non-responsive to medical treatment. ^{927–930}	IIa	C
ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SCA, sudden cardiac arrest; SCD, sudden cardiac death; VA, ventricular arrhythmia; VF, ventricular fibrillation.		
^a Class of recommendation.		
^b Level of evidence.		

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology. ^{222,911}	I	B
Clinical testing (history, ECG and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered. ^{222,278,916}	IIb	B
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered. ^{249,278,914,915}	IIb	B
Secondary prevention of SCD and treatment of VA		
ICD implantation is recommended in idiopathic VF. ^{352,917–919}	I	B
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF. ^{912,913,918,922–924,926}	IIa	C
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF. ^{923,924}	IIa	B
Catheter ablation by experienced electrophysiologists should be considered in idiopathic VF patients with recurrent episodes of VF triggered by a similar PVC non-responsive to medical treatment. ^{927–930}	IIa	C
ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SCA, sudden cardiac arrest; SCD, sudden cardiac death; VA, ventricular arrhythmia; VF, ventricular fibrillation.		
^a Class of recommendation.		
^b Level of evidence.		

© ESC 2022

Recommendation Table 40

[Open in new tab](#)

Recommendations for the management of patients with idiopathic ventricular fibrillation

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology. ^{222,911}	I	B
Clinical testing (history, ECG and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered. ^{222,278,916}	IIb	B
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered. ^{249,278,914,915}	IIb	B
Secondary prevention of SCD and treatment of VA		
ICD implantation is recommended in idiopathic VF. ^{352,917–919}	I	B
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF. ^{912,913,918,922–924,926}	IIa	C
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF. ^{923,924}	IIa	B
Catheter ablation by experienced electrophysiologists should be considered in idiopathic VF patients with recurrent episodes of VF triggered by a similar PVC non-responsive to medical treatment. ^{927–930}	IIa	C

© ESC 2022

ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SCA, sudden cardiac arrest; SCD, sudden cardiac death; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Diagnostic evaluation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology. ^{222,911}	I	B
Clinical testing (history, ECG and high precordial lead ECG, exercise test, echocardiogram) of first-degree family members of idiopathic VF patients may be considered. ^{222,278,916}	IIb	B
In idiopathic VF patients, genetic testing of genes related to channelopathy and cardiomyopathy may be considered. ^{249,278,914,915}	IIb	B
Secondary prevention of SCD and treatment of VA		
ICD implantation is recommended in idiopathic VF. ^{352,917–919}	I	B
Isoproterenol infusion, verapamil, or quinidine for acute treatment of an electrical storm or recurrent ICD discharges should be considered in idiopathic VF. ^{912,913,918,922–924,926}	IIa	C
Quinidine should be considered for chronic therapy to suppress an electrical storm or recurrent ICD discharges in idiopathic VF. ^{923,924}	IIa	B
Catheter ablation by experienced electrophysiologists should be considered in idiopathic VF patients with recurrent episodes of VF triggered by a similar PVC non-responsive to medical treatment. ^{927–930}	IIa	C
ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PVC, premature ventricular complex; SCA, sudden cardiac arrest; SCD, sudden cardiac death; VA, ventricular arrhythmia; VF, ventricular fibrillation. ^a Class of recommendation. ^b Level of evidence.		

© ESC 2022

7.2.2. Long QT syndrome (including acquired long QT syndrome)

LQTS is characterized by a prolonged QT interval and VAs mainly triggered by adrenergic activation. The mean age at presentation is 14 years. The annual rate of SCD in asymptomatic patients with untreated LQTS has been estimated to be less than 0.5%,⁸² while it increases to around 5% in those with history of syncope.⁹³¹

Rare variants in 17 genes⁹³² have been associated with LQTS. However, the causality for several of the identified genes has been questioned.¹⁶⁶ The undisputed genes are those causing LQT1, LQT2 and LQTS3: *KCNQ1*, *KCNH2* and *SCN5A*, respectively, with gene-specific triggers being exercise (LQTS1), emotional stress (LQTS2) and sleep (LQTS3). Genetic screening identifies a mutation in 75% of LQTS cases and three main genes account for 90% of positively genotyped cases.¹⁷⁸ The subtypes of LQTS may be grouped as follows:

1. Autosomal-dominant LQTS (prevalence: 1 in 2500) without extra-cardiac manifestation.
2. Autosomal-dominant LQTS with extra-cardiac manifestation, comprising:
 - Andersen-Tawil Syndrome (LQT7), increasingly considered its own entity.^{933,934}
 - Timothy Syndrome (LQT8), characterized by prolonged QT, syndactyly, cardiac malformations, autism spectrum disorder and dysmorphism.⁹³⁵
3. Autosomal-recessive LQTS (Jervell and Lange-Nielsen Syndrome), combining extreme QT prolongation with congenital deafness.⁹³⁶

This panel of experts has confirmed the diagnostic criteria proposed in the previous version of the guidelines: a QTc ≥ 480 ms or a LQTS risk score >3 (Table 10) for clinical diagnosis (Figures 29 and 30). In the presence of arrhythmic syncope or CA, a QTc ≥ 460 ms is sufficient to consider a diagnosis of LQTS. A challenge for the clinician is establishing the duration of the QT interval in patients with broad QRS complexes (e.g. in the presence of ventricular pacing or ventricular conduction defects). In this setting, a formula has been proposed that adjusts QT by QRS duration.⁹³⁸ While measuring the QT, the patient moving briskly from recumbent to orthostatic position may be helpful for the diagnosis of LQTS.^{232,939} Epinephrine challenge is not recommended as a routine diagnostic tool, as reproducibility is modest.¹³⁷ Patients with a clinical diagnosis of LQTS are recommended to undergo genetic counselling and testing in specialized centres to receive genotype-specific management and permit identification of relatives at risk. Relatives with a mutation but without QT prolongation still receive a diagnosis of LQTS, as they are at risk of experiencing VA, although less frequently than phenotype-positive patients.⁹⁴⁰




Figure 29

Long QT syndrome electrocardiograms and torsade-de-pointes ventricular tachycardia.

(A) ECG characteristics in the three major LQTS phenotypes. (B) Example of Torsade-de-pointes in a male patient with a *SCN5A* (c.1238C>A, p.A413E) mutation. LQT, Long QT.

[Open in new tab](#)[Download slide](#)




Figure 30

Brisk standing electrocardiogram changes and T wave alternans in Long QT syndrome patients.

(A) ECG changes during brisk standing test in a male LQTS patient with *KCNH2* (p.S818L) mutation, increased heart rate is associated with less adaptation of QT interval. (B) T wave alternans in a male patient with *CACNA1C* (p.G406R) mutation.

[Open in new tab](#)[Download slide](#)

Table 10

[Open in new tab](#)

Modified long QT syndrome diagnostic score²⁴³

Findings			Points
ECG	QTc	≥480 ms	3.5
		≥460 ms	2
		≥450 ms (in males)	1
		≥480 ms during 4th minute of recovery from exercise stress test	1
	Torsade de pointes		2
	T wave alternans		1
	Notched T wave in 3 leads		1
	Low heart rate for age		0.5
Clinical history	Syncope	With stress	2
		Without stress	1
Family history	Family member(s) with definite LQTS		1
	Unexplained SCD at age <30 years in first-degree family		0.5
Genetic finding	Pathogenic mutation		3.5

Findings			Points
ECG	QTc	≥480 ms	3.5
		≥460 ms	2
		≥450 ms (in males)	1
		≥480 ms during 4th minute of recovery from exercise stress test	1
	Torsade de pointes		2
	T wave alternans		1
	Notched T wave in 3 leads		1
	Low heart rate for age		0.5
Clinical history	Syncope	With stress	2
		Without stress	1
Family history	Family member(s) with definite LQTS		1
	Unexplained SCD at age <30 years in first-degree family		0.5
Genetic finding	Pathogenic mutation		3.5

ECG, electrocardiogram; LQTS, long QT syndrome; SCD, sudden cardiac death.

Diagnosis of LQTS with a score >3.

Â© ESC 2022

TableÂ 10

[Open in new tab](#)

Modified long QT syndrome diagnostic score²⁴³

Findings			Points
ECG	QTc	≥ 480 ms	3.5
		≥ 460 ms	2
		≥ 450 ms (in males)	1
		≥ 480 ms during 4th minute of recovery from exercise stress test	1
	Torsade de pointes		2
	T wave alternans		1
	Notched T wave in 3 leads		1
	Low heart rate for age		0.5
Clinical history	Syncope	With stress	2
		Without stress	1
Family history	Family member(s) with definite LQTS		1
	Unexplained SCD at age <30 years in first-degree family		0.5
Genetic finding	Pathogenic mutation		3.5

Findings			Points
ECG	QTc	≥ 480 ms	3.5
		≥ 460 ms	2
		≥ 450 ms (in males)	1
		≥ 480 ms during 4th minute of recovery from exercise stress test	1
	Torsade de pointes		2
	T wave alternans		1
	Notched T wave in 3 leads		1
	Low heart rate for age		0.5
Clinical history	Syncope	With stress	2
		Without stress	1
Family history	Family member(s) with definite LQTS		1
	Unexplained SCD at age <30 years in first-degree family		0.5
Genetic finding	Pathogenic mutation		3.5

ECG, electrocardiogram; LQTS, long QT syndrome; SCD, sudden cardiac death.

Diagnosis of LQTS with a score >3.

Survivors of a CA have a high risk of recurrences, even on beta-blockers (14% within 5 years on therapy), supporting the use of ICD in CA survivors.⁹⁵² Moreover, ICD implantation is indicated when a patient experiences syncope and/or VA despite optimal pharmacological therapy, as syncopal events are associated with increased risk of CA.^{953,954} Women with LQTS, particularly LQT2, have an increased risk of arrhythmia both during pregnancy and especially first year post-partum.⁹⁵⁵ Silent carriers of mutations present a low, but not negligible, risk of cardiac events, and the use of beta-blockers should be considered in this group of patients.⁹⁵⁶

[Figure 31](#) illustrates the algorithm for the management of patients with long QT syndrome.



Algorithm for the management of patients with long QT syndrome.

ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQT, long QT; N, No; VA, ventricular arrhythmia; Y, Yes. ^aGeneral recommendations: avoidance of QT-prolonging drugs (<http://www.crediblemeds.org>), correction of electrolyte abnormalities (hypokalaemia, hypomagnesaemia, and hypocalcaemia), avoidance of genotype-specific triggers for arrhythmias (strenuous swimming in LQT1, exposure to loud noises in LQT2). ^bPreferred beta-blockers: nadolol and propranolol.

[Open in new tab](#)[Download slide](#)

Recommendation Table 41

[Open in new tab](#)

Recommendations for the management of patients with long QT syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that LQTS is diagnosed with either QTc $\geq$ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3 .	I	C
In patients with clinically diagnosed LQTS, genetic testing and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
The LQTS diagnosis should be considered in the presence of a QTc $\geq$ 460 ms and $<$ 480 ms in repeated 12-lead ECGs in patients with an arrhythmic syncope in the absence of secondary causes for QT prolongation. ^{952,962,963}	IIa	C
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS. ¹³⁷	III	C
General recommendations to prevent SCD		
The following is recommended in LQTS: • Avoid QT-prolonging drugs.^c • Avoid and correct electrolyte abnormalities. • Avoid genotype-specific triggers for arrhythmias.⁹⁴³	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events. ^{940,945,946}	I	B
Mexiletine is indicated in LQT3 patients with a prolonged QT interval. ⁹⁴⁸	I	C
Beta-blockers should be considered in patients with a pathogenic mutation and a normal QTc interval. ⁸²	IIa	B
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA. ^{952,953,962,963}	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ^d while receiving beta-blockers and genotype-specific therapies.	I	C

LCSD is indicated in patients with symptomatic ^d LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific drugs with an ICD and experiences multiple shocks or syncope due to VA. ^{541,957–959}	I	C
Either ICD implantation or LCSD should be considered in patients with symptomatic ^d LQTS, when beta-blockers and genotype-specific therapies are not tolerated or contraindicated at the therapeutic dose.	IIa	C
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval. ⁹⁴⁰	IIa	C
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients). ^{82,940,947,948}	IIb	B
Invasive electrophysiologic study is not recommended in LQTS. ⁹⁶¹	III	C

CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; SCD, sudden cardiac death; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

^c<http://www.crediblemeds.org>

^dArrhythmic syncope or haemodynamically non-tolerated VA.

© ESC 2022

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3 .	I	C
In patients with clinically diagnosed LQTS, genetic testing and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
The LQTS diagnosis should be considered in the presence of a QTc ≥ 460 ms and <480 ms in repeated 12-lead ECGs in patients with an arrhythmic syncope in the absence of secondary causes for QT prolongation. ^{952,962,963}	IIa	C

Routine diagnostic testing with epinephrine challenge is not recommended in LQTS. ¹³⁷	III	C
General recommendations to prevent SCD		
The following is recommended in LQTS: • Avoid QT-prolonging drugs. ^c • Avoid and correct electrolyte abnormalities. • Avoid genotype-specific triggers for arrhythmias. ⁹⁴³	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events. ^{940,945,946}	I	B
Mexiletine is indicated in LQT3 patients with a prolonged QT interval. ⁹⁴⁸	I	C
Beta-blockers should be considered in patients with a pathogenic mutation and a normal QTc interval. ⁸²	IIa	B
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA. ^{952,953,962,963}	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ^d while receiving beta-blockers and genotype-specific therapies.	I	C
LCSD is indicated in patients with symptomatic ^d LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific drugs with an ICD and experiences multiple shocks or syncope due to VA. ^{541,957–959}	I	C
Either ICD implantation or LCSD should be considered in patients with symptomatic ^d LQTS, when beta-blockers and genotype-specific therapies are not tolerated or contraindicated at the therapeutic dose.	IIa	C
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval. ⁹⁴⁰	IIa	C
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients). ^{82,940,947,948}	IIb	B
Invasive electrophysiologic study is not recommended in LQTS. ⁹⁶¹	III	C
© ESC 2022		
CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; SCD, sudden cardiac death; VA, ventricular arrhythmia.		
^a Class of recommendation.		
^b Level of evidence.		
^c http://www.crediblemeds.org		
^d Arrhythmic syncope or haemodynamically non-tolerated VA.		

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3 .	I	C
In patients with clinically diagnosed LQTS, genetic testing and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
The LQTS diagnosis should be considered in the presence of a QTc ≥ 460 ms and <480 ms in repeated 12-lead ECGs in patients with an arrhythmic syncope in the absence of secondary causes for QT prolongation. ^{952,962,963}	IIa	C
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS. ¹³⁷	III	C
General recommendations to prevent SCD		
The following is recommended in LQTS: • Avoid QT-prolonging drugs.^c • Avoid and correct electrolyte abnormalities. • Avoid genotype-specific triggers for arrhythmias.⁹⁴³	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events. ^{940,945,946}	I	B
Mexiletine is indicated in LQT3 patients with a prolonged QT interval. ⁹⁴⁸	I	C
Beta-blockers should be considered in patients with a pathogenic mutation and a normal QTc interval. ⁸²	IIa	B
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA. ^{952,953,962,963}	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ^d while receiving beta-blockers and genotype-specific therapies.	I	C

LCSD is indicated in patients with symptomatic ^d LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific drugs with an ICD and experiences multiple shocks or syncope due to VA. ^{541,957–959}	I	C
Either ICD implantation or LCSD should be considered in patients with symptomatic ^d LQTS, when beta-blockers and genotype-specific therapies are not tolerated or contraindicated at the therapeutic dose.	IIa	C
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval. ⁹⁴⁰	IIa	C
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients). ^{82,940,947,948}	IIb	B
Invasive electrophysiologic study is not recommended in LQTS. ⁹⁶¹	III	C

CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; SCD, sudden cardiac death; VA, ventricular arrhythmia.

^aClass of recommendation.

^bLevel of evidence.

^c<http://www.crediblemeds.org>

^dArrhythmic syncope or haemodynamically non-tolerated VA.

© ESC 2022

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3 .	I	C
In patients with clinically diagnosed LQTS, genetic testing and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
The LQTS diagnosis should be considered in the presence of a QTc ≥ 460 ms and <480 ms in repeated 12-lead ECGs in patients with an arrhythmic syncope in the absence of secondary causes for QT prolongation. ^{952,962,963}	IIa	C

Routine diagnostic testing with epinephrine challenge is not recommended in LQTS. ¹³⁷	III	C
General recommendations to prevent SCD		
The following is recommended in LQTS: • Avoid QT-prolonging drugs.^c • Avoid and correct electrolyte abnormalities. • Avoid genotype-specific triggers for arrhythmias.⁹⁴³	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events. ^{940,945,946}	I	B
Mexiletine is indicated in LQT3 patients with a prolonged QT interval. ⁹⁴⁸	I	C
Beta-blockers should be considered in patients with a pathogenic mutation and a normal QTc interval. ⁸²	IIa	B
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA. ^{952,953,962,963}	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ^d while receiving beta-blockers and genotype-specific therapies.	I	C
LCSD is indicated in patients with symptomatic ^d LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific drugs with an ICD and experiences multiple shocks or syncope due to VA. ^{541,957–959}	I	C
Either ICD implantation or LCSD should be considered in patients with symptomatic ^d LQTS, when beta-blockers and genotype-specific therapies are not tolerated or contraindicated at the therapeutic dose.	IIa	C
In LQTS, it should be considered to calculate the arrhythmic risk before initiation of therapy based on the genotype and the duration of QTc interval. ⁹⁴⁰	IIa	C
ICD implantation may be considered in asymptomatic LQTS patients with high-risk profile (according to the 1-2-3 LQTS Risk calculator) in addition to genotype-specific medical therapies (mexiletine in LQT3 patients). ^{82,940,947,948}	IIb	B
Invasive electrophysiologic study is not recommended in LQTS. ⁹⁶¹	III	C
© ESC 2022		
CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; LQTS, long QT syndrome; SCD, sudden cardiac death; VA, ventricular arrhythmia. ^aClass of recommendation. ^bLevel of evidence. ^chttp://www.crediblemeds.org ^dArrhythmic syncope or haemodynamically non-tolerated VA.		

7.2.3. Andersen-Tawil syndrome Type 1

Andersen-Tawil syndrome Type 1, also classified as LQT7, is a rare disease (1:1 000 000) characterized by three main symptoms: frequent VA (e.g. bidirectional VT), dysmorphologies and periodic paralysis.^{964,967,968} The inward rectifier current (I_{K1}) decreased by *KCNJ2* loss of function mutation⁹⁶⁸ causes an increase in U wave amplitude rather than QT prolongation.^{964,967,970} Syncope or documented VT is associated with life-threatening VA, and a study found a 5-year probability of 7.9% of life-threatening VA.⁹⁶⁷ In patients with sustained haemodynamically not-tolerated VT or CA, an ICD is recommended.^{964,967} Flecainide and/or beta-blockers seem to reduce VA,^{964,970,971} whereas amiodarone may be proarrhythmic and should only be used with an ICD.⁹⁶⁷ In patients with syncope despite medical therapy, an ICD or ILR should be discussed.⁹⁶⁷

Recommendation Table 42

[Open in new tab](#)

Recommendations for management of patients with Andersen-Tawil syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
Genetic testing is recommended in patients with suspected Andersen-Tawil syndrome. ^{964,967}	I	C
Andersen-Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • <i>KCNJ2</i> pathogenic loss of function mutation.^{964,965,967,968,972}	IIa	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with Andersen-Tawil syndrome after aborted CA or not-tolerated sustained VT. ^{964,967}	I	C
Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen-Tawil syndrome to treat VA. ^{964,970}	IIa	C
An ILR should be considered in patients with Andersen-Tawil syndrome and unexplained syncope.	IIa	C
ICD implantation may be considered in patients with Andersen-Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT. ⁹⁶⁷	IIb	C

© ESC 2022

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVCs, premature ventricular complexes; SCD, sudden cardiac death; SHD, structural heart disease; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Diagnosis		
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome. ^{964,967}	I	C
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.^{964,965,967,968,972}	IIa	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT. ^{964,967}	I	C
Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA. ^{964,970}	IIa	C
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa	C
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT. ⁹⁶⁷	IIb	C
CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVCs, premature ventricular complexes; SCD, sudden cardiac death; SHD, structural heart disease; VA, ventricular arrhythmia; VT, ventricular tachycardia. ^a Class of recommendation. ^b Level of evidence.		

© ESC 2022

Recommendation Table 42

[Open in new tab](#)

Recommendations for management of patients with Andersen–Tawil syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome. ^{964,967}	I	C
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.^{964,965,967,968,972}	IIa	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT. ^{964,967}	I	C
Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA. ^{964,970}	IIa	C
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa	C
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT. ⁹⁶⁷	IIb	C

© ESC 2022

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVCs, premature ventricular complexes; SCD, sudden cardiac death; SHD, structural heart disease; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.


Recommendations	Class ^a	Level ^b
Diagnosis		
Genetic testing is recommended in patients with suspected Andersen–Tawil syndrome. ^{964,967}	I	C
Andersen–Tawil syndrome should be considered in patients without SHD who present with at least two of the following: • Prominent U waves with or without prolongation of the QT interval • Bidirectional and/or polymorphic PVCs/VT • Dysmorphic features • Periodic paralysis • KCNJ2 pathogenic loss of function mutation.^{964,965,967,968,972}	IIa	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with Andersen–Tawil syndrome after aborted CA or not-tolerated sustained VT. ^{964,967}	I	C
Beta-blockers and/or flecainide with or without acetazolamide should be considered in patients with Andersen–Tawil syndrome to treat VA. ^{964,970}	IIa	C
An ILR should be considered in patients with Andersen–Tawil syndrome and unexplained syncope.	IIa	C
ICD implantation may be considered in patients with Andersen–Tawil syndrome who have a history of unexplained syncope or suffer from tolerated sustained VT. ⁹⁶⁷	IIb	C

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVCs, premature ventricular complexes; SCD, sudden cardiac death; SHD, structural heart disease; VA, ventricular arrhythmia; VT, ventricular tachycardia.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

7.2.4. Brugada syndrome

The type 1 Brugada ECG pattern is characterized by J point elevation of $>2\text{aPR}^{\text{a}}\text{mV}$ with coved ST elevation and T wave inversion in at least one right precordial ECG lead, V1 or V2, positioned in the second, third or fourth intercostal spaces (FigureÂ 32). It may occur either spontaneously or be induced by exposure to sodium channel-blocking drugs or fever.^{135,231,973aPROBUE978} It is mandatory to exclude other conditions that may explain the type 1 pattern, so-called phenocopies.⁹⁷⁹



FigureÂ 32

Typical examples of (A) Brugada type 1 electrocardiogram, and (B) early repolarization pattern electrocardiogram.


ECG, Electrocardiogram.

[Open in new tab](#)[Download slide](#)

BrS is diagnosed in patients without other heart disease and a spontaneous type 1 pattern, regardless of symptoms, due to its rarity in the general population and association with risk.^{231,979,980} The type 1 pattern may also be induced by administration of a sodium channel-blocking drug as a diagnostic test in patients suspected to have concealed BrS but without a spontaneous type 1 ECG.^{135,136,231,387,973,978,981a,985} However, provocation with drug or fever is less specific than previously thought, with 2a⁹⁸⁵4% prevalence in healthy subjects and higher prevalence in patients with AV nodal reentry tachycardia or an accessory pathway in one study.^{977,978,986} In the opinion of this panel of experts, an induced type 1 ECG pattern therefore requires other clinical features, such as documented PVT/VF, arrhythmic syncope, or relevant family history.

The yield of genetic testing in BrS patients is approximately 20%, with the *SCN5A* gene the only gene with evidence of association for clinical testing purposes.^{164,980} Phenotype and genotype mismatch is seen in *SCN5A* families, which is explained by variable effects of mutation severity and a polygenic risk score derived from genome-wide association studies.^{170,979,987} Recent data also support a potential for prognostication.^{988,989}

Psychotropic drugs, selected AADs, anaesthetic agents, cocaine, excessive alcohol intake and fever are potential triggers to exacerbate the type 1 pattern and trigger VF.^{231,297} The risk of recurrent VF among patients presenting with CA is 48% at 10 years. ICD implantation is therefore indicated in symptomatic BrS patients who are survivors of CA or have documented spontaneous sustained VA (*FigureÂ 33*).^{980,990a,992,994,996} Approximately one-third of BrS patients present with syncope.⁹⁹⁵ The risk of arrhythmic events in BrS patients with unexplained syncope is 4 times higher than the risk in asymptomatic patients.^{155,990a,992,994,996} Detailed patient history, including absence of prodrome or specific triggers, is essential for distinguishing arrhythmic from non-arrhythmic syncope. Nonetheless, the aetiology of syncope is difficult to determine in up to 30% of BrS patients. In small studies, arrhythmia detected by ILR changed clinical management in 20a⁹⁹⁵36% of the BrS patients with unexplained syncope.^{997a,998,999}



FigureÂ 33

Part One. Algorithm for the management of patients with Brugada pattern electrocardiogram.

[Open in new tab](#)[Download slide](#)

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern. ⁹⁷⁴⁻⁹⁷⁶	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever. ^{135,136,975,981,982}	I	C
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS. ^{164,1016}	I	C
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa	C

BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG. ^{136,973,975,978,984,985}	IIb	C
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III	C
General recommendations		
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA and/or (b) Have documented spontaneous sustained VT. ^{980,990–992}	I	C
ICD implantation should be considered in patients with type 1 Brugada pattern and an arrhythmic syncope. ^{990,992,996}	IIa	C
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope. ^{997,999}	IIa	C
Quinidine should be considered in patients with BrS who qualify for an ICD but have a contraindication, decline, or have recurrent ICD shocks. ^{922,1006,1007}	IIa	C
Isoproterenol infusion should be considered in BrS patients suffering electrical storm. ¹⁰⁰⁸	IIa	C
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy. ^{1010–1015}	IIa	C
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG. ¹⁵⁵	IIb	B
ICD implantation may be considered in selected asymptomatic BrS patients with inducible VF during PES using up to 2 extra stimuli. ¹⁵⁵	IIb	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C

© ESC 2022

BrS, Brugada syndrome; CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PES, programmed electrical stimulation; PVCs, premature ventricular complexes; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern. ⁹⁷⁴⁻⁹⁷⁶	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever. ^{135,136,975,981,982}	I	C
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS. ^{164,1016}	I	C
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa	C

BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG. ^{136,973,975,978,984,985}	IIb	C
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III	C
General recommendations		
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA and/or (b) Have documented spontaneous sustained VT. ^{980,990–992}	I	C
ICD implantation should be considered in patients with type 1 Brugada pattern and an arrhythmic syncope. ^{990,992,996}	IIa	C
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope. ^{997,999}	IIa	C
Quinidine should be considered in patients with BrS who qualify for an ICD but have a contraindication, decline, or have recurrent ICD shocks. ^{922,1006,1007}	IIa	C
Isoproterenol infusion should be considered in BrS patients suffering electrical storm. ¹⁰⁰⁸	IIa	C
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy. ^{1010–1015}	IIa	C
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG. ¹⁵⁵	IIb	B
ICD implantation may be considered in selected asymptomatic BrS patients with inducible VF during PES using up to 2 extra stimuli. ¹⁵⁵	IIb	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
BrS, Brugada syndrome; CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PES, programmed electrical stimulation; PVCs, premature ventricular complexes; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation. ^aClass of recommendation. ^bLevel of evidence.		

© ESC 2022

Recommendation Table 43

[Open in new tab](#)

Recommendations for management of patients with Brugada syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern. ⁹⁷⁴⁻⁹⁷⁶	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever. ^{135,136,975,981,982}	I	C
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS. ^{164,1016}	I	C
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa	C

BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG. ^{136,973,975,978,984,985}	IIb	C
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III	C
General recommendations		
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA and/or (b) Have documented spontaneous sustained VT. ^{980,990–992}	I	C
ICD implantation should be considered in patients with type 1 Brugada pattern and an arrhythmic syncope. ^{990,992,996}	IIa	C
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope. ^{997,999}	IIa	C
Quinidine should be considered in patients with BrS who qualify for an ICD but have a contraindication, decline, or have recurrent ICD shocks. ^{922,1006,1007}	IIa	C
Isoproterenol infusion should be considered in BrS patients suffering electrical storm. ¹⁰⁰⁸	IIa	C
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy. ^{1010–1015}	IIa	C
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG. ¹⁵⁵	IIb	B
ICD implantation may be considered in selected asymptomatic BrS patients with inducible VF during PES using up to 2 extra stimuli. ¹⁵⁵	IIb	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
© ESC 2022		

BrS, Brugada syndrome; CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PES, programmed electrical stimulation; PVCs, premature ventricular complexes; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation.

^aClass of recommendation.
^bLevel of evidence.


Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern. ⁹⁷⁴⁻⁹⁷⁶	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever. ^{135,136,975,981,982}	I	C
Genetic testing for <i>SCN5A</i> gene is recommended for probands with BrS. ^{164,1016}	I	C
BrS should be considered in patients with no other heart disease and induced type 1 Brugada pattern who have at least one of: • Arrhythmic syncope or nocturnal agonal respiration • A family history of BrS • A family history of SD (<45 years old) with a negative autopsy and circumstance suspicious for BrS.	IIa	C

BrS may be considered as a diagnosis in patients with no other heart disease who exhibit an induced type 1 Brugada ECG. ^{136,973,975,978,984,985}	IIb	C
Sodium channel blocker test is not recommended in patients with a prior type I Brugada pattern.	III	C
General recommendations		
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA and/or (b) Have documented spontaneous sustained VT. ^{980,990–992}	I	C
ICD implantation should be considered in patients with type 1 Brugada pattern and an arrhythmic syncope. ^{990,992,996}	IIa	C
Implantation of a loop recorder should be considered in BrS patients with an unexplained syncope. ^{997,999}	IIa	C
Quinidine should be considered in patients with BrS who qualify for an ICD but have a contraindication, decline, or have recurrent ICD shocks. ^{922,1006,1007}	IIa	C
Isoproterenol infusion should be considered in BrS patients suffering electrical storm. ¹⁰⁰⁸	IIa	C
Catheter ablation of triggering PVCs and/or RVOT epicardial substrate should be considered in BrS patients with recurrent appropriate ICD shocks refractory to drug therapy. ^{1010–1015}	IIa	C
PES may be considered in asymptomatic patients with a spontaneous type I BrS ECG. ¹⁵⁵	IIb	B
ICD implantation may be considered in selected asymptomatic BrS patients with inducible VF during PES using up to 2 extra stimuli. ¹⁵⁵	IIb	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
© ESC 2022		
BrS, Brugada syndrome; CA, cardiac arrest; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; PES, programmed electrical stimulation; PVCs, premature ventricular complexes; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation.		
^a Class of recommendation.		
^b Level of evidence.		

7.2.5. Early repolarization syndromes

Early repolarization syndrome (ERS) is diagnosed in a patient resuscitated from PVT or VF without any heart disease and the early repolarization pattern (ERP); J-point elevation ≥ 1 mm in ≥ 2 adjacent inferior and/or lateral ECG leads ([Figure A32](#)).^{135,231,1017,1019} However, ERP is most often a benign finding and the prevalence of ERP has been reported as 5.8% in adults and is more common in young males and athletes.^{135,231,1017,1021}

Nonetheless, ERP is over-represented in relatives of SADS cases^{282,1022} and of CA survivors.^{182,916} The diagnostic yield and utility of genetic testing is low.^{1023a[†]1025} High-risk ECG features have been proposed to increase likelihood of ERS: prominent J-waves $\geq 2a[†]2a[†]mm$, dynamic changes in J-point elevation ($>0.1a[†]2a[†]mV$) and J-waves associated with a horizontal or descending ST-segment ([FigureÂ 34](#)).^{231,1026,1027} ERP with a horizontal ST-segment was associated with arrhythmic risk in an elderly and IVF population.¹⁰²⁷ Nonetheless, ERS survivors and relatives of SADS cases also exhibit a higher prevalence of the ascending/upsloping ST segment than controls.^{282,1018} At least 40% of ERS patients with VF have subsequent episodes, with 27% suffering multiple episodes.^{231,1028,1029} Isoproterenol infusion is effective in acute suppression of recurrent ICD discharges and electrical storm.^{1030a[†]1032} AADs that block the transient outward potassium current can prevent VF.^{922,1030,1033} A retrospective multicentre study showed a reduction of recurrent VF after initiation of quinidine.¹⁰³⁰ Phosphodiesterase-3 inhibitors such as cilostazol and milrinone also reduced the recurrence of VF.¹⁰³² Ablation of a PVC trigger, usually from the Purkinje system, has an acute success rate of 87a[†]100% and may be effective in preventing recurrence in patients with drug-refractory VF.^{1010,1017} Detailed electroanatomic mapping may reveal localized structural alterations in 39% of ERS patients.¹⁰¹⁰ Ablation of these areas successfully suppressed electrical storm and may be a therapeutic option in experienced centres. Whether catheter ablation improves long-term outcomes is currently unknown.



FigureÂ 34
Management of patients with early repolarization pattern/syndrome.

ERP, early repolarization pattern; ERS, early repolarization syndrome; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; N, No; PVC, premature ventricular complex; SD, sudden death; Y, Yes. ^aERP high risk features: J waves $>2a[†]2a[†]mm$, dynamic changes in ST morphology.

[Open in new tab](#)[Download slide](#)

Data for risk stratification of patients with suspected ERS without prior CA are unavailable. In individuals with ERP and unexplained syncope, this panel has recommended that follow-up with an ILR should be considered. As the prognosis of asymptomatic subjects with ERP is good, ICD therapy is usually not indicated.^{1034a[†]1036} If, however, there is a high-risk ERP and a strong family history of unexplained juvenile SD, then ICD implantation or quinidine may be considered.

Recommendation TableÂ 44

[Open in new tab](#)

Recommendations for the management of patients with early repolarization pattern/syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads. ^{1017,1018}	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP. ^{1017,1018}	I	C
In an SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered. ^{1017,1018}	IIa	C
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^{c,1022,1037}	IIa	B
Genetic testing in ERS patients may be considered. ^{1023,1025}	IIb	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP. ^{1038,1039}	III	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA. ¹⁰¹⁷	I	B
Isoproterenol infusion should be considered for ERS patients with electrical storm. ^{1017,1030–1032}	IIa	B
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients. ^{922,1030,1033}	IIa	B
ILR should be considered in individuals with ERP and at least one risk feature ^d or arrhythmic syncope. ¹⁰²⁰	IIa	C
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment. ¹⁰¹⁰	IIa	C
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^{d,1030,1033}	IIb	C
ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^c in the presence of a family history of unexplained juvenile SD. ^{1030,1033}	IIb	C
ICD implantation is not recommended in asymptomatic patients with an isolated ERP. ^{1034,1035,1040}	III	C
CA, cardiac arrest; ECG, electrocardiogram; ERP, early repolarization pattern; ERS, early repolarization syndrome; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVC, premature ventricular complexes; PVT, ventricular tachycardia; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation. ^aClass of recommendation. ^bLevel of evidence. ^cERP high-risk features: J waves >2 mm, dynamic changes in J point and ST morphology.^{1020,1041} ^dHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.		

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads. ^{1017,1018}	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP. ^{1017,1018}	I	C
In an SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered. ^{1017,1018}	IIa	C
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^{c,1022,1037}	IIa	B
Genetic testing in ERS patients may be considered. ^{1023,1025}	IIb	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP. ^{1038,1039}	III	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA. ¹⁰¹⁷	I	B
Isoproterenol infusion should be considered for ERS patients with electrical storm. ^{1017,1030–1032}	IIa	B
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients. ^{922,1030,1033}	IIa	B
ILR should be considered in individuals with ERP and at least one risk feature ^d or arrhythmic syncope. ¹⁰²⁰	IIa	C
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment. ¹⁰¹⁰	IIa	C
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^{d,1030,1033}	IIb	C
ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^c in the presence of a family history of unexplained juvenile SD. ^{1030,1033}	IIb	C
ICD implantation is not recommended in asymptomatic patients with an isolated ERP. ^{1034,1035,1040}	III	C
CA, cardiac arrest; ECG, electrocardiogram; ERP, early repolarization pattern; ERS, early repolarization syndrome; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVC, premature ventricular complexes; PVT, ventricular tachycardia; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation. ^aClass of recommendation. ^bLevel of evidence. ^cERP high-risk features: J waves >2 mm, dynamic changes in J point and ST morphology.^{1020,1041} ^dHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.		

© ESC 2022

Recommendation Table 44

[Open in new tab](#)

Recommendations for the management of patients with early repolarization pattern/syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads. ^{1017,1018}	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP. ^{1017,1018}	I	C
In an SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered. ^{1017,1018}	IIa	C
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^{c,1022,1037}	IIa	B
Genetic testing in ERS patients may be considered. ^{1023,1025}	IIb	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP. ^{1038,1039}	III	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA. ¹⁰¹⁷	I	B
Isoproterenol infusion should be considered for ERS patients with electrical storm. ^{1017,1030–1032}	IIa	B
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients. ^{922,1030,1033}	IIa	B
ILR should be considered in individuals with ERP and at least one risk feature ^d or arrhythmic syncope. ¹⁰²⁰	IIa	C
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment. ¹⁰¹⁰	IIa	C
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^{d,1030,1033}	IIb	C
ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^c in the presence of a family history of unexplained juvenile SD. ^{1030,1033}	IIb	C
ICD implantation is not recommended in asymptomatic patients with an isolated ERP. ^{1034,1035,1040}	III	C
CA, cardiac arrest; ECG, electrocardiogram; ERP, early repolarization pattern; ERS, early repolarization syndrome; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVC, premature ventricular complexes; PVT, ventricular tachycardia; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation. ^aClass of recommendation. ^bLevel of evidence. ^cERP high-risk features: J waves >2 mm, dynamic changes in J point and ST morphology.^{1020,1041} ^dHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.		

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads. ^{1017,1018}	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP. ^{1017,1018}	I	C
In an SCD victim with a negative autopsy and medical chart review, and an ante-mortem ECG demonstrating the ERP, the diagnosis of ERS should be considered. ^{1017,1018}	IIa	C
First-degree relatives of ERS patients should be considered for clinical evaluation for ERP with additional high-risk features. ^{c,1022,1037}	IIa	B
Genetic testing in ERS patients may be considered. ^{1023,1025}	IIb	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP. ^{1038,1039}	III	C
Risk stratification, prevention of SCD and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA. ¹⁰¹⁷	I	B
Isoproterenol infusion should be considered for ERS patients with electrical storm. ^{1017,1030–1032}	IIa	B
Quinidine in addition to an ICD should be considered for recurrent VF in ERS patients. ^{922,1030,1033}	IIa	B
ILR should be considered in individuals with ERP and at least one risk feature ^d or arrhythmic syncope. ¹⁰²⁰	IIa	C
PVC ablation should be considered in ERS patients with recurrent VF episodes triggered by a similar PVC non-responsive to medical treatment. ¹⁰¹⁰	IIa	C
ICD implantation or quinidine may be considered in individuals with ERP and arrhythmic syncope and additional risk features. ^{d,1030,1033}	IIb	C
ICD implantation or quinidine may be considered in asymptomatic individuals who demonstrate a high-risk ERP ^c in the presence of a family history of unexplained juvenile SD. ^{1030,1033}	IIb	C
ICD implantation is not recommended in asymptomatic patients with an isolated ERP. ^{1034,1035,1040}	III	C
CA, cardiac arrest; ECG, electrocardiogram; ERP, early repolarization pattern; ERS, early repolarization syndrome; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; PVC, premature ventricular complexes; PVT, ventricular tachycardia; SCD, sudden cardiac death; SD, sudden death; VA, ventricular arrhythmia; VF, ventricular fibrillation. ^aClass of recommendation. ^bLevel of evidence. ^cERP high-risk features: J waves >2 mm, dynamic changes in J point and ST morphology. ^{1020,1041} ^dHigh-risk ERP: family history of unexplained SD <40 years, family history of ERS.		

© ESC 2022

7.2.6. Catecholaminergic polymorphic ventricular tachycardia

CPVT is a heritable disorder characterized by catecholamine-induced bidirectional VT and PVT in the absence of SHD or ischaemia. The disease has an estimated prevalence of 1 in 10 000 ([Figure A 35](#)). ¹³⁵




Figure 35

Management of patients with catecholaminergic polymorphic ventricular tachycardia.

CPVT, catecholaminergic polymorphic ventricular tachycardia; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; PVC, premature ventricular complex; VA, ventricular arrhythmia; VT, ventricular tachycardia. ^aGeneral recommendations: avoidance of competitive sports, avoidance of strenuous exercise, avoidance of stressful environments. ^bPreferred beta-blockers: nadolol, propranolol.

[Open in new tab](#)[Download slide](#)

There are two main genetic types: a dominant disorder due to mutations in the gene encoding for the cardiac ryanodine receptor (*RYR2*) and a recessive disorder caused by mutations in the cardiac calsequestrin gene (*CASQ2*).^{135,178} Mutations in *TRDN* and *CALM1-3* have been identified in patients with atypical forms of catecholaminergic VAs.¹⁰⁴² At the present time, however, it is unclear whether they are distinct arrhythmic entities.¹⁰⁴³ Patients with *KCNJ2* mutations causing Andersen-Tawil syndrome type 1 may sometimes exhibit bidirectional and PVT, but are distinguished by their syndromic associations.¹⁰⁴⁴

The clinical manifestations of CPVT usually occur in the first decade of life prompted by physical activity or emotional stress.¹⁰⁴⁵ Most CPVT patients have normal ECG and echocardiogram. Nonetheless, some patients present mild ECG abnormalities, such as sinus bradycardia and prominent U waves. Exercise stress test is the most important diagnostic test, because it elicits the distinguishing bidirectional or PVT that establish the diagnosis (Figure 36).¹³⁵ A diagnosis can also be made in the presence of a mutation in genes associated with CPVT. Epinephrine or isoproterenol challenge may be considered when exercise stress testing is not feasible.¹⁰⁴⁶




Figure 36

Exercise test of a patient with catecholaminergic polymorphic ventricular tachycardia.

PVC, premature ventricular complex.

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional, or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb	C
General recommendations		
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Therapeutic interventions		
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT. ^{1045,1048,1059}	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA. ^{1045,1047,1060}	I	C
Therapy with beta-blockers should be considered for genetically positive CPVT patients without phenotype. ^{1047,1050}	IIa	C
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated. ¹⁰⁵⁶	IIa	C
ICD implantation should be considered in patients with CPVT who experience arrhythmogenic syncope and/or documented bidirectional/PVT while on highest tolerated beta-blocker dose and on flecainide. ^{1047,1050}	IIa	C
Flecainide should be considered in patients with CPVT who experience recurrent syncope, polymorphic/bidirectional VT, or persistent exertional PVCs, while on beta-blockers at the highest tolerated dose. ^{1052,1053,1060}	IIa	C
PES is not recommended for stratification of SCD risk.	III	C

CA, cardiac arrest; CPVT, catecholaminergic ventricular tachycardia; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; PVT, polymorphic ventricular tachycardia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional, or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb	C
General recommendations		
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Therapeutic interventions		
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT. ^{1045,1048,1059}	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA. ^{1045,1047,1060}	I	C
Therapy with beta-blockers should be considered for genetically positive CPVT patients without phenotype. ^{1047,1050}	IIa	C
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated. ¹⁰⁵⁶	IIa	C
ICD implantation should be considered in patients with CPVT who experience arrhythmogenic syncope and/or documented bidirectional/PVT while on highest tolerated beta-blocker dose and on flecainide. ^{1047,1050}	IIa	C
Flecainide should be considered in patients with CPVT who experience recurrent syncope, polymorphic/bidirectional VT, or persistent exertional PVCs, while on beta-blockers at the highest tolerated dose. ^{1052,1053,1060}	IIa	C
PES is not recommended for stratification of SCD risk.	III	C

CA, cardiac arrest; CPVT, catecholaminergic ventricular tachycardia; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; PVT, polymorphic ventricular tachycardia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional, or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb	C
General recommendations		
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Therapeutic interventions		
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT. ^{1045,1048,1059}	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA. ^{1045,1047,1060}	I	C
Therapy with beta-blockers should be considered for genetically positive CPVT patients without phenotype. ^{1047,1050}	IIa	C
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated. ¹⁰⁵⁶	IIa	C
ICD implantation should be considered in patients with CPVT who experience arrhythmogenic syncope and/or documented bidirectional/PVT while on highest tolerated beta-blocker dose and on flecainide. ^{1047,1050}	IIa	C
Flecainide should be considered in patients with CPVT who experience recurrent syncope, polymorphic/bidirectional VT, or persistent exertional PVCs, while on beta-blockers at the highest tolerated dose. ^{1052,1053,1060}	IIa	C
PES is not recommended for stratification of SCD risk.	III	C

CA, cardiac arrest; CPVT, catecholaminergic ventricular tachycardia; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; PVT, polymorphic ventricular tachycardia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional, or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Epinephrine or isoproterenol challenge may be considered for the diagnosis of CPVT when an exercise test is not possible.	IIb	C
General recommendations		
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Therapeutic interventions		
Beta-blockers, ideally non-selective (nadolol or propranolol) are recommended in all patients with a clinical diagnosis of CPVT. ^{1045,1048,1059}	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA. ^{1045,1047,1060}	I	C
Therapy with beta-blockers should be considered for genetically positive CPVT patients without phenotype. ^{1047,1050}	IIa	C
LCSD should be considered in patients with diagnosis of CPVT when the combination of beta-blockers and flecainide at therapeutic dosage are either not effective, not tolerated, or contraindicated. ¹⁰⁵⁶	IIa	C
ICD implantation should be considered in patients with CPVT who experience arrhythmogenic syncope and/or documented bidirectional/PVT while on highest tolerated beta-blocker dose and on flecainide. ^{1047,1050}	IIa	C
Flecainide should be considered in patients with CPVT who experience recurrent syncope, polymorphic/bidirectional VT, or persistent exertional PVCs, while on beta-blockers at the highest tolerated dose. ^{1052,1053,1060}	IIa	C
PES is not recommended for stratification of SCD risk.	III	C

CA, cardiac arrest; CPVT, catecholaminergic ventricular tachycardia; ECG, electrocardiogram; ICD, implantable cardioverter defibrillator; LCSD, left cardiac sympathetic denervation; PVT, polymorphic ventricular tachycardia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

7.2.7. Short QT syndrome

Short QT syndrome (SQTS) is a rare genetic disorder characterised by a short QT interval, premature AF and VF in the context of a structurally normal heart.¹⁰⁶¹ It has been associated with gain of function mutations in *KCNH2*, *KCNQ1* and loss of function in *SLC4A*.^{1062,1063} This panel proposed two QTc cut-off threshold for diagnosis: A QTc ≤ 320 ms alone, or B a QTc ≤ 360 ms combined with a family history of SQTS, aborted CA in the absence of heart disease or pathogenic mutation^{1064,1065,1067} (Figure 37). The disease has high lethality in all age groups, including the first months of life.^{1063,1068,1069} The probability of a first CA by the age of 40 years is $>40\%$.¹⁰⁶⁸ While an ICD is used for secondary prevention,¹⁰⁶⁹ primary prevention remains contentious and is based upon prior symptoms and QTc interval.^{1063,1068,1069} Quinidine is currently the best supported AAD, but should be monitored for excessive QT prolongation, while isoprenaline may be considered in electrical storm.^{1070,1071} Drugs that shorten QT interval should be avoided, e.g. nicorandil.¹⁰⁷² Loop recorder implantation should be considered in children and young asymptomatic SQTS patients.




Figure 37
Short QT syndrome.

[Open in new tab](#)[Download slide](#)

Recommendation Table 46

[Open in new tab](#)

Recommendations for the management of patients with short QT syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that SQTS is diagnosed in the presence of a QTc ≤ 360 ms and one or more of the following: (a) a pathogenic mutation, (b) a family history of SQTS, (c) survival from a VT/VF episode in the absence of heart disease. ^{1061,1068}	I	C
Genetic testing is indicated in patients diagnosed with SQTS. ¹⁰⁶³	I	C

SQTS should be considered in the presence of a QTc $\leq$ 320 ms. ^{1064–1067,1073,1074}	IIa	C
SQTS should be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and arrhythmic syncope.	IIa	C
SQTS may be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and a family history of SD at age $<$ 40 years.	IIb	C
Risk stratification, SCD prevention and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA and/or (b) have documented spontaneous sustained VT. ¹⁰⁶³	I	C
ILR should be considered in young SQTS patients.	IIa	C
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa	C
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD. ^{1069–1071}	IIb	C
Isoproterenol may be considered in SQTS patients with an electrical storm. ¹⁰⁷⁵	IIb	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; SCD, sudden cardiac death; SD, sudden death; SQTS, short QT syndrome; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that SQTS is diagnosed in the presence of a QTc $\leq$ 360 ms and one or more of the following: (a) a pathogenic mutation, (b) a family history of SQTS, (c) survival from a VT/VF episode in the absence of heart disease. ^{1061,1068}	I	C
Genetic testing is indicated in patients diagnosed with SQTS. ¹⁰⁶³	I	C

SQTS should be considered in the presence of a QTc $\leq$ 320 ms. ^{1064–1067,1073,1074}	IIa	C
SQTS should be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and arrhythmic syncope.	IIa	C
SQTS may be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and a family history of SD at age $<$ 40 years.	IIb	C
Risk stratification, SCD prevention and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA and/or (b) have documented spontaneous sustained VT. ¹⁰⁶³	I	C
ILR should be considered in young SQTS patients.	IIa	C
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa	C
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD. ^{1069–1071}	IIb	C
Isoproterenol may be considered in SQTS patients with an electrical storm. ¹⁰⁷⁵	IIb	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; SCD, sudden cardiac death; SD, sudden death; SQTS, short QT syndrome; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 46

[Open in new tab](#)

Recommendations for the management of patients with short QT syndrome

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that SQTS is diagnosed in the presence of a QTc $\leq$ 360 ms and one or more of the following: (a) a pathogenic mutation, (b) a family history of SQTS, (c) survival from a VT/VF episode in the absence of heart disease. ^{1061,1068}	I	C
Genetic testing is indicated in patients diagnosed with SQTS. ¹⁰⁶³	I	C

SQTS should be considered in the presence of a QTc $\leq$ 320 ms. ^{1064–1067,1073,1074}	IIa	C
SQTS should be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and arrhythmic syncope.	IIa	C
SQTS may be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and a family history of SD at age $<$ 40 years.	IIb	C
Risk stratification, SCD prevention and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA and/or (b) have documented spontaneous sustained VT. ¹⁰⁶³	I	C
ILR should be considered in young SQTS patients.	IIa	C
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa	C
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD. ^{1069–1071}	IIb	C
Isoproterenol may be considered in SQTS patients with an electrical storm. ¹⁰⁷⁵	IIb	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; SCD, sudden cardiac death; SD, sudden death; SQTS, short QT syndrome; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
Diagnosis		
It is recommended that SQTS is diagnosed in the presence of a QTc $\leq$ 360 ms and one or more of the following: (a) a pathogenic mutation, (b) a family history of SQTS, (c) survival from a VT/VF episode in the absence of heart disease. ^{1061,1068}	I	C
Genetic testing is indicated in patients diagnosed with SQTS. ¹⁰⁶³	I	C

SQTS should be considered in the presence of a QTc $\leq$ 320 ms. ^{1064–1067,1073,1074}	IIa	C
SQTS should be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and arrhythmic syncope.	IIa	C
SQTS may be considered in the presence of a QTc $\geq$ 320 ms and $\leq$ 360 ms and a family history of SD at age $<$ 40 years.	IIb	C
Risk stratification, SCD prevention and treatment of VA		
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA and/or (b) have documented spontaneous sustained VT. ¹⁰⁶³	I	C
ILR should be considered in young SQTS patients.	IIa	C
ICD implantation should be considered in SQTS patients with arrhythmic syncope.	IIa	C
Quinidine may be considered in (a) SQTS patients who qualify for an ICD but present a contraindication to the ICD or refuse it, and (b) asymptomatic SQTS patients and a family history of SCD. ^{1069–1071}	IIb	C
Isoproterenol may be considered in SQTS patients with an electrical storm. ¹⁰⁷⁵	IIb	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C

CA, cardiac arrest; ICD, implantable cardioverter defibrillator; ILR, implantable loop recorder; SCD, sudden cardiac death; SD, sudden death; SQTS, short QT syndrome; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

8. Special aspects in selected populations

8.1. Pregnant patients and peri-partum cardiomyopathy

Pregnancy and the post-partum period contribute a significant risk in women with arrhythmic heart disease and is covered in the 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy. ^{1076,1077,1080} New-onset VT may present during pregnancy and risk of recurrent VT is higher in patients with previous VT and SHD. ¹⁰⁸¹ Peri-partum cardiomyopathy (PPCM) should be ruled out in the case of new-onset VT during the last 6 weeks of pregnancy or in the early post-partum period. ¹⁰⁸² A genetic contribution may be present in up to 20% of patients with PPCM (in particular titin-truncating variants) ¹⁰⁸³ and future studies should reveal if genetic testing plays a role in PPCM patients with a positive family history.

8.1.1. Electrocardioversion and implantable cardioverter defibrillator therapy in pregnancy

Cardioversion seems safe in all phases of pregnancy and case reports show neither compromise in fetal blood flow nor initiation of pre-term labour. ¹⁰⁸⁴ Foetal heart rate should be routinely controlled after cardioversion. ¹⁰⁸⁵ If an ICD is indicated, ICD implantation should be performed beyond 8 weeks of gestation with radiation protection by experienced operator teams. ¹⁰⁸⁶ Echocardiographic guidance or a 3D mapping system may be helpful in ICD implantation to avoid radiation. ^{1087,1088} In pregnant patients with existing ICD, routine ICD interrogation is recommended prior to delivery. In patients with PPCM, thresholds for early ICD implantations are higher than in other conditions because of a high rate of spontaneous recovery after delivery. ¹⁰⁸⁹ WCD have been suggested to temporarily prevent SCD during the first 38 weeks 6 months after diagnosis of PPCM with LVEF $\leq$ 35%, while awaiting recovery. ^{1090,1091}

8.1.2. Pharmacological treatment

Pharmacological treatment of arrhythmias and heart failure in pregnant women should follow treatment in non-pregnant patients with avoidance of drugs contraindicated in pregnancy, such as ACE inhibitors, ARB inhibitors/ARNI and renin inhibitors.^{1080,1082,1092,1093} The first trimester is associated with the greatest teratogenic risk. Start of pharmacological therapy is advised to begin at the latest possible point in the pregnancy and with the lowest effective dose. Drug exposure in the second and third trimesters may confer adverse effects on foetal growth and development as well as increasing the risk of pro-arrhythmia.

It is recommended to check drugs and safety data before initiation of a new drug during pregnancy, according to the 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy.¹⁰⁸⁰ From this list, AADs can be summarized as followed:

- **Well tolerated:** sotalol, oral verapamil
- **Use only if potential benefit outweighs potential risk:** bisoprolol, carvedilol, digoxin, diltiazem (possible teratogenic effects), disopyramide (uterine contractions), flecainide, lidocaine, metoprolol, nadolol, propranolol, verapamil IV, quinidine
- **Inadequate data:** Ivabradine, mexiletine, propafenone, vernakalant
- **Contraindicated:** amiodarone, atenolol, dronedarone.

In LQTS (LQT2 in particular), risk of cardiac events increases substantially in the post-partum period (up to one year after delivery).¹⁰⁹⁴ Therefore, it is important to continue beta-blocker therapy throughout pregnancy and post-partum.^{955,1094,1095} Continuation of beta-blocker treatment is recommended in LQTS and CPVT¹⁰⁹⁶ and should be considered in ARVC.¹⁰⁹⁷ No additional risk by pregnancy is known in women with BrS^{1100,1101} (ESC CardioMed chapter 53.6).¹¹⁰²

In patients with PPCM, use of bromocriptine as disease-specific therapy in addition to standard heart failure therapy has shown promising results in two clinical trials.^{1103,1104}

8.1.3. Catheter ablation

In planned pregnancies, symptomatic tachyarrhythmia should be treated by catheter ablation before pregnancy. If catheter ablation is indicated in a pregnant patient, one should avoid the procedure in the first trimester, and electroanatomical mapping-guided procedures should be preferred.^{1105,1106}

Recommendation Table 47

[Open in new tab](#)

Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy

Recommendations	Class ^a	Level ^b
Acute management of VA		
During pregnancy, electrical cardioversion is recommended for sustained VT. ¹⁰⁸⁴	I	C
For acute conversion of haemodynamically tolerated SMVT during pregnancy, a beta-blocker, sotalol, flecainide, procainamide, or overdrive ventricular pacing should be considered.	IIa	C
Long-term management of VA		
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection. ^{1087,1107}	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in women with LQTS or CPVT. ^{955,1094–1096}	I	C
Continuation of beta-blockers should be considered during pregnancy in women with ARVC. ^{1097–1099}	IIa	C
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa	C
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs. ¹¹⁰⁵	IIa	C

© ESC 2022

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; CPVT, catecholaminergic polymorphic ventricular tachycardia; ICD, implantable cardioverter defibrillator; LQTS, long QT syndrome; SMVT, sustained monomorphic VT; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Acute management of VA		
During pregnancy, electrical cardioversion is recommended for sustained VT. ¹⁰⁸⁴	I	C
For acute conversion of haemodynamically tolerated SMVT during pregnancy, a beta-blocker, sotalol, flecainide, procainamide, or overdrive ventricular pacing should be considered.	IIa	C
Long-term management of VA		
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection. ^{1087,1107}	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in women with LQTS or CPVT. ^{955,1094–1096}	I	C
Continuation of beta-blockers should be considered during pregnancy in women with ARVC. ^{1097–1099}	IIa	C
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa	C
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs. ¹¹⁰⁵	IIa	C

© ESC 2022

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; CPVT, catecholaminergic polymorphic ventricular tachycardia; ICD, implantable cardioverter defibrillator; LQTS, long QT syndrome; SMVT, sustained monomorphic VT; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendation Table 47

[Open in new tab](#)

Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy

Recommendations	Class ^a	Level ^b
Acute management of VA		
During pregnancy, electrical cardioversion is recommended for sustained VT. ¹⁰⁸⁴	I	C
For acute conversion of haemodynamically tolerated SMVT during pregnancy, a beta-blocker, sotalol, flecainide, procainamide, or overdrive ventricular pacing should be considered.	IIa	C
Long-term management of VA		
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection. ^{1087,1107}	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in women with LQTS or CPVT. ^{955,1094–1096}	I	C
Continuation of beta-blockers should be considered during pregnancy in women with ARVC. ^{1097–1099}	IIa	C
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa	C
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs. ¹¹⁰⁵	IIa	C

© ESC 2022

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; CPVT, catecholaminergic polymorphic ventricular tachycardia; ICD, implantable cardioverter defibrillator; LQTS, long QT syndrome; SMVT, sustained monomorphic VT; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Acute management of VA		
During pregnancy, electrical cardioversion is recommended for sustained VT. ¹⁰⁸⁴	I	C
For acute conversion of haemodynamically tolerated SMVT during pregnancy, a beta-blocker, sotalol, flecainide, procainamide, or overdrive ventricular pacing should be considered.	IIa	C
Long-term management of VA		
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection. ^{1087,1107}	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in women with LQTS or CPVT. ^{955,1094–1096}	I	C
Continuation of beta-blockers should be considered during pregnancy in women with ARVC. ^{1097–1099}	IIa	C
Oral metoprolol, propranolol, or verapamil should be considered for long-term management of idiopathic sustained VT during pregnancy.	IIa	C
Catheter ablation using non-fluoroscopic mapping systems should be considered, preferably after the first trimester, in women with highly symptomatic recurrent SMVT refractory or who are intolerant to AADs. ¹¹⁰⁵	IIa	C

AAD, anti-arrhythmic drug; ARVC, arrhythmogenic right ventricular cardiomyopathy; CPVT, catecholaminergic polymorphic ventricular tachycardia; ICD, implantable cardioverter defibrillator; LQTS, long QT syndrome; SMVT, sustained monomorphic VT; VA, ventricular arrhythmia; VT, ventricular tachycardia.

^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

8.2. Heart transplantation

Patients listed for HTX are exposed to an SCD risk and have a high incidence of VA. Data from large registries suggest a survival benefit from ICDs.^{1108a,1109,1110,1111} The only randomized study including patients listed for HTX was prematurely stopped because of low enrolment.¹¹¹² There are no data regarding the role of WCD in patients listed for HTX. The median expected time on the waiting list, 8a,1109,1110,1111 16 months, needs to be considered.^{1108,1110} However, this panel shares the opinion that a WCD may be an alternative for an ICD in selected patients waiting for HTX.^{371,1113}

In post-HTX patients, SCD is responsible for around 10% of deaths.^{1114,1115} Transplant rejection and allograft vasculopathy are associated with SCD^{1114,1116,1117}; therefore, ICD implantation may be appropriate in selected high-risk patients.¹¹¹⁸

Recommendation Table 48

[Open in new tab](#)

Recommendations for the prevention of sudden cardiac death before and after heart transplantation

Recommendations	Class ^a	Level ^b
Prior to heart transplant		
In patients awaiting heart transplantation, ICD implantation for primary prevention should be considered. ^{1108,1111,1112}	IIa	C
In patients awaiting heart transplantation, WCD may be considered. ^{1109,1113,1119}	IIb	C
Post heart transplant		
In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered. ^{1114,1116}	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Prior to heart transplant		
In patients awaiting heart transplantation, ICD implantation for primary prevention should be considered. ^{1108,1111,1112}	IIa	C
In patients awaiting heart transplantation, WCD may be considered. ^{1109,1113,1119}	IIb	C
Post heart transplant		
In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered. ^{1114,1116}	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendation Table 48

[Open in new tab](#)

Recommendations for the prevention of sudden cardiac death before and after heart transplantation

Recommendations	Class ^a	Level ^b
Prior to heart transplant		
In patients awaiting heart transplantation, ICD implantation for primary prevention should be considered. ^{1108,1111,1112}	IIa	C
In patients awaiting heart transplantation, WCD may be considered. ^{1109,1113,1119}	IIb	C
Post heart transplant		
In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered. ^{1114,1116}	IIb	C

© ESC 2022

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

Recommendations	Class ^a	Level ^b
Prior to heart transplant		
In patients awaiting heart transplantation, ICD implantation for primary prevention should be considered. ^{1108,1111,1112}	IIa	C
In patients awaiting heart transplantation, WCD may be considered. ^{1109,1113,1119}	IIb	C
Post heart transplant		
In selected transplanted patients with cardiac allograft vasculopathy or treated rejection, ICD implantation may be considered. ^{1114,1116}	IIb	C

ICD, implantable cardioverter defibrillator; WCD, wearable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

8.3. Sudden cardiac death in athletes

The incidence of SCD in athletes increases with age.^{4,1120} In apparently healthy athletes (>35 years), the estimated incidence of SCD ranges from 2 to 6.3 per 100 000 participant-years. In comparison, in young competitive athletes (≤35 years) the incidence of fatal events is significantly lower, 0.4±0.3 per 100 000 participant-years.^{46,47,1120} Women athletes are at low risk of SCD; on average, 1 in 14 SCD in athletes occurs in women.¹¹²¹

Pre-participation cardiovascular evaluation offers the potential to identify athletes at-risk for cardiovascular disease before onset of symptoms.¹¹²²⁻¹¹²⁶ The evaluation protocol needs to be adapted to the age of the athlete to account for age-specific cardiovascular disease.¹¹²⁵ Pre-participation evaluation, including medical history, physical examination, and ECG, appears effective in identifying cardiovascular disease in young athletes (≤35 years of age) by identification of relevant symptoms (e.g. exertional syncope) or ECG abnormalities consistent with inheritable cardiomyopathies or channelopathies.¹¹²⁷⁻¹¹³⁰ Although echocardiography can increase the sensitivity of screening for SHD, it is unfeasible as a routine test in mass screening. More cardiovascular diseases are identified by serial (annual) evaluations of adolescent athletes.^{1129,1131} The prevalence of false-positive results strongly depends on the criteria used to define an ECG as abnormal.^{1132,1133} Additional tests, such as echocardiography, 24-hour Holter monitoring, stress testing, and CMR, are requested for athletes who had positive findings at the initial evaluation. Athletes diagnosed with clinically relevant cardiovascular disease are managed according to available ESC Guidelines.^{4,1134-1136}

In middle-aged/senior athletes, the most common cause of SCD is CAD.^{1125,1137} Before engaging in a vigorous physical activity, asymptomatic middle-aged/senior athletes should be evaluated using risk score systems such as the ESC SCORE2.^{4,65}

Excellent rates of survival with favourable neurological outcome after CA has been reported in sports centres equipped with AED.^{1137,1138} This justifies the efforts for implementing emergency programs for SCD prevention, with distribution of AED in sports arenas and training of coaches and staff to perform CPR and defibrillation.¹¹³⁹

Recommendation Table 49

[Open in new tab](#)

Recommendations for risk stratification and prevention of sudden cardiac death in athletes

Recommendations	Class ^a	Level ^b
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended. ^{1123,1133,1135}	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED. ^{93,1137}	I	C
Pre-participation cardiovascular evaluation of competitive athletes should be considered. ^{46,1122,1123,1127}	IIa	C
It should be considered that cardiovascular evaluation of young (<35 years) competitive athletes includes history, physical examination, and 12-lead ECG. ^{1123,1126,1130,1140}	IIa	C
The cardiovascular risk of middle-aged and elderly individuals should be evaluated before engaging in strenuous sports through established scores such as the SCORE2 risk chart. ^{46,1141,1142}	IIa	C

© ESC 2022

AED, automated external defibrillator; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; ECG, electrocardiogram; SCD, sudden cardiac death.

^aClass of recommendation.

^bLevel of evidence.

Recommendations	Class ^a	Level ^b
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended. ^{1123,1133,1135}	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED. ^{93,1137}	I	C
Pre-participation cardiovascular evaluation of competitive athletes should be considered. ^{46,1122,1123,1127}	IIa	C
It should be considered that cardiovascular evaluation of young (<35 years) competitive athletes includes history, physical examination, and 12-lead ECG. ^{1123,1126,1130,1140}	IIa	C
The cardiovascular risk of middle-aged and elderly individuals should be evaluated before engaging in strenuous sports through established scores such as the SCORE2 risk chart. ^{46,1141,1142}	IIa	C

© ESC 2022

AED, automated external defibrillator; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; ECG, electrocardiogram; SCD, sudden cardiac death.

^aClass of recommendation.

^bLevel of evidence.

[Open in new tab](#)

Recommendations for risk stratification and prevention of sudden cardiac death in athletes

Recommendations	Class ^a	Level ^b
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended. ^{1123,1133,1135}	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED. ^{93,1137}	I	C
Pre-participation cardiovascular evaluation of competitive athletes should be considered. ^{46,1122,1123,1127}	IIa	C
It should be considered that cardiovascular evaluation of young (<35 years) competitive athletes includes history, physical examination, and 12-lead ECG. ^{1123,1126,1130,1140}	IIa	C
The cardiovascular risk of middle-aged and elderly individuals should be evaluated before engaging in strenuous sports through established scores such as the SCORE2 risk chart. ^{46,1141,1142}	IIa	C

AED, automated external defibrillator; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; ECG, electrocardiogram; SCD, sudden cardiac death.

^aClass of recommendation.

^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended. 1123,1133,1135	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED. 93,1137	I	C
Pre-participation cardiovascular evaluation of competitive athletes should be considered. 46,1122,1123,1127	IIa	C
It should be considered that cardiovascular evaluation of young (<35 years) competitive athletes includes history, physical examination, and 12-lead ECG. 1123,1126,1130,1140	IIa	C
The cardiovascular risk of middle-aged and elderly individuals should be evaluated before engaging in strenuous sports through established scores such as the SCORE2 risk chart. 46,1141,1142	IIa	C

AED, automated external defibrillator; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; ECG, electrocardiogram; SCD, sudden cardiac death.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

8.4. Wolff-Parkinson-White syndrome

In patients with Wolff-Parkinson-White (WPW) syndrome, the most common arrhythmia is AV re-entry tachycardia (AVRT; 80%), followed by AF (20%). SCD secondary to pre-excited AF resulting in VF is the most feared manifestation of WPW syndrome. The risk of CA/VF in untreated WPW patients has been estimated at 0.9–2.4 per 1000 person-year. [1143,1144](#) The management of WPW patients has recently been reviewed in the 2019 ESC Guidelines for the management of patients with SVT [302](#) and updated with a particular focus on athletes in 2020. [1145](#) In patients with ventricular pre-excitation and symptomatic AVRT, catheter ablation is recommended (class I). In asymptomatic patients with ventricular pre-excitation, both invasive (class IIa) and non-invasive (class IIb) assessment are options for risk stratification for SCD. While catheter ablation is recommended for asymptomatic accessory pathways with high-risk features (class I), clinical follow-up (class IIa), or catheter ablations (class IIb) are options based on informed patient choice. This decision should take into account the location of the accessory pathway, the local ablation experience and the fact that symptomatic arrhythmias will frequently develop during follow-up. [1146](#) For the paediatric population, SVT due to WPW can usually be managed pharmacologically, and accessory pathways often lose antegrade conduction in the first years of life. [1147](#) In children with asymptomatic accessory pathways, risk stratification is not recommended before the age of 8 years. [1148](#)

8.5. Prevention of sudden cardiac death in the elderly

Age is a strong risk factor for death. In several studies, advanced age was one of the factors associated with a reduced expected benefit from ICD treatment. In patients with ischaemic cardiomyopathy enrolled in MADIT-II trial, a risk scheme composed of 5 clinical factors including age >70 years was predictive of the absence of long-term benefit conferred by the ICD. [1149](#) In a recent analysis of ICD patients enrolled in four MADIT studies, age <75 years was a predictor of VT/VF whereas age ≥75 years was a predictor of non-arrhythmic mortality. [365](#) Consistently, in the recent randomized DANISH trial in patients with non-ischaemic heart failure, the association between the ICD and survival decreased linearly with increasing age, while an age cut-off at ≥70 years yielded the highest survival. [647](#) Contemporary non-randomized data corroborate these findings. In the EU-CERT-ICD prospective cohort study in patients with CAD or cardiomyopathies with indication for primary prevention ICD implantation, no ICD benefit was

observed in patients aged ≥75 years.³⁵⁷ In contrast, there was a significant survival benefit in patients aged <75 years. Further retrospective data are in line with this observation.¹¹⁵⁰

Biological age may vary in part according to comorbidities. Indeed, comorbidities significantly influence survival of ICD recipients and a high comorbidity index is associated with less survival gain in primary or secondary prevention ICD patients.^{1149,1151} Therefore, a simple age cut-off cannot guide the decision on ICD implantation adequately and the ICD indication in elderly patients should be based on a personalized assessment considering the overall condition and comorbidities.

Recommendation Table 50

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in the elderly

Recommendations	Class ^a	Level ^b
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered. ^{647,1150,1152}	IIb	B

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered. ^{647,1150,1152}	IIb	B

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendation Table 50

[Open in new tab](#)

Recommendations for implantable cardioverter defibrillator implantation in the elderly

Recommendations	Class ^a	Level ^b
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered. ^{647,1150,1152}	IIb	B

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

Recommendations	Class ^a	Level ^b
In elderly patients in whom a benefit from the defibrillator is not expected due to the patient's age and comorbidities, omission of ICD implantation for primary prevention may be considered. ^{647,1150,1152}	IIb	B

ICD, implantable cardioverter defibrillator.
^aClass of recommendation.
^bLevel of evidence.

© ESC 2022

9. Key messages

9.1. General aspects

- Increased availability of public access defibrillators and community training in basic life support are key elements to improve survival of out-of-hospital cardiac arrest victims.
- Risk calculators for SCD and VA implemented in clinical practice need to meet agreed high standards for the development, external validation, and reporting of prediction models.
- Patients with genetic cardiomyopathies and arrhythmia syndromes require genetic testing as a routine part of their care.
- Genetic testing and counselling require access to an expert multidisciplinary team.
- A systematic workup of cardiac arrest survivors requires a multimodal approach.
- A comprehensive autopsy is recommended in all cases of sudden death under 50 years and is desirable in all sudden death victims.
- Clinical and genetic evaluation of SADS decedents and their families leads to a diagnosis of genetic heart disease in a substantial proportion of families.
- An electrical storm refractory to drug treatment requires the availability of advanced catheter ablation techniques, mechanical circulatory support, and autonomic modulation.
- When considering ICD therapy benefit, competing risk factors for non-arrhythmic death and the patient's wishes and quality of life need to be taken into account.

9.2. Structural heart disease

- Catheter ablation is recommended in CAD patients with recurrent symptomatic SMVT despite chronic amiodarone therapy.
- Catheter ablation is the first-line treatment for PVC-induced cardiomyopathy.
- In HND/DCM patients, indication for primary prevention ICD implantation should not be restricted to an LVEF $\leq 35\%$. Additional risk factors (e.g. CMR and genetics) are important to consider.
- Patients with an *LMNA* mutation require specific risk stratification for SCD.
- ARVC patients have a high rate of appropriate ICD interventions which may not necessarily be classified as lifesaving.
- A validated risk calculator (HCM Risk-Kids score) is useful to assess the risk for SCD in HCM patients younger than 16 years.
- Myotonic dystrophy patients with palpitations suspected of arrhythmia, syncope or aborted sudden death need to be evaluated by invasive electrophysiological study.
- In patients with repaired tetralogy of Fallot and monomorphic VT, catheter ablation is the preferred treatment.

9.3. Primary electrical disease

- Nadolol or propranolol are the preferred beta-blockers in LQTS and CPVT patients.
- In asymptomatic LQTS patients the arrhythmic risk (1-2-3 Risk calculator) may be useful to calculate.
- A type 1 Brugada ECG pattern provoked by sodium channel blocker test in the absence of other findings does not diagnose the BrS.
- SCD risk stratification in asymptomatic BrS patients with a spontaneous type 1 pattern remains controversial.
- Routine catheter ablation is not recommended in asymptomatic BrS patients.
- The diagnosis of idiopathic VF requires exclusion of an underlying structural, channelopathic, or metabolic aetiology.
- ERP can be a benign finding and is distinct from ERS.
- Left cardiac denervation plays an important role in the management of CPVT and LQTS patients.

10. Gaps in evidence

10.1. General aspects

- Accurate screening tests to detect heart conditions associated with SCD in asymptomatic individuals in the general population are required.
- In patients with SHD, the optimal time interval between repeated non-invasive and invasive prognostic tests, in case of a negative test, is unknown.
- Improved assessment of genetic variants of uncertain significance and likely pathogenic variants is needed.
- The utility of polygenic risk scores in patients at risk of SCD requires investigation.

10.2. Structural heart disease ~~and~~ general

- The long-term safety and efficacy of S-ICDs is unknown.
- The role of primary preventive ICD therapy in patients with SHD and mildly reduced or preserved ejection fraction has not been studied systematically.
- The optimal techniques to undertake VT substrate mapping and ablation in SHD remain to be determined.
- The role of ICD implantation in end-stage heart failure patients supported by current-generation, continuous-flow LVADs is unclear.

10.3. Idiopathic premature ventricular complexes/ventricular tachycardia

- The beneficial role of catheter ablation or anti-arrhythmic drug treatment in patients with asymptomatic, frequent PVCs and preserved cardiac function needs to be determined.

10.4. Coronary artery disease

- It is unknown which patients with chronic CAD and severely impaired LVEF are at low risk for SCD.
- The role of LGE-CMR for risk stratification for SCD in chronic CAD is unclear.
- RCTs are needed to determine the role of ICDs after successful VT ablation in chronic CAD with mildly reduced or preserved LVEF.

10.5. Cardiomyopathies

- It is unknown if PVC-induced cardiomyopathy is a diagnosis on its own or if an underlying predisposition is needed.
- The predictive value of LGE-CMR findings (e.g. pattern and amount of LGE) for *individual* risk stratification for SCD in patients with cardiac sarcoidosis, HCM, and DCM/HNDC is unclear.
- Studies are needed to determine the role of PES in patients with cardiac sarcoidosis and DCM/HNDC who have a mildly reduced or preserved cardiac function and LGE on CMR.

- Prospective data on the association between intensity and duration of exercise and manifestation and severity of the phenotype in healthy ARVC mutation carriers are lacking.
- The beneficial role of ICDs after successful ablation in ARVC patients who present with haemodynamically tolerated VT needs to be studied.
- Data on clinical outcome, predictors for arrhythmic events, and indication for treatment, including ICD therapy, in patients with biventricular and left-dominant arrhythmogenic cardiomyopathy are required.

10.6. Valvular heart disease

- There is a lack of knowledge to identify patients with MVP at risk for VA and SCD.

10.7. Congenital heart disease

- There is a lack of knowledge regarding the absolute risk for VA and SCD in CHD that have undergone repair with contemporary surgical approaches.

10.8. Primary electrical disease

- Robust evidence is required to support the prophylactic use of the ICD on top of medical therapy with beta-blockers and gene-specific therapy in LQTS patients.
- More data are required to determine the role of LCSD and ICD in high-risk LQTS patients who do not tolerate medical therapy.
- Improved diagnostic and risk stratification tools for asymptomatic Brugada patients and suspected early repolarization syndrome are needed.
- The role of endo-epicardial mapping to identify localized structural alterations potentially related to IVF and targeted catheter ablation needs to be further studied.
- Long-term data are required on the efficacy of the ICD vs. no ICD in CPVT survivors of cardiac arrest.
- It is poorly understood why women are at low risk of sport-related SCD.

11. What to do and what not to do messages from the Guidelines

[Open in new tab](#)

Recommendations	Class ^a	Level ^b
Recommendations for public basic life support and access to automated external defibrillator		
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ²	I	B
Prompt CPR by bystanders is recommended at OHCA.	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I	B
Recommendations for genetic testing		
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD.	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework.	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team.	I	C
It is recommended that Class III variants (of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C
Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmia		
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
Recommendations for evaluation of sudden cardiac arrest survivors		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team.	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing.	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors.	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precardial lead ECG) as well as continuous cardiac monitoring are recommended.	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause.	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause.	I	B
Recommendations for evaluation of sudden death victims		
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority.	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files.	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those under 50 years of age.	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained.	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death.	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended.	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50 years of age) and/or the circumstances and/or family history support a primary electrical disease.	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedents		
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended.	III	B
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years or if there are other circumstantial data or family history to suggest heritable disease.	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation.	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard- and high-precordial lead ECG, echocardiography, and exercise testing.	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood.	I	C
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes.	III	C
Recommendations for treatment of reversible conditions		
Withdrawal of offending agents is recommended whenever drug-induced VA are suspected.	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ⁹ is recommended in patients with VA.	I	C
Recommendations for the acute treatment of sustained ventricular tachycardia and electrical storm		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT.	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C
In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended.	I	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism.	III	B
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated.	I	B

Intravenous magnesium with supplementation of potassium is recommended in patients with TdP.	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs.	I	B
Recommendations for treatment with heart failure medication		
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I	A
Recommendations for implantable cardioverter defibrillation implantation (general aspects)		
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C
Recommendations for secondary prevention of sudden cardiac death		
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes.	I	A
Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillator		
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator.	I	C
Recommendations for optimization of device programming		
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality.	I	A
In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing.	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals).	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients.	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones.	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m.	I	B
It is recommended to activate lead failure alerts.	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks.	I	B
Programming of burst ATP as a first attempt is recommended over ramp ATP.	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone	I	B
Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy		
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies.	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming.	I	C
Recommendation for psychosocial management after implantable cardioverter defibrillator implantation		
Assessment of psychological status and treatment of distress is recommended in ICD patients.	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression.	I	C
Recommendations for prevention of implantable cardioverter defibrillator complications		
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications.	I	A
Recommendations for end-of-life issues in implantable cardioverter defibrillator carriers		
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration.	I	C
Recommendations for treatment of ventricular arrhythmia in acute coronary syndrome and vasospasm		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated.	I	B
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS.	III	B
Recommendations for risk stratification and treatment of ventricular arrhythmia early after myocardial infarction		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI.	I	B
In patients with pre-discharge LVEF $\leq 40\%$, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in chronic coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT.	I	A
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended.	III	A
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI.	I	A
In patients with CAD and recurrent, symptomatic SMVT or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I	B
Recommendations for sudden cardiac death prevention in patients with coronary anomalies		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia.	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs or angina when other causes have been excluded.	I	C
Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia		
Regular assessment of ventricular function of patients with PVC burden $>10\%$ and normal ventricular function is indicated.	I	C
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.*	I	B
Beta-blockers or non-dihydropyridine CCB are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I	C
Catheter ablation is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated.	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs.	III	C
Recommendations for the management of patients with premature ventricular complex-induced or aggravated cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	I	C

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy		
Genetic testing (including at least LMNA, PLN, RBM20, and RPLNC genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I	B
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a LMNA mutation.	III	C
ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/VF or experience haemodynamically not-tolerated SMVT.	I	B
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: - the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or - there is family history of DCM/HNDCM, or premature unexpected SD.	I	C
Recommendations for diagnostic, risk stratification, sudden cardiac death prevention and treatment of ventricular arrhythmia in arrhythmogenic right ventricular cardiomyopathy		
In patients with suspected ARVC, CMR is recommended.	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC.	I	B
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF.	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In a first-degree relative of a patient with ARVC, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in hypertrophic cardiomyopathy		
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I	B
Genetic counselling and testing are recommended in HCM patients.	I	B
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals or when there is a change in clinical status.	I	C
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF.	I	B
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in neuromuscular diseases		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease.	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders.	I	C
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA, VT, or surviving a CA.	I	C
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III	C
In symptomatic patients with BBR-VT, catheter ablation is recommended.	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in myocarditis		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VA are referred to a specialized centre.	I	C
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in cardiac sarcoidosis		
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF ≤35%.	I	B
ICD implantation is recommended in patients with cardiac sarcoidosis who (a) have documented sustained VT, or (2) aborted CA.	I	B
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in valvular heart disease		
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention.	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in congenital heart disease		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF ≤35% despite ≥3 months of OMT, ICD implantation is indicated.	I	C
In patients with CHD presenting with sustained VAs, evaluation for residual or new anatomical abnormalities is recommended.	I	B
In patients with CHD with not-tolerated VT/aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes.	I	C
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended.	I	C
Recommendations for management of patients with idiopathic ventricular fibrillation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I	B
ICD implantation is recommended in idiopathic VF.	I	B
Recommendations for management of patients with long QT syndrome		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3.	I	C
In patients with clinically diagnosed long QT syndrome, genetic testing, and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III	C
The following is recommended in LQTS: - Avoid QT-prolonging drugs.¹ - Avoid and correct electrolyte abnormalities. - Avoid genotype-specific triggers for arrhythmias.	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I	B
Mexiletine is indicated in LQTS patients with a prolonged QT interval.	I	C
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA.	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ⁴ while receiving beta-blockers and genotype-specific therapies.	I	C

LCSD is indicated in patients with symptomatic ⁴ LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific therapies with an ICD and experiences multiple shocks or syncope due to VA.	I	C
Invasive electrophysiologic study is not recommended in LQTS.	III	C
Recommendations for management of patients with Andersen-Tawil syndrome		
Genetic testing is recommended in patients with suspected Andersen-Tawil syndrome.	I	C
ICD implantation is recommended in patients with Andersen-Tawil syndrome after aborted CA not tolerated sustained VT.	I	C
Recommendations for management of patients with Brugada syndrome		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern.	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever.	I	C
Genetic testing for SCN5A gene is recommended for probands with BrS.	I	C
Sodium channel blocker test is not recommended in patients with a prior type 1 Brugada pattern.	III	C
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA, and/or (b) Have documented spontaneous sustained VT.	I	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
Recommendations for management of patients with early repolarization pattern/syndrome		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III	C
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I	B
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III	C
Recommendations for management of patients with catecholaminergic polymorphic ventricular tachycardia		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Beta-blockers, ideally non-selective (nadolol or propranolol), are recommended in all patients with a clinical diagnosis of CPVT.	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA.	I	C
PES is not recommended for stratification of SCD risk.	III	C
Recommendations for management of patients with short QT syndrome		
It is recommended that SQTS is diagnosed in the presence of a QTc ≤ 360 ms and one or more of the following: (a) a pathogenic mutation; (b) a family history of SQTS; (c) survival from a VT/VF episode in the absence of heart disease.	I	C
Genetic testing is indicated in patients diagnosed with SQTS.	I	C
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA, and/or (b) have documented spontaneous sustained VT.	I	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C
Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy		
During pregnancy, electrical cardioversion is recommended for sustained VT.	I	C
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection.	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in patients with LQTS or CPVT.	I	C
Recommendations for risk stratification and prevention of sudden cardiac death in athletes		
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended.	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED.	I	C

AAD, anti-arrhythmic drug; ACE-I, angiotensin-converting-enzyme inhibitor; AED, automated external defibrillator; ARB, angiotensin receptor blocker; ARNI, angiotensin receptor neprilysin inhibitor; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CIED, cardiac implantable electronic device; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HNDCH, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; CAD, coronary artery disease; IAR, implantable loop recorder; LCSD, left cardiac sympathetic denervation; LGE, late gadolinium enhancement; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OPT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SADS, sudden arrhythmic death syndrome; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SQT, short QT syndrome; STEMI, ST-elevation myocardial infarction; SVT, supraventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

⁴Class of recommendation.
⁵Level of evidence.
⁶Shopping malls, stadiums, public transport stations, casinos.
⁷List not exhaustive.
⁸LQTS C for VT/PVCs from left fascicles.
<http://www.crediblemed.org>
⁹Arrhythmic syncope or haemodynamically non-tolerated VA

© ESC 2022

Recommendations	Class ^a	Level ^b
Recommendations for public basic life support and access to automated external defibrillator		
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ²	I	B
Prompt CPR by bystanders is recommended at OHCA.	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I	B
Recommendations for genetic testing		
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD.	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework.	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team.	I	C
It is recommended that Class III variants (of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C
Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmia		
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
Recommendations for evaluation of sudden cardiac arrest survivors		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team.	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing.	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors.	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG) as well as continuous cardiac monitoring are recommended.	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause.	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause.	I	B
Recommendations for evaluation of sudden death victims		
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority.	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files.	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those under 50 years of age.	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained.	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death.	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended.	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50 years of age) and/or the circumstances and/or family history support a primary electrical disease.	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedents		
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended.	III	B
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years or if there are other circumstantial data or family history to suggest heritable disease.	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation.	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard- and high-precordial lead ECG, echocardiography, and exercise testing.	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood.	I	C
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes.	III	C
Recommendations for treatment of reversible conditions		
Withdrawal of offending agents is recommended whenever drug-induced VA are suspected.	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ⁹ is recommended in patients with VA.	I	C
Recommendations for the acute treatment of sustained ventricular tachycardia and electrical storm		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT.	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C
In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended.	I	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism.	III	B
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated.	I	B

Intravenous magnesium with supplementation of potassium is recommended in patients with TdP.	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs.	I	B
Recommendations for treatment with heart failure medication		
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I	A
Recommendations for implantable cardioverter defibrillation implantation (general aspects)		
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C
Recommendations for secondary prevention of sudden cardiac death		
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes.	I	A
Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillator		
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator.	I	C
Recommendations for optimization of device programming		
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality.	I	A
In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing.	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals).	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients.	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones.	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m.	I	B
It is recommended to activate lead failure alerts.	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks.	I	B
Programming of burst ATP as a first attempt is recommended over ramp ATP.	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone	I	B
Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy		
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies.	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming.	I	C
Recommendation for psychosocial management after implantable cardioverter defibrillator implantation		
Assessment of psychological status and treatment of distress is recommended in ICD patients.	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression.	I	C
Recommendations for prevention of implantable cardioverter defibrillator complications		
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications.	I	A
Recommendations for end-of-life issues in implantable cardioverter defibrillator carriers		
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration.	I	C
Recommendations for treatment of ventricular arrhythmia in acute coronary syndrome and vasospasm		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated.	I	B
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS.	III	B
Recommendations for risk stratification and treatment of ventricular arrhythmia early after myocardial infarction		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI.	I	B
In patients with pre-discharge LVEF $\leq 40\%$, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in chronic coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT.	I	A
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended.	III	A
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI.	I	A
In patients with CAD and recurrent, symptomatic SMVT or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I	B
Recommendations for sudden cardiac death prevention in patients with coronary anomalies		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia.	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs or angina when other causes have been excluded.	I	C
Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia		
Regular assessment of ventricular function of patients with PVC burden $>10\%$ and normal ventricular function is indicated.	I	C
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.*	I	B
Beta-blockers or non-dihydropyridine CCB are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I	C
Catheter ablation is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated.	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs.	III	C
Recommendations for the management of patients with premature ventricular complex-induced or aggravated cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	I	C

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy		
Genetic testing (including at least LMNA, PLN, RBM20, and RPLNC genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I	B
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a LMNA mutation.	III	C
ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/Vf or experience haemodynamically not-tolerated SMVT.	I	B
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: • the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or • there is family history of DCM/HNDCM, or premature unexpected SD.	I	C
Recommendations for diagnostic, risk stratification, sudden cardiac death prevention and treatment of ventricular arrhythmia in arrhythmogenic right ventricular cardiomyopathy		
In patients with suspected ARVC, CMR is recommended.	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC.	I	B
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF.	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In a first-degree relative of a patient with ARVC, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in hypertrophic cardiomyopathy		
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I	B
Genetic counselling and testing are recommended in HCM patients.	I	B
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals or when there is a change in clinical status.	I	C
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF.	I	B
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in neuromuscular diseases		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease.	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders.	I	C
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA, VT, or surviving a CA.	I	C
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III	C
In symptomatic patients with BBR-VT, catheter ablation is recommended.	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in myocarditis		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VA are referred to a specialized centre.	I	C
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in cardiac sarcoidosis		
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF ≤35%.	I	B
ICD implantation is recommended in patients with cardiac sarcoidosis who (a) have documented sustained VT, or (2) aborted CA.	I	B
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in valvular heart disease		
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention.	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in congenital heart disease		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF ≤35% despite ≥3 months of OMT, ICD implantation is indicated.	I	C
In patients with CHD presenting with sustained VAs, evaluation for residual or new anatomical abnormalities is recommended.	I	B
In patients with CHD with not-tolerated VT/aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes.	I	C
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended.	I	C
Recommendations for management of patients with idiopathic ventricular fibrillation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I	B
ICD implantation is recommended in idiopathic VF.	I	B
Recommendations for management of patients with long QT syndrome		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3.	I	C
In patients with clinically diagnosed long QT syndrome, genetic testing, and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III	C
The following is recommended in LQTS: • Avoid QT-prolonging drugs.¹ • Avoid and correct electrolyte abnormalities. • Avoid genotype-specific triggers for arrhythmias.	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I	B
Mexiletine is indicated in LQTS patients with a prolonged QT interval.	I	C
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA.	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ⁴ while receiving beta-blockers and genotype-specific therapies.	I	C

LCSD is indicated in patients with symptomatic ⁴ LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific therapies with an ICD and experiences multiple shocks or syncope due to VA.	I	C
Invasive electrophysiologic study is not recommended in LQTS.	III	C
Recommendations for management of patients with Andersen-Tawil syndrome		
Genetic testing is recommended in patients with suspected Andersen-Tawil syndrome.	I	C
ICD implantation is recommended in patients with Andersen-Tawil syndrome after aborted CA not tolerated sustained VT.	I	C
Recommendations for management of patients with Brugada syndrome		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern.	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever.	I	C
Genetic testing for SCN5A gene is recommended for probands with BrS.	I	C
Sodium channel blocker test is not recommended in patients with a prior type 1 Brugada pattern.	III	C
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA, and/or (b) Have documented spontaneous sustained VT.	I	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
Recommendations for management of patients with early repolarization pattern/syndrome		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III	C
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I	B
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III	C
Recommendations for management of patients with catecholaminergic polymorphic ventricular tachycardia		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Beta-blockers, ideally non-selective (nadolol or propranolol), are recommended in all patients with a clinical diagnosis of CPVT.	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA.	I	C
PES is not recommended for stratification of SCD risk.	III	C
Recommendations for management of patients with short QT syndrome		
It is recommended that SQTS is diagnosed in the presence of a QTc ≤ 360 ms and one or more of the following: (a) a pathogenic mutation; (b) a family history of SQTS; (c) survival from a VT/VF episode in the absence of heart disease.	I	C
Genetic testing is indicated in patients diagnosed with SQTS.	I	C
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA, and/or (b) have documented spontaneous sustained VT.	I	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C
Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy		
During pregnancy, electrical cardioversion is recommended for sustained VT.	I	C
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection.	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in patients with LQTS or CPVT.	I	C
Recommendations for risk stratification and prevention of sudden cardiac death in athletes		
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended.	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED.	I	C

AAD, anti-arrhythmic drug; ACE-I, angiotensin-converting-enzyme inhibitor; AED, automated external defibrillator; ARB, angiotensin receptor blocker; ARNI, angiotensin receptor neprilysin inhibitor; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CIED, cardiac implantable electronic device; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HNDCH, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; CAD, coronary artery disease; IAR, implantable loop recorder; LCSD, left cardiac sympathetic denervation; LGE, late gadolinium enhancement; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OPT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SADS, sudden arrhythmic death syndrome; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SQT, short QT syndrome; STEMI, ST-elevation myocardial infarction; SVT, supraventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

⁴Class of recommendation.
⁵Level of evidence.
⁶Shopping malls, stadiums, public transport stations, casinos.
⁷List not exhaustive.
⁸LQTS C for VT/PVCs from left fascicles.
<http://www.crediblemed.org>
⁹Arrhythmic syncope or haemodynamically non-tolerated VA

[Open in new tab](#)

Recommendations	Class ^a	Level ^b
Recommendations for public basic life support and access to automated external defibrillator		
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ²	I	B
Prompt CPR by bystanders is recommended at OHCA.	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I	B
Recommendations for genetic testing		
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD.	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework.	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team.	I	C
It is recommended that Class III variants (of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C
Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmia		
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
Recommendations for evaluation of sudden cardiac arrest survivors		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team.	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing.	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors.	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG) as well as continuous cardiac monitoring are recommended.	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause.	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause.	I	B
Recommendations for evaluation of sudden death victims		
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority.	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files.	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those under 50 years of age.	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained.	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death.	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended.	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50 years of age) and/or the circumstances and/or family history support a primary electrical disease.	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedents		
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended.	III	B
Familial evaluation of SADS decedents is recommended:		
• for first-degree relatives	I	B
• for relatives who must carry a mutation based on analysis of the family history	I	B
• for relatives with suspicious symptoms	I	B
• when the decedent's age is <50 years or if there are other circumstantial data or family history to suggest heritable disease.	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation.	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard- and high-precordial lead ECG, echocardiography, and exercise testing.	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood.	I	C
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes.	III	C
Recommendations for treatment of reversible conditions		
Withdrawal of offending agents is recommended whenever drug-induced VA are suspected.	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ⁹ is recommended in patients with VA.	I	C
Recommendations for the acute treatment of sustained ventricular tachycardia and electrical storm		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT.	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C
In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended.	I	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism.	III	B
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated.	I	B

Intravenous magnesium with supplementation of potassium is recommended in patients with TdP.	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs.	I	B
Recommendations for treatment with heart failure medication		
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I	A
Recommendations for implantable cardioverter defibrillation implantation (general aspects)		
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C
Recommendations for secondary prevention of sudden cardiac death		
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes.	I	A
Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillator		
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator.	I	C
Recommendations for optimization of device programming		
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality.	I	A
In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing.	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals).	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients.	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones.	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m.	I	B
It is recommended to activate lead failure alerts.	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks.	I	B
Programming of burst ATP as a first attempt is recommended over ramp ATP.	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone	I	B
Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy		
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies.	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming.	I	C
Recommendation for psychosocial management after implantable cardioverter defibrillator implantation		
Assessment of psychological status and treatment of distress is recommended in ICD patients.	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression.	I	C
Recommendations for prevention of implantable cardioverter defibrillator complications		
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications.	I	A
Recommendations for end-of-life issues in implantable cardioverter defibrillator carriers		
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration.	I	C
Recommendations for treatment of ventricular arrhythmia in acute coronary syndrome and vasospasm		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated.	I	B
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS.	III	B
Recommendations for risk stratification and treatment of ventricular arrhythmia early after myocardial infarction		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI.	I	B
In patients with pre-discharge LVEF $\leq 40\%$, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in chronic coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT.	I	A
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended.	III	A
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI.	I	A
In patients with CAD and recurrent, symptomatic SMVT or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I	B
Recommendations for sudden cardiac death prevention in patients with coronary anomalies		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia.	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs or angina when other causes have been excluded.	I	C
Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia		
Regular assessment of ventricular function of patients with PVC burden $>10\%$ and normal ventricular function is indicated.	I	C
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.*	I	B
Beta-blockers or non-dihydropyridine CCB are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I	C
Catheter ablation is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated.	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs.	III	C
Recommendations for the management of patients with premature ventricular complex-induced or aggravated cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	I	C

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy		
Genetic testing (including at least LMNA, PLN, RBM20, and RPLNC genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I	B
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a LMNA mutation.	III	C
ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/Vf or experience haemodynamically not-tolerated SMVT.	I	B
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: - the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or - there is family history of DCM/HNDCM, or premature unexpected SD.	I	C
Recommendations for diagnostic, risk stratification, sudden cardiac death prevention and treatment of ventricular arrhythmia in arrhythmogenic right ventricular cardiomyopathy		
In patients with suspected ARVC, CMR is recommended.	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC.	I	B
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF.	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In a first-degree relative of a patient with ARVC, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in hypertrophic cardiomyopathy		
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I	B
Genetic counselling and testing are recommended in HCM patients.	I	B
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals or when there is a change in clinical status.	I	C
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF.	I	B
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in neuromuscular diseases		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease.	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders.	I	C
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA, VT, or surviving a CA.	I	C
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III	C
In symptomatic patients with BBR-VT, catheter ablation is recommended.	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in myocarditis		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VA are referred to a specialized centre.	I	C
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in cardiac sarcoidosis		
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF ≤35%.	I	B
ICD implantation is recommended in patients with cardiac sarcoidosis who (a) have documented sustained VT, or (2) aborted CA.	I	B
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in valvular heart disease		
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention.	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in congenital heart disease		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF ≤35% despite ≥3 months of OMT, ICD implantation is indicated.	I	C
In patients with CHD presenting with sustained VAs, evaluation for residual or new anatomical abnormalities is recommended.	I	B
In patients with CHD with not-tolerated VT/aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes.	I	C
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended.	I	C
Recommendations for management of patients with idiopathic ventricular fibrillation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I	B
ICD implantation is recommended in idiopathic VF.	I	B
Recommendations for management of patients with long QT syndrome		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3.	I	C
In patients with clinically diagnosed long QT syndrome, genetic testing, and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III	C
The following is recommended in LQTS: - Avoid QT-prolonging drugs.¹ - Avoid and correct electrolyte abnormalities. - Avoid genotype-specific triggers for arrhythmias.	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I	B
Mexiletine is indicated in LQTS patients with a prolonged QT interval.	I	C
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA.	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ⁴ while receiving beta-blockers and genotype-specific therapies.	I	C

LCSD is indicated in patients with symptomatic ⁴ LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific therapies with an ICD and experiences multiple shocks or syncope due to VA.	I	C
Invasive electrophysiologic study is not recommended in LQTS.	III	C
Recommendations for management of patients with Andersen-Tawil syndrome		
Genetic testing is recommended in patients with suspected Andersen-Tawil syndrome.	I	C
ICD implantation is recommended in patients with Andersen-Tawil syndrome after aborted CA not tolerated sustained VT.	I	C
Recommendations for management of patients with Brugada syndrome		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern.	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever.	I	C
Genetic testing for SCN5A gene is recommended for probands with BrS.	I	C
Sodium channel blocker test is not recommended in patients with a prior type 1 Brugada pattern.	III	C
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA, and/or (b) Have documented spontaneous sustained VT.	I	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
Recommendations for management of patients with early repolarization pattern/syndrome		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III	C
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I	B
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III	C
Recommendations for management of patients with catecholaminergic polymorphic ventricular tachycardia		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Beta-blockers, ideally non-selective (nadolol or propranolol), are recommended in all patients with a clinical diagnosis of CPVT.	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA.	I	C
PES is not recommended for stratification of SCD risk.	III	C
Recommendations for management of patients with short QT syndrome		
It is recommended that SQTS is diagnosed in the presence of a QTc ≤ 360 ms and one or more of the following: (a) a pathogenic mutation; (b) a family history of SQTS; (c) survival from a VT/VF episode in the absence of heart disease.	I	C
Genetic testing is indicated in patients diagnosed with SQTS.	I	C
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA, and/or (b) have documented spontaneous sustained VT.	I	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C
Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy		
During pregnancy, electrical cardioversion is recommended for sustained VT.	I	C
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection.	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in patients with LQTS or CPVT.	I	C
Recommendations for risk stratification and prevention of sudden cardiac death in athletes		
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended.	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED.	I	C

AAD, anti-arrhythmic drug; ACE-I, angiotensin-converting-enzyme inhibitor; AED, automated external defibrillator; ARB, angiotensin receptor blocker; ARNI, angiotensin receptor neprilysin inhibitor; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CIED, cardiac implantable electronic device; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HNDCH, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; CAD, coronary artery disease; IAR, implantable loop recorder; LCSD, left cardiac sympathetic denervation; LGE, late gadolinium enhancement; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; OHCA, out-of-hospital cardiac arrest; OPT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SADS, sudden arrhythmic death syndrome; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SQT, short QT syndrome; STEMI, ST-elevation myocardial infarction; SVT, supraventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

⁴Class of recommendation.
⁵Level of evidence.
⁶Shopping malls, stadiums, public transport stations, casinos.
⁷List not exhaustive.
⁸LQTS C for VT/PVCs from left fascicles.
<http://www.crediblemed.org>
⁹Arrhythmic syncope or haemodynamically non-tolerated VA

© ESC 2022

Recommendations	Class ^a	Level ^b
Recommendations for public basic life support and access to automated external defibrillator		
It is recommended that public access defibrillation be available at sites where cardiac arrest is more likely to occur. ²	I	B
Prompt CPR by bystanders is recommended at OHCA.	I	B
It is recommended to promote community training in basic life support to increase bystander CPR rate and AED use.	I	B
Recommendations for genetic testing		
Genetic testing is recommended when a condition is diagnosed in a living or deceased individual with a likely genetic basis and a risk of VA and SCD.	I	B
When a putative causative variant is first identified, evaluation for pathogenicity is recommended using an internationally accepted framework.	I	C
When a Class IV or Class V variant has been identified in a living or deceased individual with a condition that carries a risk of VA and SCD, genetic testing of first-degree and symptomatic relatives and obligate carriers is recommended.	I	C
It is recommended that genetic testing and counselling on its potential consequences should be undertaken by an expert multidisciplinary team.	I	C
It is recommended that Class III variants (of uncertain significance) and Class IV variants should be evaluated for segregation in families where possible, and the variant re-evaluated periodically.	I	C
It is not recommended to undertake genetic testing in index patients with insufficient evidence of a genetic disease.	III	C
Recommendations for evaluation of patients presenting with newly documented ventricular arrhythmia		
In patients with newly documented VA (frequent PVCs, NSVT, SMVT), a baseline 12-lead ECG, recording of the VA on 12-lead ECG, whenever possible, and an echocardiogram are recommended as first-line evaluation.	I	C
Recommendations for evaluation of sudden cardiac arrest survivors		
The investigation of a SCA survivor without obvious extra-cardiac cause is recommended to be overseen by a multidisciplinary team.	I	B
In electrically unstable patients after SCA, with suspicion of ongoing myocardial ischaemia, a coronary angiogram is indicated.	I	C
In SCA survivors, collection of blood samples at presentation is recommended for potential toxicology and genetic testing.	I	B
Retrieval of recordings from CIEDs and wearable monitors is recommended for all SCA survivors.	I	B
In SCA survivors, repeated 12-lead ECGs during stable rhythm (including high precordial lead ECG) as well as continuous cardiac monitoring are recommended.	I	B
Echocardiography is recommended for evaluation of cardiac structure and function in all SCA survivors.	I	C
Coronary imaging and CMR with LGE are recommended for evaluation of cardiac structure and function in all SCA survivors without a clear underlying cause.	I	B
Sodium channel blocker test and exercise testing is recommended in SCA survivors without a clear underlying cause.	I	B
Recommendations for evaluation of sudden death victims		
Investigation of unexpected SD, especially in case of suspicion of inherited disease, should be made a public health priority.	I	B
In cases of SD, it is recommended to collect a detailed description of circumstances of death, symptoms prior to death, the family history, and to review prior medical files.	I	B
A comprehensive autopsy is recommended, ideally, in all cases of unexpected SD, and always in those under 50 years of age.	I	B
In cases of SCD, it is recommended to retain samples suitable for DNA extraction and consult a cardiac pathologist when an inherited cause is suspected or the cause of death unexplained.	I	B
Toxicology screens are recommended in SD cases with uncertain cause of death.	I	B
For SCD where the cause is known or suspected to be heritable, genetic testing targeted to the cause is recommended.	I	B
Following SADS, post-mortem genetic testing targeted to primary electrical disease is recommended when the decedent is young (<50 years of age) and/or the circumstances and/or family history support a primary electrical disease.	I	B
When an autopsy diagnoses possible heritable cardiac disease, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
In non-autopsied cases of SD where inherited cardiac disease is suspected, it is recommended to refer first-degree relatives for cardiac assessment in a specialized clinic.	I	B
Recommendations for evaluation of relatives of sudden arrhythmic death syndrome decedents		
Following SADS, hypothesis-free post-mortem genetic testing using exome or genome sequencing is not recommended.	III	B
Familial evaluation of SADS decedents is recommended: • for first-degree relatives • for relatives who must carry a mutation based on analysis of the family history • for relatives with suspicious symptoms • when the decedent's age is <50 years or if there are other circumstantial data or family history to suggest heritable disease.	I	B
Familial evaluation of SADS decedents is recommended to include genetic testing when post-mortem genetic testing in a SADS decedent detects a pathogenic mutation.	I	B
Baseline familial evaluation of SADS decedents is recommended to include taking a medical history and performing physical examination, standard- and high-precordial lead ECG, echocardiography, and exercise testing.	I	B
In SADS families without a diagnosis after clinical evaluation, follow-up is recommended for children of decedents until they reach adulthood.	I	C
In SADS families without a diagnosis after clinical evaluation, follow-up is not recommended for asymptomatic adults who can be discharged with advice to return if they develop symptoms or if the family history changes.	III	C
Recommendations for treatment of reversible conditions		
Withdrawal of offending agents is recommended whenever drug-induced VA are suspected.	I	B
Investigation for reversible causes (e.g. electrolyte imbalances, ischaemia, hypoxaemia, fever) ⁹ is recommended in patients with VA.	I	C
Recommendations for the acute treatment of sustained ventricular tachycardia and electrical storm		
DC cardioversion is recommended as the first-line treatment for patients with haemodynamically not-tolerated SMVT.	I	B
DC cardioversion is recommended as the first-line treatment for patients presenting with tolerated SMVT provided that the anaesthetic/sedation risk is low.	I	C
In patients presenting with a haemodynamically tolerated idiopathic VT, treatment with intravenous beta-blocker (RVOT VT) or verapamil (fascicular VT) is recommended.	I	C
Intravenous verapamil is not recommended in broad QRS complex tachycardia of unknown mechanism.	III	B
Mild to moderate sedation is recommended in patients with electrical storm to alleviate psychological distress and reduce sympathetic tone.	I	C
Antiarrhythmic therapy with beta-blockers (non-selective preferred) in combination with intravenous amiodarone is recommended in patients with SHD and electrical storm unless contraindicated.	I	B

Intravenous magnesium with supplementation of potassium is recommended in patients with TdP.	I	C
Isoproterenol or transvenous pacing to increase heart rate is recommended in patients with acquired LQT syndrome and recurrent TdP despite correction of precipitating conditions and magnesium.	I	C
Catheter ablation is recommended in patients presenting with incessant VT or electrical storm due to SMVT refractory to AADs.	I	B
Recommendations for treatment with heart failure medication		
Optimal medical treatment including ACE-I/ARB/ARNIs, MRAs, beta-blockers, and SGLT2 inhibitors is indicated in all heart failure patients with reduced EF.	I	A
Recommendations for implantable cardioverter defibrillation implantation (general aspects)		
Implantation of a cardioverter defibrillator is only recommended in patients who have an expectation of good quality survival >1 year.	I	C
It is not recommended to implant an ICD in patients with incessant VAs until the VA is controlled.	III	C
Recommendations for secondary prevention of sudden cardiac death		
ICD implantation is recommended in patients with documented VF or haemodynamically not-tolerated VT in the absence of reversible causes.	I	A
Recommendations for adding cardiac resynchronization therapy to implantable cardioverter defibrillator		
When an ICD is indicated, it is recommended to evaluate whether the patient could benefit from CRT-defibrillator.	I	C
Recommendations for optimization of device programming		
Optimization of ICD programming is indicated to avoid inappropriate and unnecessary therapies and to reduce mortality.	I	A
In single- or dual-chamber ICD patients without bradycardia pacing indications, it is recommended to minimize ventricular pacing.	I	A
Programming of prolonged detection settings is indicated (duration criteria of at least 6–12 s or 30 intervals).	I	A
It is recommended to program the slowest tachycardia therapy zone limit ≥ 188 b.p.m. in primary prevention ICD patients.	I	A
In patients with SHD, programming of at least one ATP therapy is recommended in all tachyarrhythmias zones.	I	A
It is recommended to program algorithms for SVT vs. VT discrimination for tachycardias with rates up to 230 b.p.m.	I	B
It is recommended to activate lead failure alerts.	I	B
Remote monitoring is recommended to reduce the incidence of inappropriate shocks.	I	B
Programming of burst ATP as a first attempt is recommended over ramp ATP.	I	B
For S-ICDs, a dual detection zone configuration is recommended with activation of discrimination algorithm in the lower conditional shock zone	I	B
Recommendations for concomitant treatment to avoid inappropriate implantable cardioverter defibrillator therapy		
Catheter ablation is recommended for ICD patients with recurrent SVT resulting in inappropriate ICD therapies.	I	C
Pharmacological treatment or catheter ablation is recommended in patients with AF-related inappropriate ICD therapies despite optimal ICD programming.	I	C
Recommendation for psychosocial management after implantable cardioverter defibrillator implantation		
Assessment of psychological status and treatment of distress is recommended in ICD patients.	I	C
Communication between patient and physician/healthcare professional is recommended to address ICD-related concerns and to discuss quality-of-life issues before ICD implantation and during disease progression.	I	C
Recommendations for prevention of implantable cardioverter defibrillator complications		
Single-chamber ICD is recommended over dual-chamber ICD in primary prevention patients without current or expected indication for atrial or AV sequential pacing due to a lower risk of device-related complications.	I	A
Recommendations for end-of-life issues in implantable cardioverter defibrillator carriers		
Informed discussion with patient and family about ICD deactivation options and shared decision-making is indicated prior to implantation and in case of significant health status deterioration.	I	C
Recommendations for treatment of ventricular arrhythmia in acute coronary syndrome and vasospasm		
Intravenous beta-blocker treatment is indicated for patients with recurrent PVT/VF during STEMI unless contraindicated.	I	B
Prophylactic treatment with AADs (other than beta-blockers) is not recommended in ACS.	III	B
Recommendations for risk stratification and treatment of ventricular arrhythmia early after myocardial infarction		
Early (before discharge) assessment of LVEF is recommended in all patients with acute MI.	I	B
In patients with pre-discharge LVEF $\leq 40\%$, re-evaluation of LVEF 6–12 weeks after MI is recommended to assess the potential need for primary prevention ICD implantation.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in chronic coronary artery disease		
In patients with syncope and previous STEMI, PES is indicated when syncope remains unexplained after non-invasive evaluation.	I	C
ICD therapy is recommended in patients with CAD, symptomatic heart failure (NYHA class II–III), and LVEF $\leq 35\%$ despite ≥ 3 months of OMT.	I	A
In patients with CAD, prophylactic treatment with AADs other than beta-blockers is not recommended.	III	A
ICD implantation is recommended in patients without ongoing ischaemia with documented VF or haemodynamically not-tolerated VT occurring later than 48 h after MI.	I	A
In patients with CAD and recurrent, symptomatic SMVT or ICD shocks for SMVT despite chronic amiodarone therapy, catheter ablation is recommended in preference to escalating AAD therapy.	I	B
Recommendations for sudden cardiac death prevention in patients with coronary anomalies		
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test in patients with anomalous aortic origin of a coronary artery with an interarterial course to confirm/exclude myocardial ischaemia.	I	C
Cardiac stress imaging during physical exercise is recommended in addition to cardiopulmonary exercise test after surgery in patients with anomalous aortic origin of a coronary artery with a history of aborted CA.	I	C
Surgery is recommended in patients with anomalous aortic origin of a coronary artery with CA, syncope suspected to be due to VAs or angina when other causes have been excluded.	I	C
Recommendations for the management of patients with idiopathic premature ventricular complexes/ventricular tachycardia		
Regular assessment of ventricular function of patients with PVC burden $>10\%$ and normal ventricular function is indicated.	I	C
Catheter ablation as first-line treatment is recommended for symptomatic idiopathic VT/PVCs from the RVOT or the left fascicles.*	I	B
Beta-blockers or non-dihydropyridine CCB are indicated in symptomatic patients with idiopathic VT/PVCs from an origin other than the RVOT or the left fascicles.	I	C
Catheter ablation is not recommended in children <5 years of age or <10 kg weight except when previous medical therapy fails or when VT is not haemodynamically tolerated.	III	C
Amiodarone as a first-line treatment is not recommended in patients with idiopathic VTs/PVCs.	III	C
Verapamil is not recommended in children <1 year of age with PVC/VT, particularly if they have signs of heart failure or concurrent use of other AADs.	III	C
Recommendations for the management of patients with premature ventricular complex-induced or aggravated cardiomyopathy		
In patients with a cardiomyopathy suspected to be caused by frequent and predominately monomorphic PVCs, catheter ablation is recommended.	I	C

Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in dilated cardiomyopathy/hypokinetic non-dilated cardiomyopathy		
Genetic testing (including at least LMNA, PLN, RBM20, and RPLNC genes) is recommended in patients with DCM/HNDCM and AV conduction delay at <50 years, or who have a family history of DCM/HNDCM or SCD in a first-degree relative (at age <50 years).	I	B
Participation in high-intensity exercise including competitive sports is not recommended for individuals with DCM/HNDCM and a LMNA mutation.	III	C
ICD implantation is recommended in patients with DCM/HNDCM, who survive SCA due to VT/VF or experience haemodynamically not-tolerated SMVT.	I	B
In a first-degree relative of a DCM/HNDCM patient, an ECG, and an echocardiogram are recommended if: • the index patient was diagnosed <50 years of age or has clinical features suggestive of an inherited cause, or • there is family history of DCM/HNDCM, or premature unexpected SD.	I	C
Recommendations for diagnostic, risk stratification, sudden cardiac death prevention and treatment of ventricular arrhythmia in arrhythmogenic right ventricular cardiomyopathy		
In patients with suspected ARVC, CMR is recommended.	I	B
In patients with a suspected or definite diagnosis of ARVC, genetic counselling and testing are recommended.	I	B
Avoidance of high-intensity exercise is recommended in patients with a definite diagnosis of ARVC.	I	B
ICD implantation is recommended in ARVC patients with haemodynamically not-tolerated VT or VF.	I	C
In patients with ARVC and non-sustained or sustained VAs, beta-blocker therapy is recommended.	I	C
In a first-degree relative of a patient with ARVC, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in hypertrophic cardiomyopathy		
CMR with LGE is recommended in HCM patients for diagnostic work-up.	I	B
Genetic counselling and testing are recommended in HCM patients.	I	B
It is recommended that the 5-year risk of SCD is assessed at first evaluation and at 1–3-year intervals or when there is a change in clinical status.	I	C
ICD implantation is recommended in HCM patients with haemodynamically not-tolerated VT or VF.	I	B
In a first-degree relative of a patient with HCM, ECG, and echocardiogram are recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in neuromuscular diseases		
Annual follow-up with at least a 12-lead ECG is recommended in patients with muscular dystrophies, even in the concealed phase of the disease.	I	C
It is recommended that patients with neuromuscular disorders who have VAs or ventricular dysfunction are treated in the same way for arrhythmia as patients without neuromuscular disorders.	I	C
Invasive electrophysiological evaluation is recommended in patients with myotonic dystrophy and palpitations or syncope suggestive of VA, VT, or surviving a CA.	I	C
ICD implantation is recommended in patients with myotonic dystrophy and SMVT or aborted CA not caused by BBR-VT.	I	C
In myotonic dystrophy patients, serial electrophysiological evaluation of AV conduction and arrhythmia induction is not recommended without arrhythmia suspicion or progression of ECG conduction disorders.	III	C
In symptomatic patients with BBR-VT, catheter ablation is recommended.	I	C
In patients with myotonic dystrophy undergoing ablation for BBR-VT, pacemaker/ICD implantation is recommended.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in myocarditis		
In confirmed or clinically suspected acute myocarditis, it is recommended that patients who present with life-threatening VA are referred to a specialized centre.	I	C
In patients with haemodynamically not-tolerated SMVT occurring in the chronic phase of myocarditis, an ICD implantation is recommended.	I	C
Recommendations for risk stratification, sudden cardiac death prevention, and treatment of ventricular arrhythmia in cardiac sarcoidosis		
ICD implantation is recommended in patients with cardiac sarcoidosis who have an LVEF ≤35%.	I	B
ICD implantation is recommended in patients with cardiac sarcoidosis who (a) have documented sustained VT, or (2) aborted CA.	I	B
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in valvular heart disease		
PES with standby catheter ablation is recommended in patients with aortic valve disease and SMVT to identify and ablate BBR-VT, especially if it occurs following a valve intervention.	I	C
In patients with valvular heart disease and persistent LV dysfunction after surgical correction (if possible) it is recommended that ICD implantation for primary prevention follows DCM/HNDCM recommendations.	I	C
Recommendations for sudden cardiac death prevention, and treatment of ventricular arrhythmia in congenital heart disease		
In adults with CHD with biventricular physiology and a left systemic ventricle presenting with symptomatic heart failure (NYHA II/III) and EF ≤35% despite ≥3 months of OMT, ICD implantation is indicated.	I	C
In patients with CHD presenting with sustained VAs, evaluation for residual or new anatomical abnormalities is recommended.	I	B
In patients with CHD with not-tolerated VT/aborted CA due to VF, ICD implantation is indicated after exclusion of reversible causes.	I	C
In patients with repaired TOF who present with SMVT or recurrent, symptomatic appropriate ICD therapy for SMVT, catheter ablation performed in specialized centres is recommended.	I	C
Recommendations for management of patients with idiopathic ventricular fibrillation		
It is recommended that idiopathic VF is diagnosed in a SCA survivor, preferably with documentation of VF, after exclusion of an underlying structural, channelopathic, metabolic, or toxicological aetiology.	I	B
ICD implantation is recommended in idiopathic VF.	I	B
Recommendations for management of patients with long QT syndrome		
It is recommended that LQTS is diagnosed with either QTc ≥ 480 ms in repeated 12-lead ECGs with or without symptoms or LQTS diagnostic score >3.	I	C
In patients with clinically diagnosed long QT syndrome, genetic testing, and genetic counselling are recommended.	I	C
It is recommended that LQTS is diagnosed in the presence of a pathogenic mutation, irrespective of the QT duration.	I	C
Routine diagnostic testing with epinephrine challenge is not recommended in LQTS.	III	C
The following is recommended in LQTS: • Avoid QT-prolonging drugs.¹ • Avoid and correct electrolyte abnormalities. • Avoid genotype-specific triggers for arrhythmias.	I	C
Beta-blockers, ideally non-selective beta-blockers (nadolol or propranolol), are recommended in LQTS patients with documented QT interval prolongation, to reduce risk of arrhythmic events.	I	B
Mexiletine is indicated in LQTS patients with a prolonged QT interval.	I	C
ICD implantation in addition to beta-blockers is recommended in LQTS patients with CA.	I	B
ICD implantation is recommended in patients with LQTS who are symptomatic ⁴ while receiving beta-blockers and genotype-specific therapies.	I	C

LCSD is indicated in patients with symptomatic ⁴ LQTS when: (a) ICD therapy is contraindicated or declined; (b) patient is on beta-blockers and genotype-specific therapies with an ICD and experiences multiple shocks or syncope due to VA.	I	C
Invasive electrophysiologic study is not recommended in LQTS.	III	C
Recommendations for management of patients with Andersen-Tawil syndrome		
Genetic testing is recommended in patients with suspected Andersen-Tawil syndrome.	I	C
ICD implantation is recommended in patients with Andersen-Tawil syndrome after aborted CA not tolerated sustained VT.	I	C
Recommendations for management of patients with Brugada syndrome		
It is recommended that BrS is diagnosed in patients with no other heart disease and a spontaneous type 1 Brugada ECG pattern.	I	C
It is recommended that BrS is diagnosed in patients with no other heart disease who have survived a CA due to VF or PVT and exhibit a type 1 Brugada ECG induced by sodium channel blocker challenge or during fever.	I	C
Genetic testing for SCN5A gene is recommended for probands with BrS.	I	C
Sodium channel blocker test is not recommended in patients with a prior type 1 Brugada pattern.	III	C
The following is recommended in all patients with BrS: (a) Avoidance of drugs that may induce ST-segment elevation in right precordial leads (http://www.brugadadrugs.org). (b) Avoidance of cocaine, cannabis, and excessive alcohol intake. (c) Treatment of fever with antipyretic drugs.	I	C
ICD implantation is recommended in patients with BrS who: (a) Are survivors of an aborted CA, and/or (b) Have documented spontaneous sustained VT.	I	C
Catheter ablation in asymptomatic BrS patients is not recommended.	III	C
Recommendations for management of patients with early repolarization pattern/syndrome		
It is recommended that the ERP is diagnosed as J-point elevation of ≥ 1 mm in two adjacent inferior and/or lateral ECG leads.	I	C
It is recommended that the ERS is diagnosed in a patient resuscitated from unexplained VF/PVT in the presence of ERP.	I	C
Clinical evaluation is not recommended routinely in asymptomatic subjects with ERP.	III	C
ICD implantation is recommended in patients with a diagnosis of ERS who have survived a CA.	I	B
ICD implantation is not recommended in asymptomatic patients with an isolated ERP.	III	C
Recommendations for management of patients with catecholaminergic polymorphic ventricular tachycardia		
It is recommended that CPVT is diagnosed in the presence of a structurally normal heart, normal ECG, and exercise- or emotion-induced bidirectional or PVT.	I	C
It is recommended that CPVT is diagnosed in patients who are carriers of a mutation in disease-causing genes.	I	C
Genetic testing and genetic counselling are indicated in patients with clinical suspicion or clinical diagnosis of CPVT.	I	C
Avoidance of competitive sports, strenuous exercise, and exposure to stressful environments is recommended in all patients with CPVT.	I	C
Beta-blockers, ideally non-selective (nadolol or propranolol), are recommended in all patients with a clinical diagnosis of CPVT.	I	C
ICD implantation combined with beta-blockers and flecainide is recommended in CPVT patients after aborted CA.	I	C
PES is not recommended for stratification of SCD risk.	III	C
Recommendations for management of patients with short QT syndrome		
It is recommended that SQTS is diagnosed in the presence of a QTc ≤ 360 ms and one or more of the following: (a) a pathogenic mutation; (b) a family history of SQTS; (c) survival from a VT/VF episode in the absence of heart disease.	I	C
Genetic testing is indicated in patients diagnosed with SQTS.	I	C
ICD implantation is recommended in patients with a diagnosis of SQTS who: (a) are survivors of an aborted CA, and/or (b) have documented spontaneous sustained VT.	I	C
PES is not recommended for SCD risk stratification in SQTS patients.	III	C
Recommendations for the prevention of sudden cardiac death and management of ventricular arrhythmia during pregnancy		
During pregnancy, electrical cardioversion is recommended for sustained VT.	I	C
If ICD implantation is indicated during pregnancy, implantation is recommended with optimal radiation protection.	I	C
Continuation of beta-blockers is recommended during pregnancy and post-partum in patients with LQTS or CPVT.	I	C
Recommendations for risk stratification and prevention of sudden cardiac death in athletes		
In athletes with positive medical history, abnormal physical examination, or ECG alterations, further investigations including echocardiography and/or CMR to confirm (or exclude) an underlying disease are recommended.	I	C
It is recommended that athletes diagnosed with a cardiovascular disease associated with SCD are managed according to current guidelines for sports eligibility.	I	C
It is recommended that staff at sporting facilities are trained in CPR and in the use of AED.	I	C

AAD, anti-arrhythmic drug; ACE-I, angiotensin-converting-enzyme inhibitor; AED, automated external defibrillator; ARB, angiotensin receptor blocker; ARNI, angiotensin receptor neprilysin inhibitor; ARVC, arrhythmogenic right ventricular cardiomyopathy; ATP, anti-tachycardia pacing; AV, atrioventricular; BBR-VT, bundle branch re-entry; BrS, Brugada syndrome; CA, cardiac arrest; CAD, coronary artery disease; CCB, calcium channel blocker; CHD, congenital heart disease; CIED, cardiac implantable electronic device; CMR, cardiac magnetic resonance; CPR, cardiopulmonary resuscitation; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRT, cardiac resynchronization therapy; DC, direct current; DCM, dilated cardiomyopathy; ECG, electrocardiogram; EF, ejection fraction; ERP, early repolarization pattern; ERS, early repolarization syndrome; HCM, hypertrophic cardiomyopathy; HNCDCM, hypokinetic non-dilated cardiomyopathy; ICD, implantable cardioverter defibrillator; CAD, coronary artery disease; IAR, implantable loop recorder; LCSD, left cardiac sympathetic denervation; LGE, late gadolinium enhancement; LQTS, long QT syndrome; LV, left ventricular; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NSVT, non-sustained ventricular tachycardia; NYHA, New York Heart Association; GHCA, out-of-hospital cardiac arrest; OHIT, optimal medical therapy; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RVOT, right ventricular outflow tract; SADS, sudden arrhythmic death syndrome; SCA, sudden cardiac arrest; SCD, sudden cardiac death; SD, sudden death; SGLT2, sodium-glucose co-transporter 2; SHD, structural heart disease; SMVT, sustained monomorphic ventricular tachycardia; SQT, short QT syndrome; STEMI, ST-elevation myocardial infarction; SVT, supraventricular tachycardia; VA, ventricular arrhythmia; VF, ventricular fibrillation; VT, ventricular tachycardia.

⁴Class of recommendation.
⁵Level of evidence.
⁶Shopping malls, stadiums, public transport stations, casinos.
⁷List not exhaustive.
⁸OE C for VT/PVCs from left fascicles.
<http://www.crediblemed.org>
⁹Arrhythmic syncope or haemodynamically non-tolerated VA

12. Quality indicators

Quality indicators (QIs) are tools that may be used to evaluate care quality, including structural aspects, process, and outcomes of care.¹¹⁵³ They serve as a mechanism for enhancing adherence to guideline recommendations, through associated quality improvement initiatives and the benchmarking of care providers.^{1154,1155} As such, the role of QIs in improving care and outcomes is increasingly recognized by healthcare authorities, professional organizations, payers, and the public.¹¹⁵³

The ESC understands the need for measuring and reporting quality and outcomes of cardiovascular care, and has established methods for the development of the ESC QIs for the quantification of care and outcomes for

cardiovascular disease.¹¹⁵³ In parallel with the writing of this Clinical Practice Guideline document, a process has been initiated to develop QIs for patients with, or at risk of, VA or SCD using the ESC methodology and through the collaboration with domain experts and the European Heart Rhythm Association. These QIs, alongside their specifications and development process, will be published separately.

13. Supplementary data

[Supplementary data](#) is available at *European Heart Journal online*.

14. Data availability statement

No new data were generated or analysed in support of this research.

15. Author information

Author/task force Member Affiliations:

Marta de Riva, Cardiology, Leiden University Medical Centre, Leiden, Netherlands; **Bo Gregers Winkel**, Cardiology, Rigshospitalet, Copenhagen, Denmark, European Reference Networks for rare, low prevalence and complex diseases of the heart, ERN-GUARD HEART; **Elijah R. Behr**, Cardiovascular Clinical Academic Group, Cardiology Section, St George's Hospitals, University of London, London, United Kingdom, Department of Cardiology, St George's University Hospitals NHS Foundation Trust, London, United Kingdom, and Cardiology, Mayo Clinic Healthcare, London, United Kingdom; **Nico A. Blom**, Paediatric Cardiology, Leiden University Medical Centre, Leiden, Netherlands, and Paediatric Cardiology, Amsterdam University Medical Center, Amsterdam, Netherlands; **Philippe Charron**, APHP, Centre de Référence des Maladies Cardiaques Rares, Hôpital Pitié-Salpêtrière, Paris, France, UMR_S 1166, and ICAN Institute for Cardiometabolism, and Nutrition, Sorbonne Université, Paris, France, and European Reference Networks for rare, low prevalence and complex diseases of the heart, ERN-Guard HEART, Paris, France; **Domenico Corrado**, Department of Cardiac, Thoracic and Vascular Sciences and Public Health, University of Padova, Padova, Italy; **Nikolaos Dagres**, Department of Electrophysiology, Heart Center Leipzig at University of Leipzig, Leipzig, Germany; **Christian de Chillou**, Department of Cardiology, CHRU-Nancy, Nancy, France, and IADI, INSERM U1254, Université de Lorraine, Nancy, France; **Lars Eckardt**, Department of Cardiology II Electrophysiology, University Hospital Münster, Münster, Germany; **Tim Friede**, Department of Medical Statistics, University Medical Center Goettingen, Goettingen, Germany, and Partner Site Goettingen, DZHK (German Center for Cardiovascular Research), Goettingen, Germany; **Kristina H. Haugaa**, Department of Cardiology, Oslo University Hospital, Rikshospitalet, Oslo, Norway, and Faculty of Medicine, University of Oslo, Oslo, Norway; **Márcio Ze Hocini**, Cardiology Department, Liryc Institute, Pessac, France, Hôpital Cardiologique du Haut Lévéque, Pessac, France, and Université de Bordeaux, Bordeaux, France; **Pier D. Lambiase**, Institute of Cardiovascular Science, University College, London, United Kingdom, Barts Heart Centre, St Bartholomews Hospital, London, United Kingdom, and Heart, Vascular and Thoracic Institute, Cleveland Clinic, London, United Kingdom; **Eloi Marijon**, Cardiology Department, European Georges Pompidou Hospital, Paris, France; **Jose L. Merino**, Arrhythmia and Electrophysiology Robotic Unit, La Paz University Hospital, Universidad Autonoma, IdiPaz, Madrid, Spain, Cardiology Department, Hospital Ruber Juan Bravo, Madrid, Spain, and Cardiac Electrophysiology, Hospital Viamed Santa Elena, Madrid, Spain; **Petr Peichl**, Cardiology Department, IKEM, Prague, Czech Republic; **Silvia G. Priori**, Molecular Medicine Department, University of Pavia, Pavia, Italy, Molecular Cardiology Department, Istituti Clinici Scientifici Maugeri SpA SB, Pavia, Italy, and Molecular Cardiology Department, Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC), Madrid, Spain, European Reference Networks for rare, low prevalence and complex diseases of the heart, ERN-GUARD HEART; **Tobias Reichlin**, Department of Cardiology, Inselspital University Hospital Bern, University of Bern, Bern, Switzerland; **Jeanette Schulz-Menger**, Cardiology, ECRC, Charité Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany, Cardiology, Helios Clinics Berlin-Buch, Berlin, Germany, and DZHK Partnersite Berlin, Charité, Berlin, Germany; **Christian Sticherling**, Department of Cardiology, University Hospital Basel, University of Basel, Basel, Switzerland; **Stylianos Tzeis**, Cardiology Department, Mitera Hospital, Hygeia Group, Athens, Greece; **Axel Verstraël** (Belgium), ESC Patient Forum, Sophia Antipolis, France; and **Maurizio Volterrani**, Department of Cardiology, IRCCS San Raffaele Roma, Rome, Italy, and Professor of Exercise Science and Medicine, San Raffaele Telematic University of Rome, Rome.

16. Appendix

ESC Scientific Document Group

Includes Document Reviewers and ESC National Cardiac Societies.

Document Reviewers: Maja Cikes (CPG Review Coordinator) (Croatia), Paulus Kirchhof (CPG Review Coordinator) (Germany), Magdy Abdelhamid (Egypt), Victor Aboyans (France), Elena Arbelo (Spain), Fernando Arribas (Spain), Riccardo Asteggiano (Italy), Cristina Basso (Italy), Axel Bauer (Austria), Emanuele Bertaglia (Italy), Tor Biering-Sørensen (Denmark), Carina Blomström-Lundqvist (Sweden), Michael A. Borger (Germany), Jelena Åbergelutkienė (Lithuania), Bernard Cosyns (Belgium), Volkmar Falk (Germany), Laurent Fauchier (France), Bulent Gorennek (Turkey), Sigrun Halvorsen (Norway), Robert Hatala (Slovakia), Hein Heidbuchel (Belgium), Stefan Kaab (Germany), Aleksandra Konradi (Russian Federation), Konstantinos C. Koskinas (Switzerland), Dipak Kotecha (United Kingdom), Ulf Landmesser (Germany), Basil S. Lewis (Israel), Ales Linhart (Czech Republic), Maja-Lisa Løchen (Norway), Lars H. Lund (Sweden), Andreas Metzner (Germany), Richard Mindham (United Kingdom), Jens Cosedis Nielsen (Denmark), Tone M. Norekvål (Norway), Monica Patten (Germany), Eva Prescott (Denmark), Amina Rakisheva (Kazakhstan), Carol Ann Remme (Netherlands), Ivo Roca-Luque (Spain), Andrea Sarkozy (Belgium), Daniel Scherr (Austria), Marta Sitges (Spain), Rhian M. Touyz (Canada/United Kingdom), Nicolas Van Mieghem (Netherlands), Vedran Velagic (Croatia), Sami Viskin (Israel), and Paul G. A. Volders (Netherlands).

ESC National Cardiac Societies actively involved in the review process of the 2022 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death.

Algeria: Algerian Society of Cardiology, Brahim Kichou; **Armenia:** Armenian Cardiologists Association, Mihran Martirosyan; **Austria:** Austrian Society of Cardiology, Daniel Scherr; **Azerbaijan:** Azerbaijan Society of Cardiology, Farid Aliyev; **Belgium:** Belgian Society of Cardiology, Rik Willems; **Bosnia and Herzegovina:** Association of Cardiologists of Bosnia and Herzegovina, Nabil Naser; **Bulgaria:** Bulgarian Society of Cardiology, Tchavdar Shalganov; **Croatia:** Croatian Cardiac Society, Davor Milicic; **Cyprus:** Cyprus Society of Cardiology, Theodoros Christophides; **Czech Republic:** Czech Society of Cardiology, Josef Kautzner; **Denmark:** Danish Society of Cardiology, Jim Hansen; **Egypt:** Egyptian Society of Cardiology, Lamyaa Allam; **Estonia:** Estonian Society of Cardiology, Priit Kampus; **Finland:** Finnish Cardiac Society, Juhani Junttila; **France:** French Society of Cardiology, Christophe Leclercq; **Georgia:** Georgian Society of Cardiology, Kakhaber Etsadashvili; **Germany:** German Cardiac Society, Daniel Steven; **Greece:** Hellenic Society of Cardiology, Konstantinos Gatzoulis; **Hungary:** Hungarian Society of Cardiology, László Gellér; **Iceland:** Icelandic Society of Cardiology, David O. Arnar; **Ireland:** Irish Cardiac Society, Joseph Galvin, European Reference Networks for rare, low prevalence and complex diseases of the heart, ERN-GUARD HEART; **Israel:** Israel Heart Society, Moti Haim; **Italy:** Italian Federation of Cardiology, Carlo Pappone; **Kosovo (Republic of):** Kosovo Society of Cardiology, Shpend Elezi; **Kyrgyzstan:** Kyrgyz Society of Cardiology, Alina Kerimkulova; **Latvia:** Latvian Society of Cardiology, Oskars Kalejs; **Lebanon:** Lebanese Society of Cardiology, Ali Rabah; **Lithuania:** Lithuanian Society of Cardiology, Aras Puodziukynas; **Luxembourg:** Luxembourg Society of Cardiology, Carlo Dimmer; **Malta:** Maltese Cardiac Society, Mark Adrian Sammut; **Moldova (Republic of):** Moldavian Society of Cardiology, Lilia David; **Montenegro:** Montenegro Society of Cardiology, Aneta Boskovic; **Morocco:** Moroccan Society of Cardiology, Abdelhamid Moustaghfir; **Netherlands:** Netherlands Society of Cardiology, Alexander H. Maass; **North Macedonia:** North Macedonian Society of Cardiology, Lidija Poposka; **Norway:** Norwegian Society of Cardiology, Ole Christian Mjølstad; **Poland:** Polish Cardiac Society, Przemysław Mitkowski; **Portugal:** Portuguese Society of Cardiology, Leonor Parreira; **Romania:** Romanian Society of Cardiology, Dragos Cozma; **Russian Federation:** Russian Society of Cardiology, Elena Golukhova; **San Marino:** San Marino Society of Cardiology, Roberto Bini; **Serbia:** Cardiology Society of Serbia, Sinisa Stojkovic; **Slovakia:** Slovak Society of Cardiology, Peter Hlivak; **Slovenia:** Slovenian Society of Cardiology, Andrej Pernat; **Spain:** Spanish Society of Cardiology, Nicasio Perez Castellano; **Sweden:** Swedish Society of Cardiology, Pyotr G. Platonov; **Switzerland:** Swiss Society of Cardiology, Firat Duru; **Syrian Arab Republic:** Syrian Cardiovascular Association, Ahmad Rasheed Al Saadi; **Tunisia:** Tunisian Society of Cardiology and Cardio-Vascular Surgery, Sana Ouali; **Turkey:** Turkish Society of Cardiology, Sabri Demircan; **Ukraine:** Ukrainian Association of Cardiology, Oleg Sychov; and **United Kingdom of Great Britain and Northern Ireland:** British Cardiovascular Society, Alistair Slade.

ESC Clinical Practice Guidelines (CPG) Committee: Colin Baigent (Chairperson) (United Kingdom), Magdy Abdelhamid (Egypt), Victor Aboyans (France), Sotiris Antoniou (United Kingdom), Elena Arbelo (Spain), Riccardo Asteggiano (Italy), Andreas Baumbach (United Kingdom), Michael A. Borger (Germany), Jelena Åbergelutkienė (Lithuania), Maja Cikes (Croatia), Jean-Philippe Collet (France), Volkmar Falk (Germany), Laurent Fauchier (France), Chris P. Gale (United Kingdom), Sigrun Halvorsen (Norway), Bernard Iung (France), Tiny Jaarsma (Sweden), Aleksandra Konradi (Russian Federation), Konstantinos C. Koskinas (Switzerland), Dipak Kotecha (United Kingdom), Ulf Landmesser (Germany), Basil S. Lewis (Israel), Ales Linhart (Czech Republic), Maja-Lisa Løchen

(Norway), Richard Mindham (United Kingdom), Jens Cosedis Nielsen (Denmark), Steffen E. Petersen (United Kingdom), Eva Prescott (Denmark), Amina Rakisheva (Kazakhstan), Marta Sitges (Spain), and Rhian M. Touyz (Canada/United Kingdom).

References

- 1Brignole Â M, Moya Â A, de Lange Â FJ, Deharo Â J-C, Elliott Â PM, Fanciulli Â A, et al. Â 2018 ESC Guidelines for the diagnosis and management of syncope. *Eur Heart J* Â 2018;39:1883â1948.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 2Glikson Â M, Nielsen Â JC, Kronborg Â MB, Michowitz Â Y, Auricchio Â A, Barbash Â IM, et al. Â 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Eur Heart J* Â 2021;42:3427â3520.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 3Ibanez Â B, James Â S, Agewall Â S, Antunes Â MJ, Bucciarelli-Ducci Â C, Bueno Â H, et al. Â 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the task force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* Â 2018;39:119â177.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 4Pelliccia Â A, Sharma Â S, Gati Â S, BÃck Â M, BjÃrjesson Â M, Caselli Â S, et al. Â 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J* Â 2021;42:17â96.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 5Wahbi Â K, Babuty Â D, Probst Â V, Wissocque Â L, Labombarda Â F, Porcher Â R, et al. Â Incidence and predictors of sudden death, major conduction defects and sustained ventricular tachyarrhythmias in 1388 patients with myotonic dystrophy type 1. *Eur Heart J* Â 2017;38:751â758.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 6Bucci Â E, Testa Â M, Licchelli Â L, Frattari Â A, El Halabieh Â NA, Gabriele Â E, et al. Â A 34-year longitudinal study on long-term cardiac outcomes in DM1 patients with normal ECG at baseline at an Italian clinical centre. *J Neurol* Â 2018;265:885â895.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 7Chong-Nguyen Â C, Wahbi Â K, Algalarrondo Â V, BÃcane Â HM, Radvanyi-Hoffman Â H, Arnaud Â P, et al. Â Association between mutation size and cardiac involvement in myotonic dystrophy type 1: an analysis of the DM1-heart registry. *Circ Cardiovasc Genet* Â 2017;10:e001526.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 8Sabovic Â M, Medica Â I, Logar Â N, MandiÃ Â E, Zidar Â J, Peterlin Â B. Relation of CTG expansion and clinical variables to electrocardiogram conduction abnormalities and sudden death in patients with myotonic dystrophy. *Neuromuscul Disord* Â 2003;13:822â826.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 9Clarke Â NR, Kelion Â AD, Nixon Â J, Hilton-Jones Â D, Forfar Â JC. Does cytosine-thymine-guanine (CTG) expansion size predict cardiac events and electrocardiographic progression in myotonic dystrophy? Â *Heart* Â 2001;86:411â416.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 10Melacini Â P, Villanova Â C, Menegazzo Â E, Novelli Â G, Danieli Â G, Rizzoli Â G, et al. Â Correlation between cardiac involvement and CTG trinucleotide repeat length in myotonic dystrophy. *J Am Coll Cardiol* Â 1995;25:239â245.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 11Lazarus Â A, Varin Â J, Ounnoughene Â Z, Radvanyi Â H, Junien Â C, Coste Â J, et al. Â Relationships among electrophysiological findings and clinical status, heart function, and extent of DNA mutation in myotonic dystrophy. *Circulation* Â 1999;99:1041â1046.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 12Sansone Â VA, Brigonzi Â E, Schoser Â B, Villani Â S, Gaeta Â M, De Ambroggi Â G, et al. Â The frequency and severity of cardiac involvement in myotonic dystrophy type 2 (DM2): long-term outcomes. *Int J Cardiol* Â 2013;168:1147â1153.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 13Breton R, Mathieu J. Usefulness of clinical and electrocardiographic data for predicting adverse cardiac events in patients with myotonic dystrophy. *Can J Cardiol* 2009;25:e233-237. [Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 14Hermans MCE, Faber CG, Bekkers SCAM, de Die-Smulders CEM, Gerrits MM, Merckies ISJ, et al. Structural and functional cardiac changes in myotonic dystrophy type 1: a cardiovascular magnetic resonance study. *J Cardiovasc Magn Reson* 2012;14:48. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 15Schmacht L, Traber J, Grieben U, Utz W, Dieringer MA, Kellman P, et al. Cardiac involvement in myotonic dystrophy type 2 patients with preserved ejection fraction: detection by cardiovascular magnetic resonance. *Circ Cardiovasc Imaging* 2016;9:e004615. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 16Groh WJ, Groh MR, Saha C, Kincaid JC, Simmons Z, Ciafaloni E, et al. Electrocardiographic abnormalities and sudden death in myotonic dystrophy type 1. *N Engl J Med* 2008;358:2688-2697. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 17Bhakta D, Groh MR, Shen C, Pascuzzi RM, Groh WJ. Increased mortality with left ventricular systolic dysfunction and heart failure in adults with myotonic dystrophy type 1. *Am Heart J* 2010;160:1137-1141. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 18Petri H, Ahtarovski KA, Vejlsr   N, Vissing J, Witting N, K  ber L, et al. Myocardial fibrosis in patients with myotonic dystrophy type 1: a cardiovascular magnetic resonance study. *J Cardiovasc Magn Reson* 2014;16:59. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 19Myerburg RJ, Junttila MJ. Sudden cardiac death caused by coronary heart disease. *Circulation* 2012;125:1043-1052. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 20Stecker EC, Reinier K, Marijon E, Narayanan K, Teodorescu C, Uy-Evanado A, et al. Public health burden of sudden cardiac death in the United States. *Circ Arrhythm Electrophysiol* 2014;7:212-217. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 21Chen LY, Sotoodehnia N, B  kov   P, Lopez FL, Yee LM, Heckbert SR, et al. Atrial fibrillation and the risk of sudden cardiac death: the atherosclerosis risk in communities study and cardiovascular health study. *JAMA Intern Med* 2013;173:29-35. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 22Marijon E, Uy-Evanado A, Dumas F, Karam N, Reinier K, Teodorescu C, et al. Warning symptoms are associated with survival from sudden cardiac arrest. *Ann Intern Med* 2016;164:23-29. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 23Glinge C, Jabbari R, Risgaard B, Lynge TH, Engstr  m T, Albert CM, et al. Symptoms before sudden arrhythmic death syndrome: a nationwide study among the young in Denmark. *J Cardiovasc Electrophysiol* 2015;26:761-767. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 24  elsgesen FN, Lynge TH, Blanche P, Banner J, Prescott E, Jabbari R, et al. Temporal trends and sex differences in sudden cardiac death in the Copenhagen City Heart Study. *Heart* 2021;107:1303-1309. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 25Winkel BG, Holst AG, Theilade J, Kristensen IB, Thomsen JL, Ottesen GL, et al. Nationwide study of sudden cardiac death in persons aged 1-35 years. *Eur Heart J* 2011;32:983-990. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 26Pilmer CM, Kirsh JA, Hildebrandt D, Krahn AD, Gow RM. Sudden cardiac death in children and adolescents between 1 and 19 years of age. *Heart Rhythm* 2014;11:239-245. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 27Risgaard B, Winkel BG, Jabbari R, Behr ER, Ingemann-Hansen O, Thomsen JL, et al. Burden of sudden cardiac death in persons aged 1 to 49 years: nationwide study in Denmark. *Circ Arrhythm Electrophysiol* 2014;7:205-211.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

28Fishman [Â](#) GI, Chugh [Â](#) SS, DiMarco [Â](#) JP, Albert [Â](#) CM, Anderson [Â](#) ME, Bonow [Â](#) RO, et al. [Â](#) Sudden cardiac death prediction and prevention report from a national heart, lung, and blood institute and heart rhythm society workshop. *Circulation* [Â](#) 2010;122:2335[PAGETOP](#)[PAGETOP](#)2348.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

29Wong [Â](#) CX, Brown [Â](#) A, Lau [Â](#) DH, Chugh [Â](#) SS, Albert [Â](#) CM, Kalman [Â](#) JM, et al. [Â](#) Epidemiology of sudden cardiac death: global and regional perspectives. *Heart Lung Circ* [Â](#) 2019;28:6[PAGETOP](#)[PAGETOP](#)14.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

30Bougouin [Â](#) W, Lamhaut [Â](#) L, Marijon [Â](#) E, Jost [Â](#) D, Dumas [Â](#) F, Deye [Â](#) N, et al. [Â](#) Characteristics and prognosis of sudden cardiac death in Greater Paris: population-based approach from the Paris Sudden Death Expertise Center (Paris-SDEC). *Intensive Care Med* [Â](#) 2014;40:846[PAGETOP](#)[PAGETOP](#)854.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

31Deo [Â](#) R, Albert [Â](#) CM. Epidemiology and genetics of sudden cardiac death. *Circulation* [Â](#) 2012;125:620[PAGETOP](#)[PAGETOP](#)637.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

32Gerber [Â](#) Y, Jacobsen [Â](#) SJ, Frye [Â](#) RL, Weston [Â](#) SA, Killian [Â](#) JM, Roger [Â](#) VL. Secular trends in deaths from cardiovascular diseases: a 25-year community study. *Circulation* [Â](#) 2006;113:2285[PAGETOP](#)[PAGETOP](#)2292.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

33Krahn [Â](#) AD, Connolly [Â](#) SJ, Roberts [Â](#) RS, Gent [Â](#) M, ATMA Investigators. Diminishing proportional risk of sudden death with advancing age: implications for prevention of sudden death. *Am Heart J* [Â](#) 2004;147:837[PAGETOP](#)[PAGETOP](#)840.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

34Becker [Â](#) LB, Han [Â](#) BH, Meyer [Â](#) PM, Wright [Â](#) FA, Rhodes [Â](#) KV, Smith [Â](#) DW, et al. [Â](#) Racial differences in the incidence of cardiac arrest and subsequent survival. The CPR Chicago Project. *N Engl J Med* [Â](#) 1993;329:600[PAGETOP](#)[PAGETOP](#)606.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

35Zhao [Â](#) D, Post [Â](#) WS, Blasco-Colmenares [Â](#) E, Cheng [Â](#) A, Zhang [Â](#) Y, Deo [Â](#) R, et al. [Â](#) Racial differences in sudden cardiac death. *Circulation* [Â](#) 2019;139:1688[PAGETOP](#)[PAGETOP](#)1697.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

36de Vreede-Swagemakers [Â](#) JJ, Gorgels [Â](#) AP, Dubois-Arbouw [Â](#) WI, van Ree [Â](#) JW, Daemen [Â](#) MJ, Houben [Â](#) LG, et al. [Â](#) Out-of-hospital cardiac arrest in the 1990s: a population-based study in the Maastricht area on incidence, characteristics and survival. *J Am Coll Cardiol* [Â](#) 1997;30:1500[PAGETOP](#)[PAGETOP](#)1505.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

37Lyngge [Â](#) TH, Risgaard [Â](#) B, Banner [Â](#) J, Nielsen [Â](#) JL, Jespersen [Â](#) T, Stampe [Â](#) NK, et al. [Â](#) Nationwide burden of sudden cardiac death: a study of 54,028 deaths in Denmark. *Heart Rhythm* [Â](#) 2021;18:1657[PAGETOP](#)[PAGETOP](#)1665.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

38Atwood [Â](#) C, Eisenberg [Â](#) MS, Herlitz [Â](#) J, Rea [Â](#) TD. Incidence of EMS-treated out-of-hospital cardiac arrest in Europe. *Resuscitation* [Â](#) 2005;67:75[PAGETOP](#)[PAGETOP](#)80.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

39Empana [Â](#) J-P, Blom [Â](#) MT, BÅttiger [Â](#) BW, Dagres [Â](#) N, Dekker [Â](#) JM, Gislason [Â](#) G, et al. [Â](#) Determinants of occurrence and survival after sudden cardiac arrestâ€”a European perspective: The ESCAPE-NET project. *Resuscitation* [Â](#) 2018;124:7[PAGETOP](#)[PAGETOP](#)13.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

40Fox [Â](#) CS, Evans [Â](#) JC, Larson [Â](#) MG, Kannel [Â](#) WB, Levy [Â](#) D. Temporal trends in coronary heart disease mortality and sudden cardiac death from 1950 to 1999: the Framingham Heart Study. *Circulation* [Â](#) 2004;110:522[PAGETOP](#)[PAGETOP](#)527.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

41Niemeijer [Â](#) MN, van den Berg [Â](#) ME, Leening [Â](#) MJG, Hofman [Â](#) A, Franco [Â](#) OH, Deckers [Â](#) JW, et al. [Â](#) Declining incidence of sudden cardiac death from 1990s to 2010 in a general middle-aged and elderly population: the Rotterdam Study. *Heart Rhythm* [Â](#) 2015;12:123[PAGETOP](#)[PAGETOP](#)129.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

- 42Lyngøe TH, Nielsen JL, Blanche P, Gislason G, Torp-Pedersen C, Winkel BG, et al. Decline in incidence of sudden cardiac death in the young: a 10-year nationwide study of 8756 deaths in Denmark. *Europace* 2019;21:909a-917.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 43Zheng ZJ, Croft JB, Giles WH, Mensah GA. Sudden cardiac death in the United States, 1989 to 1998. *Circulation* 2001;104:2158a-2163.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 44Dudas K, Lappas G, Stewart S, Rosengren A. Trends in out-of-hospital deaths due to coronary heart disease in Sweden (1991 to 2006). *Circulation* 2011;123:46a-52.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 45Harmon KG, Asif IM, Maleszewski JJ, Owens DS, Prutkin JM, Salerno JC, et al. Incidence, cause, and comparative frequency of sudden cardiac death in national collegiate athletic association athletes: a decade in review. *Circulation* 2015;132:10a-19.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 46Risgaard B, Winkel BG, Jabbari R, Glinge C, Ingemann-Hansen O, Thomsen JL, et al. Sports-related sudden cardiac death in a competitive and a noncompetitive athlete population aged 12 to 49 years: data from an unselected nationwide study in Denmark. *Heart Rhythm* 2014;11:1673a-1681.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 47Holst AG, Winkel BG, Theilade J, Kristensen IB, Thomsen JL, Ottesen GL, et al. Incidence and etiology of sports-related sudden cardiac death in Denmark: implications for preparticipation screening. *Heart Rhythm* 2010;7:1365a-1371.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 48Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO. Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980-2006. *Circulation* 2009;119:1085a-1092.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 49Corrado D, Basso C, Thiene G. Sudden cardiac death in athletes: what is the role of screening? *Curr Opin Cardiol* 2012;27:41a-48.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 50Albert CM, Mittleman MA, Chae CU, Lee IM, Hennekens CH, Manson JE. Triggering of sudden death from cardiac causes by vigorous exertion. *N Engl J Med* 2000;343:1355a-1361.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 51Siscovick DS, Weiss NS, Fletcher RH, Lasky T. The incidence of primary cardiac arrest during vigorous exercise. *N Engl J Med* 1984;311:874a-877.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 52Marijon E, Tafflet M, Celermajer DS, Dumas F, Perier M-C, Mustafic H, et al. Sports-related sudden death in the general population. *Circulation* 2011;124:672a-681.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 53Heidbuchel H, Willems R, Jordaens L, Olshansky B, Carre F, Lozano IF, et al. Intensive recreational athletes in the prospective multinational ICD sports safety registry: results from the European cohort. *Eur J Prev Cardiol* 2019;26:764a-775.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 54Marijon E, Bougouin W, Poirier M-C, Celermajer DS, Jouven X. Incidence of sports-related sudden death in France by specific sports and sex. *JAMA* 2013;310:642a-643.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 55Marijon E, Uy-Evanado A, Reinier K, Teodorescu C, Narayanan K, Jouven X, et al. Sudden cardiac arrest during sports activity in middle age. *Circulation* 2015;131:1384a-1391.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 56Bagnall RD, Weintraub RG, Ingles J, Dufloy J, Yeates L, Lam L, et al. A prospective study of sudden cardiac death among children and young adults. *N Engl J Med* 2016;374:2441a-2452.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 57Wisten A, Forsberg H, Krantz P, Messner T. Sudden cardiac death in 15-35-year olds in Sweden during 1992-99. *J Intern Med* 2002;252:529a-536.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

58Chugh [Â](#) SS, Jui [Â](#) J, Gunson [Â](#) K, Stecker [Â](#) EC, John [Â](#) BT, Thompson [Â](#) B, et al. [Â](#) Current burden of sudden cardiac death: multiple source surveillance versus retrospective death certificate-based review in a large US. community. *J Am Coll Cardiol* [Â](#) 2004;44:1268[a](#)^{PADST}1275.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

59Winkel [Â](#) BG, Risgaard [Â](#) B, Sadjadieh [Â](#) G, Bundgaard [Â](#) H, Hauns^Â, [Â](#) S, Tfelt-Hansen [Â](#) J. Sudden cardiac death in children (13[a](#)^{PADST}18 years): symptoms and causes of death in a nationwide setting. *Eur Heart J* [Â](#) 2014;35:868[a](#)^{PADST}875.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

60Byrne [Â](#) R, Constant [Â](#) O, Smyth [Â](#) Y, Callagy [Â](#) G, Nash [Â](#) P, Daly [Â](#) K, et al. [Â](#) Multiple source surveillance incidence and aetiology of out-of-hospital sudden cardiac death in a rural population in the West of Ireland. *Eur Heart J* [Â](#) 2008;29:1418[a](#)^{PADST}1423.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

61Eckart [Â](#) RE, Shry [Â](#) EA, Burke [Â](#) AP, McNear [Â](#) JA, Appel [Â](#) DA, Castillo-Rojas [Â](#) LM, et al. [Â](#) Sudden death in young adults an autopsy-based series of a population undergoing active surveillance. *J Am Coll Cardiol* [Â](#) 2011;58:1254[a](#)^{PADST}1261.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

62Waldmann [Â](#) V, Karam [Â](#) N, Bougouin [Â](#) W, Sharifzadehgan [Â](#) A, Dumas [Â](#) F, Narayanan [Â](#) K, et al. [Â](#) Burden of coronary artery disease as a cause of sudden cardiac arrest in the young. *J Am Coll Cardiol* [Â](#) 2019;73:2118[a](#)^{PADST}2120.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

63Waldmann [Â](#) V, Karam [Â](#) N, Rischard [Â](#) J, Bougouin [Â](#) W, Sharifzadehgan [Â](#) A, Dumas [Â](#) F, et al. [Â](#) Low rates of immediate coronary angiography among young adults resuscitated from sudden cardiac arrest. *Resuscitation* [Â](#) 2020;147:34[a](#)^{PADST}42.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

64Lloyd-Jones [Â](#) DM, Wilson [Â](#) PWF, Larson [Â](#) MG, Beiser [Â](#) A, Leip [Â](#) EP, D^Â^{PADST}Agostino [Â](#) RB, et al. [Â](#) Framingham risk score and prediction of lifetime risk for coronary heart disease. *Am J Cardiol* [Â](#) 2004;94:203[a](#)^{PADST}24.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

65Visseren [Â](#) FLJ, Mach [Â](#) F, Smulders [Â](#) YM, Carballo [Â](#) D, Koskinas [Â](#) KC, B^Â^{PADST}ock [Â](#) M, et al. [Â](#) 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* [Â](#) 2021;42:3227[a](#)^{PADST}3337.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

66Jouven [Â](#) X, Desnos [Â](#) M, Guerot [Â](#) C, Ducimet^Â^{PADST}re [Â](#) P. Predicting sudden death in the population: the Paris Prospective Study I. *Circulation* [Â](#) 1999;99:1978[a](#)^{PADST}1983.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

67Dekker [Â](#) LRC, Bezzina [Â](#) CR, Henriques [Â](#) JPS, Tanck [Â](#) MW, Koch [Â](#) KT, Alings [Â](#) MW, et al. [Â](#) Familial sudden death is an important risk factor for primary ventricular fibrillation: a case-control study in acute myocardial infarction patients. *Circulation* [Â](#) 2006;114:1140[a](#)^{PADST}1145.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

68Kaikkonen [Â](#) KS, Kortelainen [Â](#) M-L, Linna [Â](#) E, Huikuri [Â](#) HV. Family history and the risk of sudden cardiac death as a manifestation of an acute coronary event. *Circulation* [Â](#) 2006;114:1462[a](#)^{PADST}1467.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

69Jabbari [Â](#) R, Engstr^Â^{PADST}m [Â](#) T, Glinge [Â](#) C, Risgaard [Â](#) B, Jabbari [Â](#) J, Winkel [Â](#) BG, et al. [Â](#) Incidence and risk factors of ventricular fibrillation before primary angioplasty in patients with first ST-elevation myocardial infarction: a nationwide study in Denmark. *J Am Heart Assoc* [Â](#) 2015;4:e001399.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

70Friedlander [Â](#) Y, Siscovick [Â](#) DS, Weinmann [Â](#) S, Austin [Â](#) MA, Psaty [Â](#) BM, Lemaitre [Â](#) RN, et al. [Â](#) Family history as a risk factor for primary cardiac arrest. *Circulation* [Â](#) 1998;97:155[a](#)^{PADST}160.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

71Deo [Â](#) R, Norby [Â](#) FL, Katz [Â](#) R, Sotoodehnia [Â](#) N, Adabag [Â](#) S, DeFilippi [Â](#) CR, et al. [Â](#) Development and validation of a sudden cardiac death prediction model for the general population. *Circulation* [Â](#) 2016;134:806[a](#)^{PADST}816.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

- 72Aro $\hat{A}$ AL, Reinier $\hat{A}$ K, Rusinaru $\hat{A}$ C, Uy-Evanado $\hat{A}$ A, Darouian $\hat{A}$ N, Phan $\hat{A}$ D, et al. $\hat{A}$ Electrical risk score beyond the left ventricular ejection fraction: prediction of sudden cardiac death in the Oregon Sudden Unexpected Death Study and the Atherosclerosis Risk in Communities Study. *Eur Heart J* $\hat{A}$ 2017;38:3017a $\hat{A}$ _{PROSTIS}3025.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 73Waks $\hat{A}$ JW, Sitlani $\hat{A}$ CM, Soliman $\hat{A}$ EZ, Kabir $\hat{A}$ M, Ghafoori $\hat{A}$ E, Biggs $\hat{A}$ ML, et al. $\hat{A}$ Global electric heterogeneity risk score for prediction of sudden cardiac death in the general population: the Atherosclerosis Risk in Communities (ARIC) and Cardiovascular Health (CHS) studies. *Circulation* $\hat{A}$ 2016;133:2222a $\hat{A}$ _{PROSTIS}2234.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 74Yoshinaga $\hat{A}$ M, Ushinohama $\hat{A}$ H, Sato $\hat{A}$ S, Tauchi $\hat{A}$ N, Horigome $\hat{A}$ H, Takahashi $\hat{A}$ H, et al. $\hat{A}$ Electrocardiographic screening of 1-month-old infants for identifying prolonged QT intervals. *Circ Arrhythm Electrophysiol* $\hat{A}$ 2013;6:932a $\hat{A}$ _{PROSTIS}938.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 75Yoshinaga $\hat{A}$ M, Kucho $\hat{A}$ Y, Sarantuya $\hat{A}$ J, Ninomiya $\hat{A}$ Y, Horigome $\hat{A}$ H, Ushinohama $\hat{A}$ H, et al. $\hat{A}$ Genetic characteristics of children and adolescents with long-QT syndrome diagnosed by school-based electrocardiographic screening programs. *Circ Arrhythm Electrophysiol* $\hat{A}$ 2014;7:107a $\hat{A}$ _{PROSTIS}112.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 76J $\hat{A}$,rgensen $\hat{A}$ T, Jacobsen $\hat{A}$ RK, Toft $\hat{A}$ U, Aadahl $\hat{A}$ M, Gl $\hat{A}$ $\frac{1}{4}$ mer $\hat{A}$ C, Pisinger $\hat{A}$ C. Effect of screening and lifestyle counselling on incidence of ischaemic heart disease in general population: Inter99 randomised trial. *BMJ* $\hat{A}$ 2014;348:g3617.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 77Boas $\hat{A}$ R, Sappler $\hat{A}$ N, von St $\hat{A}$ $\frac{1}{4}$ lpnagel $\hat{A}$ L, Klemm $\hat{A}$ M, Diken $\hat{A}$ U, Thune $\hat{A}$ JJ, et al. $\hat{A}$ Periodic repolarization dynamics identifies ICD-responders in non-ischemic cardiomyopathy: a DANISH substudy. *Circulation* $\hat{A}$ 2022;145:754a $\hat{A}$ _{PROSTIS}764.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 78Spirito $\hat{A}$ P, Bellone $\hat{A}$ P, Harris $\hat{A}$ KM, Bernabo $\hat{A}$ P, Bruzzi $\hat{A}$ P, Maron $\hat{A}$ BJ. Magnitude of left ventricular hypertrophy and risk of sudden death in hypertrophic cardiomyopathy. *N Engl J Med* $\hat{A}$ 2000;342:1778a $\hat{A}$ _{PROSTIS}1785.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 79Barsheshet $\hat{A}$ A, Goldenberg $\hat{A}$ I, O-Uchi $\hat{A}$ J, Moss $\hat{A}$ AJ, Jons $\hat{A}$ C, Shimizu $\hat{A}$ W, et al. $\hat{A}$ Mutations in cytoplasmic loops of the KCNQ1 channel and the risk of life-threatening events: implications for mutation-specific response to $\hat{I}^2$ -blocker therapy in type 1 long-QT syndrome. *Circulation* $\hat{A}$ 2012;125:1988a $\hat{A}$ _{PROSTIS}1996.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 80Wahbi $\hat{A}$ K, Ben Yaou $\hat{A}$ R, Gandjbakhch $\hat{A}$ E, Anselme $\hat{A}$ F, Gossios $\hat{A}$ T, Lakdawala $\hat{A}$ NK, et al. $\hat{A}$ Development and validation of a new risk prediction score for life-threatening ventricular tachyarrhythmias in laminopathies. *Circulation* $\hat{A}$ 2019;140:293a $\hat{A}$ _{PROSTIS}302.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 81Cadrin-Tourigny $\hat{A}$ J, Bosman $\hat{A}$ LP, Nozza $\hat{A}$ A, Wang $\hat{A}$ W, Tadros $\hat{A}$ R, Bhonsale $\hat{A}$ A, et al. $\hat{A}$ A new prediction model for ventricular arrhythmias in arrhythmogenic right ventricular cardiomyopathy. *Eur Heart J* $\hat{A}$ 2019;40:1850a $\hat{A}$ _{PROSTIS}1858.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 82Priori $\hat{A}$ SG, Schwartz $\hat{A}$ PJ, Napolitano $\hat{A}$ C, Bloise $\hat{A}$ R, Ronchetti $\hat{A}$ E, Grillo $\hat{A}$ M, et al. $\hat{A}$ Risk stratification in the long-QT syndrome. *N Engl J Med* $\hat{A}$ 2003;348:1866a $\hat{A}$ _{PROSTIS}1874.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 83Miron $\hat{A}$ A, Lafreniere-Roula $\hat{A}$ M, Steve Fan $\hat{A}$ C-P, Armstrong $\hat{A}$ KR, Dragulescu $\hat{A}$ A, Papaz $\hat{A}$ T, et al. $\hat{A}$ A validated model for sudden cardiac death risk prediction in pediatric hypertrophic cardiomyopathy. *Circulation* $\hat{A}$ 2020;142:217a $\hat{A}$ _{PROSTIS}229.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 84Norrish $\hat{A}$ G, Ding $\hat{A}$ T, Field $\hat{A}$ E, Zi $\hat{A}$ $\frac{3}{4}$ lkowska $\hat{A}$ L, Olivotto $\hat{A}$ I, Limongelli $\hat{A}$ G, et al. $\hat{A}$ Development of a novel risk prediction model for sudden cardiac death in childhood hypertrophic cardiomyopathy (HCM risk-kids). *JAMA Cardiol* $\hat{A}$ 2019;4:918a $\hat{A}$ _{PROSTIS}927.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) $\hat{A}$
- 85O $\hat{A}$ _{PROSTIS}gciMahony $\hat{A}$ C, Jichi $\hat{A}$ F, Pavlou $\hat{A}$ M, Monserrat $\hat{A}$ L, Anastasakis $\hat{A}$ A, Rapezzi $\hat{A}$ C, et al. $\hat{A}$ A novel clinical risk prediction model for sudden cardiac death in hypertrophic cardiomyopathy (HCM risk-SCD). *Eur*

- Heart J 2014;35:2010â€¦2020.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 86Steyerberg Â EW, Vergouwe Â Y. Towards better clinical prediction models: seven steps for development and an ABCD for validation. Eur Heart J 2014;35:1925â€¦1931.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 87Collins Â GS, Reitsma Â JB, Altman Â DG, Moons Â KGM, TRIPOD Group. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD statement. Circulation 2015;131:211â€¦219.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 88Alba Â AC, Agoritsas Â T, Walsh Â M, Hanna Â S, Iorio Â A, Devereaux Â PJ, et al. Discrimination and calibration of clinical prediction models: users' guides to the medical literature. JAMA 2017;318:1377â€¦1384.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 89Yan Â S, Gan Â Y, Jiang Â N, Wang Â R, Chen Â Y, Luo Â Z, et al. The global survival rate among adult out-of-hospital cardiac arrest patients who received cardiopulmonary resuscitation: a systematic review and meta-analysis. Crit Care 2020;24:61.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 90Hallstrom Â AP, Ornato Â JP, Weisfeldt Â M, Travers Â A, Christenson Â J, McBurnie Â MA, et al. Public-access defibrillation and survival after out-of-hospital cardiac arrest. N Engl J Med 2004;351:637â€¦646.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 91Nakashima Â T, Noguchi Â T, Tahara Â Y, Nishimura Â K, Yasuda Â S, Onozuka Â D, et al. Public-access defibrillation and neurological outcomes in patients with out-of-hospital cardiac arrest in Japan: a population-based cohort study. Lancet 2019;394:2255â€¦2262.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 92Pollack Â RA, Brown Â SP, Rea Â T, Aufderheide Â T, Barbic Â D, Buick Â JE, et al. Impact of bystander automated external defibrillator use on survival and functional outcomes in shockable observed public cardiac arrests. Circulation 2018;137:2104â€¦2113.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 93Kragholm Â K, Wissenberg Â M, Mortensen Â RN, Hansen Â SM, Malta Hansen Â C, Thorsteinsson Â K, et al. Bystander efforts and 1-year outcomes in out-of-hospital cardiac arrest. N Engl J Med 2017;376:1737â€¦1747.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 94Kitamura Â T, Kiyohara Â K, Sakai Â T, Matsuyama Â T, Hatakeyama Â T, Shimamoto Â T, et al. Public-access defibrillation and out-of-hospital cardiac arrest in Japan. N Engl J Med 2016;375:1649â€¦1659.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 95Hasselqvist-Ax Â I, Riva Â G, Herlitz Â J, Rosenqvist Â M, Hollenberg Â J, Nordberg Â P, et al. Early cardiopulmonary resuscitation in out-of-hospital cardiac arrest. N Engl J Med 2015;372:2307â€¦2315.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 96Marijon Â E, Bougouin Â W, Celermajer Â DS, Perier Â M-C, Benameur Â N, Lamhaut Â L, et al. Major regional disparities in outcomes after sudden cardiac arrest during sports. Eur Heart J 2013;34:3632â€¦3640.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 97Karam Â N, Narayanan Â K, Bougouin Â W, Benameur Â N, Beganton Â F, Jost Â D, et al. Major regional differences in Automated External Defibrillator placement and Basic Life Support training in France: further needs for coordinated implementation. Resuscitation 2017;118:49â€¦54.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 98Mpotos Â N, De Wever Â B, Cleymans Â N, Raemaekers Â J, Loeys Â T, Herregods Â L, et al. Repetitive sessions of formative self-testing to refresh CPR skills: a randomised non-inferiority trial. Resuscitation 2014;85:1282â€¦1286.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 99Gundry Â JW, Comess Â KA, DeRook Â FA, Jorgenson Â D, Bardy Â GH. Comparison of naive sixth-grade children with trained professionals in the use of an automated external defibrillator. Circulation 1999;100:1703â€¦1707.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

100Blewer ^ AL, Putt ^ ME, Becker ^ LB, Riegel ^ BJ, Li ^ J, Leary ^ M, et al. ^ Video-only cardiopulmonary resuscitation education for high-risk families before hospital discharge: a multicenter pragmatic trial. *Circ Cardiovasc Qual Outcomes* ^ 2016;9:740a-748.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

101Ringh ^ M, Rosenqvist ^ M, Hollenberg ^ J, Jonsson ^ M, Fredman ^ D, Nordberg ^ P, et al. ^ Mobile-phone dispatch of laypersons for CPR in out-of-hospital cardiac arrest. *N Engl J Med* ^ 2015;372:2316a-2325.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

102Stroop ^ R, Kerner ^ T, Strickmann ^ B, Hensel ^ M. Mobile phone-based alerting of CPR-trained volunteers simultaneously with the ambulance can reduce the resuscitation-free interval and improve outcome after out-of-hospital cardiac arrest: a German, population-based cohort study. *Resuscitation* ^ 2020;147:57a-64.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

103Lee ^ SY, Shin ^ SD, Lee ^ YJ, Song ^ KJ, Hong ^ KJ, Ro ^ YS, et al. ^ Text message alert system and resuscitation outcomes after out-of-hospital cardiac arrest: a before-and-after population-based study. *Resuscitation* ^ 2019;138:198a-207.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

104Fordyce ^ CB, Hansen ^ CM, Kragholm ^ K, Dupre ^ ME, Jollis ^ JG, Roettig ^ ML, et al. ^ Association of public health initiatives with outcomes for out-of-hospital cardiac arrest at home and in public locations. *JAMA Cardiol* ^ 2017;2:1226a-1235.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

105Andelius ^ L, Malta Hansen ^ C, Lippert ^ FK, Karlsson ^ L, Torp-Pedersen ^ C, Kjær Ersbøll ^ A, et al. ^ Smartphone activation of citizen responders to facilitate defibrillation in out-of-hospital cardiac arrest. *J Am Coll Cardiol* ^ 2020;76:43a-53.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

106Ranthe ^ MF, Winkel ^ BG, Andersen ^ EW, Risgaard ^ B, Wohlfahrt ^ J, Bundgaard ^ H, et al. ^ Risk of cardiovascular disease in family members of young sudden cardiac death victims. *Eur Heart J* ^ 2013;34:503a-511.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

107Patton ^ KK, Sotoodehnia ^ N, DeFilippi ^ C, Siscovick ^ DS, Gottdiener ^ JS, Kronmal ^ RA. N-terminal pro-B-type natriuretic peptide is associated with sudden cardiac death risk: the Cardiovascular Health Study. *Heart Rhythm* ^ 2011;8:228a-233.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

108Everett ^ BM, Moorthy ^ MV, Tikkanen ^ JT, Cook ^ NR, Albert ^ CM. Markers of myocardial stress, myocardial injury, and subclinical inflammation and the risk of sudden death. *Circulation* ^ 2020;142:1148a-1158.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

109Berger ^ R, Huelsman ^ M, Strecker ^ K, Bojic ^ A, Moser ^ P, Stanek ^ B, et al. ^ B-type natriuretic peptide predicts sudden death in patients with chronic heart failure. *Circulation* ^ 2002;105:2392a-2397.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

110Scott ^ PA, Barry ^ J, Roberts ^ PR, Morgan ^ JM. Brain natriuretic peptide for the prediction of sudden cardiac death and ventricular arrhythmias: a meta-analysis. *Eur J Heart Fail* ^ 2009;11:958a-966.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

111Ahmad ^ T, Fiuzat ^ M, Neely ^ B, Neely ^ ML, Pencina ^ MJ, Kraus ^ WE, et al. ^ Biomarkers of myocardial stress and fibrosis as predictors of mode of death in patients with chronic heart failure. *JACC Heart Fail* ^ 2014;2:260a-268.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

112Goldberger ^ JJ, Cain ^ ME, Hohnloser ^ SH, Kadish ^ AH, Knight ^ BP, Lauer ^ MS, et al. ^ American Heart Association/American College of Cardiology Foundation/Heart Rhythm Society scientific statement on noninvasive risk stratification techniques for identifying patients at risk for sudden cardiac death: a scientific statement from the American Heart Association Council on Clinical Cardiology Committee on Electrocardiography and Arrhythmias and Council on Epidemiology and Prevention. *Circulation* ^ 2008;118:1497a-1518.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) ^

- 113Varma Â N, Cygankiewicz Â I, Turakhia Â MP, Heidebuchel Â H, Hu Â Y-F, Chen Â LY, et al. Â 2021
ISHNE/HRS/EHRA/APHR expert collaborative statement on mHealth in arrhythmia management: digital
medical tools for heart rhythm professionals: from the international society for holter and noninvasive
electrocardiology/heart rhythm society/European heart rhythm association/Asia-Pacific heart rhythm society.
Circ Arrhythm Electrophysiol Â 2021;14:e009204.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 114Solbiati Â M, Casazza Â G, Dipaola Â F, Barbic Â F, Caldato Â M, Montano Â N, et al. Â The diagnostic
yield of implantable loop recorders in unexplained syncope: a systematic review and meta-analysis. Int J Cardiol
Â 2017;231:170â176.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 115Gatzoulis Â KA, Arsenos Â P, Trachanas Â K, Dilaveris Â P, Antoniou Â C, Tsiachris Â D, et al. Â Signal-
averaged electrocardiography: past, present, and future. J Arrhythmia Â 2018;34:222â229.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 116Marcus Â FI, McKenna Â WJ, Sherrill Â D, Basso Â C, Bauce Â B, Bluemke Â DA, et al. Â Diagnosis of
arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the task force criteria. Eur
Heart J Â 2010;31:806â814.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 117Giudicessi Â JR, Ackerman Â MJ. Exercise testing oversights underlie missed and delayed diagnosis of
catecholaminergic polymorphic ventricular tachycardia in young sudden cardiac arrest survivors. Heart Rhythm
Â 2019;16:1232â1239.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 118Roston Â TM, Kallas Â D, Davies Â B, Franciosi Â S, De Souza Â AM, Laksman Â ZW, et al. Â Burst
exercise testing can unmask arrhythmias in patients with incompletely penetrant catecholaminergic polymorphic
ventricular tachycardia. JACC Clin Electrophysiol Â 2021;7:437â441.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 119Sy Â RW, van der Werf Â C, Chattha Â IS, Chockalingam Â P, Adler Â A, Healey Â JS, et al. Â Derivation
and validation of a simple exercise-based algorithm for prediction of genetic testing in relatives of LQTS
probands. Circulation Â 2011;124:2187â2194.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 120Binder Â T. Transthoracic echocardiography and the standard examination of specific cardiac structures. ESC
CardioMed. 3rd ed. Oxford University Press; 2022. p425â431.
- 121Haugaa Â KH, Basso Â C, Badano Â LP, Bucciarelli-Ducci Â C, Cardim Â N, Gaemperli Â O, et al.
Â Comprehensive multi-modality imaging approach in arrhythmogenic cardiomyopathy: an expert
consensus document of the European Association of Cardiovascular Imaging. Eur Heart J Cardiovasc Imaging
Â 2017;18:237â253.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 122Haugaa Â KH, Grenne Â BL, Eek Â CH, Ersboll Â M, Valeur Â N, Svendsen Â JH, et al. Â Strain
echocardiography improves risk prediction of ventricular arrhythmias after myocardial infarction. JACC
Cardiovasc Imaging Â 2013;6:841â850.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 123Haugaa Â KH, Hasselberg Â NE, Edvardsen Â T. Mechanical dispersion by strain echocardiography: a predictor
of ventricular arrhythmias in subjects with lamin A/C mutations. JACC Cardiovasc Imaging
Â 2015;8:1043â106.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 124Ersboll Â M, Valeur Â N, Andersen Â MJ, Mogensen Â UM, Vinther Â M, Svendsen Â JH, et al. Â Early
echocardiographic deformation analysis for the prediction of sudden cardiac death and life-threatening
arrhythmias after myocardial infarction. JACC Cardiovasc Imaging Â 2013;6:851â860.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 125Biering-Sørensen Â T, Knappe Â D, Pouleur Â A-C, Claggett Â B, Wang Â PJ, Moss Â AJ, et al.
Â Regional longitudinal deformation improves prediction of ventricular tachyarrhythmias in patients with heart
failure with reduced ejection fraction: a MADIT-CRT substudy (Multicenter Automatic Defibrillator
Implantation Trial-Cardiac Resynchronization Therapy). Circ Cardiovasc Imaging Â 2017;10:e005096.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 126Garg R, Friedrich MG. Cardiac myopathies. ESC CardioMed. 3rd ed. Oxford University Press; 2022, p510a-515.
- 127Borgquist R, Haugaa KH, Gilljam T, Bundgaard H, Hansen J, Eschen O, et al. The diagnostic performance of imaging methods in ARVC using the 2010 task force criteria. Eur Heart J Cardiovasc Imaging. 2014;15:1219a-1225.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 128Elliott PM, Anastakis A, Borger MA, Borggrefe M, Cecchi F, Charron P, et al. 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: the task force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). Eur Heart J. 2014;35:2733a-2779.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 129Di Marco A, Anguera I, Schmitt M, Klem I, Neilan TG, White JA, et al. Late gadolinium enhancement and the risk for ventricular arrhythmias or sudden death in dilated cardiomyopathy: systematic review and meta-analysis. JACC Heart Fail. 2017;5:28a-38.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 130Dejaard LA, Skjellvåg ET, Lie A, Helle-Strøm M, Stokke MK, Hegbom F, et al. The mitral annulus disjunction arrhythmic syndrome. J Am Coll Cardiol. 2018;72:1600a-1609.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 131Basso C, Iliceto S, Thiene G, Perazzolo Marra M. Mitral valve prolapse, ventricular arrhythmias, and sudden death. Circulation. 2019;140:952a-964.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 132Achenbach S. Technology of cardiac computed tomography. ESC CardioMed. 3rd ed. Oxford University Press; 2022, p537a-541.
- 133Meijboom WB, van Mieghem CAG, Mollet NR, Pugliese F, Weustink AC, van Pelt N, et al. 64-Slice computed tomography coronary angiography in patients with high, intermediate, or low pretest probability of significant coronary artery disease. J Am Coll Cardiol. 2007;50:1469a-1475.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 134Douglas PS, Hoffmann U, Patel MR, Mark DB, Al-Khalidi HR, Cavanaugh B, et al. Outcomes of anatomical versus functional testing for coronary artery disease. N Engl J Med. 2015;372:1291a-1300.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 135Priori SG, Wilde AA, Horie M, Cho Y, Behr ER, Berul C, et al. Executive summary: HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes. Europace. 2013;15:1389a-1406.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 136Govindan M, Batchvarov VN, Raju H, Shanmugam N, Bizrah M, Bastiaenen R, et al. Utility of high and standard right precordial leads during ajmaline testing for the diagnosis of Brugada syndrome. Heart. 2010;96:1904a-1908.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 137Churet M, Luttuo K, Hocini M, Haïssaguerre M, Sacher F, Duchateau J. Diagnostic reproducibility of epinephrine drug challenge interpretation in suspected long QT syndrome. J Cardiovasc Electrophysiol. 2019;30:896a-901.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 138Rolf S. The ajmaline challenge in Brugada syndrome: diagnostic impact, safety, and recommended protocol. Eur Heart J. 2003;24:1104a-1112.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 139Tadros R, Nannenberg EA, Lieve KV, Korić M, Milosavljević D, Lahrouchi N, Lekanne Deprez RH, et al. Yield and pitfalls of ajmaline testing in the evaluation of unexplained cardiac arrest and sudden unexplained death: single-center experience with 482 families. JACC Clin Electrophysiol. 2017;3:1400a-1408.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 140Meregalli PG, Ruijter JM, Hofman N, Bezzina CR, Wilde AAM, Tan HL. Diagnostic value of flecainide testing in unmasking SCN5A-related Brugada syndrome. J Cardiovasc Electrophysiol.

Â 2006;17:857â864.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

141Krahn Â AD, Healey Â JS, Chauhan Â VS, Birnie Â DH, Champagne Â J, Sanatani Â S, et al. Â Epinephrine infusion in the evaluation of unexplained cardiac arrest and familial sudden death: from the cardiac arrest survivors with preserved Ejection Fraction Registry. *Circ Arrhythm Electrophysiol* Â 2012;5:933â940.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

142Sueda Â S, Kohno Â H, Ochi Â T, Uraoka Â T. Overview of the acetylcholine spasm provocation test. *Clin Cardiol* Â 2015;38:430â438.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

143Takagi Â Y, Yasuda Â S, Takahashi Â J, Tsunoda Â R, Ogata Â Y, Seki Â A, et al. Â Clinical implications of provocation tests for coronary artery spasm: safety, arrhythmic complications, and prognostic impact: multicentre registry study of the Japanese Coronary Spasm Association. *Eur Heart J* Â 2013;34:258â267.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

144Garratt Â CJ, Antoniou Â A, Griffith Â MJ, Ward Â DE, Camm Â AJ. Use of intravenous adenosine in sinus rhythm as a diagnostic test for latent preexcitation. *Am J Cardiol* Â 1990;65:868â873.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

145Bourke Â JP, Richards Â DA, Ross Â DL, Wallace Â EM, McGuire Â MA, Uther Â JB. Routine programmed electrical stimulation in survivors of acute myocardial infarction for prediction of spontaneous ventricular tachyarrhythmias during follow-up: results, optimal stimulation protocol and cost-effective screening. *J Am Coll Cardiol* Â 1991;18:780â788.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

146Gatzoulis Â KA, Tsiachris Â D, Arsenos Â P, Archontakis Â S, Dilaveris Â P, Voulitis Â A, et al. Â Prognostic value of programmed ventricular stimulation for sudden death in selected high risk patients with structural heart disease and preserved systolic function. *Int J Cardiol* Â 2014;176:1449â1451.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

147Gatzoulis Â KA, Voulitis Â A-I, Tsiachris Â D, Salourou Â M, Archontakis Â S, Dilaveris Â P, et al. Â Primary prevention of sudden cardiac death in a nonischemic dilated cardiomyopathy population: reappraisal of the role of programmed ventricular stimulation. *Circ Arrhythm Electrophysiol* Â 2013;6:504â512.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

148Brilakis Â ES, Shen Â WK, Hammill Â SC, Hodge Â DO, Rea Â RF, Lexvold Â NY, et al. Â Role of programmed ventricular stimulation and implantable cardioverter defibrillators in patients with idiopathic dilated cardiomyopathy and syncope. *Pacing Clin Electrophysiol* Â 2001;24:1623â1630.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

149Brembilla-Perrot Â B, Suty-Selton Â C, Beurrier Â D, Houriez Â P, Nippert Â M, de la Chaise Â AT, et al. Â Differences in mechanisms and outcomes of syncope in patients with coronary disease or idiopathic left ventricular dysfunction as assessed by electrophysiologic testing. *J Am Coll Cardiol* Â 2004;44:594â601.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

150Schmitt Â C, Barthel Â P, Ndrepepa Â G, Schrieck Â J, Plewan Â A, SchÃ¶mig Â A, et al. Â Value of programmed ventricular stimulation for prophylactic internal cardioverter-defibrillator implantation in postinfarction patients preselected by noninvasive risk stratifiers. *J Am Coll Cardiol* Â 2001;37:1901â1907.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

151Gatzoulis Â KA, Tsiachris Â D, Arsenos Â P, Antoniou Â C-K, Dilaveris Â P, Sideris Â S, et al. Â Arrhythmic risk stratification in post-myocardial infarction patients with preserved ejection fraction: the PRESERVE EF study. *Eur Heart J* Â 2019;40:2940â2949.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

152Denis Â A, Sacher Â F, Derval Â N, Lim Â HS, Cochet Â H, Shah Â AJ, et al. Â Diagnostic value of isoproterenol testing in arrhythmogenic right ventricular cardiomyopathy. *Circ Arrhythm Electrophysiol* Â 2014;7:590â597.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

153Merino Â JL, Carmona Â JR, FernÃ¡ndez-Lozano Â I, Peinado Â R, Basterra Â N, Sobrino Â JA. Mechanisms of sustained ventricular tachycardia in myotonic dystrophy: implications for catheter ablation. *Circulation* Â 1998;98:541â546.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 154Pezawas Â T, Stix Â G, Kastner Â J, Wolzt Â M, Mayer Â C, Moertl Â D, et al. Â Unexplained syncope in patients with structural heart disease and no documented ventricular arrhythmias: value of electrophysiologically guided implantable cardioverter defibrillator therapy. *Europace* Â 2003;5:305â312.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 155Sroubek Â J, Probst Â V, Mazzanti Â A, Delise Â P, Hevia Â JC, Ohkubo Â K, et al. Â Programmed ventricular stimulation for risk stratification in the Brugada syndrome: a pooled analysis. *Circulation* Â 2016;133:622â630.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 156Nayyar Â S, Downar Â E, Bhaskaran Â AP, MassÃ© Â S, Nanthakumar Â K. Signature signal strategy: electrogram-based ventricular tachycardia mapping. *Heart Rhythm* Â 2020;17:2000â2009.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 157Lazarus Â A, Varin Â J, Babuty Â D, Anselme Â F, Coste Â J, Duboc Â D. Long-term follow-up of arrhythmias in patients with myotonic dystrophy treated by pacing: a multicenter diagnostic pacemaker study. *J Am Coll Cardiol* Â 2002;40:1645â1652.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 158Tandri Â H, Asimaki Â A, Abraham Â T, Dalal Â D, Tops Â L, Jain Â R, et al. Â Prolonged RV endocardial activation duration: a novel marker of arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Heart Rhythm* Â 2009;6:769â775.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 159Gomes Â J, Finlay Â M, Ahmed Â AK, Ciaccio Â EJ, Asimaki Â A, Saffitz Â JE, et al. Â Electrophysiological abnormalities precede overt structural changes in arrhythmogenic right ventricular cardiomyopathy due to mutations in desmoplakin-A combined murine and human study. *Eur Heart J* Â 2012;33:1942â1953.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 160Orini Â M, Graham Â AJ, Srinivasan Â NT, Campos Â FO, Hanson Â BM, Chow Â A, et al. Â Evaluation of the reentry vulnerability index to predict ventricular tachycardia circuits using high-density contact mapping. *Heart Rhythm* Â 2020;17:576â583.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 161Jackson Â N, Gizurason Â S, Viswanathan Â K, King Â B, MassÃ© Â S, Kusha Â M, et al. Â Decrement evoked potential mapping: basis of a mechanistic strategy for ventricular tachycardia ablation. *Circ Arrhythm Electrophysiol* Â 2015;8:1433â1442.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 162Avella Â A, d'Amati Â G, Pappalardo Â A, Re Â F, Silenzi Â PF, Laurenzi Â F, et al. Â Diagnostic value of endomyocardial biopsy guided by electroanatomic voltage mapping in arrhythmogenic right ventricular cardiomyopathy/dysplasia. *J Cardiovasc Electrophysiol* Â 2008;19:1127â1134.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 163Wilde Â AAM, Behr Â ER. Genetic testing for inherited cardiac disease. *Nat Rev Cardiol* Â 2013;10:571â583.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 164Hosseini Â SM, Kim Â R, Udupa Â S, Costain Â G, Jobling Â R, Liston Â E, et al. Â Reappraisal of reported genes for sudden arrhythmic death: evidence-based evaluation of gene validity for Brugada syndrome. *Circulation* Â 2018;138:1195â1205.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 165Ingles Â J, Goldstein Â J, Thaxton Â C, Caleshu Â C, Corty Â EW, Crowley Â SB, et al. Â Evaluating the clinical validity of hypertrophic cardiomyopathy genes. *Circ Genom Precis Med* Â 2019;12:e002460.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 166Adler Â A, Novelli Â V, Amin Â AS, Abiusi Â E, Care Â M, Nannenberg Â EA, et al. Â An international, multicentered, evidence-based reappraisal of genes reported to cause congenital long QT syndrome. *Circulation* Â 2020;141:418â428.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 167Risgaard Â B, Jabbari Â R, Refsgaard Â L, Holst Â AG, HaunsÃ¸ Â S, Sadjadieh Â A, et al. Â High prevalence of genetic variants previously associated with Brugada syndrome in new exome data. *Clin Genet* Â 2013;84:489â495.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 168James Â CA, Jongbloed Â JDH, Hershberger Â RE, Morales Â A, Judge Â DP, Syrris Â P, et al. Â An international evidence based reappraisal of genes associated with arrhythmogenic right ventricular cardiomyopathy (ARVC) using the ClinGen framework. *Circ Genom Precis Med* Â 2021;14:e003273.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 169Lahrouchi Â N, Tadros Â R, Crotti Â L, Mizusawa Â Y, Postema Â PG, Beekman Â L, et al. Â Transethnic genome-wide association study provides insights in the genetic architecture and heritability of long QT syndrome. *Circulation* Â 2020;142:324â338.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 170Wijeyeratne Â YD, Tanck Â MW, Mizusawa Â Y, Batchvarov Â V, Barc Â J, Crotti Â L, et al. Â SCN5A mutation type and a genetic risk score associate variably with Brugada syndrome phenotype in SCN5A families. *Circ Genom Precis Med* Â 2020;13:e002911.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 171Harper Â AR, Goel Â A, Grace Â C, Thomson Â KL, Petersen Â SE, Xu Â X, et al. Â Common genetic variants and modifiable risk factors underpin hypertrophic cardiomyopathy susceptibility and expressivity. *Nat Genet* Â 2021;53:135â142.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 172Tadros Â R, Francis Â C, Xu Â X, Vermeer Â AMC, Harper Â AR, Huurman Â R, et al. Â Shared genetic pathways contribute to risk of hypertrophic and dilated cardiomyopathies with opposite directions of effect. *Nat Genet* Â 2021;53:128â134.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 173Tadros Â R, Tan Â HL, ESCAPE-NET Investigators, El Mathari Â S, Kors Â JA, Postema Â PG, et al. Â Predicting cardiac electrical response to sodium-channel blockade and Brugada syndrome using polygenic risk scores. *Eur Heart J* Â 2019;40:3097â3107.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 174Ware Â JS, Roberts Â AM, Cook Â SA. Next generation sequencing for clinical diagnostics and personalised medicine: implications for the next generation cardiologist. *Heart* Â 2012;98:276â281.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 175Ashar Â FN, Mitchell Â RN, Albert Â CM, Newton-Cheh Â C, Brody Â JA, MÃ¼ller-Nurasyid Â M, et al. Â A comprehensive evaluation of the genetic architecture of sudden cardiac arrest. *Eur Heart J* Â 2018;39:3961â3969.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 176Richards Â S, Aziz Â N, Bale Â S, Bick Â D, Das Â S, Gastier-Foster Â J, et al. Â Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* Â 2015;17:405â424.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 177Ingles Â J, Lind Â JM, Phongsavan Â P, Semsarian Â C. Psychosocial impact of specialized cardiac genetic clinics for hypertrophic cardiomyopathy. *Genet Med* Â 2008;10:117â120.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 178Ackerman Â MJ, Priori Â SG, Willems Â S, Berul Â C, Brugada Â R, Calkins Â H, et al. Â HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies: this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Europace* Â 2011;13:1077â1109.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 179Conte Â G, Wilde Â A, Behr Â ER, Scherr Â D, Lenarczyk Â R, Gandjbakhkh Â E, et al. Â Importance of dedicated units for the management of patients with inherited arrhythmia syndromes. *Circ Genom Precis Med* Â 2021;14:e003313.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 180Minier Â M, Probst Â V, Berthome Â P, Tixier Â R, Briand Â J, Geoffroy Â O, et al. Â Age at diagnosis of Brugada syndrome: influence on clinical characteristics and risk of arrhythmia. *Heart Rhythm* Â 2020;17:743â749.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 181Wong Â LCH, Roses-Noguer Â F, Till Â JA, Behr Â ER. Cardiac evaluation of pediatric relatives in sudden arrhythmic death syndrome: a 2-center experience. *Circ Arrhythm Electrophysiol* Â 2014;7:800â806.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

182Mellor [Â](#) GJ, Blom [Â](#) LJ, Groeneveld [Â](#) SA, Winkel [Â](#) BG, Ensam [Â](#) B, Bargehr [Â](#) J, et al. [Â](#) Familial evaluation in idiopathic ventricular fibrillation: diagnostic yield and significance of J Wave Syndromes. *Circ Arrhythm Electrophysiol* [Â](#) 2021;14:e009089.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

183Lahrouchi [Â](#) N, Raju [Â](#) H, Lodder [Â](#) EM, Papatheodorou [Â](#) E, Ware [Â](#) JS, Papadakis [Â](#) M, et al. [Â](#) Utility of post-mortem genetic testing in cases of sudden arrhythmic death syndrome. *J Am Coll Cardiol* [Â](#) 2017;69:2134a-2145.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

184Faber [Â](#) TS, Gradingier [Â](#) R, Treusch [Â](#) S, Morkel [Â](#) C, Brachmann [Â](#) J, Bode [Â](#) C, et al. [Â](#) Incidence of ventricular tachyarrhythmias during permanent pacemaker therapy in low-risk patients results from the German multicentre EVENTS study. *Eur Heart J* [Â](#) 2007;28:2238a-2242.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

185Middlekauff [Â](#) HR, Stevenson [Â](#) WG, Stevenson [Â](#) LW, Saxon [Â](#) LA. Syncope in advanced heart failure: high risk of sudden death regardless of origin of syncope. *J Am Coll Cardiol* [Â](#) 1993;21:110a-116.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

186Noda [Â](#) T, Shimizu [Â](#) W, Taguchi [Â](#) A, Aiba [Â](#) T, Satomi [Â](#) K, Suyama [Â](#) K, et al. [Â](#) Malignant entity of idiopathic ventricular fibrillation and polymorphic ventricular tachycardia initiated by premature extrasystoles originating from the right ventricular outflow tract. *J Am Coll Cardiol* [Â](#) 2005;46:1288a-1294.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

187Belhassen [Â](#) B, Viskin [Â](#) S. Idiopathic ventricular tachycardia and fibrillation. *J Cardiovasc Electrophysiol* [Â](#) 1993;4:356a-368.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

188Zareba [Â](#) W, Platonov [Â](#) P. ECG patterns related to arrhythmias and sudden death: channelopathies, early repolarization, and pre-excitation. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p382a-389.

189Voigt [Â](#) J-U. Left ventricular function, heart failure, and resynchronization therapy. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p450a-454.

190Edvardsen [Â](#) T. Cardiomyopathies, myocarditis, and the transplanted heart. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p456a-460.

191Cygankiewicz [Â](#) I. Ambulatory ECG monitoring. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p393a-400.

192Gray [Â](#) B, Kirby [Â](#) A, Kabunga [Â](#) P, Freedman [Â](#) SB, Yeates [Â](#) L, Kanthan [Â](#) A, et al. [Â](#) Twelve-lead ambulatory electrocardiographic monitoring in Brugada syndrome: potential diagnostic and prognostic implications. *Heart Rhythm* [Â](#) 2017;14:866a-874.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

193Karogiannis [Â](#) N, Senior [Â](#) R. Stress echocardiography. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p431a-434.

194Nucifora [Â](#) G, Muser [Â](#) D, Masci [Â](#) PG, Barison [Â](#) A, Rebellato [Â](#) L, Piccoli [Â](#) G, et al. [Â](#) Prevalence and prognostic value of concealed structural abnormalities in patients with apparently idiopathic ventricular arrhythmias of left versus right ventricular origin: a magnetic resonance imaging study. *Circ Arrhythm Electrophysiol* [Â](#) 2014;7:456a-462.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

195Muser [Â](#) D, Nucifora [Â](#) G, Muser [Â](#) D, Nucifora [Â](#) G, Pieroni [Â](#) M, Castro [Â](#) SA, et al. [Â](#) Prognostic value of nonischemic ringlike left ventricular scar in patients with apparently idiopathic nonsustained ventricular arrhythmias. *Circulation* [Â](#) 2021;143:1359a-1373.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

196Josephson [Â](#) ME, Callans [Â](#) DJ. Using the twelve-lead electrocardiogram to localize the site of origin of ventricular tachycardia. *Heart Rhythm* [Â](#) 2005;2:443a-446.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

197Corrado [Â](#) D, Basso [Â](#) C, Leoni [Â](#) L, Tokajuk [Â](#) B, Baucé [Â](#) B, Frigo [Â](#) G, et al. [Â](#) Three-dimensional electroanatomic voltage mapping increases accuracy of diagnosing arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Circulation* [Â](#) 2005;111:3042a-3050.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

198Casella Â M, Dello Russo Â A, Bergonti Â M, Catto Â V, Conte Â E, Sommariva Â E, et al. Â Diagnostic yield of electroanatomic voltage mapping in guiding endomyocardial biopsies. *Circulation* Â 2020;142:1249â1260.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

199Haanschoten Â DM, Adiyaman Â A, [PROSTIS](#) Hart Â NA, Jager Â PL, Elvan Â A. Value of 3D mapping-guided endomyocardial biopsy in cardiac sarcoidosis: case series and narrative review on the value of electro-anatomic mapping-guided endomyocardial biopsies. *Eur J Clin Invest* Â 2021;51:e13497.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

200Tung Â R, Bauer Â B, Schelbert Â H, Lynch Â JP, Auerbach Â M, Gupta Â P, et al. Â Incidence of abnormal positron emission tomography in patients with unexplained cardiomyopathy and ventricular arrhythmias: the potential role of occult inflammation in arrhythmogenesis. *Heart Rhythm* Â 2015;12:2488â2498.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

201Palmisano Â A, Vignale Â D, Peretto Â G, Busnardo Â E, Calcagno Â C, Campochiaro Â C, et al. Â Hybrid FDG-PET/MR or FDG-PET/CT to detect disease activity in patients with persisting arrhythmias after myocarditis. *JACC Cardiovasc Imaging* Â 2021;14:288â292.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

202Venlet Â J, Piers Â SRD, Jongbloed Â JDH, Androulakis Â AFA, Naruse Â Y, den Uijl Â DW, et al. Â Isolated subepicardial right ventricular outflow tract scar in athletes with ventricular tachycardia. *J Am Coll Cardiol* Â 2017;69:497â507.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

203Geri Â G, Dumas Â F, Bougouin Â W, Varenne Â O, Daviaud Â F, PÃne Â F, et al. Â Immediate percutaneous coronary intervention is associated with improved short- and long-term survival after out-of-hospital cardiac arrest. *Circ Cardiovasc Interv* Â 2015;8:e002303.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

204Vyas Â A, Chan Â PS, Cram Â P, Nallamothu Â BK, McNally Â B, Girotra Â S. Early coronary angiography and survival after out-of-hospital cardiac arrest. *Circ Cardiovasc Interv* Â 2015;8:e002321.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

205Kalarus Â Z, Svendsen Â JH, Capodanno Â D, Dan Â G-A, De Maria Â E, Gorenek Â B, et al. Â Cardiac arrhythmias in the emergency settings of acute coronary syndrome and revascularization: an European Heart Rhythm Association (EHRA) consensus document, endorsed by the European Association of Percutaneous Cardiovascular Interventions (EAPCI), and European Acute Cardiovascular Care Association (ACCA). *Europace* Â 2019;21:1603â1604.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

206Noc Â M, Fajadet Â J, Lassen Â JF, Kala Â P, McCarthy Â P, Olivecrona Â GK, et al. Â Invasive coronary treatment strategies for out-of-hospital cardiac arrest: a consensus statement from the European association for percutaneous cardiovascular interventions (EAPCI)/stent for life (SFL) groups. *EuroIntervention* Â 2014;10:31â37.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

207Verma Â BR, Sharma Â V, Shekhar Â S, Kaur Â M, Khubber Â S, Bansal Â A, et al. Â Coronary angiography in patients with out-of-hospital cardiac arrest without ST-segment elevation: a systematic review and meta-analysis. *JACC Cardiovasc Interv* Â 2020;13:2193â2205.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

208Barbarawi Â M, Zayed Â Y, Kheiri Â B, Barbarawi Â O, Al-Abdoun Â A, Dhillon Â H, et al. Â Optimal timing of coronary intervention in patients resuscitated from cardiac arrest without ST-segment elevation myocardial infarction (NSTEMI): a systematic review and meta-analysis. *Resuscitation* Â 2019;144:137â144.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

209Meng-Chang Â Y, Meng-Jun Â W, Xiao-Yan Â X, Peng Â KL, Peng Â YG, RuRong Â W. Coronary angiography or not after cardiac arrest without ST segment elevation: a systematic review and meta-analysis. *Medicine (Baltimore)* Â 2020;99:e22197.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) [Â](#)

210Khan Â MS, Shah Â SMM, Mubashir Â A, Khan Â AR, Fatima Â K, Schenone Â AL, et al. Â Early coronary angiography in patients resuscitated from out of hospital cardiac arrest without ST-segment elevation: a

- systematic review and meta-analysis. *Resuscitation* 2017;121:127â–134.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 211Welsford Â M, Bossard Â M, Shortt Â C, Pritchard Â J, Natarajan Â MK, Belley-Côté Â EP. Does early coronary angiography improve survival after out-of-hospital cardiac arrest? A systematic review with meta-analysis. *Can J Cardiol* 2018;34:180â–194.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 212Chelly Â J, Mongardon Â N, Dumas Â F, Varenne Â O, Spaulding Â C, Vignaux Â O, et al. Â Benefit of an early and systematic imaging procedure after cardiac arrest: insights from the PROCAT (Parisian Region Out of Hospital Cardiac Arrest) registry. *Resuscitation* 2012;83:1444â–1450.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 213Stiles Â MK, Wilde Â AAM, Abrams Â DJ, Ackerman Â MJ, Albert Â CM, Behr Â ER, et al. Â 2020 APHRS/HRS expert consensus statement on the investigation of decedents with sudden unexplained death and patients with sudden cardiac arrest, and of their families. *Heart Rhythm* 2021;18:e1â–e50.
[OpenURL Placeholder Text](#) Â
- 214Tseng Â ZH, Olgin Â JE, Vittinghoff Â E, Ursell Â PC, Kim Â AS, Sporer Â K, et al. Â Prospective countywide surveillance and autopsy characterization of sudden cardiac death: POST SCD study. *Circulation* 2018;137:2689â–2700.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 215Fellmann Â F, van El Â CG, Charron Â P, Michaud Â K, Howard Â HC, Boers Â SN, et al. Â European recommendations integrating genetic testing into multidisciplinary management of sudden cardiac death. *Eur J Hum Genet* 2019;27:1763â–1773.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 216Yamamoto Â T, Takayama Â M, Sato Â N, Yodogawa Â K, Iwasaki Â Y-K, Kato Â K, et al. Â Inappropriate analyses of automated external defibrillators used during in-hospital ventricular fibrillation. *Circ J* 2008;72:679â–681.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 217Sweeney Â MO, Ruetz Â LL, Belk Â P, Mullen Â TJ, Johnson Â JW, Sheldon Â T. Bradycardia pacing-induced short-long-short sequences at the onset of ventricular tachyarrhythmias: a possible mechanism of proarrhythmia? *J Am Coll Cardiol* 2007;50:614â–622.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 218Tseng Â ZH, Hayward Â RM, Clark Â NM, Mulvanny Â CG, Colburn Â BJ, Ursell Â PC, et al. Â Sudden death in patients with cardiac implantable electronic devices. *JAMA Intern Med* 2015;175:1342â–1350.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 219Lacour Â P, Buschmann Â C, Storm Â C, Nee Â J, Parwani Â AS, Huemer Â M, et al. Â Cardiac implantable electronic device interrogation at forensic autopsy: an underestimated resource? *Circulation* 2018;137:2730â–2740.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 220Curcio Â A, Mazzanti Â A, Bloise Â R, Monteforte Â N, Indolfi Â C, Priori Â SG, et al. Â Clinical presentation and outcome of Brugada syndrome diagnosed with the new 2013 criteria. *J Cardiovasc Electrophysiol* 2016;27:937â–943.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 221Haïssaguerre Â M, Shah Â DC, Jaïs Â P, Shoda Â M, Kautzner Â J, Arentz Â T, et al. Â Role of Purkinje conducting system in triggering of idiopathic ventricular fibrillation. *Lancet* 2002;359:677â–678.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 222Krahn Â AD, Healey Â JS, Chauhan Â V, Birnie Â DH, Simpson Â CS, Champagne Â J, et al. Â Systematic assessment of patients with unexplained cardiac arrest: cardiac arrest survivors with preserved ejection fraction registry (CASPER). *Circulation* 2009;120:278â–285.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 223van der Werf Â C, Hofman Â N, Tan Â HL, van Dessel Â PF, Alders Â M, van der Wal Â AC, et al. Â Diagnostic yield in sudden unexplained death and aborted cardiac arrest in the young: the experience of a tertiary referral center in The Netherlands. *Heart Rhythm* 2010;7:1383â–1389.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 224Eckart Â RE, Scoville Â SL, Campbell Â CL, Shry Â EA, Stajduhar Â KC, Potter Â RN, et al. Â Sudden death in young adults: a 25-year review of autopsies in military recruits. *Ann Intern Med* Â 2004;141:829â834.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 225Kunadian Â V, Chieffo Â A, Camici Â PG, Berry Â C, Escaned Â J, Maas Â AHEM, et al. Â An EAPCI expert consensus document on ischaemia with non-obstructive coronary arteries in collaboration with European Society of Cardiology Working Group on Coronary Pathophysiology & Microcirculation Endorsed by Coronary Vasomotor Disorders International Study Group. *Eur Heart J* Â 2020;41:3504â3520.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 226Rodrigues Â P, Joshi Â A, Williams Â H, Westwood Â M, Petersen Â SE, Zemrak Â F, et al. Â Diagnosis and prognosis in sudden cardiac arrest survivors without coronary artery disease: utility of a clinical approach using cardiac magnetic resonance imaging. *Circ Cardiovasc Imaging* Â 2017;10:e006709.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 227White Â JA, Fine Â NM, Gula Â L, Yee Â R, Skanes Â A, Klein Â G, et al. Â Utility of cardiovascular magnetic resonance in identifying substrate for malignant ventricular arrhythmias. *Circ Cardiovasc Imaging* Â 2012;5:12â20.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 228Basso Â C, Perazzolo Marra Â M, Rizzo Â S, De Lazzari Â M, Giorgi Â B, Cipriani Â A, et al. Â Arrhythmic mitral valve prolapse and sudden cardiac death. *Circulation* Â 2015;132:556â566.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 229Obeyesekere Â MN, Klein Â GJ, Modi Â S, Leong-Sit Â P, Gula Â LJ, Yee Â R, et al. Â How to perform and interpret provocative testing for the diagnosis of Brugada syndrome, long-QT syndrome, and catecholaminergic polymorphic ventricular tachycardia. *Circ Arrhythm Electrophysiol* Â 2011;4:958â964.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 230Cheung Â CC, Mellor Â G, Deyell Â MW, Ensam Â B, Batchvarov Â V, Papadakis Â M, et al. Â Comparison of ajmaline and procainamide provocation tests in the diagnosis of Brugada syndrome. *JACC Clin Electrophysiol* Â 2019;5:504â512.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 231Antzelevitch Â C, Yan Â G-X, Ackerman Â MJ, Borggrefe Â M, Corrado Â D, Guo Â J, et al. Â J-Wave syndromes expert consensus conference report: emerging concepts and gaps in knowledge. *Europace* Â 2017;19:665â694.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 232Adler Â A, van der Werf Â C, Postema Â PG, Rosso Â R, Bhuiyan Â ZA, Kalman Â JM, et al. Â The phenomenon of âQT stunningâ: the abnormal QT prolongation provoked by standing persists even as the heart rate returns to normal in patients with long QT syndrome. *Heart Rhythm* Â 2012;9:901â908.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 233Dionne Â A, Fournier Â A, Dahdah Â N, Abrams Â D, Khairy Â P, Abadir Â S. Dynamic QT interval changes from supine to standing in healthy children. *Can J Cardiol* Â 2018;34:66â72.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 234Foo Â FS, Stiles Â MK, Heaven Â D. Unmasking latent preexcitation of a right-sided accessory pathway with intravenous adenosine after unexplained sudden cardiac arrest. *J Arrhythm* Â 2020;36:939â941.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 235Marjamaa Â A, Hiippala Â A, Arrhenius Â B, Lahtinen Â AM, Kontula Â K, Toivonen Â L, et al. Â Intravenous epinephrine infusion test in diagnosis of catecholaminergic polymorphic ventricular tachycardia. *J Cardiovasc Electrophysiol* Â 2012;23:194â199.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 236Ackerman Â MJ, Khositseth Â A, Tester Â DJ, Hejlik Â JB, Shen Â W-K, Porter Â CJ. Epinephrine-induced QT interval prolongation: a gene-specific paradoxical response in congenital long QT syndrome. *Mayo Clin Proc* Â 2002;77:413â421.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 237Shimizu Â W, Noda Â T, Takaki Â H, Nagaya Â N, Satomi Â K, Kurita Â T, et al. Â Diagnostic value of epinephrine test for genotyping LQT1, LQT2, and LQT3 forms of congenital long QT syndrome. *Heart Rhythm* Â 2004;1:276â283.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 238Magnano Â AR, Talathoti Â N, Hallur Â R, Bloomfield Â DM, Garan Â H. Sympathomimetic infusion and cardiac repolarization: the normative effects of epinephrine and isoproterenol in healthy subjects. *J Cardiovasc Electrophysiol* Â 2006;17:983â^{PADIST}989.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 239Vyas Â H, Hejlik Â J, Ackerman Â MJ. Epinephrine QT stress testing in the evaluation of congenital long-QT syndrome: diagnostic accuracy of the paradoxical QT response. *Circulation* Â 2006;113:1385â^{PADIST}1392.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 240Waldmann Â V, Bougouin Â W, Karam Â N, Narayanan Â K, Sharifzadehgan Â A, Spaulding Â C, et al. Â Coronary vasospasm-related sudden cardiac arrest in the community. *J Am Coll Cardiol* Â 2018;72:814â^{PADIST}815.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 241Etienne Â P, Huchet Â F, Gaborit Â N, Barc Â J, Thollet Â A, Kyndt Â F, et al. Â Mental stress test: a rapid, simple, and efficient test to unmask long QT syndrome. *Europace* Â 2018;20:2014â^{PADIST}2020.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 242Kop Â WJ, Krantz Â DS, Nearing Â BD, Gottdiener Â JS, Quigley Â JF, Oâ^{PADIST}Callahan Â M, et al. Â Effects of acute mental stress and exercise on T-wave alternans in patients with implantable cardioverter defibrillators and controls. *Circulation* Â 2004;109:1864â^{PADIST}1869.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 243Schwartz Â PJ, Lia Â C. QTc behavior during exercise and genetic testing for the long-QT syndrome. *Circulation* Â 2011;124:2181â^{PADIST}2184.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 244Wang Â YS, Scheinman Â MM, Chien Â WW, Cohen Â TJ, Lesh Â MD, Griffin Â JC. Patients with supraventricular tachycardia presenting with aborted sudden death: incidence, mechanism and long-term follow-up. *J Am Coll Cardiol* Â 1991;18:1711â^{PADIST}1719.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 245Roberts Â JD, Gollob Â MH, Young Â C, Connors Â SP, Gray Â C, Wilton Â SB, et al. Â Bundle branch re-entrant ventricular tachycardia: novel genetic mechanisms in a life-threatening arrhythmia. *JACC Clin Electrophysiol* Â 2017;3:276â^{PADIST}288.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 246Santangeli Â P, Hamilton-Craig Â C, Dello Russo Â A, Pieroni Â M, Casella Â M, Pelargonio Â G, et al. Â Imaging of scar in patients with ventricular arrhythmias of right ventricular origin: cardiac magnetic resonance versus electroanatomic mapping. *J Cardiovasc Electrophysiol* Â 2011;22:1359â^{PADIST}1366.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 247Haâ^{ss}aguerre Â M, Duchateau Â J, Dubois Â R, Hocini Â M, Cheniti Â G, Sacher Â F, et al. Â Idiopathic ventricular fibrillation: role of Purkinje system and microstructural myocardial abnormalities. *JACC Clin Electrophysiol* Â 2020;6:591â^{PADIST}608.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 248Haâ^{ss}aguerre Â M, Hocini Â M, Cheniti Â G, Duchateau Â J, Sacher Â F, Puyo Â S, et al. Â Localized structural alterations underlying a subset of unexplained sudden cardiac death. *Circ Arrhythm Electrophysiol* Â 2018;11:e006120.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 249Mellor Â G, Laksman Â ZWM, Tadros Â R, Roberts Â JD, Gerull Â B, Simpson Â CS, et al. Â Genetic testing in the evaluation of unexplained cardiac arrest: from the CASPER (Cardiac Arrest Survivors With Preserved Ejection Fraction Registry). *Circ Cardiovasc Genet* Â 2017;10:e001686.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 250Burns Â C, Bagnall Â RD, Lam Â L, Semsarian Â C, Ingles Â J. Multiple gene variants in hypertrophic cardiomyopathy in the era of next-generation sequencing. *Circ Cardiovasc Genet* Â 2017;10:e001666.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 251Karam Â N, Jabre Â P, Narayanan Â K, Sharifzadehgan Â A, Perier Â M-C, Tennenbaum Â J, et al. Â Psychological support and medical screening of first-degree relatives of sudden cardiac arrest victims. *JACC Clin Electrophysiol* Â 2020;6:586â^{PADIST}587.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 252Behr Â E, Wood Â DA, Wright Â M, Syrris Â P, Sheppard Â MN, Casey Â A, et al. Â Cardiological assessment of first-degree relatives in sudden arrhythmic death syndrome. *Lancet* Â 2003;362:1457â1459.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 253Tan Â HL, Hofman Â N, van Langen Â IM, van der Wal Â AC, Wilde Â AAM. Sudden unexplained death: heritability and diagnostic yield of cardiological and genetic examination in surviving relatives. *Circulation* Â 2005;112:207â213.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 254Lemkes Â JS, Janssens Â GN, van der Hoeven Â NW, Jewbali Â LSD, Dubois Â EA, Meuwissen Â M, et al. Â Coronary angiography after cardiac arrest without ST-segment elevation. *N Engl J Med* Â 2019;380:1397â1407.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 255Kern Â KB, Radsel Â P, Jentzer Â JC, Seder Â DB, Lee Â KS, Lotun Â K, et al. Â Randomized pilot clinical trial of early coronary angiography versus no early coronary angiography after cardiac arrest without ST-segment elevation: the PEARL study. *Circulation* Â 2020;142:2002â2012.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 256Desch Â S, Freund Â A, Akin Â I, Behnes Â M, Preusch Â MR, Zelniker Â TA, et al. Â Angiography after out-of-hospital cardiac arrest without ST-segment elevation. *N Engl J Med* Â 2021;385:2544â2553.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 257Arnaout Â M, Mongardon Â N, Deye Â N, Legriel Â S, Dumas Â F, Sauneuf Â B, et al. Â Out-of-hospital cardiac arrest from brain cause: epidemiology, clinical features, and outcome in a multicenter cohort. *Crit Care Med* Â 2015;43:453â460.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 258Horner Â JM, Horner Â MM, Ackerman Â MJ. The diagnostic utility of recovery phase QTc during treadmill exercise stress testing in the evaluation of long QT syndrome. *Heart Rhythm* Â 2011;8:1698â1704.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 259Perrin Â MJ, Angaran Â P, Laksman Â Z, Zhang Â H, Porepa Â LF, Rutberg Â J, et al. Â Exercise testing in asymptomatic gene carriers exposes a latent electrical substrate of arrhythmogenic right ventricular cardiomyopathy. *J Am Coll Cardiol* Â 2013;62:1772â1779.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 260Waldmann Â V, Bougouin Â W, Karam Â N, Dumas Â F, Sharifzadehgan Â A, Gandjbakhch Â E, et al. Â Characteristics and clinical assessment of unexplained sudden cardiac arrest in the real-world setting: focus on idiopathic ventricular fibrillation. *Eur Heart J* Â 2018;39:1981â1987.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 261Nakao Â K, Ohgushi Â M, Yoshimura Â M, Morooka Â K, Okumura Â K, Ogawa Â H, et al. Â Hyperventilation as a specific test for diagnosis of coronary artery spasm. *Am J Cardiol* Â 1997;80:545â549.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 262Behr Â ER, Casey Â A, Sheppard Â M, Wright Â M, Bowker Â TJ, Davies Â MJ, et al. Â Sudden arrhythmic death syndrome: a national survey of sudden unexplained cardiac death. *Heart* Â 2007;93:601â605.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 263Basso Â C, Aguilera Â B, Banner Â J, Cohle Â S, d'Amati Â G, de Gouveia Â RH, et al. Â Guidelines for autopsy investigation of sudden cardiac death: 2017 update from the Association for European Cardiovascular Pathology. *Virchows Arch* Â 2017;471:691â705.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 264de Noronha Â SV, Behr Â ER, Papadakis Â M, Ohta-Ogo Â K, Banya Â W, Wells Â J, et al. Â The importance of specialist cardiac histopathological examination in the investigation of young sudden cardiac deaths. *Europace* Â 2014;16:899â907.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 265Gulino Â SP, Burns Â K, Gunther Â WM, MacLeod Â H. Improving forensic pathologic investigation of sudden death in the young. *Acad Forensic Pathol* Â 2018;8:347â391.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 266Hansen Â BL, Jacobsen Â EM, Kjerrumgaard Â A, Tfelt-Hansen Â J, Winkel Â BG, Bundgaard Â H, et al. Â Diagnostic yield in victims of sudden cardiac death and their relatives. *Europace* Â 2020;22:964â971.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

267Bjune Â T, Risgaard Â B, Kruckow Â L, Glinge Â C, Ingemann-Hansen Â O, Leth Â PM, et al. Â Post-mortem toxicology in young sudden cardiac death victims: a nationwide cohort study. *Europace* Â 2018;20:614â621.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

268Allan Â KS, Morrison Â LJ, Pinter Â A, Tu Â JV, Dorian Â P, Rescu Epistery Investigators. Â Presumed cardiac arrest in children and young adults: a misnomer? Â *Resuscitation* Â 2017;117:73â79.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

269Lahrouchi Â N, Raju Â H, Lodder Â EM, Papatheodorou Â S, Miles Â C, Ware Â JS, et al. Â The yield of postmortem genetic testing in sudden death cases with structural findings at autopsy. *Eur J Hum Genet* Â 2020;28:17â22.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

270Tester Â DJ, Medeiros-Domingo Â A, Will Â ML, Haglund Â CM, Ackerman Â MJ. Cardiac channel molecular autopsy: insights from 173 consecutive cases of autopsy-negative sudden unexplained death referred for postmortem genetic testing. *Mayo Clin Proc* Â 2012;87:524â539.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

271Papadakis Â M, Raju Â H, Behr Â ER, De Noronha Â SV, Spath Â N, Kouloubinis Â A, et al. Â Sudden cardiac death with autopsy findings of uncertain significance: potential for erroneous interpretation. *Circ Arrhythm Electrophysiol* Â 2013;6:588â596.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

272Jacobsen Â EM, Hansen Â BL, Kjerrumgaard Â A, Tfelt-Hansen Â J, Hassager Â C, Kjaergaard Â J, et al. Â Diagnostic yield and long-term outcome of nonischemic sudden cardiac arrest survivors and their relatives: results from a tertiary referral center. *Heart Rhythm* Â 2020;17:1679â1686.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

273Kjerrumgaard Â A, Jacobsen Â EM, Hansen Â BL, Tfelt-Hansen Â J, Winkel Â BG, Christensen Â AH, et al. Â Diagnostic findings and follow-up outcomes in relatives to young non-autopsied sudden death victims. *Int J Cardiol* Â 2020;318:61â66.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

274Shanks Â GW, Tester Â DJ, Ackerman Â JP, Simpson Â MA, Behr Â ER, White Â SM, et al. Â Importance of variant interpretation in whole-exome molecular autopsy: population-based case series. *Circulation* Â 2018;137:2705â2715.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

275Narula Â N, Tester Â DJ, Paulmichl Â A, Maleszewski Â JJ, Ackerman Â MJ. Post-mortem Whole exome sequencing with gene-specific analysis for autopsy-negative sudden unexplained death in the young: a case series. *Pediatr Cardiol* Â 2015;36:768â778.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

276Miles Â CJ, Behr Â ER. The role of genetic testing in unexplained sudden death. *Transl Res* Â 2016;168:59â73.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

277Behr Â ER, Dalageorgou Â C, Christiansen Â M, Syrris Â P, Hughes Â S, Tome Esteban Â MT, et al. Â Sudden arrhythmic death syndrome: familial evaluation identifies inheritable heart disease in the majority of families. *Eur Heart J* Â 2008;29:1670â1680.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

278Kumar Â S, Peters Â S, Thompson Â T, Morgan Â N, Maccicoca Â I, Trainer Â A, et al. Â Familial cardiological and targeted genetic evaluation: low yield in sudden unexplained death and high yield in unexplained cardiac arrest syndromes. *Heart Rhythm* Â 2013;10:1653â1660.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

279McGorrian Â C, Constant Â O, Harper Â N, O'Connell Â C, Codd Â M, Keelan Â E, et al. Â Family-based cardiac screening in relatives of victims of sudden arrhythmic death syndrome. *Europace* Â 2013;15:1050â1058.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

280Mellor Â G, Raju Â H, de Noronha Â SV, Papadakis Â M, Sharma Â S, Behr Â ER, et al. Â Clinical characteristics and circumstances of death in the sudden arrhythmic death syndrome. *Circ Arrhythm*

- Electrophysiol 2014;7:1078â1083.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 281Papadakis M, Papatheodorou E, Mellor G, Raju H, Bastiaenen R, Wijeyeratne Y, et al. The diagnostic yield of Brugada syndrome after sudden death with normal autopsy. *J Am Coll Cardiol* 2018;71:1204â1214.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 282Mellor G, Nelson CP, Robb C, Raju H, Wijeyeratne Y, Hengstenberg C, et al. The prevalence and significance of the early repolarization pattern in sudden arrhythmic death syndrome families. *Circ Arrhythm Electrophysiol* 2016;9:e003960.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 283Nunn LM, Lopes LR, Syrris P, Murphy C, Plagnol V, Firman E, et al. Diagnostic yield of molecular autopsy in patients with sudden arrhythmic death syndrome using targeted exome sequencing. *Europace* 2016;18:888â896.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 284van der Werf C, Stiekema L, Tan HL, Hofman N, Alders M, van der Wal AC, et al. Low rate of cardiac events in first-degree relatives of diagnosis-negative young sudden unexplained death syndrome victims during follow-up. *Heart Rhythm* 2014;11:1728â1732.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 285Madhavan M, Friedman PA, Lennon RJ, Prasad A, White RD, Sriram CS, et al. Implantable cardioverter-defibrillator therapy in patients with ventricular fibrillation out of hospital cardiac arrest secondary to acute coronary syndrome. *J Am Heart Assoc* 2015;4:e001255.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 286Wyse DG, Friedman PL, Brodsky MA, Beckman KJ, Carlson MD, Curtis AB, et al. Life-threatening ventricular arrhythmias due to transient or correctable causes: high risk for death in follow-up. *J Am Coll Cardiol* 2001;38:1718â1724.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 287Goyal A, Spertus JA, Gosch K, Venkitachalam L, Jones PG, Van den Berghe G, et al. Serum potassium levels and mortality in acute myocardial infarction. *JAMA* 2012;307:157â164.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 288Hoppe LK, Muhlack DC, Koenig W, Carr PR, Brenner H, SchÃ¼ttker B. Association of abnormal serum potassium levels with arrhythmias and cardiovascular mortality: a systematic review and meta-analysis of observational studies. *Cardiovasc Drugs Ther* 2018;32:197â212.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 289Ravn Jacobsen M, Jabbari R, Glinge C, KjÃ¦r Stampe N, Butt JH, Blanche P, et al. Potassium disturbances and risk of ventricular fibrillation among patients with ST-segment-elevation myocardial infarction. *J Am Heart Assoc* 2020;9:e014160.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 290Ahmed W, Flynn MA, Alpert MA. Cardiovascular complications of weight reduction diets. *Am J Med Sci* 2001;321:280â284.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 291Stephenson V. In pharmacy nursingâa dynamic new field. *Nurs RSA* 1988;3:51â53.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 292Dumas F, Cariou A, Manzo-Silberman S, Grimaldi D, Vivien B, Rosencher J, et al. Immediate percutaneous coronary intervention is associated with better survival after out-of-hospital cardiac arrest: insights from the PROCAT (Parisian Region Out of hospital Cardiac Arrest) registry. *Circ Cardiovasc Interv* 2010;3:200â207.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 293Gibbs C, Thalamus J, Haldal K, Holla L, Haugaa KH, Hysing J. Predictors of mortality in high-risk patients with QT prolongation in a community hospital. *Europace* 2018;20:f99âf107.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 294Simpson TF, Salazar JW, Vittinghoff E, Probert J, Iwahashi A, Olgin JE, et al. Association of QT-prolonging medications with risk of autopsy-defined causes of sudden death. *JAMA Intern Med* 2020;180:698â706.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

295Tzivoni [Â](#) D, Banai [Â](#) S, Schuger [Â](#) C, Benhorin [Â](#) J, Keren [Â](#) A, Gottlieb [Â](#) S, et al. [Â](#) Treatment of torsade de pointes with magnesium sulfate. *Circulation* [Â](#) 1988;77:392â397.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

296Ladejobi [Â](#) A, Pasupula [Â](#) DK, Adhikari [Â](#) S, Javed [Â](#) A, Durrani [Â](#) AF, Patil [Â](#) S, et al. [Â](#) Implantable defibrillator therapy in cardiac arrest survivors with a reversible cause. *Circ Arrhythm Electrophysiol* [Â](#) 2018;11:e005940.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

297Postema [Â](#) PG, Wolpert [Â](#) C, Amin [Â](#) AS, Probst [Â](#) V, Borggrefe [Â](#) M, Roden [Â](#) DM, et al. [Â](#) Drugs and Brugada syndrome patients: review of the literature, recommendations, and an up-to-date website (www.brugadadrugs.org). *Heart Rhythm* [Â](#) 2009;6:1335â1341.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

298Schupp [Â](#) T, Behnes [Â](#) M, von Zworowsky [Â](#) M, Kim [Â](#) S-H, Weidner [Â](#) K, Rusnak [Â](#) J, et al. [Â](#) Hypokalemia but not hyperkalemia is associated with recurrences of ventricular tachyarrhythmias in ICD recipients. *Clin Lab* [Â](#) 2020;66.

[Google Scholar](#) [OpenURL Placeholder Text](#) [Â](#)

299Gupta [Â](#) A, Pasupula [Â](#) DK, Bhonsale [Â](#) A, Kancharla [Â](#) K, Wang [Â](#) NC, Adelstein [Â](#) E, et al. [Â](#) Implantable cardioverter-defibrillator therapy in device recipients who survived a cardiac arrest associated with a reversible cause. *J Cardiovasc Electrophysiol* [Â](#) 2018;29:1413â1417.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

300Marill [Â](#) KA, Wolfram [Â](#) S, Desouza [Â](#) IS, Nishijima [Â](#) DK, Kay [Â](#) D, Setnik [Â](#) GS, et al. [Â](#) Adenosine for wide-complex tachycardia: efficacy and safety. *Crit Care Med* [Â](#) 2009;37:2512â2518.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

301Lerman [Â](#) BB, Belardinelli [Â](#) L, West [Â](#) GA, Berne [Â](#) RM, DiMarco [Â](#) JP. Adenosine-sensitive ventricular tachycardia: evidence suggesting cyclic AMP-mediated triggered activity. *Circulation* [Â](#) 1986;74:270â280.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

302Brugada [Â](#) J, Katritsis [Â](#) DG, Arbelo [Â](#) E, Arribas [Â](#) F, Bax [Â](#) JJ, Blomstr  m-Lundqvist [Â](#) C, et al. [Â](#) 2019 ESC Guidelines for the management of patients with supraventricular tachycardiaThe task force for the management of patients with supraventricular tachycardia of the European Society of Cardiology (ESC). *Eur Heart J* [Â](#) 2020;41:655â720.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

303Ortiz [Â](#) M, Mart  n [Â](#) A, Arribas [Â](#) F, Coll-Vinent [Â](#) B, Del Arco [Â](#) C, Peinado [Â](#) R, et al. [Â](#) Randomized comparison of intravenous procainamide vs. intravenous amiodarone for the acute treatment of tolerated wide QRS tachycardia: the PROCAMIO study. *Eur Heart J* [Â](#) 2017;38:1329â1335.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) [Â](#)

304Gill [Â](#) JS, Mehta [Â](#) D, Ward [Â](#) DE, Camm [Â](#) AJ. Efficacy of flecainide, sotalol, and verapamil in the treatment of right ventricular tachycardia in patients without overt cardiac abnormality. *Br Heart J* [Â](#) 1992;68:392â397.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

305Manz [Â](#) M, Mletzko [Â](#) R, Jung [Â](#) W, L  deritz [Â](#) B. Electrophysiological and haemodynamic effects of lidocaine and ajmaline in the management of sustained ventricular tachycardia. *Eur Heart J* [Â](#) 1992;13:1123â1128.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

306Buxton [Â](#) AE, Marchlinski [Â](#) FE, Doherty [Â](#) JU, Cassidy [Â](#) DM, Vassallo [Â](#) JA, Flores [Â](#) BT, et al. [Â](#) Repetitive, monomorphic ventricular tachycardia: clinical and electrophysiologic characteristics in patients with and patients without organic heart disease. *Am J Cardiol* [Â](#) 1984;54:997â1002.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

307Griffith [Â](#) MJ, Garratt [Â](#) CJ, Rowland [Â](#) E, Ward [Â](#) DE, Camm [Â](#) AJ. Effects of intravenous adenosine on verapamil-sensitive âidiopathicâ ventricular tachycardia. *Am J Cardiol* [Â](#) 1994;73:759â764.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

308Buxton [Â](#) AE, Marchlinski [Â](#) FE, Doherty [Â](#) JU, Flores [Â](#) B, Josephson [Â](#) ME. Hazards of intravenous verapamil for sustained ventricular tachycardia. *Am J Cardiol* [Â](#) 1987;59:1107â1110.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

- 309Rankin A AC, Rae A AP, Cobbe A SM. Misuse of intravenous verapamil in patients with ventricular tachycardia. *Lancet* 1987;2:472a-474.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 310Pedersen A CT, Kay A GN, Kalman A J, Borggrefe A M, Della-Bella A P, Dickfeld A T, et al. A EHRA/HRS/APHRS expert consensus on ventricular arrhythmias. *Europace* 2014;16:1257a-1283.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 311Cronin A EM, Bogun A FM, Maury A P, Peichl A P, Chen A M, Namboodiri A N, et al. 2019 HRS/EHRA/APHRS/LAHS expert consensus statement on catheter ablation of ventricular arrhythmias. *Europace* 2019;21:1143a-1144.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 312Kowligi A GN, Cha A Y-M. Management of ventricular electrical storm: a contemporary appraisal. *Europace* 2020;22:1768a-1780.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 313Guerra A F, Shkoko A M, Scappini A L, Flori A M, Capucci A A. Role of electrical storm as a mortality and morbidity risk factor and its clinical predictors: a meta-analysis. *Europace* 2014;16:347a-353.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 314Noda A T, Kurita A T, Nitta A T, Chiba A Y, Furushima A H, Matsumoto A N, et al. A Significant impact of electrical storm on mortality in patients with structural heart disease and an implantable cardiac defibrillator. *Int J Cardiol* 2018;255:85a-91.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 315Soar A J, Maconochie A I, Wyckoff A MH, Olasveengen A TM, Singletary A EM, Greif A R, et al. 2019 International consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. *Circulation* 2019;140:e826a-e880.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 316Eifling A M, Razavi A M, Massumi A A. The evaluation and management of electrical storm. *Tex Heart Inst J* 2011;38:111a-121.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 317Chatzidou A S, Kontogiannis A C, Tsilimigras A DI, Georgiopoulos A G, Kosmopoulos A M, Papadopoulou A E, et al. A Propranolol versus metoprolol for treatment of electrical storm in patients with implantable cardioverter-defibrillator. *J Am Coll Cardiol* 2018;71:1897a-1906.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 318Connolly A SJ, Dorian A P, Roberts A RS, Gent A M, Bailin A S, Fain A ES, et al. A Comparison of beta-blockers, amiodarone plus beta-blockers, or sotalol for prevention of shocks from implantable cardioverter defibrillators: the OPTIC Study: a randomized trial. *JAMA* 2006;295:165a-171.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 319Shiga A T, Ikeda A T, Shimizu A W, Kinugawa A K, Sakamoto A A, Nagai A R, et al. A Efficacy and safety of landiolol in patients with ventricular tachyarrhythmias with or without renal impairment a subanalysis of the J-Land II study. *Circ Rep* 2020;2:440a-445.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 320Kanamori A K, Aoyagi A T, Mikamo A T, Tsutsui A K, Kunishima A T, Inaba A H, et al. A Successful treatment of refractory electrical storm with landiolol after more than 100 electrical defibrillations. *Int Heart J* 2015;56:555a-557.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 321Gorgels A AP, van den Dool A A, Hofs A A, Mulleneers A R, Smeets A JL, Vos A MA, et al. A Comparison of procainamide and lidocaine in terminating sustained monomorphic ventricular tachycardia. *Am J Cardiol* 1996;78:43a-46.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 322MartÃ-Carvajal A AJ, Simancas-Racines A D, Anand A V, Bangdiwala A S. Prophylactic lidocaine for myocardial infarction. *Cochrane Database Syst Rev* 2015;2015:CD008553.
[Google Scholar](#) [OpenURL Placeholder Text](#)

- 323Viskin A S, Chorin A E, Viskin A D, Hochstadt A A, Halkin A A, Tovia-Brodie O, et al. A Quinidine-responsive polymorphic ventricular tachycardia in patients with coronary heart disease. *Circulation* A 2019;139:2304a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 324Viskin A S, Hochstadt A A, Chorin A E, Viskin A D, Havakuk A O, Khoury S, et al. A Quinidine-responsive out-of-hospital polymorphic ventricular tachycardia in patients with coronary heart disease. *Europace* A 2020;22:265a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 325Martins A RP, Urien A J-M, Barbarot A N, Rieul A G, Sellal A J-M, Borella A L, et al. A Effectiveness of deep sedation for patients with intractable electrical storm refractory to antiarrhythmic drugs. *Circulation* A 2020;142:1599a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 326Fudim A M, Boortz-Marx A R, Ganesh A A, Waldron A NH, Qadri A YJ, Patel A CB, et al. A Stellate ganglion blockade for the treatment of refractory ventricular arrhythmias: a systematic review and meta-analysis. *J Cardiovasc Electrophysiol* A 2017;28:1460a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 327Do A DH, Bradfield A J, Ajijola A OA, Vaseghi A M, Le A J, Rahman A S, et al. A Thoracic epidural anesthesia can be effective for the short-term management of ventricular tachycardia storm. *J Am Heart Assoc* A 2017;6:e007080. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 328Vaseghi A M, Barwad A P, Malavassi Corrales A FJ, Tandri A H, Mathuria A N, Shah A R, et al. A Cardiac sympathetic denervation for refractory ventricular arrhythmias. *J Am Coll Cardiol* A 2017;69:3070a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 329Gatzoulis A KA, Andrikopoulos A GK, Apostolopoulos A T, Sotiropoulos A E, Zervopoulos A G, Antoniou A J, et al. A Electrical storm is an independent predictor of adverse long-term outcome in the era of implantable defibrillator therapy. *Europace* A 2005;7:184a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 330Carbucicchio A C, Santamaria A M, Trevisi A N, Maccabelli A G, Giraldi A F, Fassini A G, et al. A Catheter ablation for the treatment of electrical storm in patients with implantable cardioverter-defibrillators: short- and long-term outcomes in a prospective single-center study. *Circulation* A 2008;117:462a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 331Vergara A P, Tung A R, Vaseghi A M, Brombin A C, Frankel A DS, Di Biase A L, et al. A Successful ventricular tachycardia ablation in patients with electrical storm reduces recurrences and improves survival. *Heart Rhythm* A 2018;15:48a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 332Komatsu A Y, Hocini A M, Nogami A A, Maury A P, Peichl A P, Iwasaki A Y, et al. A Catheter ablation of refractory ventricular fibrillation storm after myocardial infarction: a multicenter study. *Circulation* A 2019;139:2315a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 333Knecht A S, Sacher A F, Wright A M, Hocini A M, Nogami A A, Arentz A T, et al. A Long-term follow-up of idiopathic ventricular fibrillation ablation: a multicenter study. *J Am Coll Cardiol* A 2009;54:522a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 334Peichl A P, Cihák A R, Kozeluhová A M, Wichterle A D, Vancura A V, Kautzner A J. Catheter ablation of arrhythmic storm triggered by monomorphic ectopic beats in patients with coronary artery disease. *J Interv Card Electrophysiol* A 2010;27:51a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 335Le Pennec-Prigent A S, Flecher A E, Auffret A V, Leurent A G, Daubert A J-C, Leclercq A C, et al. A Effectiveness of extracorporeal life support for patients with cardiogenic shock due to intractable arrhythmic storm. *Crit Care Med* A 2017;45:e281a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 336Mariani A S, Napp A LC, Lo Coco A V, Delnoij A TSR, Luermans A JGLM, Ter Bekke A RMA, et al. A Mechanical circulatory support for life-threatening arrhythmia: a systematic review. *Int J Cardiol* A 2020;308:42a. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A

- 337Muser Â D, Liang Â JJ, Castro Â SA, Hayashi Â T, Enriquez Â A, Troutman Â GS, et al. Â Outcomes with prophylactic use of percutaneous left ventricular assist devices in high-risk patients undergoing catheter ablation of scar-related ventricular tachycardia: a propensity-score matched analysis. *Heart Rhythm* Â 2018;15:1500â1506.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 338Mathuria Â N, Wu Â G, Rojas-Delgado Â F, Shuraih Â M, Razavi Â M, Civitello Â A, et al. Â Outcomes of pre-emptive and rescue use of percutaneous left ventricular assist device in patients with structural heart disease undergoing catheter ablation of ventricular tachycardia. *J Interv Card Electrophysiol* Â 2017;48:27â34.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 339Patel Â KK, Spertus Â JA, Khariton Â Y, Tang Â Y, Curtis Â LH, Chan Â PS, et al. Â Association between prompt defibrillation and epinephrine treatment with long-term survival after in-hospital cardiac arrest. *Circulation* Â 2018;137:2041â2051.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 340Vaseghi Â M, Gima Â J, Kanaan Â C, Ajijola Â OA, Marmureanu Â A, Mahajan Â A, et al. Â Cardiac sympathetic denervation in patients with refractory ventricular arrhythmias or electrical storm: Intermediate and long-term follow-up. *Heart Rhythm* Â 2014;11:360â366.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 341Ponikowski Â P, Voors Â AA, Anker Â SD, Bueno Â H, Cleland Â JGF, Coats Â AJS, et al. Â 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the task force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J* Â 2016;37:2129â2200.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 342McDonagh Â TA, Metra Â M, Adamo Â M, Gardner Â RS, Baumbach Â A, Böhm Â M, et al. Â 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* Â 2021;42:3599â3726.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 343AlJaroudi Â WA, Refaat Â MM, Habib Â RH, Al-Shaar Â L, Singh Â M, Gutmann Â R, et al. Â Effect of angiotensin-converting enzyme inhibitors and receptor blockers on appropriate implantable cardiac defibrillator shock in patients with severe systolic heart failure (from the GRADE Multicenter Study). *Am J Cardiol* Â 2015;115:924â931.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 344Pitt Â B, Remme Â W, Zannad Â F, Neaton Â J, Martinez Â F, Roniker Â B, et al. Â Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* Â 2003;348:1309â1321.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 345McMurray Â JJV, Packer Â M, Desai Â AS, Gong Â J, Lefkowitz Â MP, Rizkala Â AR, et al. Â Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* Â 2014;371:993â1004.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 346Packer Â M, Anker Â SD, Butler Â J, Filippatos Â G, Pocock Â SJ, Carson Â P, et al. Â Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med* Â 2020;383:1413â1424.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 347Desai Â AS, McMurray Â JJV, Packer Â M, Swedberg Â K, Rouleau Â JL, Chen Â F, et al. Â Effect of the angiotensin-receptor-neprilysin inhibitor LCZ696 compared with enalapril on mode of death in heart failure patients. *Eur Heart J* Â 2015;36:1990â1997.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 348Hindricks Â G, Potpara Â T, Dagres Â N, Arbelo Â E, Bax Â JJ, Blomström-Lundqvist Â C, et al. Â 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The task force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J* Â 2021;42:373â498.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 349Connolly Â SJ, Gent Â M, Roberts Â RS, Dorian Â P, Roy Â D, Sheldon Â RS, et al. Â Canadian implantable defibrillator study (CIDS): a randomized trial of the implantable cardioverter defibrillator against amiodarone. *Circulation* Â 2000;101:1297â1302.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

350Antiarrhythmics versus Implantable Defibrillators (AVID) Investigators. A comparison of antiarrhythmic-drug therapy with implantable defibrillators in patients resuscitated from near-fatal ventricular arrhythmias. *N Engl J Med* 1997;337:1576-1583.

[Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

351Kuck KH, Cappato R, Siebels J, Rappel R. Randomized comparison of antiarrhythmic drug therapy with implantable defibrillators in patients resuscitated from cardiac arrest: the Cardiac Arrest Study Hamburg (CASH). *Circulation* 2000;102:748-754.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

352Connolly SJ, Hallstrom AP, Cappato R, Schron EB, Kuck KH, Zipes DP, et al. Meta-analysis of the implantable cardioverter defibrillator secondary prevention trials. AVID, CASH and CIDS studies. Antiarrhythmics vs Implantable Defibrillator study. Cardiac Arrest Study Hamburg. Canadian Implantable Defibrillator Study. *Eur Heart J* 2000;21:2071-2078.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

353Moss AJ, Hall WJ, Cannom DS, Daubert JP, Higgins SL, Klein H, et al. Improved survival with an implanted defibrillator in patients with coronary disease at high risk for ventricular arrhythmia. *N Engl J Med* 1996;335:1933-1940.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

354Moss AJ, Zareba W, Hall WJ, Klein H, Wilber DJ, Cannom DS, et al. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med* 2002;346:877-883.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

355Buxton AE, Lee KL, Fisher JD, Josephson ME, Prystowsky EN, Hafley G. A randomized study of the prevention of sudden death in patients with coronary artery disease. Multicenter Unsustained Tachycardia Trial Investigators. *N Engl J Med* 1999;341:1882-1890.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

356Bardy GH, Lee KL, Mark DB, Poole JE, Packer DL, Boineau R, et al. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med* 2005;352:225-237.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

357Zabel M, Willems R, Lubinski A, Bauer A, Brugada J, Conen D, et al. Clinical effectiveness of primary prevention implantable cardioverter-defibrillators: results of the EU-CERT-ICD controlled multicentre cohort study. *Eur Heart J* 2020;41:3437-3447.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

358Schrage B, Uijl A, Benson L, Westermann D, Ståhlberg M, Stolfo D, et al. Association between use of primary-prevention implantable cardioverter-defibrillators and mortality in patients with heart failure: a prospective propensity score-matched analysis from the Swedish Heart Failure Registry. *Circulation* 2019;140:1530-1539.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

359Kjaer L, Thune JJ, Nielsen JC, Haarbø J, Videbæk L, Korup E, et al. Defibrillator implantation in patients with nonischemic systolic heart failure. *N Engl J Med* 2016;375:1221-1230.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

360Jukema JW, Timal RJ, Rotmans JI, Hensen LCR, Buiten MS, de Bie MK, et al. Prophylactic use of implantable cardioverter-defibrillators in the prevention of sudden cardiac death in dialysis patients. *Circulation* 2019;139:2628-2638.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

361Sticherling C, Arendacka B, Svendsen JH, Wijers S, Friede T, Stockinger J, et al. Sex differences in outcomes of primary prevention implantable cardioverter-defibrillator therapy: combined registry data from eleven European countries. *Europace* 2018;20:963-970.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

362Junttila MJ, Pelli A, Kenttö TV, Friede T, Willems R, Bergau L, et al. Appropriate shocks and mortality in patients with versus without diabetes with prophylactic implantable cardioverter defibrillators. *Diabetes Care* 2020;43:196-200.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [↗](#)

- 363Koller Â MT, Schaer Â B, Wolbers Â M, Sticherling Â C, Bucher Â HC, Osswald Â S. Death without prior appropriate implantable cardioverter-defibrillator therapy: a competing risk study. *Circulation* Â 2008;117:1918â1926.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 364Cleland Â JGF, Halliday Â BP, Prasad Â SK. Selecting patients with nonischemic dilated cardiomyopathy for ICDs: myocardial function, fibrosis, and whatâs attached? Â *J Am Coll Cardiol* Â 2017;70:1228â1231.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 365Younis Â A, Goldberger Â JJ, Kutyla Â V, Zareba Â W, Polonsky Â B, Klein Â H, et al. Â Predicted benefit of an implantable cardioverter-defibrillator: the MADIT-ICD benefit score. *Eur Heart J* Â 2021;42:1676â1684.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 366Knops Â RE, Olde Nordkamp Â LRA, Delnoy Â P-PH, Boersma Â LVA, Kuschyk Â J, El-Chami Â MF, et al. Â Subcutaneous or transvenous defibrillator therapy. *N Engl J Med* Â 2020;383:526â536.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 367Cleland Â JGF, Daubert Â J-C, Erdmann Â E, Freemantle Â N, Gras Â D, Kappenberger Â L, et al. Â The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med* Â 2005;352:1539â1549.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 368Bristow Â MR, Saxon Â LA, Boehmer Â J, Krueger Â S, Kass Â DA, De Marco Â T, et al. Â Cardiac-resynchronization therapy with or without an implantable defibrillator in advanced chronic heart failure. *N Engl J Med* Â 2004;350:2140â2150.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 369Moss Â AJ, Hall Â WJ, Cannom Â DS, Klein Â H, Brown Â MW, Daubert Â JP, et al. Â Cardiac-resynchronization therapy for the prevention of heart-failure events. *N Engl J Med* Â 2009;361:1329â1338.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 370Masri Â A, Altibi Â AM, Erqou Â S, Zmaili Â MA, Saleh Â A, Al-Adham Â R, et al. Â Wearable cardioverter-defibrillator therapy for the prevention of sudden cardiac death: a systematic review and meta-analysis. *JACC Clin Electrophysiol* Â 2019;5:152â161.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 371Garcia Â R, Combes Â N, Defaye Â P, Narayanan Â K, Guedon-Moreau Â L, Boveda Â S, et al. Â Wearable cardioverter-defibrillator in patients with a transient risk of sudden cardiac death: the WEARIT-France cohort study. *Europace* Â 2021;23:73â81.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 372Olgin Â JE, Pletcher Â MJ, Vittinghoff Â E, Wranicz Â J, Malik Â R, Morin Â DP, et al. Â Wearable cardioverter-defibrillator after myocardial infarction. *N Engl J Med* Â 2018;379:1205â1215.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 373Scott Â PA, Silberbauer Â J, McDonagh Â TA, Murgatroyd Â FD. Impact of prolonged implantable cardioverter-defibrillator arrhythmia detection times on outcomes: a meta-analysis. *Heart Rhythm* Â 2014;11:828â835.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 374Tan Â VH, Wilton Â SB, Kuriachan Â V, Sumner Â GL, Exner Â DV. Impact of programming strategies aimed at reducing nonessential implantable cardioverter defibrillator therapies on mortality: a systematic review and meta-analysis. *Circ Arrhythm Electrophysiol* Â 2014;7:164â170.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 375Saeed Â M, Hanna Â I, Robotis Â D, Styperek Â R, Polosajian Â L, Khan Â A, et al. Â Programming implantable cardioverter-defibrillators in patients with primary prevention indication to prolong time to first shock: results from the PROVIDE study. *J Cardiovasc Electrophysiol* Â 2014;25:52â59.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 376Wilkoff Â BL, Fauchier Â L, Stiles Â MK, Morillo Â CA, Al-Khatib Â SM, Almendral Â J, et al. Â 2015 HRS/EHRA/APHRS/SOLAECE expert consensus statement on optimal implantable cardioverter-defibrillator programming and testing. *Europace* Â 2016;18:159â183.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 377Stiles Â MK, Fauchier Â L, Morillo Â CA, Wilkoff Â BL, ESC Scientific Document Group. 2019 HRS/EHRA/APHRS/LAHRs focused update to 2015 expert consensus statement on optimal implantable cardioverter-defibrillator programming and testing. *Europace* Â 2019;21:1442â1443.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 378Barsheshet Â A, Moss Â AJ, McNitt Â S, Jons Â C, Glikson Â M, Klein Â HU, et al. Â Long-term implications of cumulative right ventricular pacing among patients with an implantable cardioverter-defibrillator. *Heart Rhythm* Â 2011;8:212â218.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 379Wilkoff Â BL, Cook Â JR, Epstein Â AE, Greene Â HL, Hallstrom Â AP, Hsia Â H, et al. Â Dual-chamber pacing or ventricular backup pacing in patients with an implantable defibrillator: the Dual Chamber and VVI Implantable Defibrillator (DAVID) Trial. *JAMA* Â 2002;288:3115â3123.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 380Olshansky Â B, Day Â JD, Moore Â S, Gering Â L, Rosenbaum Â M, McGuire Â M, et al. Â Is dual-chamber programming inferior to single-chamber programming in an implantable cardioverter-defibrillator? Results of the INTRINSIC RV (Inhibition of Unnecessary RV Pacing With AVSH in ICDs) study. *Circulation* Â 2007;115:9â16.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 381Hindricks Â G, KÄ¼hl Â M, Dagres Â N. The implantable cardioverter defibrillator, conclusions on sudden cardiac death, and future perspective. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p2370â2376.
- 382Gasparini Â M, Proclemer Â A, Klersy Â C, Kloppe Â A, Ferrer Â JBM, Hersi Â A, et al. Â Effect of long-detection interval vs standard-detection interval for implantable cardioverter-defibrillators on antitachycardia pacing and shock delivery: the ADVANCE III randomized clinical trial. *JAMA* Â 2013;309:1903â1911.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 383Moss Â AJ, Schuger Â C, Beck Â CA, Brown Â MW, Cannom Â DS, Daubert Â JP, et al. Â Reduction in inappropriate therapy and mortality through ICD programming. *N Engl J Med* Â 2012;367:2275â2283.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 384Wilkoff Â BL, Ousdigian Â KT, Sterns Â LD, Wang Â ZJ, Wilson Â RD, Morgan Â JM, et al. Â A comparison of empiric to physician-tailored programming of implantable cardioverter-defibrillators: results from the prospective randomized multicenter EMPIRIC trial. *J Am Coll Cardiol* Â 2006;48:330â339.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 385Wilkoff Â BL, Williamson Â BD, Stern Â RS, Moore Â SL, Lu Â F, Lee Â SW, et al. Â Strategic programming of detection and therapy parameters in implantable cardioverter-defibrillators reduces shocks in primary prevention patients: results from the PREPARE (Primary Prevention Parameters Evaluation) study. *J Am Coll Cardiol* Â 2008;52:541â550.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 386Gilliam Â FR, Hayes Â DL, Boehmer Â JP, Day Â J, Heidenreich Â PA, Seth Â M, et al. Â Real world evaluation of dual-zone ICD and CRT-D programming compared to single-zone programming: the ALTITUDE REDUCES study. *J Cardiovasc Electrophysiol* Â 2011;22:1023â1029.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 387Hernandez-Ojeda Â J, Arbelo Â E, Borrás Â R, Berne Â P, Tolosana Â JM, Gomez-Juanatey Â A, et al. Â Patients with Brugada syndrome and implanted cardioverter-defibrillators: long-term follow-up. *J Am Coll Cardiol* Â 2017;70:1991â2002.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 388Gold Â MR, Weiss Â R, Theuns Â DAMJ, Smith Â W, Leon Â A, Knight Â BP, et al. Â Use of a discrimination algorithm to reduce inappropriate shocks with a subcutaneous implantable cardioverter-defibrillator. *Heart Rhythm* Â 2014;11:1352â1358.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 389Mesquita Â J, Cavaco Â D, Ferreira Â A, Lopes Â N, Santos Â PG, Carvalho Â MS, et al. Â Effectiveness of subcutaneous implantable cardioverter-defibrillators and determinants of inappropriate shock delivery. *Int J Cardiol* Â 2017;232:176â180.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 390Gold Â MR, Lambiase Â PD, El-Chami Â MF, Knops Â RE, Aasbo Â JD, Bongiorni Â MG, et al. Â Primary results from the understanding outcomes with the S-ICD in primary prevention patients with low ejection

- fraction (UNTOUCHED) trial. *Circulation* 2021;143:731-741. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 391 Wathen MS, DeGroot PJ, Sweeney MO, Stark AJ, Otterness MF, Adkisson WO, et al. Prospective randomized multicenter trial of empirical antitachycardia pacing versus shocks for spontaneous rapid ventricular tachycardia in patients with implantable cardioverter-defibrillators: Pacing Fast Ventricular Tachycardia Reduces Shock Therapies (PainFREE Rx II) trial results. *Circulation* 2004;110:2591-2596. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 392 Gulizia MM, Piraino L, Scherillo M, Puntrello C, Vasco C, Scianaro MC, et al. A randomized study to compare ramp versus burst antitachycardia pacing therapies to treat fast ventricular tachyarrhythmias in patients with implantable cardioverter defibrillators: the PITAGORA ICD trial. *Circ Arrhythm Electrophysiol* 2009;2:146-153. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 393 Saxon LA, Hayes DL, Gilliam FR, Heidenreich PA, Day J, Seth M, et al. Long-term outcome after ICD and CRT implantation and influence of remote device follow-up: the ALTITUDE survival study. *Circulation* 2010;122:2359-2367. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 394 Varma N, Piccini JP, Snell J, Fischer A, Dalal N, Mittal S. The relationship between level of adherence to automatic wireless remote monitoring and survival in pacemaker and defibrillator patients. *J Am Coll Cardiol* 2015;65:2601-2610. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 395 Guédon-Moreau L, Kouakam C, Klug D, Marquié C, Brigadeau F, Boulé S, et al. Decreased delivery of inappropriate shocks achieved by remote monitoring of ICD: a substudy of the ECOST trial. *J Cardiovasc Electrophysiol* 2014;25:763-770. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 396 Varma N, Michalski J, Epstein AE, Schweikert R. Automatic remote monitoring of implantable cardioverter-defibrillator lead and generator performance: the Lumos-T Safely Reduces Routine Office Device Follow-Up (TRUST) trial. *Circ Arrhythm Electrophysiol* 2010;3:428-436. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 397 Ploux S, Swerdlow CD, Strik M, Welte N, Klotz N, Ritter P, et al. Towards eradication of inappropriate therapies for ICD lead failure by combining comprehensive remote monitoring and lead noise alerts. *J Cardiovasc Electrophysiol* 2018;29:1125-1134. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 398 Ellenbogen KA, Gunderson BD, Stromberg KD, Swerdlow CD. Performance of Lead Integrity Alert to assist in the clinical diagnosis of implantable cardioverter defibrillator lead failures: analysis of different implantable cardioverter defibrillator leads. *Circ Arrhythm Electrophysiol* 2013;6:1169-1177. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 399 Swerdlow CD, Gunderson BD, Ousdigian KT, Abeyratne A, Sachanandani H, Ellenbogen KA. Downloadable software algorithm reduces inappropriate shocks caused by implantable cardioverter-defibrillator lead fractures: a prospective study. *Circulation* 2010;122:1449-1455. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 400 Ruwald MH, Abu-Zeitone A, Jons C, Ruwald A-C, McNitt S, Kutyifa V, et al. Impact of carvedilol and metoprolol on inappropriate implantable cardioverter-defibrillator therapy: the MADIT-CRT trial (Multicenter Automatic Defibrillator Implantation With Cardiac Resynchronization Therapy). *J Am Coll Cardiol* 2013;62:1343-1350. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 401 Miyazaki S, Taniguchi H, Kusa S, Komatsu Y, Ichihara N, Takagi T, et al. Catheter ablation of atrial tachyarrhythmias causing inappropriate implantable cardioverter-defibrillator shocks. *Europace* 2015;17:289-294. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 402 Mainigi SK, Almuti K, Figueredo VM, Guttenplan NA, Aouthmany A, Smukler J, et al. Usefulness of radiofrequency ablation of supraventricular tachycardia to decrease inappropriate shocks from implantable cardioverter-defibrillators. *Am J Cardiol* 2012;109:231-237. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 403Kosiuk A J, Nedios S, Darma A, Rolf S, Richter S, Arya A, et al. Impact of single atrial fibrillation catheter ablation on implantable cardioverter defibrillator therapies in patients with ischaemic and non-ischaemic cardiomyopathies. *Europace* 2014;16:1322â1326.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 404Kirchhof P, Camm AJ, Goette A, Brandes A, Eckardt L, Elvan A, et al. Early rhythm-control therapy in patients with atrial fibrillation. *N Engl J Med* 2020;383:1305â1316.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 405Gasparini M, Kloppe A, Lunati M, Anselme F, Landolina M, Martinez-Ferrer JB, et al. Atrioventricular junction ablation in patients with atrial fibrillation treated with cardiac resynchronization therapy: positive impact on ventricular arrhythmias, implantable cardioverter-defibrillator therapies and hospitalizations: atrioventricular junction ablation in CRT patients with AF. *Eur J Heart Fail* 2018;20:1472â1481.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 406Gasparini M, Galimberti P. Rate control: ablation and device therapy (ablate and pace). *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p2159â2162.
- 407Kitamura T, Fukamizu S, Kawamura I, Hojo R, Aoyama Y, Komiyama K, et al. Long-term efficacy of catheter ablation for paroxysmal atrial fibrillation in patients with Brugada syndrome and an implantable cardioverter-defibrillator to prevent inappropriate shock therapy. *Heart Rhythm* 2016;13:1455â1459.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 408Magyar-Russell G, Thombs BD, Cai JX, Baveja T, Kuhl EA, Singh PP, et al. The prevalence of anxiety and depression in adults with implantable cardioverter defibrillators: a systematic review. *J Psychosom Res* 2011;71:223â231.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 409Tzeis S, Kolb C, Baumert J, Reents T, Zrenner B, Deisenhofer I, et al. Effect of depression on mortality in implantable cardioverter defibrillator recipientsâfindings from the prospective LICAD study. *Pacing Clin Electrophysiol* 2011;34:991â997.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 410Andersen CM, Theuns DAMJ, Johansen JB, Pedersen SS. Anxiety, depression, ventricular arrhythmias and mortality in patients with an implantable cardioverter defibrillator: 7 yearsâfollow-up of the MIDAS cohort. *Gen Hosp Psychiatry* 2020;66:154â160.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 411Berg SK, Thygesen LC, Svendsen JH, Christensen AV, Zwisler A-D. Anxiety predicts mortality in ICD patients: results from the cross-sectional national CopenHeartICD survey with register follow-up. *Pacing Clin Electrophysiol* 2014;37:1641â1650.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 412Thygesen I, Moser DK, Strimling A, Dekker RA, Chung ML. Concerns about implantable cardioverter-defibrillator shocks mediate the relationship between actual shocks and psychological distress. *Europace* 2016;18:828â835.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 413Pedersen SS, van Domburg RT, Theuns DAMJ, Jordaens L, Erdman RAM. Concerns about the implantable cardioverter defibrillator: a determinant of anxiety and depressive symptoms independent of experienced shocks. *Am Heart J* 2005;149:664â669.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 414Frizelle DJ, Lewin B, Kaye G, Moniz-Cook ED. Development of a measure of the concerns held by people with implanted cardioverter defibrillators: the ICDC. *Br J Health Psychol* 2006;11:293â301.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 415Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361â370.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 416Frydensberg VS, Johansen JB, Moller S, Riahi S, Wehberg S, Haarbo J, et al. Anxiety and depression symptoms in Danish patients with an implantable cardioverter-defibrillator: prevalence and association with indication and sex up to 2 years of follow-up (data from the national DEFIB-WOMEN study).

- Europace 2020;22:1830-1840.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 417Hoogwegt MT, Kupper N, Theuns DAMJ, Zijlstra WP, Jordaens L, Pedersen SS.
 Undertreatment of anxiety and depression in patients with an implantable cardioverter-defibrillator: impact on health status. *Health Psychol* 2012;31:745-753.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 418Lane DA, Aguinaga L, Blomström-Lundqvist C, Boriani G, Dan G-A, Hills MT, et al.
 Cardiac tachyarrhythmias and patient values and preferences for their management: the European Heart Rhythm Association (EHRA) consensus document endorsed by the Heart Rhythm Society (HRS), Asia Pacific Heart Rhythm Society (APHRS), and Sociedad Latinoamericana de Estimulación Cardíaca y Electrofisiología (SOLEACE). *Europace* 2015;17:1747-1769.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 419Dunbar SB, Dougherty CM, Sears SF, Carroll DL, Goldstein NE, Mark DB, et al.
 Educational and psychological interventions to improve outcomes for recipients of implantable cardioverter defibrillators and their families: a scientific statement from the American Heart Association. *Circulation* 2012;126:2146-2172.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 420Sears SF, Sowell LDV, Kuhl EA, Kovacs AH, Serber ER, Handberg E, et al.
 The ICD shock and stress management program: a randomized trial of psychosocial treatment to optimize quality of life in ICD patients. *Pacing Clin Electrophysiol* 2007;30:858-864.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 421Berg SK, Rasmussen TB, Herning M, Svendsen JH, Christensen AV, Thygesen LC.
 Cognitive behavioural therapy significantly reduces anxiety in patients with implanted cardioverter defibrillator compared with usual care: findings from the Screen-ICD randomised controlled trial. *Eur J Prev Cardiol* 2020;27:258-268.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 422Schulz SM, Ritter O, Zniwa R, Nordbeck P, Wacker C, Jack M, et al.
 Efficacy of a web-based intervention for improving psychosocial well-being in patients with implantable cardioverter-defibrillators: the randomized controlled ICD-FORUM trial. *Eur Heart J* 2020;41:1203-1211.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 423van den Broek KC, Tekle FB, Habibović M, Alings M, van der Voort PH, Denollet J.
 Emotional distress, positive affect, and mortality in patients with an implantable cardioverter defibrillator. *Int J Cardiol* 2013;165:327-332.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 424Hauptman PJ, Chibnall JT, Guild C, Armbrecht ES.
 Patient perceptions, physician communication, and the implantable cardioverter-defibrillator. *JAMA Intern Med* 2013;173:571-577.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 425Cikes M, Jakus N, Claggett B, Brugts JJ, Timmermans P, Pouleur A-C, et al.
 Cardiac implantable electronic devices with a defibrillator component and all-cause mortality in left ventricular assist device carriers: results from the PCHF-VAD registry. *Eur J Heart Fail* 2019;21:1129-1141.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 426Galand V, Flacher E, Auffret V, Boulé S, Vincentelli A, Dambrin C, et al.
 Predictors and clinical impact of late ventricular arrhythmias in patients with continuous-flow left ventricular assist devices. *JACC Clin Electrophysiol* 2018;4:1166-1175.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 427Nakahara S, Chien C, Gelow J, Dalouk K, Henrikson CA, Mudd J, et al.
 Ventricular arrhythmias after left ventricular assist device. *Circ Arrhythm Electrophysiol* 2013;6:648-654.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 428Clerkin KJ, Topkara VK, Demmer RT, Dizon JM, Yuzefpolskaya M, Fried JA, et al.
 Implantable cardioverter-defibrillators in patients with a continuous-flow left ventricular assist device: an analysis of the INTERMACS registry. *JACC Heart Fail* 2017;5:916-926.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 429Oz MC, Rose EA, Slater J, Kuiper JJ, Catanese KA, Levin HR.
 Malignant ventricular arrhythmias are well tolerated in patients receiving long-term left ventricular assist devices. *J Am Coll Cardiol*

- Â 1994;24:1688â^{PADSTS}1691.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 430Potapov Â EV, Antonides Â C, Crespo-Leiro Â MG, Combes Â A, FÃrber Â G, Hannan Â MM, et al. Â 2019 EACTS Expert Consensus on long-term mechanical circulatory support. *Eur J Cardiothorac Surg* Â 56:230â^{PADSTS}270.
[Crossref](#) [Search ADS](#) [PubMed](#) Â
- 431Makki Â N, Mesubi Â O, Steyers Â C, Olshansky Â B, Abraham Â WT. Meta-analysis of the relation of ventricular arrhythmias to all-cause mortality after implantation of a left ventricular assist device. *Am J Cardiol* Â 2015;116:1385â^{PADSTS}1390.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 432Yoruk Â A, Sherazi Â S, Massey Â HT, Kutyla Â V, McNitt Â S, Hallinan Â W, et al. Â Predictors and clinical relevance of ventricular tachyarrhythmias in ambulatory patients with a continuous flow left ventricular assist device. *Heart Rhythm* Â 2016;13:1052â^{PADSTS}1056.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 433Bedi Â M, Kormos Â R, Winowich Â S, McNamara Â DM, Mathier Â MA, Murali Â S. Ventricular arrhythmias during left ventricular assist device support. *Am J Cardiol* Â 2007;99:1151â^{PADSTS}1153.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 434Brenyo Â A, Rao Â M, Koneru Â S, Hallinan Â W, Shah Â S, Massey Â HT, et al. Â Risk of mortality for ventricular arrhythmia in ambulatory LVAD patients. *J Cardiovasc Electrophysiol* Â 2012;23:515â^{PADSTS}520.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 435Vakil Â K, Kazmirczak Â F, Sathnur Â N, Adabag Â S, Cantillon Â DJ, Kiehl Â EL, et al. Â Implantable cardioverter-defibrillator use in patients with left ventricular assist devices: a systematic review and meta-analysis. *JACC Heart Fail* Â 2016;4:772â^{PADSTS}779.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 436Refaat Â MM, Tanaka Â T, Kormos Â RL, McNamara Â D, Teuteberg Â J, Winowich Â S, et al. Â Survival benefit of implantable cardioverter-defibrillators in left ventricular assist device-supported heart failure patients. *J Card Fail* Â 2012;18:140â^{PADSTS}145.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 437Cantillon Â DJ, Tarakji Â KG, Kumbhani Â DJ, Smedira Â NG, Starling Â RC, Wilkoff Â BL. Improved survival among ventricular assist device recipients with a concomitant implantable cardioverter-defibrillator. *Heart Rhythm* Â 2010;7:466â^{PADSTS}471.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 438Joyce Â E, Starling Â RC. HFrEF other treatment: ventricular assist devices. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p1884â^{PADSTS}1889.
- 439Younes Â A, Al-Kindi Â SG, Alajaji Â W, Mackall Â JA, Oliveira Â GH. Presence of implantable cardioverter-defibrillators and wait-list mortality of patients supported with left ventricular assist devices as bridge to heart transplantation. *Int J Cardiol* Â 2017;231:211â^{PADSTS}215.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 440Agrawal Â S, Garg Â L, Nanda Â S, Sharma Â A, Bhatia Â N, Manda Â Y, et al. Â The role of implantable cardioverter-defibrillators in patients with continuous flow left ventricular assist devices â^{PADSTS} a meta-analysis. *Int J Cardiol* Â 2016;222:379â^{PADSTS}384.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 441BlomstrÃm-Lundqvist Â C, Traykov Â V, Erba Â PA, Burri Â H, Nielsen Â JC, Bongiorni Â MG, et al. Â European Heart Rhythm Association (EHRA) international consensus document on how to prevent, diagnose, and treat cardiac implantable electronic device infections-endorsed by the Heart Rhythm Society (HRS), the Asia Pacific Heart Rhythm Society (APHRS), the Latin American Heart Rhythm Society (LAHRS), International Society for Cardiovascular Infectious Diseases (ISCVID) and the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). *Europace* Â 2020;22:515â^{PADSTS}549.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 442Burri Â H, Starck Â C, Auricchio Â A, Biffi Â M, Burri Â M, Dâ^{PADSTS}Avila Â A, et al. Â EHRA expert consensus statement and practical guide on optimal implantation technique for conventional pacemakers and implantable cardioverter-defibrillators: endorsed by the Heart Rhythm Society (HRS), the Asia Pacific Heart Rhythm Society (APHRS), and the Latin-American Heart Rhythm Society (LAHRS). *Europace*

Â 2021;23:983â1008.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

443Tarakji Â KG, Mittal Â S, Kennergren Â C, Corey Â R, Poole Â JE, Schloss Â E, et al. Â Antibacterial envelope to prevent cardiac implantable device infection. *N Engl J Med* Â 2019;380:1895â1905.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

444Atti Â V, Turagam Â MK, Garg Â J, Koerber Â S, Angirekula Â A, Gopinathannair Â R, et al. Â Subclavian and axillary vein access versus cephalic vein cutdown for cardiac implantable electronic device implantation: a meta-analysis. *JACC Clin Electrophysiol* Â 2020;6:661â671.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

445Benz Â AP, Vamos Â M, Erath Â JW, Hohnloser Â SH. Cephalic vs. subclavian lead implantation in cardiac implantable electronic devices: a systematic review and meta-analysis. *Europace* Â 2019;21:121â129.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

446Chan Â N-Y, Kwong Â N-P, Cheong Â A-P. Venous access and long-term pacemaker lead failure: comparing contrast-guided axillary vein puncture with subclavian puncture and cephalic cutdown. *Europace* Â 2017;19:1193â1197.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

447Defaye Â P, Boveda Â S, Klug Â D, Beganton Â F, Piot Â O, Narayanan Â K, et al. Â Dual- vs. single-chamber defibrillators for primary prevention of sudden cardiac death: long-term follow-up of the DÃ©fibrillateur Automatique Implantable-PrÃ©vention Primaire registry. *Europace* Â 2017;19:1478â1484.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

448Dewland Â TA, Pellegrini Â CN, Wang Â Y, Marcus Â GM, Keung Â E, Varosy Â PD. Dual-chamber implantable cardioverter-defibrillator selection is associated with increased complication rates and mortality among patients enrolled in the NCDR implantable cardioverter-defibrillator registry. *J Am Coll Cardiol* Â 2011;58:1007â1013.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

449Friedman Â PA, Bradley Â D, Koestler Â C, Slusser Â J, Hodge Â D, Bailey Â K, et al. Â A prospective randomized trial of single- or dual-chamber implantable cardioverter-defibrillators to minimize inappropriate shock risk in primary sudden cardiac death prevention. *Europace* Â 2014;16:1460â1468.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

450Chen Â B-W, Liu Â Q, Wang Â X, Dang Â A-M. Are dual-chamber implantable cardioverter-defibrillators really better than single-chamber ones? A systematic review and meta-analysis. *J Interv Card Electrophysiol* Â 2014;39:273â280.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

451Epstein Â LM, Love Â CJ, Wilkoff Â BL, Chung Â MK, Hackler Â JW, Bongiorno Â MG, et al. Â Superior vena cava defibrillator coils make transvenous lead extraction more challenging and riskier. *J Am Coll Cardiol* Â 2013;61:987â989.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

452Larsen Â JM, HjortshÃ¸j Â SP, Nielsen Â JC, Johansen Â JB, Petersen Â HH, Haarlo Â J, et al. Â Single-coil and dual-coil defibrillator leads and association with clinical outcomes in a complete Danish nationwide ICD cohort. *Heart Rhythm* Â 2016;13:706â712.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

453Kumar Â KR, Mandleywala Â SN, Madias Â C, Weinstock Â J, Rowin Â EJ, Maron Â BJ, et al. Â Single coil implantable cardioverter defibrillator leads in patients with hypertrophic cardiomyopathy. *Am J Cardiol* Â 2020;125:1896â1900.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

454Friedman Â PA, Rasmussen Â MJ, Grice Â S, Trusty Â J, Glikson Â M, Stanton Â MS. Defibrillation thresholds are increased by right-sided implantation of totally transvenous implantable cardioverter defibrillators. *Pacing Clin Electrophysiol* Â 1999;22:1186â1192.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

455Stoevelaar Â R, Brinkman-Stoppelenburg Â A, Bhagwandien Â RE, van Bruchem-Visser Â RL, Theuns Â DA, van der Heide Â A, et al. Â The incidence and impact of implantable cardioverter defibrillator shocks in the last phase of life: an integrated review. *Eur J Cardiovasc Nurs* Â 2018;17:477â485.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 456Kapa S, Mueller PS, Hayes DL, Asirvatham SJ. Perspectives on withdrawing pacemaker and implantable cardioverter-defibrillator therapies at end of life: results of a survey of medical and legal professionals and patients. *Mayo Clin Proc* 2010;85:981-990.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 457Padeletti L, Arnar DO, Boncinelli L, Brachman J, Camm JA, Daubert JC, et al. EHRA Expert Consensus Statement on the management of cardiovascular implantable electronic devices in patients nearing end of life or requesting withdrawal of therapy. *Europace* 2010;12:1480-1489.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 458Stoevelaar R, Brinkman-Stoppelenburg A, van Driel AG, Theuns DA, Bhagwandien RE, van Bruchem-Visser RL, et al. Trends in time in the management of the implantable cardioverter defibrillator in the last phase of life: a retrospective study of medical records. *Eur J Cardiovasc Nurs* 2019;18:449-457.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 459Kirkpatrick JN, Gottlieb M, Sehgal P, Patel R, Verdino RJ. Deactivation of implantable cardioverter defibrillators in terminal illness and end of life care. *Am J Cardiol* 2012;109:914-919.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 460Stevenson WG, Khan H, Sager P, Saxon LA, Middlekauff HR, Natterson PD, et al. Identification of reentry circuit sites during catheter mapping and radiofrequency ablation of ventricular tachycardia late after myocardial infarction. *Circulation* 1993;88:1647-1670.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 461de Bakker JM, van Capelle FJ, Janse MJ, Tasseron S, Vermeulen JT, de Jonge N, et al. Slow conduction in the infarcted human heart. A zigzag course of activation. *Circulation* 1993;88:915-926.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 462de Chillou C, Lacroix D, Klug D, Magnin-Poull I, Marquié C, Messier M, et al. Isthmus characteristics of reentrant ventricular tachycardia after myocardial infarction. *Circulation* 2002;105:726-731.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 463Hsia HH, Callans DJ, Marchlinski FE. Characterization of endocardial electrophysiological substrate in patients with nonischemic cardiomyopathy and monomorphic ventricular tachycardia. *Circulation* 2003;108:704-710.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 464Soejima K, Stevenson WG, Sapp JL, Selwyn AP, Couper G, Epstein LM. Endocardial and epicardial radiofrequency ablation of ventricular tachycardia associated with dilated cardiomyopathy. *J Am Coll Cardiol* 2004;43:1834-1842.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 465Miljoen H, State S, Dechillou C, Magninpoull I, Dotto P, Andronache M, et al. Electroanatomic mapping characteristics of ventricular tachycardia in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Europace* 2005;7:516-524.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 466Priori SG, Blomström-Lundqvist C, Mazzanti A, Blom N, Borggrefe M, Camm J, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: The task force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J* 2015;36:2793-2867.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 467Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, et al. 2017 AHA/ACC/HRS Guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: executive summary. *J Am Coll Cardiol* 2018;72:1677-1749.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 468Towbin JA, McKenna WJ, Abrams DJ, Ackerman MJ, Calkins H, Darrieux FCC, et al. 2019 HRS expert consensus statement on evaluation, risk stratification, and management of arrhythmogenic cardiomyopathy: executive summary. *Heart Rhythm* 2019;16:e373-407.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 469Moss AJ, Greenberg H, Case RB, Zareba W, Hall WJ, Brown MW, et al. Long-term clinical course of patients after termination of ventricular tachyarrhythmia by an implanted defibrillator. *Circulation* 2004;110:3760-3765.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 470Poole JE, Johnson GW, Hellkamp AS, Anderson J, Callans DJ, Raitt MH, et al. Prognostic importance of defibrillator shocks in patients with heart failure. *N Engl J Med* 2008;359:1009-1017.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 471Sapp JL, Wells GA, Parkash R, Stevenson WG, Blier L, Sarrazin J-F, et al. Ventricular tachycardia ablation versus escalation of antiarrhythmic drugs. *N Engl J Med* 2016;375:1111-1121.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 472Piccini JP, Berger JS, Olshansky S, Connor CM. Amiodarone for the prevention of sudden cardiac death: a meta-analysis of randomized controlled trials. *Eur Heart J* 2009;30:1245-1253.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 473Palaniswamy C, Kolte D, Harikrishnan P, Khera S, Aronow WS, Mujib M, et al. Catheter ablation of postinfarction ventricular tachycardia: ten-year trends in utilization, in-hospital complications, and in-hospital mortality in the United States. *Heart Rhythm* 2014;11:2056-2063.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 474Caceres J, Jazayeri M, McKinnie J, Avitall B, Denker ST, Tchou P, et al. Sustained bundle branch reentry as a mechanism of clinical tachycardia. *Circulation* 1989;79:256-270.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 475Blanck Z, Dhala A, Deshpande S, Sra J, Jazayeri M, Akhtar M. Bundle branch reentrant ventricular tachycardia: cumulative experience in 48 patients. *J Cardiovasc Electrophysiol* 1993;4:253-262.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 476Chen H, Shi L, Yang B, Ju W, Zhang F, Yang G, et al. Electrophysiological characteristics of bundle branch reentry ventricular tachycardia in patients without structural heart disease. *Circ Arrhythm Electrophysiol* 2018;11:e006049.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 477Pathak RK, Fahed J, Santangeli P, Hyman MC, Liang JJ, Kubala M, et al. Long-term outcome of catheter ablation for treatment of bundle branch re-entrant tachycardia. *JACC Clin Electrophysiol* 2018;4:331-338.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 478Stevenson WG, Wilber DJ, Natale A, Jackman WM, Marchlinski FE, Talbert T, et al. Irrigated radiofrequency catheter ablation guided by electroanatomic mapping for recurrent ventricular tachycardia after myocardial infarction: the multicenter thermocool ventricular tachycardia ablation trial. *Circulation* 2008;118:2773-2782.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 479Della Bella P, Baratto F, Tsiachris D, Trevisi N, Vergara P, Bisceglia C, et al. Management of ventricular tachycardia in the setting of a dedicated unit for the treatment of complex ventricular arrhythmias: long-term outcome after ablation. *Circulation* 2013;127:1359-1368.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 480Maury P, Baratto F, Zeppenfeld K, Klein G, Delacretaz E, Sacher F, et al. Radio-frequency ablation as primary management of well-tolerated sustained monomorphic ventricular tachycardia in patients with structural heart disease and left ventricular ejection fraction over 30%. *Eur Heart J* 2014;35:1479-1485.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 481Tung R, Vaseghi M, Frankel DS, Vergara P, Di Biase L, Nagashima K, et al. Freedom from recurrent ventricular tachycardia after catheter ablation is associated with improved survival in patients with structural heart disease: an International VT Ablation Center Collaborative Group study. *Heart Rhythm* 2015;12:1997-2007.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 482Santangeli P, Zado ES, Supple GE, Haqqani HM, Garcia FC, Tschabrunn CM, et al. Long-term outcome with catheter ablation of ventricular tachycardia in patients with arrhythmogenic right ventricular

- cardiomyopathy. *Circ Arrhythm Electrophysiol* 2015;8:1413a-1421.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 483 Marchlinski FE, Haffajee CI, Beshai JF, Dickfeld T-ML, Gonzalez MD, Hsia HH, et al. Long-term success of irrigated radiofrequency catheter ablation of sustained ventricular tachycardia. *J Am Coll Cardiol* 2016;67:674a-683.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 484 Reddy VY, Reynolds MR, Neuzil P, Richardson AW, Taborsky M, Jongnarangsin K, et al. Prophylactic catheter ablation for the prevention of defibrillator therapy. *N Engl J Med* 2007;357:2657a-2665.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 485 Kuck K-H, Schaumann A, Eckardt L, Willems S, Ventura R, Delacortaz E, et al. Catheter ablation of stable ventricular tachycardia before defibrillator implantation in patients with coronary heart disease (VTACH): a multicentre randomised controlled trial. *Lancet* 2010;375:31a-40.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 486 Anter E, Kleber AG, Rottmann M, Leshem E, Barkagan M, Tschabrunn CM, et al. Infarct-related ventricular tachycardia. *JACC Clin Electrophysiol* 2018;4:1033a-1048.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 487 Marchlinski FE, Callans DJ, Gottlieb CD, Zado E. Linear ablation lesions for control of unmappable ventricular tachycardia in patients with ischemic and nonischemic cardiomyopathy. *Circulation* 2000;101:1288a-1296.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 488 de Chillou C, Groben L, Magnin-Poull I, Andronache M, Abbas MM, Zhang N, et al. Localizing the critical isthmus of postinfarct ventricular tachycardia: the value of pace-mapping during sinus rhythm. *Heart Rhythm* 2014;11:175a-181.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 489 Jaïs P, Maury P, Khairy P, Sacher F, Nault I, Komatsu Y, et al. Elimination of local abnormal ventricular activities: a new end point for substrate modification in patients with scar-related ventricular tachycardia. *Circulation* 2012;125:2184a-2196.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 490 Berruezo A, Fernandez-Armenta J. Lines, circles, channels, and clouds: looking for the best design for substrate-guided ablation of ventricular tachycardia. *Europace* 2014;16:943a-945.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 491 Di Biase L, Burkhardt JD, Lakkireddy D, Carbucicchio C, Mohanty S, Mohanty P, et al. Ablation of stable VTs versus substrate ablation in ischemic cardiomyopathy. *J Am Coll Cardiol* 2015;66:2872a-2882.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 492 Berruezo A, Fernandez-Armenta J, Andreu D, Penela D, Herczku C, Evertz R, et al. Scar dechanneling: new method for scar-related left ventricular tachycardia substrate ablation. *Circ Arrhythm Electrophysiol* 2015;8:326a-336.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 493 Haïssaguerre M, Shoda M, Jaïs P, Nogami A, Shah DC, Kautzner J, et al. Mapping and ablation of idiopathic ventricular fibrillation. *Circulation* 2002;106:962a-967.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 494 Shirai Y, Liang JJ, Santangeli P, Arkles JS, Schaller RD, Supple GE, et al. Comparison of the ventricular tachycardia circuit between patients with ischemic and nonischemic cardiomyopathies: detailed characterization by entrainment. *Circ Arrhythm Electrophysiol* 2019;12:e007249.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 495 Bhaskaran A, Tung R, Stevenson WG, Kumar S. Catheter ablation of VT in non-ischaemic cardiomyopathies: endocardial, epicardial and intramural approaches. *Heart Lung Circ* 2019;28:84a-101.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 496 Tung R, Raiman M, Liao H, Zhan X, Chung FP, Nagel R, et al. Simultaneous endocardial and epicardial delineation of 3D reentrant ventricular tachycardia. *J Am Coll Cardiol* 2020;75:884a-897.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 497Dinov Â B, Fiedler Â L, SchÃ¶nauer Â R, Bollmann Â A, Rolf Â S, Piorkowski Â C, et al. Â Outcomes in catheter ablation of ventricular tachycardia in dilated nonischemic cardiomyopathy compared with ischemic cardiomyopathy: results from the Prospective Heart Centre of Leipzig VT (HELP-VT) Study. *Circulation* Â 2014;129:728â736.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 498Proietti Â R, Essebag Â V, Beardsall Â J, Hache Â P, Pantano Â A, Wulffhart Â Z, et al. Â Substrate-guided ablation of haemodynamically tolerated and intolerated ventricular tachycardia in patients with structural heart disease: effect of cardiomyopathy type and acute success on long-term outcome. *Europace* Â 2015;17:461â467.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 499Ebert Â M, Richter Â S, Dinov Â B, Zeppenfeld Â K, Hindricks Â G. Evaluation and management of ventricular tachycardia in patients with dilated cardiomyopathy. *Heart Rhythm* Â 2019;16:624â631.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 500Proietti Â R, Lichelli Â L, Lellouche Â N, Dhanjal Â T. The challenge of optimising ablation lesions in catheter ablation of ventricular tachycardia. *J Arrhythmia* Â 2021;37:140â147.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 501Tokuda Â M, Sobieszczyk Â P, Eisenhauer Â AC, Kojodjojo Â P, Inada Â K, Koplan Â BA, et al. Â Transcoronary ethanol ablation for recurrent ventricular tachycardia after failed catheter ablation: an update. *Circ Arrhythm Electrophysiol* Â 2011;4:889â896.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 502Kreidieh Â B, RodrÃ­guez-MaÃ±ero Â M, Schurmann Â P, Ibarra-Cortez Â SH, Dave Â AS, ValderrÃ¡bano Â M. Retrograde coronary venous ethanol infusion for ablation of refractory ventricular tachycardia. *Circ Arrhythm Electrophysiol* Â 2016;9:e004352.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 503Nguyen Â DT, Tzou Â WS, Sandhu Â A, Gianni Â C, Anter Â E, Tung Â R, et al. Â Prospective multicenter experience with cooled radiofrequency ablation using high impedance irrigant to target deep myocardial substrate refractory to standard ablation. *JACC Clin Electrophysiol* Â 2018;4:1176â1185.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 504Stevenson Â WG, Tedrow Â UB, Reddy Â V, AbdelWahab Â A, Dukkupati Â S, John Â RM, et al. Â Infusion needle radiofrequency ablation for treatment of refractory ventricular arrhythmias. *J Am Coll Cardiol* Â 2019;73:1413â1425.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 505Igarashi Â M, Nogami Â A, Fukamizu Â S, Sekiguchi Â Y, Nitta Â J, Sakamoto Â N, et al. Â Acute and long-term results of bipolar radiofrequency catheter ablation of refractory ventricular arrhythmias of deep intramural origin. *Heart Rhythm* Â 2020;17:1500â1507.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 506Della Bella Â P, Peretto Â G, Paglino Â G, Bisceglia Â C, Radinovic Â A, Sala Â S, et al. Â Bipolar radiofrequency ablation for ventricular tachycardias originating from the interventricular septum: safety and efficacy in a pilot cohort study. *Heart Rhythm* Â 2020;17:2111â2118.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 507Cuculich Â PS, Schill Â MR, Kashani Â R, Mutic Â S, Lang Â A, Cooper Â D, et al. Â Noninvasive cardiac radiation for ablation of ventricular tachycardia. *N Engl J Med* Â 2017;377:2325â2336.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 508Robinson Â CG, Samson Â PP, Moore Â KMS, Hugo Â GD, Knutson Â N, Mutic Â S, et al. Â Phase I/II trial of electrophysiology-guided noninvasive cardiac radioablation for ventricular tachycardia. *Circulation* Â 2019;139:313â321.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 509Anter Â E, Hutchinson Â MD, Deo Â R, Haqqani Â HM, Callans Â DJ, Gerstenfeld Â EP, et al. Â Surgical ablation of refractory ventricular tachycardia in patients with nonischemic cardiomyopathy. *Circ Arrhythm Electrophysiol* Â 2011;4:494â500.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 510FernÃ¡ndez-Armenta Â J, Berrueto Â A, Andreu Â D, CamarÃ¡ O, Silva Â E, Serra Â L, et al. Â Three-dimensional architecture of scar and conducting channels based on high resolution ce-CMR: insights for

- ventricular tachycardia ablation. *Circ Arrhythm Electrophysiol* 2013;6:528â–537.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 511Mahida S, Sacher F, Dubois R, Sermesant M, Bogun F, Haïssaguerre M, et al. Cardiac imaging in patients with ventricular tachycardia. *Circulation* 2017;136:2491â–2507.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 512Andreu D, Penela D, Acosta J, Fernandez-Armenta J, Perea RJ, Soto-Iglesias D, et al. Cardiac magnetic resonanceâ–aided scar dechanneling: influence on acute and long-term outcomes. *Heart Rhythm* 2017;14:1121â–1128.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 513Kuo L, Liang JJ, Nazarian S, Marchlinski FE. Multimodality imaging to guide ventricular tachycardia ablation in patients with non-ischaemic cardiomyopathy. *Arrhythm Electrophysiol Rev* 2020;8:255â–264.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 514Roca-Luque I, Van Breukelen A, Alarcon F, Garre P, Tolosana JM, Borras R, et al. Ventricular scar channel entrances identified by new wideband cardiac magnetic resonance sequence to guide ventricular tachycardia ablation in patients with cardiac defibrillators. *Europace* 2020;22:598â–606.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 515Betensky BP, Marchlinski FE. Outcomes of catheter ablation of ventricular tachycardia in the setting of structural heart disease. *Curr Cardiol Rep* 2016;18:68.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 516Dukkipati SR, Koruth JS, Choudry S, Miller MA, Whang W, Reddy VY. Catheter ablation of ventricular tachycardia in structural heart disease. *J Am Coll Cardiol* 2017;70:2924â–2941.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 517Zeppenfeld K. Ventricular tachycardia ablation in nonischemic cardiomyopathy. *JACC Clin Electrophysiol* 2018;4:1123â–1140.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 518Guandalini GS, Liang JJ, Marchlinski FE. Ventricular tachycardia ablation. *JACC Clin Electrophysiol* 2019;5:1363â–1383.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 519Peichl P, Wichterle D, Pavlu L, Cihak R, Aldhoon B, Kautzner J. Complications of catheter ablation of ventricular tachycardia: a single-center experience. *Circ Arrhythm Electrophysiol* 2014;7:684â–690.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 520Katz DF, Turakhia MP, Sauer WH, Tzou WS, Heath RR, Zipse MM, et al. Safety of ventricular tachycardia ablation in clinical practice: findings from 9699 hospital discharge records. *Circ Arrhythm Electrophysiol* 2015;8:362â–370.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 521Cheung JW, Yeo I, Ip JE, Thomas G, Liu CF, Markowitz SM, et al. Outcomes, costs, and 30-day readmissions after catheter ablation of myocardial infarctâ–associated ventricular tachycardia in the real world: nationwide readmissions database 2010 to 2015. *Circ Arrhythm Electrophysiol* 2018;11:e006754.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 522Hendriks AA, Akca F, Dabiri Abkenari L, Khan M, Bhagwandien R, Yap S-C, et al. Safety and clinical outcome of catheter ablation of ventricular arrhythmias using contact force sensing: consecutive case series. *J Cardiovasc Electrophysiol* 2015;26:1224â–1229.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 523Nogami A. Purkinje-related arrhythmias part I: monomorphic ventricular tachycardias. *Pacing Clin Electrophysiol* 2011;34:624â–650.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 524Lerman BB. Mechanism, diagnosis, and treatment of outflow tract tachycardia. *Nat Rev Cardiol* 2015;12:597â–608.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 525Kobayashi Y. Idiopathic ventricular premature contraction and ventricular tachycardia: distribution of the origin, diagnostic algorithm, and catheter ablation. *J Nippon Med Sch* 2018;85:87â–94.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

526Tada Â H, Ito Â S, Naito Â S, Kurosaki Â K, Kubota Â S, Sugiyasu Â A, et al. Â Idiopathic ventricular arrhythmia arising from the mitral annulus: a distinct subgroup of idiopathic ventricular arrhythmias. *J Am Coll Cardiol* Â 2005;45:877â886.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

527Wasmer Â K, KÄbbe Â J, Decherer Â DG, Bittner Â A, Pott Â C, MÄnnig Â G, et al. Â Ventricular arrhythmias from the mitral annulus: patient characteristics, electrophysiological findings, ablation, and prognosis. *Heart Rhythm* Â 2013;10:783â788.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

528Tada Â H, Tadokoro Â K, Ito Â S, Naito Â S, Hashimoto Â T, Kaseno Â K, et al. Â Idiopathic ventricular arrhythmias originating from the tricuspid annulus: prevalence, electrocardiographic characteristics, and results of radiofrequency catheter ablation. *Heart Rhythm* Â 2007;4:7â16.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

529Yamada Â T, Doppalapudi Â H, McElderry Â HT, Okada Â T, Murakami Â Y, Inden Â Y, et al. Â Electrocardiographic and electrophysiological characteristics in idiopathic ventricular arrhythmias originating from the papillary muscles in the left ventricle: relevance for catheter ablation. *Circ Arrhythm Electrophysiol* Â 2010;3:324â331.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

530Macias Â C, Nakamura Â K, Tung Â R, Boyle Â NG, Kalyanam Â S, Bradfield Â JS. Importance of delayed enhanced cardiac MRI in idiopathic RVOT-VT: differentiating mimics including early stage ARVC and cardiac sarcoidosis. *J Atr Fibrillation* Â 2014;7:1097.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) [Â](#)

531Heeger Â C-H, Hayashi Â K, Kuck Â K-H, Ouyang Â F. Catheter ablation of idiopathic ventricular arrhythmias arising from the cardiac outflow tractsârecent insights and techniques for the successful treatment of common and challenging cases. *Circ J* Â 2016;80:1073â1086.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

532Pathak Â RK, Ariyaratna Â N, Garcia Â FC, Sanders Â P, Marchlinski Â FE. Catheter ablation of idiopathic ventricular arrhythmias. *Heart Lung Circ* Â 2019;28:102â109.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

533Yamada Â T, McElderry Â HT, Doppalapudi Â H, Murakami Â Y, Yoshida Â Y, Yoshida Â N, et al. Â Idiopathic ventricular arrhythmias originating from the aortic root prevalence, electrocardiographic and electrophysiologic characteristics, and results of radiofrequency catheter ablation. *J Am Coll Cardiol* Â 2008;52:139â147.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

534Van Herendaël Â H, Garcia Â F, Lin Â D, Riley Â M, Bala Â R, Cooper Â J, et al. Â Idiopathic right ventricular arrhythmias not arising from the outflow tract: prevalence, electrocardiographic characteristics, and outcome of catheter ablation. *Heart Rhythm* Â 2011;8:511â518.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

535Latchamsetty Â R, Yokokawa Â M, Morady Â F, Kim Â HM, Mathew Â S, Tilz Â R, et al. Â Multicenter outcomes for catheter ablation of idiopathic premature ventricular complexes. *JACC Clin Electrophysiol* Â 2015;1:116â123.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

536Liu Â Y, Fang Â Z, Yang Â B, Kojodjojo Â P, Chen Â H, Ju Â W, et al. Â Catheter ablation of fascicular ventricular tachycardia: long-term clinical outcomes and mechanisms of recurrence. *Circ Arrhythm Electrophysiol* Â 2015;8:1443â1451.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

537Hayashi Â T, Liang Â JJ, Shirai Â Y, Kuo Â L, Muser Â D, Kubala Â M, et al. Â Trends in successful ablation sites and outcomes of ablation for idiopathic outflow tract ventricular arrhythmias. *JACC Clin Electrophysiol* Â 2020;6:221â230.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

538FarrÃ© Â J, Wellens Â HJ. Philippe Coumel: a founding father of modern arrhythmology. *Europace* Â 2004;6:464â465.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

- 539Neira Â V, Enriquez Â A, Simpson Â C, Baranchuk Â A. Update on long QT syndrome. *J Cardiovasc Electrophysiol* Â 2019;30:3068â^{PADSTs}3078.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 540Winbo Â A, Paterson Â DJ. The brain-heart connection in sympathetically triggered inherited arrhythmia syndromes. *Heart Lung Circ* Â 2020;29:529â^{PADSTs}537.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 541Schwartz Â PJ, Priori Â SG, Cerrone Â M, Spazzolini Â C, Odero Â A, Napolitano Â C, et al. Â Left cardiac sympathetic denervation in the management of high-risk patients affected by the long-QT syndrome. *Circulation* Â 2004;109:1826â^{PADSTs}1833.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 542Surman Â TL, Stuklis Â RG, Chan Â JC. Thoracoscopic sympathectomy for long QT syndrome. Literature review and case study. *Heart Lung Circ* Â 2019;28:486â^{PADSTs}494.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 543Orvin Â K, Eisen Â A, Goldenberg Â I, Gottlieb Â S, Kornowski Â R, Matetzky Â S, et al. Â Outcome of contemporary acute coronary syndrome complicated by ventricular tachyarrhythmias. *Europace* Â 2016;18:219â^{PADSTs}226.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 544Demirel Â F, Rasoul Â S, Elvan Â A, Ottervanger Â JP, Dambrink Â J-HE, Gosselink Â ATM, et al. Â Impact of out-of-hospital cardiac arrest due to ventricular fibrillation in patients with ST-elevation myocardial infarction admitted for primary percutaneous coronary intervention: Impact of ventricular fibrillation in STEMI patients. *Eur Heart J Acute Cardiovasc Care* Â 2015;4:16â^{PADSTs}23.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 545Mehta Â RH, Starr Â AZ, Lopes Â RD, Hochman Â JS, Widimsky Â P, Pieper Â KS, et al. Â Incidence of and outcomes associated with ventricular tachycardia or fibrillation in patients undergoing primary percutaneous coronary intervention. *JAMA* Â 2009;301:1779â^{PADSTs}1789.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 546Demidova Â MM, Carlson Â J, Erlinge Â D, Platonov Â PG. Predictors of ventricular fibrillation at reperfusion in patients with acute ST-elevation myocardial infarction treated by primary percutaneous coronary intervention. *Am J Cardiol* Â 2015;115:417â^{PADSTs}422.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 547Cheng Â Y-J, Li Â Z-Y, Yao Â F-J, Xu Â X-J, Ji Â C-C, Chen Â X-M, et al. Â Early repolarization is associated with a significantly increased risk of ventricular arrhythmias and sudden cardiac death in patients with structural heart diseases. *Heart Rhythm* Â 2017;14:1157â^{PADSTs}1164.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 548Dumas Â F, Bougouin Â W, Geri Â G, Lamhaut Â L, Rosencher Â J, PÃˆne Â F, et al. Â Emergency percutaneous coronary intervention in post-cardiac arrest patients without ST-segment elevation pattern: insights from the PROCAT II registry. *JACC Cardiovasc Interv* Â 2016;9:1011â^{PADSTs}1018.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 549Chatterjee Â S, Chaudhuri Â D, Vedanthan Â R, Fuster Â V, Ibanez Â B, Bangalore Â S, et al. Â Early intravenous beta-blockers in patients with acute coronary syndromeâ^{PADSTs}a meta-analysis of randomized trials. *Int J Cardiol* Â 2013;168:915â^{PADSTs}921.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 550Roolvink Â V, IbÃ±ez Â B, Ottervanger Â JP, Pizarro Â G, van Royen Â N, Mateos Â A, et al. Â Early intravenous beta-blockers in patients with ST-segment elevation myocardial infarction before primary percutaneous coronary intervention. *J Am Coll Cardiol* Â 2016;67:2705â^{PADSTs}2715.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 551Piccini Â JP, Hranitzky Â PM, Kilaru Â R, Rouleau Â J-L, White Â HD, Aylward Â PE, et al. Â Relation of mortality to failure to prescribe beta blockers acutely in patients with sustained ventricular tachycardia and ventricular fibrillation following acute myocardial infarction (from the VALsartan In Acute myocardial iNfarcTion trial [VALIANT] Registry). *Am J Cardiol* Â 2008;102:1427â^{PADSTs}1432.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 552Nademanee Â K, Taylor Â R, Bailey Â WE, Rieders Â DE, Kosar Â EM. Treating electrical storm: sympathetic blockade versus advanced cardiac life support-guided therapy. *Circulation*

Â 2000;102:742â747.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

553Bundgaard Â JS, Jacobsen Â PK, Grand Â J, Lindholm Â MG, Hassager Â C, Pehrson Â S, et al. Â Deep sedation as temporary bridge to definitive treatment of ventricular arrhythmia storm. *Eur Heart J Acute Cardiovasc Care* Â 2020;9:657â664.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â

554Piccini Â JP, Schulte Â PJ, Pieper Â KS, Mehta Â RH, White Â HD, Van de Werf Â F, et al. Â Antiarrhythmic drug therapy for sustained ventricular arrhythmias complicating acute myocardial infarction. *Crit Care Med* Â 2011;39:78â83.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

555Dorian Â P, Cass Â D, Schwartz Â B, Cooper Â R, Gelaznikas Â R, Barr Â A. Amiodarone as compared with lidocaine for shock-resistant ventricular fibrillation. *N Engl J Med* Â 2002;346:884â890.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

556Echt Â DS, Liebson Â PR, Mitchell Â LB, Peters Â RW, Obias-Manno Â D, Barker Â AH, et al. Â Mortality and morbidity in patients receiving encainide, flecainide, or placebo. The Cardiac Arrhythmia Suppression Trial. *N Engl J Med* Â 1991;324:781â788.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

557Baudry Â G, Sonnevile Â R, Waintraub Â X, Lebreton Â G, Deguillard Â C, Mertens Â E, et al. Â Extracorporeal membrane oxygenation to support life-threatening drug-refractory electrical storm. *Crit Care Med* Â 2020;48:e856âe863.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

558Demidova Â MM, Smith Â JG, HÅljljer Â C-J, Holmqvist Â F, Erlinge Â D, Platonov Â PG. Prognostic impact of early ventricular fibrillation in patients with ST-elevation myocardial infarction treated with primary PCI. *Eur Heart J Acute Cardiovasc Care* Â 2012;1:302â311.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â

559Podolecki Â T, Lenarczyk Â R, Kowalczyk Â J, Jedrzejczyk-Patej Â E, Chodor Â P, Mazurek Â M, et al. Â Prognostic significance of complex ventricular arrhythmias complicating ST-segment elevation myocardial infarction. *Am J Cardiol* Â 2018;121:805â809.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

560Bougouin Â W, Marijon Â E, Puymirat Â E, Defaye Â P, Celermajer Â DS, Le Heuzey Â J-Y, et al. Â Incidence of sudden cardiac death after ventricular fibrillation complicating acute myocardial infarction: a 5-year cause-of-death analysis of the FAST-MI 2005 registry. *Eur Heart J* Â 2014;35:116â122.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

561Liang Â JJ, Hodge Â DO, Mehta Â RA, Russo Â AM, Prasad Â A, Cha Â Y-M. Outcomes in patients with sustained ventricular tachyarrhythmias occurring within 48 h of acute myocardial infarction: when is ICD appropriate? Â *Europace* Â 2014;16:1759â1766.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

562Ahn Â J-M, Lee Â KH, Yoo Â S-Y, Cho Â Y-R, Suh Â J, Shin Â E-S, et al. Â Prognosis of variant angina manifesting as aborted sudden cardiac death. *J Am Coll Cardiol* Â 2016;68:137â145.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

563Sueda Â S, Kohno Â H. Optimal medications and appropriate implantable cardioverter-defibrillator shocks in aborted sudden cardiac death due to coronary spasm. *Intern Med* Â 2018;57:1361â1369.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

564Rodr guez-Ma ero Â M, Oloriz Â T, le Polain de Waroux Â J-B, Burri Â H, Kreidieh Â B, de Asmundis Â C, et al. Â Long-term prognosis of patients with life-threatening ventricular arrhythmias induced by coronary artery spasm. *Europace* Â 2018;20:851â858.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

565Solomon Â SD, Zelenkofske Â S, McMurray Â JJV, Finn Â PV, Velazquez Â E, Ertl Â G, et al. Â Sudden death in patients with myocardial infarction and left ventricular dysfunction, heart failure, or both. *N Engl J Med* Â 2005;352:2581â2588.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

566Busk Â M, Maeng Â M, Kristensen Â SD, Thuesen Â L, Krusell Â LR, Mortensen Â LS, et al. Â Timing, causes, and predictors of death after three yearsâ follow-up in the Danish Multicenter Randomized Study of

- Fibrinolysis versus Primary Angioplasty in Acute Myocardial Infarction (DANAMI-2) trial. *Am J Cardiol* 2009;104:210a-215.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 567St John Sutton M, Pfeffer MA, Plappert T, Rouleau JL, Moy LA, Dagenais GR, et al. Quantitative two-dimensional echocardiographic measurements are major predictors of adverse cardiovascular events after acute myocardial infarction. The protective effects of captopril. *Circulation* 1994;89:68a-75.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 568S  holm H, L  nberg J, Andersen MJ, Vejlsr  p N, Engstr  m T, M  ller JE, et al. Repeated echocardiography after first ever ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention: is it necessary? *Eur Heart J Acute Cardiovasc Care* 2015;4:528a-536.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 569Steinbeck G, Andresen D, Seidl K, Brachmann J, Hoffmann E, Wojciechowski D, et al. Defibrillator implantation early after myocardial infarction. *N Engl J Med* 2009;361:1427a-1436.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 570Hohnloser SH, Kuck KH, Dorian P, Roberts RS, Hampton JR, Hatala R, et al. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med* 2004;351:2481a-2488.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 571Exner DV, Kavanagh KM, Slawnych MP, Mitchell LB, Ramadan D, Aggarwal SG, et al. Noninvasive risk assessment early after a myocardial infarction the REFINe study. *J Am Coll Cardiol* 2007;50:2275a-2284.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 572Zaman S, Narayan A, Thiagalingam A, Sivagangabalan G, Thomas S, Ross DL, et al. Long-term arrhythmia-free survival in patients with severe left ventricular dysfunction and no inducible ventricular tachycardia after myocardial infarction. *Circulation* 2014;129:848a-854.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 573Daubert MA, White JA, Al-Khalidi HR, Velazquez EJ, Rao SV, Crowley AL, et al. Cardiac remodeling after large ST-elevation myocardial infarction in the current therapeutic era. *Am Heart J* 2020;223:87a-97.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 574Chew DS, Heikki H, Schmidt G, Kavanagh KM, Dommasch M, Bloch Thomsen PE, et al. Change in left ventricular ejection fraction following first myocardial infarction and outcome. *JACC Clin Electrophysiol* 2018;4:672a-682.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 575B  nsch D, Oyang F, Antz M, Arentz T, Weber R, Val-Mejias JE, et al. Successful catheter ablation of electrical storm after myocardial infarction. *Circulation* 2003;108:3011a-3016.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 576Altmann DR, Mutschelknauss M, Ehl N, Koller M, Schaer B, J  rg L, et al. Prevalence of severely impaired left ventricular ejection fraction after reperfused ST-elevation myocardial infarction. *Swiss Med Wkly* 2013;143:w13869.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 577Shen L, Jhund PS, Petrie MC, Claggett BL, Barlera S, Cleland JGF, et al. Declining risk of sudden death in heart failure. *N Engl J Med* 2017;377:41a-51.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 578Cairns JA, Connolly SJ, Roberts R, Gent M. Randomised trial of outcome after myocardial infarction in patients with frequent or repetitive ventricular premature depolarisations: CAMIAT. Canadian Amiodarone Myocardial Infarction Arrhythmia Trial Investigators. *Lancet* 1997;349:675a-682.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 579Julian DG, Camm AJ, Frangin G, Janse MJ, Munoz A, Schwartz PJ, et al. Randomised trial of effect of amiodarone on mortality in patients with left-ventricular dysfunction after recent myocardial infarction: EMIAT. European Myocardial Infarct Amiodarone Trial Investigators. *Lancet* 1997;349:667a-674.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 580Clemens A M, Peichl A P, Wichterle A D, Pavl A L, Fliuhk A R, Aldhoon A B, et al. Catheter ablation of ventricular tachycardia as the first-line therapy in patients with coronary artery disease and preserved left ventricular systolic function: long-term results: VT ablation in patients with preserved LV function. *J Cardiovasc Electrophysiol* 2015;26:1105a-1110.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 581Pacífico A, Hohnloser SH, Williams JH, Tao B, Saksena S, Henry PD, et al. Prevention of implantable-defibrillator shocks by treatment with sotalol. D,l-Sotalol Implantable Cardioverter-Defibrillator Study Group. *N Engl J Med* 1999;340:1855a-1862.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 582Willems S, Tilz RR, Steven D, Kuck K-H, Wegscheider K, Gellera L, et al. Preventive or deferred ablation of ventricular tachycardia in patients with ischemic cardiomyopathy and implantable defibrillator (BERLIN VT): a multicenter randomized trial. *Circulation* 2020;141:1057a-1067.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 583Kuck K-H, Tilz RR, Deneke T, Hoffmann BA, Ventura R, Hansen PS, et al. Impact of substrate modification by catheter ablation on implantable cardioverter-defibrillator interventions in patients with unstable ventricular arrhythmias and coronary artery disease: results from the multicenter randomized controlled SMS (substrate modification study). *Circ Arrhythm Electrophysiol* 2017;10:e004422.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 584Olshansky B, Hahn EA, Hartz VL, Prater SP, Mason JW. Clinical significance of syncope in the electrophysiologic study versus electrocardiographic monitoring (ESVEM) trial. The ESVEM Investigators. *Am Heart J* 1999;137:878a-886.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 585Molossi A, Agrawal H, Mery CM, Krishnamurthy R, Masand P, Sexson Tejtel SK, et al. Outcomes in anomalous aortic origin of a coronary artery following a prospective standardized approach. *Circ Cardiovasc Interv* 2020;13:e008445.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 586Krasuski RA, Magyar D, Hart S, Kalahasti V, Lorber R, Hobbs R, et al. Long-term outcome and impact of surgery on adults with coronary arteries originating from the opposite coronary cusp. *Circulation* 2011;123:154a-162.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 587Jegatheeswaran A, Devlin PJ, McCrindle BW, Williams WG, Jacobs ML, Blackstone EH, et al. Features associated with myocardial ischemia in anomalous aortic origin of a coronary artery: a congenital heart surgeons' society study. *J Thorac Cardiovasc Surg* 2019;158:822a-834.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 588Jegatheeswaran A, Devlin PJ, Williams WG, Brothers JA, Jacobs ML, DeCampi WM, et al. Outcomes after anomalous aortic origin of a coronary artery repair: a congenital heart surgeons' society study. *J Thorac Cardiovasc Surg* 2020;160:757a-771.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 589Hoffmayer KS, Bhavé PD, Marcus GM, James CA, Tichnell C, Chopra N, et al. An electrocardiographic scoring system for distinguishing right ventricular outflow tract arrhythmias in patients with arrhythmogenic right ventricular cardiomyopathy from idiopathic ventricular tachycardia. *Heart Rhythm* 2013;10:477a-482.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 590Yokokawa M, Siontis KC, Kim HM, Stojanovska J, Latchamsetty R, Crawford T, et al. Value of cardiac magnetic resonance imaging and programmed ventricular stimulation in patients with frequent premature ventricular complexes undergoing radiofrequency ablation. *Heart Rhythm* 2017;14:1695a-1701.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 591Muser D, Santangeli P, Castro SA, Casado Arroyo R, Maeda S, Benhayon DA, et al. Risk stratification of patients with apparently idiopathic premature ventricular contractions: a multicenter international CMR registry. *JACC Clin Electrophysiol* 2020;6:722a-735.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 592Kjekshus J, Bathen J, Orning OM, Storstein L. A double-blind, crossover comparison of flecainide acetate and disopyramide phosphate in the treatment of ventricular premature complexes. *Am J Cardiol* 1984;53:72B-78B.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

593Hamon [Â](#) D, Swid [Â](#) MA, Rajendran [Â](#) PS, Liu [Â](#) A, Boyle [Â](#) NG, Shivkumar [Â](#) K, et al. [Â](#) Premature ventricular contraction diurnal profiles predict distinct clinical characteristics and beta-blocker responses. *J Cardiovasc Electrophysiol* [Â](#) 2019;30:836a-843.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

594Primeau [Â](#) R, Agha [Â](#) A, Giorgi [Â](#) C, Shenasa [Â](#) M, Nadeau [Â](#) R. Long term efficacy and toxicity of amiodarone in the treatment of refractory cardiac arrhythmias. *Can J Cardiol* [Â](#) 1989;5:98a-104.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) [Â](#)

595Ling [Â](#) Z, Liu [Â](#) Z, Su [Â](#) L, Zipunnikov [Â](#) V, Wu [Â](#) J, Du [Â](#) H, et al. [Â](#) Radiofrequency ablation versus antiarrhythmic medication for treatment of ventricular premature beats from the right ventricular outflow tract: prospective randomized study. *Circ Arrhythm Electrophysiol* [Â](#) 2014;7:237a-243.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

596Baksienė [Â](#) D, Sileikienė [Â](#) R, Sileikis [Â](#) V, Kazakevicius [Â](#) T, Zabiela [Â](#) V, Zebienė [Â](#) M, et al. [Â](#) Idiopathic ventricular tachycardia in children: curative therapy with radiofrequency ablation. *Medicina (Kaunas)* [Â](#) 2007;43:803a-807.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

597Blauxfox [Â](#) AD, Felix [Â](#) GL, Saul [Â](#) JP, Pediatric Catheter Ablation Registry. Radiofrequency catheter ablation in infants $\leq$ 18 months old: when is it done and how do they fare?: Short-term data from the pediatric ablation registry. *Circulation* [Â](#) 2001;104:2803a-2808.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

598Lapage [Â](#) MJ, Bradley [Â](#) DJ, Dick [Â](#) M. Verapamil in infants: an exaggerated fear? *Â Pediatr Cardiol* [Â](#) 2013;34:1532a-1534.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

599Lee [Â](#) AKY, Andrade [Â](#) J, Hawkins [Â](#) NM, Alexander [Â](#) G, Bennett [Â](#) MT, Chakrabarti [Â](#) S, et al. [Â](#) Outcomes of untreated frequent premature ventricular complexes with normal left ventricular function. *Heart* [Â](#) 2019;105:1408a-1413.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

600Baman [Â](#) TS, Lange [Â](#) DC, Ilg [Â](#) KJ, Gupta [Â](#) SK, Liu [Â](#) T-Y, Alguire [Â](#) C, et al. [Â](#) Relationship between burden of premature ventricular complexes and left ventricular function. *Heart Rhythm* [Â](#) 2010;7:865a-869.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

601van Huls van Taxis [Â](#) CFB, Piers [Â](#) SRD, de Riva Silva [Â](#) M, Dekkers [Â](#) OM, Pijnappels [Â](#) DA, Schalijs [Â](#) MJ, et al. [Â](#) Fatigue as presenting symptom and a high burden of premature ventricular contractions are independently associated with increased ventricular wall stress in patients with normal left ventricular function. *Circ Arrhythm Electrophysiol* [Â](#) 2015;8:1452a-1459.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

602Sharma [Â](#) N, Cortez [Â](#) D, Imundo [Â](#) JR. High burden of premature ventricular contractions in structurally normal hearts: to worry or not in pediatric patients? *Â Ann Noninvasive Electrocardiol* [Â](#) 2019;24:e12663.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

603Niawano [Â](#) S, Wakisaka [Â](#) Y, Niawano [Â](#) H, Fukaya [Â](#) H, Kurokawa [Â](#) S, Kiryu [Â](#) M, et al. [Â](#) Prognostic significance of frequent premature ventricular contractions originating from the ventricular outflow tract in patients with normal left ventricular function. *Heart* [Â](#) 2009;95:1230a-1237.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

604Krittayaphong [Â](#) R, Sriratanasathavorn [Â](#) C, Dumavibhat [Â](#) C, Pumprueg [Â](#) S, Boonyapisit [Â](#) W, Pooranawattanukul [Â](#) S, et al. [Â](#) Electrocardiographic predictors of long-term outcomes after radiofrequency ablation in patients with right-ventricular outflow tract tachycardia. *Europace* [Â](#) 2006;8:601a-606.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

605Komatsu [Â](#) Y, Nogami [Â](#) A, Kurosaki [Â](#) K, Morishima [Â](#) I, Masuda [Â](#) K, Ozawa [Â](#) T, et al. [Â](#) Fascicular ventricular tachycardia originating from papillary muscles: Purkinje network involvement in the reentrant circuit. *Circ Arrhythm Electrophysiol* [Â](#) 2017;10:e004549.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

606Kirk [Â](#) CR, Gibbs [Â](#) JL, Thomas [Â](#) R, Radley-Smith [Â](#) R, Qureshi [Â](#) SA. Cardiovascular collapse after verapamil in supraventricular tachycardia. *Arch Dis Child* [Â](#) 1987;62:1265a-1266.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

- 607Duffe DF, Shen WK, Smith HC. Suppression of frequent premature ventricular contractions and improvement of left ventricular function in patients with presumed idiopathic dilated cardiomyopathy. *Mayo Clin Proc* 1998;73:430â€433.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 608Chugh SS, Shen WK, Luria DM, Smith HC. First evidence of premature ventricular complex-induced cardiomyopathy: a potentially reversible cause of heart failure. *J Cardiovasc Electrophysiol* 2000;11:328â€329.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 609Penela D, Van Huls Van Taxis C, Van Huls Vans Taxis C, Aguinaga L, Fernandez-Armenta J, Mont L, et al. Neurohormonal, structural, and functional recovery pattern after premature ventricular complex ablation is independent of structural heart disease status in patients with depressed left ventricular ejection fraction: a prospective multicenter study. *J Am Coll Cardiol* 2013;62:1195â€1202.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 610Penela D, Acosta J, Aguinaga L, Tercedor L, Ordoez A, Fernandez-Armenta J, et al. Ablation of frequent PVC in patients meeting criteria for primary prevention ICD implant: safety of withholding the implant. *Heart Rhythm* 2015;12:2434â€2442.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 611Voskoboinik A, Hadjis A, Alhede C, Im SI, Park H, Moss J, et al. Predictors of adverse outcome in patients with frequent premature ventricular complexes: the ABC-VT risk score. *Heart Rhythm* 2020;17:1066â€1074.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 612Lee A, Denman R, Haqqani HM. Ventricular ectopy in the context of left ventricular systolic dysfunction: risk factors and outcomes following catheter ablation. *Heart Lung Circ* 2019;28:379â€388.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 613Sadron Blaye-Felice M, Hamon D, Sacher F, Pascale P, Rollin A, Duparc A, et al. Premature ventricular contraction-induced cardiomyopathy: related clinical and electrophysiologic parameters. *Heart Rhythm* 2016;13:103â€110.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 614Penela D, Fernandez-Armenta J, Aguinaga L, Tercedor L, Ordoez A, Bisbal F, et al. Clinical recognition of pure premature ventricular complex-induced cardiomyopathy at presentation. *Heart Rhythm* 2017;14:1864â€1870.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 615Aquaro GD, Pingitore A, Strata E, Di Bella G, Molinaro S, Lombardi M. Cardiac magnetic resonance predicts outcome in patients with premature ventricular complexes of left bundle branch block morphology. *J Am Coll Cardiol* 2010;56:1235â€1243.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 616Oebel S, Dinov B, Arya A, Hilbert S, Sommer P, Bollmann A, et al. ECG morphology of premature ventricular contractions predicts the presence of myocardial fibrotic substrate on cardiac magnetic resonance imaging in patients undergoing ablation. *J Cardiovasc Electrophysiol* 2017;28:1316â€1323.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 617Mountantonakis SE, Frankel DS, Gerstenfeld EP, Dixit S, Lin D, Hutchinson MD, et al. Reversal of outflow tract ventricular premature depolarization-induced cardiomyopathy with ablation: effect of residual arrhythmia burden and preexisting cardiomyopathy on outcome. *Heart Rhythm* 2011;8:1608â€1614.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 618Zang M, Zhang T, Mao J, Zhou S, He B. Beneficial effects of catheter ablation of frequent premature ventricular complexes on left ventricular function. *Heart* 2014;100:787â€793.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 619Wijnmaalen AP, Delgado V, Schalij MJ, van Huls van Taxis CFB, Holman ER, Bax JJ, et al. Beneficial effects of catheter ablation on left ventricular and right ventricular function in patients with frequent premature ventricular contractions and preserved ejection fraction. *Heart* 2010;96:1275â€1280.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 620Bogun F, Crawford T, Reich S, Koelling TM, Armstrong W, Good E, et al. Radiofrequency ablation of frequent, idiopathic premature ventricular complexes: comparison with a control group without

- intervention. *Heart Rhythm* 2007;4:863â867.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 621Sarrazin J-F, Labounty T, Kuhne M, Crawford T, Armstrong WF, Desjardins B, et al. Impact of radiofrequency ablation of frequent post-infarction premature ventricular complexes on left ventricular ejection fraction. *Heart Rhythm* 2009;6:1543â1549.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 622El Kadri M, Yokokawa M, Labounty T, Mueller G, Crawford T, Good E, et al. Effect of ablation of frequent premature ventricular complexes on left ventricular function in patients with nonischemic cardiomyopathy. *Heart Rhythm* 2015;12:706â713.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 623Lakkireddy D, Di Biase L, Ryschon K, Boria M, Swarup V, Reddy YM, et al. Radiofrequency ablation of premature ventricular ectopy improves the efficacy of cardiac resynchronization therapy in nonresponders. *J Am Coll Cardiol* 2012;60:1531â1539.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 624Singh SN, Fletcher RD, Fisher SG, Singh BN, Lewis HD, Deedwania PC, et al. Amiodarone in patients with congestive heart failure and asymptomatic ventricular arrhythmia. Survival trial of antiarrhythmic therapy in congestive heart failure. *N Engl J Med* 1995;333:77â82.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 625Zhong L, Lee Y-H, Huang X-M, Asirvatham SJ, Shen W-K, Friedman PA, et al. Relative efficacy of catheter ablation vs antiarrhythmic drugs in treating premature ventricular contractions: a single-center retrospective study. *Heart Rhythm* 2014;11:187â193.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 626Hyman MC, Mustin D, Supple G, Schaller RD, Santangeli P, Arkles J, et al. Class IC antiarrhythmic drugs for suspected premature ventricular contraction-induced cardiomyopathy. *Heart Rhythm* 2018;15:159â163.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 627Laurent G, Saal S, Amarouch MY, B  ziau DM, Marsman RFJ, Faivre L, et al. Multifocal ectopic Purkinje-related premature contractions: a new SCN5A-related cardiac channelopathy. *J Am Coll Cardiol* 2012;60:144â156.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 628Calloe K, Broendberg AK, Christensen AH, Pedersen LN, Olesen MS, de Los Angeles Tejada M, et al. Multifocal atrial and ventricular premature contractions with an increased risk of dilated cardiomyopathy caused by a Nav1.5 gain-of-function mutation (G213D). *Int J Cardiol* 2018;257:160â167.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 629Mann SA, Castro ML, Ohanian M, Guo G, Zodgekar P, Sheu A, et al. R222Q SCN5A mutation is associated with reversible ventricular ectopy and dilated cardiomyopathy. *J Am Coll Cardiol* 2012;60:1566â1573.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 630Beckermann TM, McLeod K, Murday V, Potet F, George AL. Novel SCN5A mutation in amiodarone-responsive multifocal ventricular ectopy-associated cardiomyopathy. *Heart Rhythm* 2014;11:1446â1453.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 631Doisne N, Waldmann V, Redheuil A, Waintraub X, Fressart V, Ader F, et al. A novel gain-of-function mutation in SCN5A responsible for multifocal ectopic Purkinje-related premature contractions. *Hum Mutat* 2020;41:850â859.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 632Elliott P, Andersson B, Arbustini E, Bilinska Z, Cecchi F, Charron P, et al. Classification of the cardiomyopathies: a position statement from the European Society Of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* 2008;29:270â276.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 633Codd MB, Sugrue DD, Gersh BJ, Melton LJ. Epidemiology of idiopathic dilated and hypertrophic cardiomyopathy. A population-based study in Olmsted County, Minnesota, 1975â1984. *Circulation*

- 1989;80:564â572.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 634Weintraub Â RG, Semsarian Â C, Macdonald Â P. Dilated cardiomyopathy. *Lancet* Â 2017;390:400â414.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 635Beggs Â SAS, Jhund Â PS, Jackson Â CE, McMurray Â JJV, Gardner Â RS. Non-ischaemic cardiomyopathy, sudden death and implantable defibrillators: a review and meta-analysis. *Heart* Â 2018;104:144â150.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 636Towbin Â JA, Lowe Â AM, Colan Â SD, Sleeper Â LA, Orav Â EJ, Clunie Â S, et al. Incidence, causes, and outcomes of dilated cardiomyopathy in children. *JAMA* Â 2006;296:1867â1876.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 637Pahl Â E, Sleeper Â LA, Canter Â CE, Hsu Â DT, Lu Â M, Webber Â SA, et al. Incidence of and risk factors for sudden cardiac death in children with dilated cardiomyopathy: a report from the Pediatric Cardiomyopathy Registry. *J Am Coll Cardiol* Â 2012;59:607â615.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 638Bharucha Â T, Lee Â KJ, Daubeney Â PEF, Nugent Â AW, Turner Â C, Sholler Â GF, et al. Sudden death in childhood cardiomyopathy: results from a long-term national population-based study. *J Am Coll Cardiol* Â 2015;65:2302â2310.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 639Pinto Â YM, Elliott Â PM, Arbustini Â E, Adler Â Y, Anastakis Â A, BÃhm Â M, et al. Proposal for a revised definition of dilated cardiomyopathy, hypokinetic non-dilated cardiomyopathy, and its implications for clinical practice: a position statement of the ESC working group on myocardial and pericardial diseases. *Eur Heart J* Â 2016;37:1850â1858.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 640Asselbergs Â FW, Sammani Â A, Elliott Â P, Gimeno Â JR, Tavazzi Â L, Tendera Â M, et al. Differences between familial and sporadic dilated cardiomyopathy: ESC EORP Cardiomyopathy & Myocarditis registry. *ESC Heart Fail* Â 2021;8:95â105.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 641Ader Â F, De Groote Â P, RÃant Â P, Rooryck-Thambo Â C, Dupin-Deguine Â D, Rambaud Â C, et al. FLNC pathogenic variants in patients with cardiomyopathies: prevalence and genotype-phenotype correlations. *Clin Genet* Â 2019;96:317â329.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 642Kayvanpour Â E, Sedaghat-Hamedani Â F, Amr Â A, Lai Â A, Haas Â J, Holzer Â DB, et al. Genotype-phenotype associations in dilated cardiomyopathy: meta-analysis on more than 8000 individuals. *Clin Res Cardiol* Â 2017;106:127â139.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 643Ortiz-Genga Â MF, Cuenca Â S, Dal Ferro Â M, Zorio Â E, Salgado-Aranda Â R, Climent Â V, et al. Truncating FLNC mutations are associated with high-risk dilated and arrhythmogenic cardiomyopathies. *J Am Coll Cardiol* Â 2016;68:2440â2451.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 644van den Hoogenhof Â MMG, Beqqali Â A, Amin Â AS, van der Made Â I, Aufiero Â S, Khan Â MAF, et al. RBM20 mutations induce an arrhythmogenic dilated cardiomyopathy related to disturbed calcium handling. *Circulation* Â 2018;138:1330â1342.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 645Gigli Â M, Merlo Â M, Graw Â SL, Barbati Â G, Rowland Â TJ, Slavov Â DB, et al. Genetic risk of arrhythmic phenotypes in patients with dilated cardiomyopathy. *J Am Coll Cardiol* Â 2019;74:1480â1490.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 646HeliÃ Â T, Elliott Â P, Koskenvuo Â JW, Gimeno Â JR, Tavazzi Â L, Tendera Â M, et al. ESC EORP cardiomyopathy registry: real-life practice of genetic counselling and testing in adult cardiomyopathy patients. *ESC Heart Fail* Â 2020;7:3013â3021.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 647Elming Â MB, Nielsen Â JC, Haarbo Â J, VidebÃk Â L, Korup Â E, Signorovitch Â J, et al. Age and outcomes of primary prevention implantable cardioverter-defibrillators in patients with nonischemic systolic heart failure. *Circulation* Â 2017;136:1772â1780.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

648Kristensen [Â](#) SL, Levy [Â](#) WC, Shadman [Â](#) R, Nielsen [Â](#) JC, Haarto [Â](#) J, Videbæk [Â](#) L, et al. [Â](#) Risk models for prediction of implantable cardioverter-defibrillator benefit. *JACC Heart Fail* [Â](#) 2019;7:717-724.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

649Wolff [Â](#) G, Lin [Â](#) Y, Karathanos [Â](#) A, Brockmeyer [Â](#) M, Wolters [Â](#) S, Nowak [Â](#) B, et al. [Â](#) Implantable cardioverter/defibrillators for primary prevention in dilated cardiomyopathy post-DANISH: an updated meta-analysis and systematic review of randomized controlled trials. *Clin Res Cardiol* [Â](#) 2017;106:501-513.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

650Kadish [Â](#) A, Dyer [Â](#) A, Daubert [Â](#) JP, Quigg [Â](#) R, Estes [Â](#) NAM, Anderson [Â](#) KP, et al. [Â](#) Prophylactic defibrillator implantation in patients with nonischemic dilated cardiomyopathy. *N Engl J Med* [Â](#) 2004;350:2151-2158.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

651Klem [Â](#) I, Klein [Â](#) M, Khan [Â](#) M, Yang [Â](#) EY, Nabi [Â](#) F, Ivanov [Â](#) A, et al. [Â](#) Relationship of LVEF and myocardial scar to long-term mortality risk and mode of death in patients with nonischemic cardiomyopathy. *Circulation* [Â](#) 2021;143:1343-1358.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

652van Rijsingen [Â](#) IAW, Arbustini [Â](#) E, Elliott [Â](#) PM, Mogensen [Â](#) J, Hermans-van Ast [Â](#) JF, van der Kooi [Â](#) AJ, et al. [Â](#) Risk factors for malignant ventricular arrhythmias in lamin a/c mutation carriers a European cohort study. *J Am Coll Cardiol* [Â](#) 2012;59:493-500.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

653Thuillot [Â](#) M, Maupain [Â](#) C, Gandjbakhch [Â](#) E, Waintraub [Â](#) X, Hidden-Lucet [Â](#) F, Isnard [Â](#) R, et al. [Â](#) External validation of risk factors for malignant ventricular arrhythmias in lamin A/C mutation carriers. *Eur J Heart Fail* [Â](#) 2019;21:253-254.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

654Pasotti [Â](#) M, Klersy [Â](#) C, Pilotto [Â](#) A, Marziliano [Â](#) N, Rapezzi [Â](#) C, Serio [Â](#) A, et al. [Â](#) Long-term outcome and risk stratification in dilated cardiomyopathies. *J Am Coll Cardiol* [Â](#) 2008;52:1250-1260.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

655Skjælsvik [Â](#) ET, Hasselberg [Â](#) NE, Dejgaard [Â](#) LA, Lie [Â](#) H, Andersen [Â](#) K, Holm [Â](#) T, et al. [Â](#) Exercise is associated with impaired left ventricular systolic function in patients with lamin A/C genotype. *J Am Heart Assoc* [Â](#) 2020;9:e012937.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

656Verstraelen [Â](#) TE, van Lint [Â](#) FHM, Bosman [Â](#) LP, de Brouwer [Â](#) R, Proost [Â](#) VM, Abeln [Â](#) BGS, et al. [Â](#) Prediction of ventricular arrhythmia in phospholamban p.Arg14del mutation carriers-reaching the frontiers of individual risk prediction. *Eur Heart J* [Â](#) 2021;42:2842-2850.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

657Knight [Â](#) BP, Goyal [Â](#) R, Pelosi [Â](#) F, Flemming [Â](#) M, Horwood [Â](#) L, Morady [Â](#) F, et al. [Â](#) Outcome of patients with nonischemic dilated cardiomyopathy and unexplained syncope treated with an implantable defibrillator. *J Am Coll Cardiol* [Â](#) 1999;33:1964-1970.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

658Desai [Â](#) AS, Fang [Â](#) JC, Maisel [Â](#) WH, Baughman [Â](#) KL. Implantable defibrillators for the prevention of mortality in patients with nonischemic cardiomyopathy: a meta-analysis of randomized controlled trials. *JAMA* [Â](#) 2004;292:2874-2879.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

659Merlo [Â](#) M, Gentile [Â](#) P, Artico [Â](#) J, Cannata [Â](#) A, Paldino [Â](#) A, De Angelis [Â](#) G, et al. [Â](#) Arrhythmic risk stratification in patients with dilated cardiomyopathy and intermediate left ventricular dysfunction. *J Cardiovasc Med (Hagerstown)* [Â](#) 2019;20:343-350.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) [Â](#)

660Zecchin [Â](#) M, Di Lenarda [Â](#) A, Gregori [Â](#) D, Merlo [Â](#) M, Pivetta [Â](#) A, Vitrella [Â](#) G, et al. [Â](#) Are nonsustained ventricular tachycardias predictive of major arrhythmias in patients with dilated cardiomyopathy on optimal medical treatment? *Pacing Clin Electro* [Â](#) 2008;31:290-299.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [Â](#)

661Link [Â](#) MS, Costeas [Â](#) XF, Griffith [Â](#) JL, Colburn [Â](#) CD, Estes [Â](#) NA, Wang [Â](#) PJ. High incidence of appropriate implantable cardioverter-defibrillator therapy in patients with syncope of unknown etiology and

- inducible ventricular arrhythmias. *J Am Coll Cardiol* 1997;29:370a-375.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 662Di Marco A, Brown PF, Bradley J, Nucifora G, Claver E, de Frutos F, et al. Improved risk stratification for ventricular arrhythmias and sudden death in patients with nonischemic dilated cardiomyopathy. *J Am Coll Cardiol* 2021;77:2890a-2905.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 663Kumar S, Romero J, Mehta NK, Fujii A, Kapur S, Baldinger SH, et al. Long-term outcomes after catheter ablation of ventricular tachycardia in patients with and without structural heart disease. *Heart Rhythm* 2016;13:1957a-1963.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 664Muser D, Santangeli P, Castro SA, Pathak RK, Liang JJ, Hayashi T, et al. Long-term outcome after catheter ablation of ventricular tachycardia in patients with nonischemic dilated cardiomyopathy. *Circ Arrhythm Electrophysiol* 2016;9:e004328.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 665Kumar S, Androulakis AFA, Sella J-M, Maury P, Gandjbakhch E, Waintraub X, et al. Multicenter experience with catheter ablation for ventricular tachycardia in lamin A/C cardiomyopathy. *Circ Arrhythm Electrophysiol* 2016;9:e004357.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 666Oloriz T, Silberbauer J, Maccabelli G, Mizuno H, Baratto F, Kirubakaran S, et al. Catheter ablation of ventricular arrhythmia in nonischemic cardiomyopathy: antero-septal versus inferolateral scar subtypes. *Circ Arrhythm Electrophysiol* 2014;7:414a-423.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 667Francone M. Role of cardiac magnetic resonance in the evaluation of dilated cardiomyopathy: diagnostic contribution and prognostic significance. *ISRN Radiol* 2014;2014:365404.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 668Goldberger JJ, Subramanian H, Patel T, Cunnane R, Kadish AH. Sudden cardiac death risk stratification in patients with nonischemic dilated cardiomyopathy. *J Am Coll Cardiol* 2014;63:1879a-1889.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 669Ebert M, Wijnmaalen AP, de Riva M, Trines SA, Androulakis AFA, Glashan CA, et al. Prevalence and prognostic impact of pathogenic variants in patients with dilated cardiomyopathy referred for ventricular tachycardia ablation. *JACC Clin Electrophysiol* 2020;6:1103a-1114.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 670Corrado D, Link MS, Calkins H. Arrhythmogenic right ventricular cardiomyopathy. *N Engl J Med* 2017;376:1489a-1490.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 671Quarta G, Muir A, Pantazis A, Syrris P, Gehmlich K, Garcia-Pavia P, et al. Familial evaluation in arrhythmogenic right ventricular cardiomyopathy: impact of genetics and revised task force criteria. *Circulation* 2011;123:2701a-2709.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 672Bhonsale A, Groeneweg JA, James CA, Dooijes D, Tichnell C, Jongbloed JDH, et al. Impact of genotype on clinical course in arrhythmogenic right ventricular dysplasia/cardiomyopathy-associated mutation carriers. *Eur Heart J* 2015;36:847a-855.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 673Rigato I, Bauce B, Rampazzo A, Zorzi A, Pilichou K, Mazzotti E, et al. Compound and digenic heterozygosity predicts lifetime arrhythmic outcome and sudden cardiac death in desmosomal gene-related arrhythmogenic right ventricular cardiomyopathy. *Circ Cardiovasc Genet* 2013;6:533a-542.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 674te Riele ASJM, James CA, Groeneweg JA, Sawant AC, Kammers K, Murray B, et al. Approach to family screening in arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Eur Heart J* 2016;37:755a-763.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 675Chivulescu A M, Lie A H, Popescu BA, Skulstad H, Edvardsen T, Jurcut RO, et al. High penetrance and similar disease progression in probands and in family members with arrhythmogenic cardiomyopathy. *Eur Heart J* 2020;41:1401â1410.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 676Rastegar N, Te Riele ASJM, James CA, Bhonsale A, Murray B, Tichnell C, et al. Fibrofatty changes: incidence at cardiac MR imaging in patients with arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Radiology* 2016;280:405â412.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 677Aquaro GD, Barison A, Todiere G, Grigoratos C, Ait Ali L, Di Bella G, et al. Usefulness of combined functional assessment by cardiac magnetic resonance and tissue characterization versus task force criteria for diagnosis of arrhythmogenic right ventricular cardiomyopathy. *Am J Cardiol* 2016;118:1730â1736.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 678te Riele ASJM, Bhonsale A, James CA, Rastegar N, Murray B, Burt JR, et al. Incremental value of cardiac magnetic resonance imaging in arrhythmic risk stratification of arrhythmogenic right ventricular dysplasia/cardiomyopathy-associated desmosomal mutation carriers. *J Am Coll Cardiol* 2013;62:1761â1769.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 679Corrado D, van Tintelen PJ, McKenna WJ, Hauer RNW, Anastakis A, Asimaki A, et al. Arrhythmogenic right ventricular cardiomyopathy: evaluation of the current diagnostic criteria and differential diagnosis. *Eur Heart J* 2020;41:1414â1429.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 680Sen-Chowdhry S, Syrris P, Ward D, Asimaki A, Sevdalis E, McKenna WJ. Clinical and genetic characterization of families with arrhythmogenic right ventricular dysplasia/cardiomyopathy provides novel insights into patterns of disease expression. *Circulation* 2007;115:1710â1720.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 681Cipriani A, Baucé B, De Lazzari M, Rigato I, Bariani R, Meneghin S, et al. Arrhythmogenic right ventricular cardiomyopathy: characterization of left ventricular phenotype and differential diagnosis with dilated cardiomyopathy. *J Am Heart Assoc* 2020;9:e014628.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 682Corrado D, Perazzolo Marra M, Zorzi A, Beffagna G, Cipriani A, Lazzari MD, et al. Diagnosis of arrhythmogenic cardiomyopathy: the Padua criteria. *Int J Cardiol* 2020;319:106â114.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 683Saberniak J, Hasselberg NE, Borgquist R, Platonov PG, Sarvari SI, Smith H, et al. Vigorous physical activity impairs myocardial function in patients with arrhythmogenic right ventricular cardiomyopathy and in mutation positive family members. *Eur J Heart Fail* 2014;16:1337â1344.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 684Lie A H, Deigaard LA, Saberniak J, Rootwelt C, Stokke MK, Edvardsen T, et al. Harmful effects of exercise intensity and exercise duration in patients with arrhythmogenic cardiomyopathy. *JACC Clin Electrophysiol* 2018;4:744â753.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 685James CA, Bhonsale A, Tichnell C, Murray B, Russell SD, Tandri H, et al. Exercise increases age-related penetrance and arrhythmic risk in arrhythmogenic right ventricular dysplasia/cardiomyopathy-associated desmosomal mutation carriers. *J Am Coll Cardiol* 2013;62:1290â1297.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 686Ruwald A-C, Marcus F, Estes NAM, Link M, McNitt S, Polonsky B, et al. Association of competitive and recreational sport participation with cardiac events in patients with arrhythmogenic right ventricular cardiomyopathy: results from the North American multidisciplinary study of arrhythmogenic right ventricular cardiomyopathy. *Eur Heart J* 2015;36:1735â1743.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 687Sawant AC, te Riele ASJM, Tichnell C, Murray B, Bhonsale A, Tandri H, et al. Safety of American Heart Association-recommended minimum exercise for desmosomal mutation carriers. *Heart Rhythm* 2016;13:199â207.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 688Thiene Â G, Nava Â A, Corrado Â D, Rossi Â L, Pennelli Â N. Right ventricular cardiomyopathy and sudden death in young people. *N Engl J Med* Â 1988;318:129â133.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 689Finocchiaro Â G, Papadakis Â M, Robertus Â J-L, Dhutia Â H, Steriotis Â AK, Tome Â M, et al. Â Etiology of sudden death in sports: insights from a United Kingdom Regional Registry. *J Am Coll Cardiol* Â 2016;67:2108â2115.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 690Marcus Â GM, Glidden Â DV, Polonsky Â B, Zareba Â W, Smith Â LM, Cannom Â DS, et al. Â Efficacy of antiarrhythmic drugs in arrhythmogenic right ventricular cardiomyopathy: a report from the North American ARVC Registry. *J Am Coll Cardiol* Â 2009;54:609â615.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 691Hulot Â J-S, Jouven Â X, Empana Â J-P, Frank Â R, Fontaine Â G. Natural history and risk stratification of arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circulation* Â 2004;110:1879â1884.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 692Wang Â W, Cadrin-Tourigny Â J, Bhonsale Â A, Tichnell Â C, Murray Â B, Monfredi Â O, et al. Â Arrhythmic outcome of arrhythmogenic right ventricular cardiomyopathy patients without implantable defibrillators. *J Cardiovasc Electrophysiol* Â 2018;29:1396â1402.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 693Pinamonti Â B, Dragos Â AM, Pyxaras Â SA, Merlo Â M, Pivetta Â A, Barbati Â G, et al. Â Prognostic predictors in arrhythmogenic right ventricular cardiomyopathy: results from a 10-year registry. *Eur Heart J* Â 2011;32:1105â1113.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 694Brun Â F, Groeneweg Â JA, Gear Â K, Sinagra Â G, van der Heijden Â J, Mestroni Â L, et al. Â Risk stratification in arrhythmic right ventricular cardiomyopathy without implantable cardioverter-defibrillators. *JACC Clin Electrophysiol* Â 2016;2:558â564.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 695Bhonsale Â A, James Â CA, Tichnell Â C, Murray Â B, Gagarin Â D, Philips Â B, et al. Â Incidence and predictors of implantable cardioverter-defibrillator therapy in patients with arrhythmogenic right ventricular dysplasia/cardiomyopathy undergoing implantable cardioverter-defibrillator implantation for primary prevention. *J Am Coll Cardiol* Â 2011;58:1485â1496.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 696Corrado Â D, Calkins Â H, Link Â MS, Leoni Â L, Favale Â S, Bevilacqua Â M, et al. Â Prophylactic implantable defibrillator in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia and no prior ventricular fibrillation or sustained ventricular tachycardia. *Circulation* Â 2010;122:1144â1152.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 697Platonov Â PG, Haugaa Â KH, Bundgaard Â H, Svensson Â A, Gilljam Â T, Hansen Â J, et al. Â Primary prevention of sudden cardiac death with implantable cardioverter-defibrillator therapy in patients with arrhythmogenic right ventricular cardiomyopathy. *Am J Cardiol* Â 2019;123:1156â1162.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 698Link Â MS, Laidlaw Â D, Polonsky Â B, Zareba Â W, McNitt Â S, Gear Â K, et al. Â Ventricular arrhythmias in the North American multidisciplinary study of ARVC: predictors, characteristics, and treatment. *J Am Coll Cardiol* Â 2014;64:119â125.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 699Schinkel Â AFL. Implantable cardioverter defibrillators in arrhythmogenic right ventricular dysplasia/cardiomyopathy: patient outcomes, incidence of appropriate and inappropriate interventions, and complications. *Circ Arrhythm Electrophysiol* Â 2013;6:562â568.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 700Corrado Â D, Leoni Â L, Link Â MS, Della Bella Â P, Gaita Â F, Curnis Â A, et al. Â Implantable cardioverter-defibrillator therapy for prevention of sudden death in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Circulation* Â 2003;108:3084â3091.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 701Mazzanti Â A, Ng Â K, Faragli Â A, Maragna Â R, Chiodaroli Â E, Orphanou Â N, et al. Â Arrhythmogenic right ventricular cardiomyopathy: clinical course and predictors of arrhythmic risk. *J Am Coll Cardiol*

- Â 2016;68:2540â²⁵⁵⁰.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 702Cadrin-Tourigny Â J, Bosman Â LP, Wang Â W, Tadros Â R, Bhonsale Â A, Bourfiss Â M, et al. Â Sudden cardiac death prediction in arrhythmogenic right ventricular cardiomyopathy: a multinational collaboration. *Circ Arrhythm Electrophysiol* Â 2021;14:e008509.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 703Bosman Â LP, Sammani Â A, James Â CA, Cadrin-Tourigny Â J, Calkins Â H, van Tintelen Â JP, et al. Â Predicting arrhythmic risk in arrhythmogenic right ventricular cardiomyopathy: a systematic review and meta-analysis. *Heart Rhythm* Â 2018;15:1097â¹¹⁰⁷.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 704Saguner Â AM, Vecchiati Â A, Baldinger Â SH, RÃ¼eger Â S, Medeiros-Domingo Â A, Mueller-Burri Â AS, et al. Â Different prognostic value of functional right ventricular parameters in arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Circ Cardiovasc Imaging* Â 2014;7:230â²³⁹.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 705Saguner Â AM, Medeiros-Domingo Â A, Schwyzer Â MA, On Â C-J, Haegeli Â LM, Wolber Â T, et al. Â Usefulness of inducible ventricular tachycardia to predict long-term adverse outcomes in arrhythmogenic right ventricular cardiomyopathy. *Am J Cardiol* Â 2013;111:250â²⁵⁷.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 706Orgeron Â GM, Te Riele Â A, Tichnell Â C, Wang Â W, Murray Â B, Bhonsale Â A, et al. Â Performance of the 2015 international task force consensus statement risk stratification algorithm for implantable cardioverter-defibrillator placement in arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circ Arrhythm Electrophysiol* Â 2018;11:e005593.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 707Orgeron Â GM, James Â CA, Te Riele Â A, Tichnell Â C, Murray Â B, Bhonsale Â A, et al. Â Implantable cardioverter-defibrillator therapy in arrhythmogenic right ventricular dysplasia/cardiomyopathy: predictors of appropriate therapy, outcomes, and complications. *J Am Heart Assoc* Â 2017;6:e006242.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 708Wichter Â T, Borggreffe Â M, Haverkamp Â W, Chen Â X, Breithardt Â G. Efficacy of antiarrhythmic drugs in patients with arrhythmogenic right ventricular disease. Results in patients with inducible and noninducible ventricular tachycardia. *Circulation* Â 1992;86:29â³⁷.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 709Mahida Â S, Venlet Â J, Saguner Â AM, Kumar Â S, Baldinger Â SH, AbdelWahab Â A, et al. Â Ablation compared with drug therapy for recurrent ventricular tachycardia in arrhythmogenic right ventricular cardiomyopathy: results from a multicenter study. *Heart Rhythm* Â 2019;16:536â⁵⁴³.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 710Ermakov Â S, Gerstenfeld Â EP, Svetlichnaya Â Y, Scheinman Â MM. Use of flecainide in combination antiarrhythmic therapy in patients with arrhythmogenic right ventricular cardiomyopathy. *Heart Rhythm* Â 2017;14:564â⁵⁶⁹.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 711Canpolat Â U, KabakÃ§i Â G, Aytemir Â K, Dural Â M, AÃ¼hiner Â L, Yorgun Â H, et al. Â Fragmented QRS complex predicts the arrhythmic events in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia: frQRS and outcomes in ARVC/D. *J Cardiovasc Electrophysiol* Â 2013;24:1260â¹²⁶⁶.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 712Martin Â A, Crawford Â J, Skinner Â JR, Smith Â W. High arrhythmic burden but low mortality during long-term follow-up in arrhythmogenic right ventricular cardiomyopathy. *Heart Lung Circ* Â 2016;25:275â²⁸¹.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 713Santangeli Â P, Dello Russo Â A, Pieroni Â M, Casella Â M, Di Biase Â L, Burkhardt Â JD, et al. Â Fragmented and delayed electrograms within fibrofatty scar predict arrhythmic events in arrhythmogenic right ventricular cardiomyopathy: Results from a prospective risk stratification study. *Heart Rhythm* Â 2012;9:1200â¹²⁰⁶.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 714Berrueto Â A, FernÃ¡ndez-Armenta Â J, Mont Â L, Zeljko Â H, Andreu Â D, Herczku Â C, et al. Â Combined endocardial and epicardial catheter ablation in arrhythmogenic right ventricular dysplasia

- incorporating scar dechanneling technique. *Circ Arrhythm Electrophysiol* 2012;5:111â121.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 715Ommen Â SR, Mital Â S, Burke Â MA, Day Â SM, Deswal Â A, Elliott Â P, et al. 2020 AHA/ACC guideline for the diagnosis and treatment of patients with hypertrophic cardiomyopathy: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2020;142:e533âe557.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 716Chan Â RH, Maron Â BJ, Olivetto Â I, Pencina Â MJ, Assenza Â GE, Haas Â T, et al. Prognostic value of quantitative contrast-enhanced cardiovascular magnetic resonance for the evaluation of sudden death risk in patients with hypertrophic cardiomyopathy. *Circulation* 2014;130:484â495.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 717He Â D, Ye Â M, Zhang Â L, Jiang Â B. Prognostic significance of late gadolinium enhancement on cardiac magnetic resonance in patients with hypertrophic cardiomyopathy. *Heart Lung* 2018;47:122â126.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 718Weissler-Snir Â A, Dorian Â P, Rakowski Â H, Care Â M, Spears Â D. Primary prevention implantable cardioverter-defibrillators in hypertrophic cardiomyopathyâare there predictors of appropriate therapy? *Â Heart Rhythm* 2021;18:63â70.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 719Khanna Â S, Wen Â I, Bhat Â A, Chen Â HHL, Gan Â GCH, Pathan Â F, et al. The role of multi-modality imaging in the diagnosis of cardiac amyloidosis: a focused update. *Front Cardiovasc Med* 2020;7:590557.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 720Pradella Â S, Grazzini Â G, De Amicis Â C, Letteriello Â M, Acquafresca Â M, Miele Â V. Cardiac magnetic resonance in hypertrophic and dilated cardiomyopathies. *Radiol Med* 2020;125:1056â1071.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 721Rosmini Â S, Biagini Â E, OâMahony Â C, Bulluck Â H, Ruozzi Â N, Lopes Â LR, et al. Relationship between aetiology and left ventricular systolic dysfunction in hypertrophic cardiomyopathy. *Heart* 2017;103:300â306.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 722Ho Â CY, Day Â SM, Ashley Â EA, Michels Â M, Pereira Â AC, Jacoby Â D, et al. Genotype and lifetime burden of disease in hypertrophic cardiomyopathy: insights from the Sarcomeric Human Cardiomyopathy Registry (SHaRe). *Circulation* 2018;138:1387â1398.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 723Kim Â HY, Park Â JE, Lee Â S-C, Jeon Â E-S, On Â YK, Kim Â SM, et al. Genotype-related clinical characteristics and myocardial fibrosis and their association with prognosis in hypertrophic cardiomyopathy. *J Clin Med* 2020;9:1671.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 724Maron Â BJ, Maron Â MS, Semsarian Â C. Double or compound sarcomere mutations in hypertrophic cardiomyopathy: a potential link to sudden death in the absence of conventional risk factors. *Heart Rhythm* 2012;9:57â63.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 725Wang Â J, Wang Â Y, Zou Â Y, Sun Â K, Wang Â Z, Ding Â H, et al. Malignant effects of multiple rare variants in sarcomere genes on the prognosis of patients with hypertrophic cardiomyopathy. *Eur J Heart Fail* 2014;16:950â957.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 726Semsarian Â C, Ingles Â J, Maron Â MS, Maron Â BJ. New perspectives on the prevalence of hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2015;65:1249â1254.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 727Maron Â BJ, Rowin Â EJ, Casey Â SA, Link Â MS, Lesser Â JR, Chan Â RHM, et al. Hypertrophic cardiomyopathy in adulthood associated with low cardiovascular mortality with contemporary management strategies. *J Am Coll Cardiol* 2015;65:1915â1928.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 728OâMahony Â C, Jichi Â F, Ommen Â SR, Christiaans Â I, Arbustini Â E, Garcia-Pavia Â P, et al. International External Validation Study of the 2014 European Society of Cardiology Guidelines on Sudden

Cardiac Death Prevention in Hypertrophic Cardiomyopathy (EVIDENCE-HCM). *Circulation* 2018;137:1015-1023.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Vriesendorp PA, Schinkel AFL, Liebrechts M, Theuns DAMJ, van Cleemput J, ten Cate FJ, et al. Validation of the 2014 European Society of Cardiology guidelines risk prediction model for the primary prevention of sudden cardiac death in hypertrophic cardiomyopathy. *Circ Arrhythm Electrophysiol* 2015;8:829-835.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Lorenzini M, Anastasiou Z, O'Leary Mahony C, Guttman OP, Gimeno JR, Monserrat L, et al. Mortality among referral patients with hypertrophic cardiomyopathy vs the general European population. *JAMA Cardiol* 2020;5:73-80.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Maron BJ, Rowin EJ, Casey SA, Haas TS, Chan RHM, Udelson JE, et al. Risk stratification and outcome of patients with hypertrophic cardiomyopathy >=60 years of age. *Circulation* 2013;127:585-593.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Gimeno JR, Tomé-Esteban M, Lofiego C, Hurtado J, Pantazis A, Mist B, et al. Exercise-induced ventricular arrhythmias and risk of sudden cardiac death in patients with hypertrophic cardiomyopathy. *Eur Heart J* 2009;30:2599-2605.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Pelliccia A, Lemme E, Maestrini V, Di Paolo FM, Pisicchio C, Di Gioia G, et al. Does sport participation worsen the clinical course of hypertrophic cardiomyopathy? Clinical outcome of hypertrophic cardiomyopathy in athletes. *Circulation* 2018;137:531-533.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Dejaard LA, Haland TF, Lie OH, Ribe M, Bjune T, Leren IS, et al. Vigorous exercise in patients with hypertrophic cardiomyopathy. *Int J Cardiol* 2018;250:157-163.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Monserrat L, Elliott PM, Gimeno JR, Sharma S, Penas-Lado M, McKenna WJ. Non-sustained ventricular tachycardia in hypertrophic cardiomyopathy: an independent marker of sudden death risk in young patients. *J Am Coll Cardiol* 2003;42:873-879.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Rowin EJ, Maron BJ, Carrick RT, Patel PP, Koethe B, Wells S, et al. Outcomes in patients with hypertrophic cardiomyopathy and left ventricular systolic dysfunction. *J Am Coll Cardiol* 2020;75:3033-3043.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Rowin EJ, Maron BJ, Haas TS, Garberich RF, Wang W, Link MS, et al. Hypertrophic cardiomyopathy with left ventricular apical aneurysm: implications for risk stratification and management. *J Am Coll Cardiol* 2017;69:761-773.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Sadoul N, Prasad K, Elliott PM, Bannerjee S, Frenneaux MP, McKenna WJ. Prospective prognostic assessment of blood pressure response during exercise in patients with hypertrophic cardiomyopathy. *Circulation* 1997;96:2987-2991.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Elliott PM, Poloniecki J, Dickie S, Sharma S, Monserrat L, Varnava A, et al. Sudden death in hypertrophic cardiomyopathy: identification of high risk patients. *J Am Coll Cardiol* 2000;36:2212-2218.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Behr ER, Elliott P, McKenna WJ. Role of invasive EP testing in the evaluation and management of hypertrophic cardiomyopathy. *Card Electrophysiol Rev* 2002;6:482-486.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

Gatzoulis KA, Georgopoulos S, Antoniou C-K, Anastakis A, Dilaveris P, Arsenos P, et al. Programmed ventricular stimulation predicts arrhythmic events and survival in hypertrophic cardiomyopathy. *Int J Cardiol* 2018;254:175-181.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 742Norrish A G, Qu A C, Field A E, Cervi A E, Khraiche A D, Klaassen A S, et al. External validation of the HCM Risk-Kids model for predicting sudden cardiac death in childhood hypertrophic cardiomyopathy. *Eur J Prev Cardiol* 2022;29:678â686.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 743Petryka-Mazurkiewicz A J, Ziolkowska A L, Kowalczyk-Domagala M, Mazurkiewicz A L, Boruc A A, Spiwak M, et al. LGE for risk stratification in primary prevention in children with HCM. *JACC Cardiovasc Imaging* 2020;13:2684â2686.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 744Elliott PM, Sharma A S, Varnava A A, Poloniecki A J, Rowland E, McKenna WJ. Survival after cardiac arrest or sustained ventricular tachycardia in patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol* 1999;33:1596â1601.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 745Cecchi A F, Maron BJ, Epstein SE. Long-term outcome of patients with hypertrophic cardiomyopathy successfully resuscitated after cardiac arrest. *J Am Coll Cardiol* 1989;13:1283â1288.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 746Maron BJ, Spirito P, Shen W-K, Haas TS, Formisano F, Link MS, et al. Implantable cardioverter-defibrillators and prevention of sudden cardiac death in hypertrophic cardiomyopathy. *JAMA* 2007;298:405â412.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 747Melacini P, Maron BJ, Bobbo F, Basso C, Tokajuk B, Zucchetto M, et al. Evidence that pharmacological strategies lack efficacy for the prevention of sudden death in hypertrophic cardiomyopathy. *Heart* 2007;93:708â710.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 748McKenna WJ, Oakley CM, Krikler DM, Goodwin JF. Improved survival with amiodarone in patients with hypertrophic cardiomyopathy and ventricular tachycardia. *Br Heart J* 1985;53:412â416.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 749Link MS, Bockstall K, Weinstock J, Alsheikh-Ali AA, Semsarian C, Estes NAM, et al. Ventricular tachyarrhythmias in patients with hypertrophic cardiomyopathy and defibrillators: triggers, treatment, and implications. *J Cardiovasc Electrophysiol* 2017;28:531â537.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 750Dallaglio PD, di Marco A, Moreno Weidmann Z, Perez L, Alzueta J, Garc a-Alberola A, et al. Antitachycardia pacing for shock prevention in patients with hypertrophic cardiomyopathy and ventricular tachycardia. *Heart Rhythm* 2020;17:1084â1091.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 751Adduci C, Semprini L, Palano F, Musumeci MB, Volpe M, Autore C, et al. Safety and efficacy of anti-tachycardia pacing in patients with hypertrophic cardiomyopathy implanted with an ICD. *Pacing Clin Electrophysiol* 2019;42:610â616.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 752Vaseghi M, Hu TY, Tung R, Vergara P, Frankel DS, Di Biase L, et al. Outcomes of catheter ablation of ventricular tachycardia based on etiology in nonischemic heart disease: an international ventricular tachycardia ablation center collaborative study. *JACC Clin Electrophysiol* 2018;4:1141â1150.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 753Igarashi M, Nogami A, Kurosaki K, Hanaki Y, Komatsu Y, Fukamizu S, et al. Radiofrequency catheter ablation of ventricular tachycardia in patients with hypertrophic cardiomyopathy and apical aneurysm. *JACC Clin Electrophysiol* 2018;4:339â350.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 754Dukkipati SR, d  s  s Avila A, Soejima K, Bala R, Inada K, Singh S, et al. Long-term outcomes of combined epicardial and endocardial ablation of monomorphic ventricular tachycardia related to hypertrophic cardiomyopathy. *Circ Arrhythm Electrophysiol* 2011;4:185â194.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 755Ross SB, Singer ES, Driscoll E, Nowak N, Yeates L, Puranik R, et al. Genetic architecture of left ventricular noncompaction in adults. *Hum Genome Var* 2020;7:33.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 756Weir-McCall Â JR, Yeap Â PM, Papagiorcopulo Â C, Fitzgerald Â K, Gandy Â SJ, Lambert Â M, et al. Â Left ventricular noncompaction: anatomical phenotype or distinct cardiomyopathy? Â J Am Coll Cardiol Â 2016;68:2157â2165.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 757Aung Â N, Doimo Â S, Ricci Â F, Sanghvi Â MM, Pedrosa Â C, Woodbridge Â SP, et al. Â Prognostic significance of left ventricular noncompaction: systematic review and meta-analysis of observational studies. Circ Cardiovasc Imaging Â 2020;13:e009712.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 758Grigoratos Â C, Barison Â A, Ivanov Â A, Andreini Â D, Amzulescu Â M-S, Mazurkiewicz Â L, et al. Â Meta-analysis of the prognostic role of late gadolinium enhancement and global systolic impairment in left ventricular noncompaction. JACC Cardiovasc Imaging Â 2019;12:2141â2151.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 759Richard Â P, Ader Â F, Roux Â M, Donal Â E, Eicher Â J-C, Aoutil Â N, et al. Â Targeted panel sequencing in adult patients with left ventricular non-compaction reveals a large genetic heterogeneity. Clin Genet Â 2019;95:356â367.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 760Seferović Â PM, Polovina Â M, Bauersachs Â J, Arad Â M, Gal Â TB, Lund Â LH, et al. Â Heart failure in cardiomyopathies: a position paper from the Heart Failure Association of the European Society of Cardiology. Eur J Heart Fail Â 2019;21:553â576.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 761Baig Â S, Edward Â NC, Kotecha Â D, Liu Â B, Nordin Â S, Kozor Â R, et al. Â Ventricular arrhythmia and sudden cardiac death in Fabry disease: a systematic review of risk factors in clinical practice. Europace Â 2018;20:f153âf161.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 762Sperry Â BW, Ikram Â A, Hachamovitch Â R, Valent Â J, Vranian Â MN, Phelan Â D, et al. Â Efficacy of chemotherapy for light-chain amyloidosis in patients presenting with symptomatic heart failure. J Am Coll Cardiol Â 2016;67:2941â2948.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 763Kristen Â AV, Dengler Â TJ, Hegenbart Â U, Schonland Â SO, Goldschmidt Â H, Sack Â F-U, et al. Â Prophylactic implantation of cardioverter-defibrillator in patients with severe cardiac amyloidosis and high risk for sudden cardiac death. Heart Rhythm Â 2008;5:235â240.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 764Merino Â JL, Peinado Â R. Arrhythmias associated with neuromuscular disorders. Card Electrophysiol Rev Â 2002;6:132â135.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 765Punnoose Â AR, Kaltman Â JR, Pastor Â W, McCarter Â R, He Â J, Spurney Â CF. Cardiac disease burden and risk of mortality in hospitalized muscular dystrophy patients. Pediatr Cardiol Â 2016;37:1290â1296.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 766Wahbi Â K, Meune Â C, Porcher Â R, Bâcane Â HM, Lazarus Â A, Laforêt Â P, et al. Â Electrophysiological study with prophylactic pacing and survival in adults with myotonic dystrophy and conduction system disease. JAMA Â 2012;307:1292â1301.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 767Lallemand Â B, Clementy Â N, Bernard-Brunet Â A, Pierre Â B, Corcia Â P, Fauchier Â L, et al. Â The evolution of infrahisian conduction time in myotonic dystrophy patients: clinical implications. Heart Â 2012;98:291â296.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 768Nogami Â A. Bundle branch reentry tachycardia. ESC CardioMed. 3rd ed. Oxford University Press; 2022, pp. 2270â2275.
- 769Sanna Â T, Dello Russo Â A, Toniolo Â D, Vytopil Â M, Pelargonio Â G, De Martino Â G, et al. Â Cardiac features of Emery-Dreifuss muscular dystrophy caused by lamin A/C gene mutations. Eur Heart J Â 2003;24:2227â2236.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 770Menon Â SC, Etheridge Â SP, Liesemer Â KN, Williams Â RV, Bardsley Â T, Heywood Â MC, et al. Â Predictive value of myocardial delayed enhancement in Duchenne muscular dystrophy. *Pediatr Cardiol* Â 2014;35:1279â1285.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 771Florian Â A, Ludwig Â A, Engelen Â M, Waltenberger Â J, R  sch Â S, Sechtem Â U, et al. Â Left ventricular systolic function and the pattern of late-gadolinium-enhancement independently and additively predict adverse cardiac events in muscular dystrophy patients. *J Cardiovasc Magn Reson* Â 2014;16:81.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 772Prystowsky Â EN, Pritchett Â EL, Roses Â AD, Gallagher Â J. The natural history of conduction system disease in myotonic muscular dystrophy as determined by serial electrophysiologic studies. *Circulation* Â 1979;60:1360â1364.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 773Trachtenberg Â BH, Hare Â JM. Inflammatory cardiomyopathic syndromes. *Circ Res* Â 2017;121:803â818.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 774Hulsmans Â M, Clauss Â S, Xiao Â L, Aguirre Â AD, King Â KR, Hanley Â A, et al. Â Macrophages facilitate electrical conduction in the heart. *Cell* Â 2017;169:510â522.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 775Youker Â KA, Assad-Kottner Â C, Cordero-Reyes Â AM, Trevino Â AR, Flores-Arredondo Â JH, Barrios Â R, et al. Â High proportion of patients with end-stage heart failure regardless of aetiology demonstrates anti-cardiac antibody deposition in failing myocardium: humoral activation, a potential contributor of disease progression. *Eur Heart J* Â 2014;35:1061â1068.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 776Adamo Â L, Rocha-Resende Â C, Prabhu Â SD, Mann Â DL. Reappraising the role of inflammation in heart failure. *Nat Rev Cardiol* Â 2020;17:269â285.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 777Sagar Â S, Liu Â PP, Cooper Â LT. Myocarditis. *Lancet* Â 2012;379:738â747.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 778Ammirati Â E, Cipriani Â M, Moro Â C, Raineri Â C, Pini Â D, Sormani Â P, et al. Â Clinical presentation and outcome in a contemporary cohort of patients with acute myocarditis: multicenter Lombardy registry. *Circulation* Â 2018;138:1088â1099.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 779Fabre Â A. Sudden adult death syndrome and other non-ischaemic causes of sudden cardiac death. *Heart* Â 2005;92:316â320.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 780Lyng   Â TH, Nielsen Â TS, Gregers Winkel Â B, Tfelt-Hansen Â J, Banner Â J. Sudden cardiac death caused by myocarditis in persons aged 1  49 years: a nationwide study of 14 294 deaths in Denmark. *Forensic Sci Res* Â 2019;4:247â256.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 781Maron Â BJ, Udelson Â JE, Bonow Â RO, Nishimura Â RA, Ackerman Â MJ, Estes Â NAM, et al. Â Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: task force 3: hypertrophic cardiomyopathy, arrhythmogenic right ventricular cardiomyopathy and other cardiomyopathies, and myocarditis. *J Am Coll Cardiol* Â 2015;66:2362â2371.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 782Kindermann Â I, Barth Â C, Mahfoud Â F, Ukena Â C, Lenski Â M, Yilmaz Â A, et al. Â Update on myocarditis. *J Am Coll Cardiol* Â 2012;59:779â792.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 783Caforio Â ALP, Pankuweit Â S, Arbustini Â E, Basso Â C, Gimeno-Blanes Â J, Felix Â SB, et al. Â Current state of knowledge on aetiology, diagnosis, management, and therapy of myocarditis: a position statement of the European Society of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* Â 2013;34:2636â2648.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 784Singh Â V, Mendirichaga Â R, Savani Â GT, Rodriguez Â A, Blumer Â V, Elmariah Â S, et al. Â Comparison of utilization trends, indications, and complications of endomyocardial biopsy in native versus

- donor hearts (from the Nationwide Inpatient Sample 2002 to 2014). *Am J Cardiol* 2018;121:356-363.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 785Aoyama N, Izumi T, Hiramori K, Isobe M, Kawana M, Hiroe M, et al. National survey of fulminant myocarditis in Japan: therapeutic guidelines and long-term prognosis of using percutaneous cardiopulmonary support for fulminant myocarditis (special report from a scientific committee). *Circ J* 2002;66:133-144.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 786Shah Z, Mohammed M, Vuddanda V, Ansari MW, Masoomi R, Gupta K. National trends, gender, management, and outcomes of patients hospitalized for myocarditis. *Am J Cardiol* 2019;124:131-136.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 787Anderson BR, Silver ES, Richmond ME, Liberman L. Usefulness of arrhythmias as predictors of death and resource utilization in children with myocarditis. *Am J Cardiol* 2014;114:1400-1405.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 788Kandolin R, Lehtonen J, Salmenkivi K, Räsänen-Sokolowski A, Lommi J, Kupari M. Diagnosis, treatment, and outcome of giant-cell myocarditis in the era of combined immunosuppression. *Circ Heart Fail* 2013;6:15-22.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 789Cooper LT, Berry GJ, Shabetai R. Idiopathic giant-cell myocarditis: a natural history and treatment. *N Engl J Med* 1997;336:1860-1866.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 790McMurray JJV, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the task force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2012;33:1787-1847.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 791Blauwet LA, Cooper LT. Myocarditis. *Prog Cardiovasc Dis* 2010;52:274-288.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 792JCS Joint Working Group. Guidelines for diagnosis and treatment of myocarditis (JCS 2009): digest version. *Circ J* 2011;75:734-743.
[Crossref](#) [Search ADS](#) [PubMed](#)
- 793Ali-Ahmed F, Dalgaard F, Al-Khatib SM. Sudden cardiac death in patients with myocarditis: Evaluation, risk stratification, and management. *Am Heart J* 2020;220:29-40.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 794Rosier L, Zouaghi A, Barrat V, Martins R, Probst V, Marijon E, et al. High risk of sustained ventricular arrhythmia recurrence after acute myocarditis. *J Clin Med* 2020;9:E848.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 795Maron BJ, Towbin JA, Thiene G, Antzelevitch C, Corrado D, Arnett D, et al. Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation* 2006;113:1807-1816.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 796D'Ambrósio A, Patti G, Manzoli A, Sinagra G, Di Lenarda A, Silvestri F, et al. The fate of acute myocarditis between spontaneous improvement and evolution to dilated cardiomyopathy: a review. *Heart* 2001;85:499-504.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 797Kasper EK, Agema WRP, Hutchins GM, Deckers JW, Hare JM, Baughman KL. The causes of dilated cardiomyopathy: a clinicopathologic review of 673 consecutive patients. *J Am Coll Cardiol* 1994;23:586-590.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 798GrÃ¼n S, Schumm J, Greulich S, Wagner A, Schneider S, Bruder O, et al. Long-term follow-up of biopsy-proven viral myocarditis. *J Am Coll Cardiol* 2012;59:1604â1615.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 799Bennett MK, Gilotra NA, Harrington C, Rao S, Dunn JM, Freitag TB, et al. Evaluation of the role of endomyocardial biopsy in 851 patients with unexplained heart failure from 2000â2009. *Circ Heart Fail* 2013;6:676â684.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 800Schumm J, Greulich S, Wagner A, GrÃ¼n S, Ong P, Bentz K, et al. Cardiovascular magnetic resonance risk stratification in patients with clinically suspected myocarditis. *J Cardiovasc Magn Reson* 2014;16:14.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 801Russo AD, Casella M, Pieroni M, Pelargonio G, Bartoletti S, Santangeli P, et al. Drug-refractory ventricular tachycardias after myocarditis: endocardial and epicardial radiofrequency catheter ablation. *Circ Arrhythm Electrophysiol* 2012;5:492â498.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 802Maccabelli G, Tsiachris D, Silberbauer J, Esposito A, Bisceglia C, Baratto F, et al. Imaging and epicardial substrate ablation of ventricular tachycardia in patients late after myocarditis. *Europace* 2014;16:1363â1372.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 803Berte B, Sacher F, Cochet H, Mahida S, Yamashita S, Lim H, et al. Postmyocarditis ventricular tachycardia in patients with epicardial-only scar: a specific entity requiring a specific approach: epicardial-only VT ablation. *J Cardiovasc Electrophysiol* 2015;26:42â50.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 804Maleszewski JJ, Orellana VM, Hodge DO, Kuhl U, Schultheiss H-P, Cooper LT. Long-term risk of recurrence, morbidity and mortality in giant cell myocarditis. *Am J Cardiol* 2015;115:1733â1738.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 805El-Assaad I, Al-Kindi SG, Oliveira GH, Boyle GJ, Aziz PF. Implantable cardioverter-defibrillator and wait-list outcomes in pediatric patients awaiting heart transplantation. *Heart Rhythm* 2015;12:2443â2448.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 806EkstrÃm K, Lehtonen J, Kandolin R, RÃisÃnen-Sokolowski A, Salmenkivi K, Kupari M. Long-term outcome and its predictors in giant cell myocarditis: giant cell myocarditis. *Eur J Heart Fail* 2016;18:1452â1458.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 807Rybicki BA, Iannuzzi MC, Frederick MM, Thompson BW, Rossman MD, Bresnitz EA, et al. Familial aggregation of sarcoidosis. A case-control etiologic study of sarcoidosis (ACCESS). *Am J Respir Crit Care Med* 2001;164:2085â2091.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 808Muchtar E, Blauwet LA, Gertz MA. Restrictive cardiomyopathy: genetics, pathogenesis, clinical manifestations, diagnosis, and therapy. *Circ Res* 2017;121:819â837.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 809Birnir DH, Sauer WH, Bogun F, Cooper JM, Culver DA, Duvernoy CS, et al. HRS expert consensus statement on the diagnosis and management of arrhythmias associated with cardiac sarcoidosis. *Heart Rhythm* 2014;11:1304â1323.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 810Birnir DH, Kandolin R, Nery PB, Kupari M. Cardiac manifestations of sarcoidosis: diagnosis and management. *Eur Heart J* 2016;38:2663â2670.
[Google Scholar](#) [OpenURL Placeholder Text](#)
- 811Trivieri MG, Spagnolo P, Birnir D, Liu P, Drake W, Kovacic JC, et al. Challenges in cardiac and pulmonary sarcoidosis: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;76:1878â1901.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 812Kandolin R, Lehtonen J, Airaksinen J, Vihinen T, Miettinen H, Ylitalo K, et al. Cardiac sarcoidosis: epidemiology, characteristics, and outcome over 25 years in a nationwide study. *Circulation*

Â 2015;131:624â632.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

813Hoogendoorn Â JC, Sramko Â M, Venlet Â J, Siontis Â KC, Kumar Â S, Singh Â R, et al.

Â Electroanatomical voltage mapping to distinguish right-sided cardiac sarcoidosis from arrhythmogenic right ventricular cardiomyopathy. *JACC Clin Electrophysiol* Â 2020;6:696â707.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

814Banba Â K, Kusano Â KF, Nakamura Â K, Morita Â H, Ogawa Â A, Ohtsuka Â F, et al. Â Relationship

between arrhythmogenesis and disease activity in cardiac sarcoidosis. *Heart Rhythm* Â 2007;4:1292â1299.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

815Takaya Â Y, Kusano Â KF, Nakamura Â K, Ito Â H. Outcomes in patients with high-degree atrioventricular

block as the initial manifestation of cardiac sarcoidosis. *Am J Cardiol* Â 2015;115:505â509.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

816Nordenswan Â H-K, Lehtonen Â J, Ekstr  m Â K, Kandolin Â R, Simonen Â P, M  yr  np     Â M, et

al. Â Outcome of cardiac sarcoidosis presenting with high-grade atrioventricular block. *Circ Arrhythm Electrophysiol* Â 2018;11.

[Google Scholar](#) [OpenURL Placeholder Text](#) Â

817Greulich Â S, Deluigi Â CC, Gloekler Â S, Wahl Â A, Z  rn Â C, Kramer Â U, et al. Â CMR imaging

predicts death and other adverse events in suspected cardiac sarcoidosis. *JACC Cardiovasc Imaging* Â 2013;6:501â511.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

818Nadel Â J, Lancefield Â T, Voskoboinik Â A, Taylor Â AJ. Late gadolinium enhancement identified with

cardiac magnetic resonance imaging in sarcoidosis patients is associated with long-term ventricular arrhythmia and sudden cardiac death. *Eur Heart J Cardiovasc Imaging* Â 2015;16:631â641.

[Google Scholar](#) [OpenURL Placeholder Text](#) Â

819Murtagh Â G, Laffin Â LJ, Beshai Â JF, Maffessanti Â F, Bonham Â CA, Patel Â AV, et al. Â Prognosis of

myocardial damage in sarcoidosis patients with preserved left ventricular ejection fraction: risk stratification using cardiovascular magnetic resonance. *Circ Cardiovasc Imaging* Â 2016;9:e003738.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

820Ekstr  m Â K, Lehtonen Â J, H  nninen Â H, Kandolin Â R, Kivist   Â S, Kupari Â M. Magnetic

resonance imaging as a predictor of survival free of life-threatening arrhythmias and transplantation in cardiac sarcoidosis. *J Am Heart Assoc* Â 2016;5:e003040.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

821Coleman Â GC, Shaw Â PW, Balfour Â PC, Gonzalez Â JA, Kramer Â CM, Patel Â AR, et al. Â Prognostic

value of myocardial scarring on CMR in patients with cardiac sarcoidosis. *JACC Cardiovasc Imaging* Â 2017;10:411â420.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

822Aizer Â A, Stern Â EH, Gomes Â JA, Teirstein Â AS, Eckart Â RE, Mehta Â D. Usefulness of programmed

ventricular stimulation in predicting future arrhythmic events in patients with cardiac sarcoidosis. *Am J Cardiol* Â 2005;96:276â282.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

823Mehta Â D, Mori Â N, Goldbarg Â SH, Lubitz Â S, Wisnivesky Â JP, Teirstein Â A. Primary prevention of

sudden cardiac death in silent cardiac sarcoidosis: role of programmed ventricular stimulation. *Circ Arrhythm Electrophysiol* Â 2011;4:43â48.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

824Okada Â DR, Smith Â J, Derakhshan Â A, Gowani Â Z, Zimmerman Â SL, Misra Â S, et al.

Â Electrophysiology study for risk stratification in patients with cardiac sarcoidosis and abnormal cardiac imaging. *Int J Cardiol Heart Vasc* Â 2019;23:100342.

[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â

825Zipse Â MM, Tzou Â WS, Schuller Â JL, Aleong Â RG, Varosy Â PD, Tompkins Â C, et al.

Â Electrophysiologic testing for diagnostic evaluation and risk stratification in patients with suspected cardiac sarcoidosis with preserved left and right ventricular systolic function. *J Cardiovasc Electrophysiol* Â 2019;30:1939â1948.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 826Blankstein Â R, Osborne Â M, Naya Â M, Waller Â A, Kim Â CK, Murthy Â VL, et al. Â Cardiac positron emission tomography enhances prognostic assessments of patients with suspected cardiac sarcoidosis. *J Am Coll Cardiol* Â 2014;63:329â336.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 827Crawford Â T, Mueller Â G, Sarsam Â S, Prasitdumrong Â H, Chaiyen Â N, Gu Â X, et al. Â Magnetic resonance imaging for identifying patients with cardiac sarcoidosis and preserved or mildly reduced left ventricular function at risk of ventricular arrhythmias. *Circ Arrhythm Electrophysiol* Â 2014;7:1109â1115.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 828Schuller Â JL, Zipse Â M, Crawford Â T, Bogun Â F, Beshai Â J, Patel Â AR, et al. Â Implantable cardioverter defibrillator therapy in patients with cardiac sarcoidosis. *J Cardiovasc Electrophysiol* Â 2012;23:925â929.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 829Betensky Â BP, Tschabrunn Â CM, Zado Â ES, Goldberg Â LR, Marchlinski Â FE, Garcia Â FC, et al. Â Long-term follow-up of patients with cardiac sarcoidosis and implantable cardioverter-defibrillators. *Heart Rhythm* Â 2012;9:884â891.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 830Kron Â J, Sauer Â W, Schuller Â J, Bogun Â F, Crawford Â T, Sarsam Â S, et al. Â Efficacy and safety of implantable cardiac defibrillators for treatment of ventricular arrhythmias in patients with cardiac sarcoidosis. *EP Europace* Â 2013;15:347â354.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 831Mohsen Â A, Jimenez Â A, Hood Â RE, Dickfeld Â T, Saliaris Â A, Shorofsky Â S, et al. Â Cardiac sarcoidosis: electrophysiological outcomes on long-term follow-up and the role of the implantable cardioverter-defibrillator. *J Cardiovasc Electrophysiol* Â 2014;25:171â176.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 832Azoulay Â L-D, Waintraub Â X, Haroche Â J, Amoura Â Z, Cohen Aubart Â F. Factors associated with implantable cardioverter defibrillators appropriate therapy in cardiac sarcoidosis: a meta-analysis: implantable cardioverter defibrillators in cardiac sarcoidosis. *Sarcoidosis Vasc Diffuse Lung Dis* Â 2020;37:17â23.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 833Smedema Â J-P, van Geuns Â R-J, Ector Â J, Heidbuchel Â H, Ainslie Â G, Crijns Â HJGM. Right ventricular involvement and the extent of left ventricular enhancement with magnetic resonance predict adverse outcome in pulmonary sarcoidosis. *ESC Heart Fail* Â 2018;5:157â171.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 834Velangi Â PS, Chen Â K-HA, Kazmirczak Â F, Okasha Â O, von Wald Â L, Roukoz Â H, et al. Â Right ventricular abnormalities on cardiovascular magnetic resonance imaging in patients with sarcoidosis. *JACC Cardiovasc Imaging* Â 2020;13:1395â1405.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 835Ise Â T, Hasegawa Â T, Morita Â Y, Yamada Â N, Funada Â A, Takahama Â H, et al. Â Extensive late gadolinium enhancement on cardiovascular magnetic resonance predicts adverse outcomes and lack of improvement in LV function after steroid therapy in cardiac sarcoidosis. *Heart* Â 2014;100:1165â1172.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 836Yodogawa Â K, Seino Â Y, Ohara Â T, Takayama Â H, Katoh Â T, Mizuno Â K. Effect of corticosteroid therapy on ventricular arrhythmias in patients with cardiac sarcoidosis: effect of corticosteroid therapy. *Ann Noninvasive Electrocardiol* Â 2011;16:140â147.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 837Naruse Â Y, Sekiguchi Â Y, Nogami Â A, Okada Â H, Yamauchi Â Y, Machino Â T, et al. Â Systematic treatment approach to ventricular tachycardia in cardiac sarcoidosis. *Circ Arrhythm Electrophysiol* Â 2014;7:407â413.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 838Segawa Â M, Fukuda Â K, Nakano Â M, Kondo Â M, Satake Â H, Hirano Â M, et al. Â Time course and factors correlating with ventricular tachyarrhythmias after introduction of steroid therapy in cardiac sarcoidosis. *Circ Arrhythm Electrophysiol* Â 2016;9:e003353.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 839Kumar Â S, Barbhaiya Â C, Nagashima Â K, Choi Â E-K, Epstein Â LM, John Â RM, et al. Â Ventricular tachycardia in cardiac sarcoidosis: characterization of ventricular substrate and outcomes of catheter ablation.

- Circ Arrhythm Electrophysiol 2015;8:872-879. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 840 Muser D, Santangeli P, Liang JJ, Castro SA, Magnani S, Hayashi T, et al. Characterization of the electroanatomic substrate in cardiac sarcoidosis. JACC Clin Electrophysiol 2018;4:291-303. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 841 Jelic D, Joel B, Good E, Morady F, Rosman H, Knight B, et al. Role of radiofrequency catheter ablation of ventricular tachycardia in cardiac sarcoidosis: report from a multicenter registry. Heart Rhythm 2009;6:189-195. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 842 Papageorgiou N, Providencia R, Bronis K, Decherer DG, Srinivasan N, Eckardt L, et al. Catheter ablation for ventricular tachycardia in patients with cardiac sarcoidosis: a systematic review. Europace 2018;20:682-691. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 843 Siontis KC, Santangeli P, Muser D, Marchlinski FE, Zeppenfeld K, Hoogendoorn JC, et al. Outcomes associated with catheter ablation of ventricular tachycardia in patients with cardiac sarcoidosis. JAMA Cardiol 2022;7:175. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 844 Hoogendoorn JC, Venlet J, Out YNJ, Man S, Kumar S, Sramko M, et al. The precordial R wave: a novel discriminator between cardiac sarcoidosis and arrhythmogenic right ventricular cardiomyopathy in patients presenting with ventricular tachycardia. Heart Rhythm 2021;18:1539-1547. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 845 Bern C. Chagas disease. N Engl J Med 2015;373:456-466. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 846 Bocchi EA, Bestetti RB, Scanavacca MI, Cunha Neto E, Issa VS. Chronic chagas heart disease management. J Am Coll Cardiol 2017;70:1510-1524. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 847 Nunes MCP, Dones W, Morillo CA, Encina JJ, Ribeiro AL. Chagas disease. J Am Coll Cardiol 2013;62:767-776. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 848 Nunes MCP, Beaton A, Acquatella H, Bern C, Bolger AF, Echeverr   LE, et al. Chagas cardiomyopathy: an update of current clinical knowledge and management: a scientific statement from the American heart association. Circulation 2018;138:e169-e209. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 849 Rassi A, Rassi A, Little WC, Xavier SS, Rassi SG, Rassi AG, et al. Development and validation of a risk score for predicting death in Chagas heart disease. N Engl J Med 2006;355:799-808. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 850 Senra T, Ianni BM, Costa ACP, Mady C, Martinelli-Filho M, Kalil-Filho R, et al. Long-term prognostic value of myocardial fibrosis in patients with chagas cardiomyopathy. J Am Coll Cardiol 2018;72:2577-2587. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 851 Rassi FM, Minohara L, Rassi A, Correia LCL, Marin-Neto JA, Rassi A, et al. Systematic review and meta-analysis of clinical outcome after implantable cardioverter-defibrillator therapy in patients with chagas heart disease. JACC Clin Electrophysiol 2019;5:1213-1223. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 852 Cardinalli-Neto A, Bestetti RB, Cordeiro JA, Rodrigues VC. Predictors of all-cause mortality for patients with chronic Chagas heart disease receiving implantable cardioverter defibrillator therapy. J Cardiovasc Electrophysiol 2007;18:1236-1240. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 853 Muratore CA, Batista Sa LA, Chiale PA, Eloy R, Tentori MC, Escudero J, et al. Implantable cardioverter defibrillators and chagas disease: results of the ICD registry Latin America. Europace 2008;11:164-168. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

854 Martinelli Â M, de Siqueira Â SF, Sternick Â EB, Rassi Â A, Costa Â R, Ramires Â JAF, et al. Â Long-term follow-up of implantable cardioverter-defibrillator for secondary prevention in chagas heart disease. *Am J Cardiol* Â 2012;110:1040â1045.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

855 Gali Â WL, Sarabanda Â AV, Baggio Â JM, Ferreira Â LG, Gomes Â GG, Marin-Neto Â JA, et al. Â Implantable cardioverter-defibrillators for treatment of sustained ventricular arrhythmias in patients with chagas heart disease: comparison with a control group treated with amiodarone alone. *Europace* Â 2014;16:674â680.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

856 Carmo Â AAL, de Sousa Â MR, Agudelo Â JF, Boersma Â E, Rocha Â MOC, Ribeiro Â ALP, et al. Â Implantable cardioverter-defibrillator in chagas heart disease: a systematic review and meta-analysis of observational studies. *Int J Cardiol* Â 2018;267:88â93.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

857 Stein Â C, Migliavaca Â CB, Colpani Â V, da Rosa Â PR, Sganzerla Â D, Giordani Â NE, et al. Â Amiodarone for arrhythmia in patients with chagas disease: a systematic review and individual patient data meta-analysis. *PLoS Negl Trop Dis* Â 2018;12:e0006742.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

858 Sosa Â E, Scanavacca Â M, D'Avila Â A, Piccioni Â J, Sanchez Â O, Velarde Â JL, et al. Â Endocardial and epicardial ablation guided by nonsurgical transthoracic epicardial mapping to treat recurrent ventricular tachycardia. *J Cardiovasc Electrophysiol* Â 1998;9:229â239.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

859 Soto-Becerra Â R, Bazan Â V, Bautista Â W, Malavassi Â F, Altamar Â J, Ramirez Â JD, et al. Â Ventricular tachycardia in the setting of chagasic cardiomyopathy: use of voltage mapping to characterize endoepicardial nonischemic scar distribution. *Circ Arrhythm Electrophysiol* Â 2017;10:e004950.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

860 Pisani Â CF, Romero Â J, Lara Â S, Hardy Â C, Chokr Â M, Sacilotto Â L, et al. Â Efficacy and safety of combined endocardial/epicardial catheter ablation for ventricular tachycardia in chagas disease: a randomized controlled study. *Heart Rhythm* Â 2020;17:1510â1518.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

861 Chizner Â MA, Pearle Â DL, deLeon Â AC. The natural history of aortic stenosis in adults. *Am Heart J* Â 1980;99:419â424.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

862 Delahaye Â JP, Gare Â JP, Viguier Â E, Delahaye Â F, De Gevigney Â G, Milon Â H. Natural history of severe mitral regurgitation. *Eur Heart J* Â 1991;12:5â9.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

863 Groves Â P. Valve disease: surgery of valve disease: late results and late complications. *Heart* Â 2001;86:715â721.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

864 Blackstone Â EH, Kirklin Â JW. Death and other time-related events after valve replacement. *Circulation* Â 1985;72:753â767.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

865 Urena Â M, Webb Â JG, Eltchaninoff Â H, Muñoz-García Â AJ, Bouleti Â C, Tamburino Â C, et al. Â Late cardiac death in patients undergoing transcatheter aortic valve replacement: incidence and predictors of advanced heart failure and sudden cardiac death. *J Am Coll Cardiol* Â 2015;65:437â448.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

866 Yang Â F, Shah Â B, Iwai Â S, Markowitz Â SM, Lerman Â BB, Stein Â KM. ICD implantation and arrhythmia-free survival in patients with depressed LV function following surgery for valvular heart disease. *Pacing Clin Electrophysiol* Â 2008;31:1419â1424.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

867 Valles Â AG, Khawaja Â FJ, Gersh Â BJ, Enriquez-Sarano Â M, Friedman Â PA, Park Â SJ, et al. Â Implantable cardioverter defibrillators in patients with valvular cardiomyopathy. *J Cardiovasc Electrophysiol* Â 2012;23:1326â1332.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

868Rodríguez-Mañero Á M, Barrio-López Á MT, Assi Á EA, Expósito-García Á V, Bertomeu-González Á V, Sánchez-Gómez Á JM, et al. Primary prevention of sudden death in patients with valvular cardiomyopathy. *Rev Esp Cardiol (Engl Ed)* 2016;69:272a-278.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

869Fischer-Rasokat Á U, Renker Á M, Liebetrau Á C, Weferling Á M, Rolf Á A, Hain Á A, et al. Long-term survival in patients with or without implantable cardioverter defibrillator after transcatheter aortic valve implantation. *J Clin Med* 2021;10:2929.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

870Nies Á RJ, Frerker Á C, Adam Á M, Kuhn Á E, Mauri Á V, Nettersheim Á FS, et al. Is there a benefit of ICD treatment in patients with persistent severely reduced systolic left ventricular function after TAVI? *Clin Res Cardiol* 2022;111:492a-501.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

871Eckart Á RE, Hruczkowski Á TW, Tedrow Á UB, Koplan Á BA, Epstein Á LM, Stevenson Á WG. Sustained ventricular tachycardia associated with corrective valve surgery. *Circulation* 2007;116:2005a-2011.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

872Liang Á JJ, Castro Á SA, Muser Á D, Briceno Á DF, Shirai Á Y, Enriquez Á A, et al. Electrophysiologic substrate, safety, procedural approaches, and outcomes of catheter ablation for ventricular tachycardia in patients after aortic valve replacement. *JACC Clin Electrophysiol* 2019;5:28a-38.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

873Delling Á FN, Vasan Á RS. Epidemiology and pathophysiology of mitral valve prolapse: new insights into disease progression, genetics, and molecular basis. *Circulation* 2014;129:2158a-2170.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

874Nalliah Á CJ, Mahajan Á R, Elliott Á AD, Haqqani Á H, Lau Á DH, Vohra Á JK, et al. Mitral valve prolapse and sudden cardiac death: a systematic review and meta-analysis. *Heart* 2019;105:144a-151.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

875Nishimura Á RA, McGoon Á MD, Shub Á C, Miller Á FA, Ilstrup Á DM, Tajik Á AJ. Echocardiographically documented mitral-valve prolapse. Long-term follow-up of 237 patients. *N Engl J Med* 1985;313:1305a-1309.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

876Perazzolo Marra Á M, Basso Á C, De Lazzari Á M, Rizzo Á S, Cipriani Á A, Giorgi Á B, et al. Morphofunctional abnormalities of mitral annulus and arrhythmic mitral valve prolapse. *Circ Cardiovasc Imaging* 2016;9:e005030.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

877Sriram Á CS, Syed Á FF, Ferguson Á ME, Johnson Á JN, Enriquez-Sarano Á M, Cetta Á F, et al. Malignant bileaflet mitral valve prolapse syndrome in patients with otherwise idiopathic out-of-hospital cardiac arrest. *J Am Coll Cardiol* 2013;62:222a-230.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

878Kitkungvan Á D, Nabi Á F, Kim Á RJ, Bonow Á RO, Khan Á MA, Xu Á J, et al. Myocardial fibrosis in patients with primary mitral regurgitation with and without prolapse. *J Am Coll Cardiol* 2018;72:823a-834.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

879Nordhues Á BD, Siontis Á KC, Scott Á CG, Nkomo Á VT, Ackerman Á MJ, Asirvatham Á SJ, et al. Bileaflet mitral valve prolapse and risk of ventricular dysrhythmias and death. *J Cardiovasc Electrophysiol* 2016;27:463a-468.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

880Essayagh Á B, Sabbag Á A, Antoine Á C, Benfari Á G, Yang Á L-T, Maalouf Á J, et al. Presentation and outcome of arrhythmic mitral valve prolapse. *J Am Coll Cardiol* 2020;76:637a-649.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

881Narasimhan Á C, Jazayeri Á MR, Sra Á J, Dhala Á A, Deshpande Á S, Biehl Á M, et al. Ventricular tachycardia in valvular heart disease: facilitation of sustained bundle-branch reentry by valve surgery. *Circulation* 1997;96:4307a-4313.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Á

- 882Moons P, Bovijn L, Budts W, Belmans A, Gewillig M. Temporal trends in survival to adulthood among patients born with congenital heart disease from 1970 to 1992 in Belgium. *Circulation* 2010;122:2264-2272.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 883Marelli AJ, Ionescu-Ittu R, Mackie AS, Guo L, Dendukuri N, Kaouache M. Lifetime prevalence of congenital heart disease in the general population from 2000 to 2010. *Circulation* 2014;130:749-756.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 884Koyak Z, Harris L, de Groot JR, Silversides CK, Oechslin EN, Bouma BJ, et al. Sudden cardiac death in adult congenital heart disease. *Circulation* 2012;126:1944-1954.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 885Gallego P, Gonzalez AE, Sanchez-Recalde A, Peinado R, Polo L, Gomez-Rubin C, et al. Incidence and predictors of sudden cardiac arrest in adults with congenital heart defects repaired before adult life. *Am J Cardiol* 2012;110:109-117.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 886Ghai A, Silversides C, Harris L, Webb GD, Siu SC, Therrien J. Left ventricular dysfunction is a risk factor for sudden cardiac death in adults late after repair of tetralogy of Fallot. *J Am Coll Cardiol* 2002;40:1675-1680.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 887Khairy P, Landzberg MJ, Gatzoulis MA, Lucron H, Lambert J, MarÅon F, et al. Value of programmed ventricular stimulation after tetralogy of fallot repair: a multicenter study. *Circulation* 2004;109:1994-2000.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 888Kapel GFL, Sacher F, Dekkers OM, Watanabe M, Blom NA, Thambo J-B, et al. Arrhythmogenic anatomical isthmuses identified by electroanatomical mapping are the substrate for ventricular tachycardia in repaired tetralogy of Fallot. *Eur Heart J* 2017;38:268-276.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#)
- 889Atallah J, Gonzalez Corcia MC, Walsh EP. Participating members of the pediatric and congenital electrophysiology society. Ventricular arrhythmia and life-threatening events in patients with repaired tetralogy of Fallot. *Am J Cardiol* 2020;132:126-132.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 890Kammeraad JAE, van Deurzen CHM, Sreeram N, Bink-Boelkens MTE, Ottenkamp J, Helbing WA, et al. Predictors of sudden cardiac death after Mustard or Senning repair for transposition of the great arteries. *J Am Coll Cardiol* 2004;44:1095-1102.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 891Schwerzmann M, Salehian O, Harris L, Siu SC, Williams WG, Webb GD, et al. Ventricular arrhythmias and sudden death in adults after a Mustard operation for transposition of the great arteries. *Eur Heart J* 2009;30:1873-1879.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 892Miyazaki A, Sakaguchi H, Ohuchi H, Matsuoka M, Komori A, Yamamoto T, et al. Efficacy of hemodynamic-based management of tachyarrhythmia after repair of tetralogy of Fallot. *Circ J* 2012;76:2855-2862.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 893Harrison DA, Harris L, Siu SC, MacLoughlin CJ, Connelly MS, Webb GD, et al. Sustained ventricular tachycardia in adult patients late after repair of tetralogy of Fallot. *J Am Coll Cardiol* 1997;30:1368-1373.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 894Sabate Rotes A, Connolly HM, Warnes CA, Ammass NM, Phillips SD, Dearani JA, et al. Ventricular arrhythmia risk stratification in patients with tetralogy of Fallot at the time of pulmonary valve replacement. *Circ Arrhythm Electrophysiol* 2015;8:110-116.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 895Khairy P, Harris L, Landzberg MJ, Fernandes SM, Barlow A, Mercier L-A, et al. Sudden death and defibrillators in transposition of the great arteries with intra-atrial baffles: a multicenter study. *Circ Arrhythm Electrophysiol* 2008;1:250-257.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 896Roca-Luque A I, Rivas G, Andara N, Dos Subir A L, Francisco Pascual J, Pérez-Rodon J, Pijuan Domenech A, et al. Intra-atrial re-entrant tachycardia in patients with congenital heart disease: factors associated with disease severity. *Europace* 2018;20:1343-1351.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 897Zeppenfeld K, Schalij MJ, Bartelings MM, Tedrow UB, Koplan BA, Soejima K, et al. Catheter ablation of ventricular tachycardia after repair of congenital heart disease: electroanatomic identification of the critical right ventricular isthmus. *Circulation* 2007;116:2241-2252.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 898Kriebel T, Saul JP, Schneider H, Sigler M, Paul T. Noncontact mapping and radiofrequency catheter ablation of fast and hemodynamically unstable ventricular tachycardia after surgical repair of tetralogy of Fallot. *J Am Coll Cardiol* 2007;50:2162-2168.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 899van Zyl M, Kapa S, Padmanabhan D, Chen FC, Mulpuru SK, Packer DL, et al. Mechanism and outcomes of catheter ablation for ventricular tachycardia in adults with repaired congenital heart disease. *Heart Rhythm* 2016;13:1449-1454.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 900Laredo M, Frank R, Waintraub X, Gandjbakhch E, Iserin L, Hascoët S, et al. Ten-year outcomes of monomorphic ventricular tachycardia catheter ablation in repaired tetralogy of Fallot. *Arch Cardiovasc Dis* 2017;110:292-302.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 901Kapel GFL, Reichlin T, Wijnmaalen AP, Piers SRD, Holman ER, Tedrow UB, et al. Re-entry using anatomically determined isthmuses: a curable ventricular tachycardia in repaired congenital heart disease. *Circ Arrhythm Electrophysiol* 2015;8:102-109.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 902Bokma JP, de Wilde KC, Vliegen HW, van Dijk AP, van Melle JP, Meijboom FJ, et al. Value of cardiovascular magnetic resonance imaging in noninvasive risk stratification in tetralogy of Fallot. *JAMA Cardiol* 2017;2:678-683.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 903Koyak Z, de Groot JR, Bouma BJ, Van Gelder IC, Budts W, Zwinderman AH, et al. Symptomatic but not asymptomatic non-sustained ventricular tachycardia is associated with appropriate implantable cardioverter therapy in tetralogy of Fallot. *Int J Cardiol* 2013;167:1532-1535.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 904Khairy P, Harris L, Landzberg MJ, Viswanathan S, Barlow A, Gatzoulis MA, et al. Implantable cardioverter-defibrillators in tetralogy of Fallot. *Circulation* 2008;117:363-370.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 905Koyak Z, de Groot JR, Van Gelder IC, Bouma BJ, van Dessel PFHM, Budts W, et al. Implantable cardioverter defibrillator therapy in adults with congenital heart disease: who is at risk of shocks? *Circ Arrhythm Electrophysiol* 2012;5:101-110.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 906Cochet H, Iriart X, Allain-Nicola A, Camaioni C, Sridi S, Nivet H, et al. Focal scar and diffuse myocardial fibrosis are independent imaging markers in repaired tetralogy of Fallot. *Eur Heart J Cardiovasc Imaging* 2019;20:990-1003.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 907Babu-Narayan SV, Kilner PJ, Li W, Moon JC, Goktekin O, Davlouros PA, et al. Ventricular fibrosis suggested by cardiovascular magnetic resonance in adults with repaired tetralogy of fallot and its relationship to adverse markers of clinical outcome. *Circulation* 2006;113:405-413.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 908Gatzoulis MA, Balaji S, Webber SA, Siu SC, Hokanson JS, Poile C, et al. Risk factors for arrhythmia and sudden cardiac death late after repair of tetralogy of Fallot: a multicentre study. *Lancet* 2000;356:975-981.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 909Bokma JP, Winter MM, Vehmeijer JT, Vliegen HW, van Dijk AP, van Melle JP, et al. QRS fragmentation is superior to QRS duration in predicting mortality in adults with tetralogy of Fallot. *Heart*

- Â 2017;103:666â671.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 910Egbe Â AC, Kothapalli Â S, Borlaug Â BA, Ammash Â NM, Najam Â M, Bajwa Â N, et al. Â Mechanism and risk factors for death in adults with tetralogy of Fallot. *Am J Cardiol* Â 2019;124:803â807.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 911Survivors of out-of-hospital cardiac arrest with apparently normal heart. Need for definition and standardized clinical evaluation. Consensus statement of the joint steering committees of the unexplained cardiac arrest registry of Europe and of the idiopathic ventricular fibrillation registry of the United States. *Circulation* Â 1997;95:265â272.
[Crossref](#) [Search ADS](#) [PubMed](#) Â
- 912Leenhardt Â A, Glaser Â E, Burguera Â M, NÃ¼rnberg Â M, Maison-Blanche Â P, Coumel Â P. Short-coupled variant of torsade de pointes. A new electrocardiographic entity in the spectrum of idiopathic ventricular tachyarrhythmias. *Circulation* Â 1994;89:206â215.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 913Eisenberg Â SJ, Scheinman Â MM, Dullet Â NK, Finkbeiner Â WE, Griffin Â JC, Eldar Â M, et al. Â Sudden cardiac death and polymorphous ventricular tachycardia in patients with normal QT intervals and normal systolic cardiac function. *Am J Cardiol* Â 1995;75:687â692.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 914Asatryan Â B, Schaller Â A, Seiler Â J, Servatius Â H, Noti Â F, Baldinger Â SH, et al. Â Usefulness of genetic testing in sudden cardiac arrest survivors with or without previous clinical evidence of heart disease. *Am J Cardiol* Â 2019;123:2031â2038.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 915Visser Â M, Dooijes Â D, van der Smagt Â JJ, van der Heijden Â JF, Doevendans Â PA, Loh Â P, et al. Â Next-generation sequencing of a large gene panel in patients initially diagnosed with idiopathic ventricular fibrillation. *Heart Rhythm* Â 2017;14:1035â1040.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 916Honarbakhsh Â S, Srinivasan Â N, Kirkby Â C, Firman Â E, Tobin Â L, Finlay Â M, et al. Â Medium-term outcomes of idiopathic ventricular fibrillation survivors and family screening: a multicentre experience. *Europace* Â 2017;19:1874â1880.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 917Meissner Â MD, Lehmann Â MH, Steinman Â RT, Mosteller Â RD, Akhtar Â M, Calkins Â H, et al. Â Ventricular fibrillation in patients without significant structural heart disease: a multicenter experience with implantable cardioverter-defibrillator therapy. *J Am Coll Cardiol* Â 1993;21:1406â1412.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 918Conte Â G, Caputo Â ML, Regoli Â F, Marcon Â S, Klersy Â C, Adjibodou Â B, et al. Â True idiopathic ventricular fibrillation in out-of-hospital cardiac arrest survivors in the Swiss Canton Ticino: prevalence, clinical features, and long-term follow-up. *Europace* Â 2017;19:259â266.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 919Stampe Â NK, Jespersen Â CB, Glinge Â C, Bundgaard Â H, Tfelt-Hansen Â J, Winkel Â BG. Clinical characteristics and risk factors of arrhythmia during follow-up of patients with idiopathic ventricular fibrillation. *J Cardiovasc Electrophysiol* Â 2020;31:2677â2686.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 920Conte Â G, Belhassen Â B, Lambiase Â P, Ciconte Â G, de Asmundis Â C, Arbelo Â E, et al. Â Out-of-hospital cardiac arrest due to idiopathic ventricular fibrillation in patients with normal electrocardiograms: results from a multicentre long-term registry. *Europace* Â 2019;21:1670â1677.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 921Blom Â LJ, Visser Â M, Christiaans Â I, Scholten Â MF, Bootsma Â M, van den Berg Â MP, et al. Â Incidence and predictors of implantable cardioverter-defibrillator therapy and its complications in idiopathic ventricular fibrillation patients. *Europace* Â 2019;21:1519â1526.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 922Malhi Â N, Cheung Â CC, Deif Â B, Roberts Â JD, Gula Â LJ, Green Â MS, et al. Â Challenge and impact of quinidine access in sudden death syndromes: a national experience. *JACC Clin Electrophysiol* Â 2019;5:376â382.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 923Belhassen Â B, Glick Â A, Viskin Â S. Excellent long-term reproducibility of the electrophysiologic efficacy of quinidine in patients with idiopathic ventricular fibrillation and Brugada syndrome. *Pacing Clin Electrophysiol* Â 2009;32:294â1301.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 924Belhassen Â B, Shapira Â I, Shoshani Â D, Paredes Â A, Miller Â H, Laniado Â S. Idiopathic ventricular fibrillation: inducibility and beneficial effects of class I antiarrhythmic agents. *Circulation* Â 1987;75:809â816.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 925Viskin Â S, Belhassen Â B. Idiopathic ventricular fibrillation. *Am Heart J* Â 1990;120:661â671.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 926Belhassen Â B, Viskin Â S, Fish Â R, Glick Â A, Setbon Â I, Eldar Â M. Effects of electrophysiologic-guided therapy with Class IA antiarrhythmic drugs on the long-term outcome of patients with idiopathic ventricular fibrillation with or without the Brugada syndrome. *J Cardiovasc Electrophysiol* Â 1999;10:1301â1312.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 927Sadek Â MM, Benhayon Â D, Sureddi Â R, Chik Â W, Santangeli Â P, Supple Â GE, et al. Â Idiopathic ventricular arrhythmias originating from the moderator band: electrocardiographic characteristics and treatment by catheter ablation. *Heart Rhythm* Â 2015;12:67â75.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 928Van Herendael Â H, Zado Â ES, Haqqani Â H, Tschabrunn Â CM, Callans Â DJ, Frankel Â DS, et al. Â Catheter ablation of ventricular fibrillation: importance of left ventricular outflow tract and papillary muscle triggers. *Heart Rhythm* Â 2014;11:566â573.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 929Santoro Â F, Di Biase Â L, Hranitzky Â P, Sanchez Â JE, Santangeli Â P, Perini Â AP, et al. Â Ventricular fibrillation triggered by PVCs from papillary muscles: clinical features and ablation. *J Cardiovasc Electrophysiol* Â 2014;25:1158â1164.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 930Nakamura Â T, Schaeffer Â B, Tanigawa Â S, Muthalaly Â RG, John Â RM, Michaud Â GF, et al. Â Catheter ablation of polymorphic ventricular tachycardia/fibrillation in patients with and without structural heart disease. *Heart Rhythm* Â 2019;16:1021â1027.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 931Moss Â AJ, Schwartz Â PJ, Crampton Â RS, Tzivoni Â D, Locati Â EH, MacCluer Â J, et al. Â The long QT syndrome. Prospective longitudinal study of 328 families. *Circulation* Â 1991;84:1136â1144.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 932Schwartz Â PJ, Ackerman Â MJ, Antzelevitch Â C, Bezzina Â CR, Borggreffe Â M, Cuneo Â BF, et al. Â Inherited cardiac arrhythmias. *Nat Rev Dis Primers* Â 2020;6:58.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 933Andersen Â ED, Krasilnikoff Â PA, Overvad Â H. Intermittent muscular weakness, extrasystoles, and multiple developmental anomalies. A new syndrome? Â *Acta Paediatr Scand* Â 1971;60:559â564.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 934Tawil Â R, Ptacek Â LJ, Pavlakis Â SG, DeVivo Â DC, Penn Â AS, Ozdemir Â C, et al. Â Andersenâ syndrome: potassium-sensitive periodic paralysis, ventricular ectopy, and dysmorphic features. *Ann Neurol* Â 1994;35:326â330.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 935Splawski Â I, Timothy Â KW, Decher Â N, Kumar Â P, Sachse Â FB, Beggs Â AH, et al. Â Severe arrhythmia disorder caused by cardiac L-type calcium channel mutations. *Proc Natl Acad Sci U S A* Â 2005;102:8089â8096; discussion 8086â8088.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 936Jervell Â A, Lange-Nielsen Â F. Congenital deaf-mutism, functional heart disease with prolongation of the Q-T interval and sudden death. *Am Heart J* Â 1957;54:59â68.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 937Schwartz Â PJ, Moss Â AJ, Vincent Â GM, Crampton Â RS. Diagnostic criteria for the long QT syndrome. An update. *Circulation* Â 1993;88:782â784.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 938Rautaharju A PM, Zhang A Z-M, Prineas A R, Heiss A G. Assessment of prolonged QT and JT intervals in ventricular conduction defects. *Am J Cardiol* A 2004;93:1017â1021.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 939Viskin A S, Postema A PG, Bhuiyan A ZA, Rosso A R, Kalman A JM, Vohra A JK, et al. A The response of the QT interval to the brief tachycardia provoked by standing: a bedside test for diagnosing long QT syndrome. *J Am Coll Cardiol* A 2010;55:1955â1961.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 940Mazzanti A A, Maragna A R, Vacanti A G, Monteforte A N, Bloise A R, Marino A M, et al. A Interplay between genetic substrate, QTc duration, and arrhythmia risk in patients with long QT syndrome. *J Am Coll Cardiol* A 2018;71:1663â1671.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 941Behr A ER, Roden A D. Drug-induced arrhythmia: pharmacogenomic prescribing? A *Eur Heart J* A 2013;34:89â95.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 942Weeke A PE, Kelleman A JS, Jespersen A CB, Theilade A J, Kanter A JK, Hansen A MS, et al. A Long-term proarrhythmic pharmacotherapy among patients with congenital long QT syndrome and risk of arrhythmia and mortality. *Eur Heart J* A 2019;40:3110â3117.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 943Schwartz A PJ, Priori A SG, Spazzolini A C, Moss A AJ, Vincent A GM, Napolitano A C, et al. A Genotype-phenotype correlation in the long-QT syndrome: gene-specific triggers for life-threatening arrhythmias. *Circulation* A 2001;103:89â95.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 944Chockalingam A P, Crotti A L, Girardengo A G, Johnson A JN, Harris A KM, van der Heijden A JF, et al. A Not all beta-blockers are equal in the management of long QT syndrome types 1 and 2: higher recurrence of events under metoprolol. *J Am Coll Cardiol* A 2012;60:2092â2099.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 945Ahn A J, Kim A HJ, Choi A J-I, Lee A KN, Shim A J, Ahn A HS, et al. A Effectiveness of beta-blockers depending on the genotype of congenital long-QT syndrome: a meta-analysis. *PLoS One* A 2017;12:e0185680.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 946Priori A SG, Napolitano A C, Schwartz A PJ, Grillo A M, Bloise A R, Ronchetti A E, et al. A Association of long QT syndrome loci and cardiac events among patients treated with beta-blockers. *JAMA* A 2004;292:1341â1344.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 947Mazzanti A A, Trancuccio A A, Kukavica A D, Pagan A E, Wang A M, Mohsin A M, et al. A Independent validation and clinical implications of the risk prediction model for long QT syndrome (1-2-3-LQTS-Risk). *Europace* A 2021;24:697â698.
[Google Scholar](#) [OpenURL Placeholder Text](#) A
- 948Mazzanti A A, Maragna A R, Faragli A A, Monteforte A N, Bloise A R, Memmi A M, et al. A Gene-specific therapy with mexiletine reduces arrhythmic events in patients with long QT syndrome type 3. *J Am Coll Cardiol* A 2016;67:1053â1058.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 949Ruan A Y, Liu A N, Bloise A R, Napolitano A C, Priori A SG. Gating properties of SCN5A mutations and the response to mexiletine in long-QT syndrome type 3 patients. *Circulation* A 2007;116:1137â1144.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 950Zhu A W, Mazzanti A A, Voelker A TL, Hou A P, Moreno A JD, Angsutararux A P, et al. A Predicting patient response to the antiarrhythmic mexiletine based on genetic variation. *Circ Res* A 2019;124:539â552.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 951Priori A SG, Napolitano A C, Schwartz A PJ, Bloise A R, Crotti A L, Ronchetti A E. The elusive link between LQT3 and Brugada syndrome: the role of flecainide challenge. *Circulation* A 2000;102:945â947.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 952Moss A AJ, Zareba A W, Hall A WJ, Schwartz A PJ, Crampton A RS, Benhorin A J, et al. A Effectiveness and limitations of beta-blocker therapy in congenital long-QT syndrome. *Circulation* A 2000;101:616â623.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A

- 953Jons A C, Moss AJ, Goldenberg I, Liu J, McNitt S, Zareba W, et al. Risk of fatal arrhythmic events in long QT syndrome patients after syncope. *J Am Coll Cardiol* 2010;55:783â788.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 954Liu JF, Jons A C, Moss AJ, McNitt S, Peterson DR, Qi M, et al. Risk factors for recurrent syncope and subsequent fatal or near-fatal events in children and adolescents with long QT syndrome. *J Am Coll Cardiol* 2011;57:941â950.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 955Seth R, Moss AJ, McNitt S, Zareba W, Andrews ML, Qi M, et al. Long QT syndrome and pregnancy. *J Am Coll Cardiol* 2007;49:1092â1098.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 956Goldenberg I, Horr S, Moss AJ, Lopes CM, Barsheshet A, McNitt S, et al. Risk for life-threatening cardiac events in patients with genotype-confirmed long-QT syndrome and normal-range corrected QT intervals. *J Am Coll Cardiol* 2011;57:51â59.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 957Jang SY, Cho Y, Kim NK, Kim C-Y, Sohn J, Roh J-H, et al. Video-assisted thoracoscopic left cardiac sympathetic denervation in patients with hereditary ventricular arrhythmias. *Pacing Clin Electrophysiol* 2017;40:232â241.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 958Waddell-Smith KE, Ertresvaag KN, Li J, Chaudhuri K, Crawford JR, Hamill JK, et al. Physical and psychological consequences of left cardiac sympathetic denervation in long-QT syndrome and catecholaminergic polymorphic ventricular tachycardia. *Circ Arrhythm Electrophysiol* 2015;8:1151â1158.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 959Anderson HN, Bos JM, Rohatgi RK, Ackerman MJ. The effect of left cardiac sympathetic denervation on exercise in patients with long QT syndrome. *JACC Clin Electrophysiol* 2019;5:1084â1090.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 960Bos JM, Bos KM, Johnson JN, Moir C, Ackerman MJ. Left cardiac sympathetic denervation in long QT syndrome: analysis of therapeutic nonresponders. *Circ Arrhythm Electrophysiol* 2013;6:705â711.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 961Bhandari AK, Shapiro WA, Morady F, Shen EN, Mason J, Scheinman MM. Electrophysiologic testing in patients with the long QT syndrome. *Circulation* 1985;71:63â71.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 962Zareba W, Moss AJ, Daubert JP, Hall WJ, Robinson JL, Andrews M. Implantable cardioverter defibrillator in high-risk long QT syndrome patients. *J Cardiovasc Electrophysiol* 2003;14:337â341.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 963Schwartz PJ, Spazzolini C, Priori SG, Crotti L, Vicentini A, Landolina M, et al. Who are the long-QT syndrome patients who receive an implantable cardioverter-defibrillator and what happens to them?: Data from the European long-QT syndrome implantable cardioverter-defibrillator (LQTS ICD) registry. *Circulation* 2010;122:1272â1282.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 964Delannoy E, Sacher F, Maury P, Mabo P, Mansourati J, Magnin I, et al. Cardiac characteristics and long-term outcome in Andersen-Tawil syndrome patients related to KCNJ2 mutation. *Europace* 2013;15:1805â1811.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 965Inoue YY, Aiba T, Kawata H, Sakaguchi T, Mitsuma W, Morita H, et al. Different responses to exercise between Andersen-Tawil syndrome and catecholaminergic polymorphic ventricular tachycardia. *Europace* 2018;20:1675â1682.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 966Krych M, Biernacka EK, Ponińska J, Kukla P, Filipecki A, Gajda R, et al. Andersen-Tawil syndrome: clinical presentation and predictors of symptomatic arrhythmiasâpossible role of polymorphisms K897T in KCNH2 and H558R in SCN5A gene. *J Cardiol* 2017;70:504â510.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 967Mazzanti A, Guz D, Trancuccio A, Pagan E, Kukavica D, Chargeishvili T, et al. Natural history and risk stratification in Andersen-Tawil syndrome type 1. *J Am Coll Cardiol* 2020;75:1772â1784. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 968Zhang L, Benson DW, Tristani-Firouzi M, Ptacek LJ, Tawil R, Schwartz PJ, et al. Electrocardiographic features in Andersen-Tawil syndrome patients with KCNJ2 mutations: characteristic T-U-wave patterns predict the KCNJ2 genotype. *Circulation* 2005;111:2720â2726. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 969Horigome H, Ishikawa Y, Kokubun N, Yoshinaga M, Sumitomo N, Lin L, et al. Multivariate analysis of TU wave complex on electrocardiogram in Andersen-Tawil syndrome with KCNJ2 mutations. *Ann Noninvasive Electrocardiol* 2020;25:e12721. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 970Miyamoto K, Aiba T, Kimura H, Hayashi H, Ohno S, Yasuoka C, et al. Efficacy and safety of flecainide for ventricular arrhythmias in patients with Andersen-Tawil syndrome with KCNJ2 mutations. *Heart Rhythm* 2015;12:596â603. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 971RadwaÅski PB, Greer-Short A, Poelzing S. Inhibition of Na⁺ channels ameliorates arrhythmias in a drug-induced model of Andersen-Tawil syndrome. *Heart Rhythm* 2013;10:255â263. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 972Tristani-Firouzi M, Jensen JL, Donaldson MR, Sansone V, Meola G, Hahn A, et al. Functional and clinical characterization of KCNJ2 mutations associated with LQT7 (Andersen syndrome). *J Clin Invest* 2002;110:381â388. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 973Antzelevitch C, Brugada P, Borggrefe M, Brugada J, Brugada R, Corrado D, et al. Brugada syndrome: report of the second consensus conference: endorsed by the Heart Rhythm Society and the European Heart Rhythm Association. *Circulation* 2005;111:659â670. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 974Savastano S, Rordorf R, Vicentini A, Petracci B, Taravelli E, Castelletti S, et al. A comprehensive electrocardiographic, molecular, and echocardiographic study of Brugada syndrome: validation of the 2013 diagnostic criteria. *Heart Rhythm* 2014;11:1176â1183. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 975Richter S, Sarkozy A, Paparella G, Henkens S, Boussy T, Chierchia G-B, et al. Number of electrocardiogram leads displaying the diagnostic coved-type pattern in Brugada syndrome: a diagnostic consensus criterion to be revised. *Eur Heart J* 2010;31:1357â1364. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 976Veltmann C, Papavassiliu T, Konrad T, Doesch C, Kuschyk J, Streitner F, et al. Insights into the location of type I ECG in patients with Brugada syndrome: correlation of ECG and cardiovascular magnetic resonance imaging. *Heart Rhythm* 2012;9:414â421. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 977Adler A, Rosso R, Chorin E, Havakuk O, Antzelevitch C, Viskin S. Risk stratification in Brugada syndrome: clinical characteristics, electrocardiographic parameters, and auxiliary testing. *Heart Rhythm* 2016;13:299â310. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 978Hasdemir C, Payzin S, Kocabas U, Sahin H, Yildirim N, Alp A, et al. High prevalence of concealed Brugada syndrome in patients with atrioventricular nodal reentrant tachycardia. *Heart Rhythm* 2015;12:1584â1594. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 979Behr ER, Ben-Haim Y, Ackerman MJ, Krahn AD, Wilde AAM. Brugada syndrome and reduced right ventricular outflow tract conduction reserve: a final common pathway? *Eur Heart J* 2021;42:1073â1081. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 980Probst V, Veltmann C, Eckardt L, Meregalli PG, Gaita F, Tan HL, et al. Long-term prognosis of patients diagnosed with Brugada syndrome: results from the FINGER Brugada syndrome registry. *Circulation* 2010;121:635â643. [Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)

- 981Amin Â AS, Meregalli Â PG, Bardai Â A, Wilde Â AAM, Tan Â HL. Fever increases the risk for cardiac arrest in the Brugada syndrome. *Ann Intern Med* Â 2008;149:216â218.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 982Adler Â A, Topaz Â G, Heller Â K, Zeltser Â D, Ohayon Â T, Rozovski Â U, et al. Â Fever-induced Brugada pattern: how common is it and what does it mean? Â *Heart Rhythm* Â 2013;10:1375â1382.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 983Priori Â SG, Napolitano Â C, Gasparini Â M, Pappone Â C, Della Bella Â P, Giordano Â U, et al. Â Natural history of Brugada syndrome: insights for risk stratification and management. *Circulation* Â 2002;105:1342â1347.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 984Rizzo Â A, Borio Â G, Sieira Â J, Van Dooren Â S, Overeinder Â I, Bala Â G, et al. Â Ajmaline testing and the Brugada syndrome. *Am J Cardiol* Â 2020;135:91â98.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 985Poli Â S, Toniolo Â M, Maiani Â M, Zanuttini Â D, Rebellato Â L, Vendramin Â I, et al. Â Management of untreatable ventricular arrhythmias during pharmacologic challenges with sodium channel blockers for suspected Brugada syndrome. *Europace* Â 2018;20:234â242.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 986Hasdemir Â C, Juang Â JJ-M, Kose Â S, Kocabas Â U, Orman Â MN, Payzin Â S, et al. Â Coexistence of atrioventricular accessory pathways and drug-induced type 1 Brugada pattern. *Pacing Clin Electrophysiol* Â 2018;41:1078â1092.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 987Probst Â V, Wilde Â AAM, Barc Â J, Sacher Â F, Babuty Â D, Mabo Â P, et al. Â SCN5A mutations and the role of genetic background in the pathophysiology of Brugada syndrome. *Circ Cardiovasc Genet* Â 2009;2:552â557.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 988Meregalli Â PG, Tan Â HL, Probst Â V, Koopmann Â TT, Tanck Â MW, Bhuiyan Â ZA, et al. Â Type of SCN5A mutation determines clinical severity and degree of conduction slowing in loss-of-function sodium channelopathies. *Heart Rhythm* Â 2009;6:341â348.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 989Ishikawa Â T, Kimoto Â H, Mishima Â H, Yamagata Â K, Ogata Â S, Aizawa Â Y, et al. Â Functionally validated SCN5A variants allow interpretation of pathogenicity and prediction of lethal events in Brugada syndrome. *Eur Heart J* Â 2021;42:2854â2863.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 990Gehi Â AK, Duong Â TD, Metz Â LD, Gomes Â JA, Mehta Â D. Risk stratification of individuals with the Brugada electrocardiogram: a meta-analysis. *J Cardiovasc Electrophysiol* Â 2006;17:577â583.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 991McNamara Â DA, Goldberger Â JJ, Berendsen Â MA, Huffman Â MD. Implantable defibrillators versus medical therapy for cardiac channelopathies. *Cochrane Database Syst Rev* Â 2015;2015:CD011168.
[Google Scholar](#) [PubMed](#) [OpenURL Placeholder Text](#) Â
- 992Priori Â SG, Gasparini Â M, Napolitano Â C, Della Bella Â P, Ottonelli Â AG, Sassone Â B, et al. Â Risk stratification in Brugada syndrome: results of the PRELUDE (PRogrammed ELectrical stimUlation preDICTive valuE) registry. *J Am Coll Cardiol* Â 2012;59:37â45.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 993Dereci Â A, Yap Â S-C, Schinkel Â AFL. Meta-analysis of clinical outcome after implantable cardioverter-defibrillator implantation in patients with Brugada syndrome. *JACC Clin Electrophysiol* Â 2019;5:141â148.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 994Conte Â G, Sieira Â J, Ciconte Â G, de Asmundis Â C, Chierchia Â G-B, Baltogiannis Â G, et al. Â Implantable cardioverter-defibrillator therapy in Brugada syndrome: a 20-year single-center experience. *J Am Coll Cardiol* Â 2015;65:879â888.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â
- 995Mascia Â G, Della Bona Â R, Ameri Â P, Canepa Â M, Porto Â I, Brignole Â M. Brugada syndrome and syncope: a systematic review. *J Cardiovasc Electrophysiol* Â 2020;31:333â338.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) Â

- 996Subramanian A M, Prabhu A MA, Harikrishnan A MS, Shekhar A SS, Pai A PG, Natarajan A K. The utility of exercise testing in risk stratification of asymptomatic patients with type 1 Brugada pattern. *J Cardiovasc Electrophysiol* A 2017;28:677a-683.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 997Kubala A M, Aïssou A L, Traullès A S, Gugenheim A A-L, Hermida A J-S. Use of implantable loop recorders in patients with Brugada syndrome and suspected risk of ventricular arrhythmia. *Europace* A 2012;14:898a-902.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 998Sakhi A R, Assaf A A, Theuns A DAMJ, Verhagen A JMA, Szili-Torok A T, Roos-Hesselink A JW, et al. A Outcome of insertable cardiac monitors in symptomatic patients with Brugada syndrome at low risk of sudden cardiac death. *Cardiology* A 2020;145:413a-420.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 999Scrocco A C, Ben-Haim A Y, Devine A B, Tome-Esteban A M, Papadakis A M, Sharma A S, et al. A Role of subcutaneous implantable loop recorder for the diagnosis of arrhythmias in Brugada syndrome: a United Kingdom single-center experience. *Heart Rhythm* A 2022;19:70a-78.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1000Sieira A J, Brugada A P. Brugada syndrome: defining the risk in asymptomatic patients. *Arrhythm Electrophysiol Rev* A 2016;5:164a-169.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1001Nishizaki A M, Sakurada A H, Yamawake A N, Ueda-Tatsumoto A A, Hiraoka A M. Low risk for arrhythmic events in asymptomatic patients with drug-induced type 1 ECG. Do patients with drug-induced Brugada type ECG have poor prognosis? (Con). *Circ J* A 2010;74:2464a-2473.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1002Conte A G, de Asmundis A C, Sieira A J, Ciconte A G, Di Giovanni A G, Chierchia A G-B, et al. A Prevalence and clinical impact of early repolarization pattern and QRS-fragmentation in high-risk patients with Brugada syndrome. *Circ J* A 2016;80:2109a-2116.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1003Kataoka A N, Mizumaki A K, Nakatani A Y, Sakamoto A T, Yamaguchi A Y, Tsujino A Y, et al. A Paced QRS fragmentation is associated with spontaneous ventricular fibrillation in patients with Brugada syndrome. *Heart Rhythm* A 2016;13:1497a-1503.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1004Probst A V, Goronflot A T, Anys A S, Tixier A R, Briand A J, Berthome A P, et al. A Robustness and relevance of predictive score in sudden cardiac death for patients with Brugada syndrome. *Eur Heart J* A 2021;42:1687a-1695.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1005Honarbakhsh A S, Providencia A R, Garcia-Hernandez A J, Martin A CA, Hunter A RJ, Lim A WY, et al. A A primary prevention clinical risk score model for patients with Brugada syndrome (BRUGADA-RISK). *JACC Clin Electrophysiol* A 2021;7:210a-222.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1006Andorin A A, Gourraud A J-B, Mansourati A J, Fouchard A S, le Marec A H, Maury A P, et al. A The QUIDAM study: hydroquinidine therapy for the management of Brugada syndrome patients at high arrhythmic risk. *Heart Rhythm* A 2017;14:1147a-1154.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1007Belhassen A B, Rahkovich A M, Michowitz A Y, Glick A A, Viskin A S. Management of Brugada syndrome: thirty-three-year experience using electrophysiologically guided therapy with class 1A antiarrhythmic drugs. *Circ Arrhythm Electrophysiol* A 2015;8:1393a-1402.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1008Ohgo A T, Okamura A H, Noda A T, Satomi A K, Suyama A K, Kurita A T, et al. A Acute and chronic management in patients with Brugada syndrome associated with electrical storm of ventricular fibrillation. *Heart Rhythm* A 2007;4:695a-700.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#) A
- 1009Nademanee A K, Raju A H, de Noronha A SV, Papadakis A M, Robinson A L, Rothery A S, et al. A Fibrosis, connexin-43, and conduction abnormalities in the Brugada syndrome. *J Am Coll Cardiol*

- 1009 Haïssaguerre M, Vassallo J, Le Breton A, et al. A 2015;66:1976â1986.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1010 Nademanee K, Haissaguerre M, Hocini M, Nogami A, Cheniti G, Duchateau J, et al. A Mapping and ablation of ventricular fibrillation associated with early repolarization syndrome. *Circulation* A 2019;140:1477â1490.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1011 Nademanee K, Veerakul G, Chandanamattha P, Chaothawee L, Ariyachaipanich A, Jirasirirojanakorn K, et al. A Prevention of ventricular fibrillation episodes in Brugada syndrome by catheter ablation over the anterior right ventricular outflow tract epicardium. *Circulation* A 2011;123:1270â1279.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1012 Zhang P, Tung R, Zhang Z, Sheng X, Liu Q, Jiang R, et al. A Characterization of the epicardial substrate for catheter ablation of Brugada syndrome. *Heart Rhythm* A 2016;13:2151â2158.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1013 Haïssaguerre M, Extramiana F, Hocini M, Cauchemez B, Jaïs P, Cabrera JA, et al. A Mapping and ablation of ventricular fibrillation associated with long-QT and Brugada syndromes. *Circulation* A 2003;108:925â928.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1014 Brugada J, Pappone C, Berruezo A, Vicedomini G, Manguso F, Ciconte G, et al. A Brugada syndrome phenotype elimination by epicardial substrate ablation. *Circ Arrhythm Electrophysiol* A 2015;8:1373â1381.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1015 Pappone C, Brugada J, Vicedomini G, Ciconte G, Manguso F, Saviano M, et al. A Electrical substrate elimination in 135 consecutive patients with Brugada syndrome. *Circ Arrhythm Electrophysiol* A 2017;10:e005053.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1016 Yamagata K, Horie M, Aiba T, Ogawa S, Aizawa Y, Ohe T, et al. A Genotype-phenotype correlation of SCN5A mutation for the clinical and electrocardiographic characteristics of probands with Brugada syndrome: a Japanese multicenter registry. *Circulation* A 2017;135:2255â2270.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1017 Haïssaguerre M, Derval N, Sacher F, Jesel L, Deisenhofer I, de Roy L, et al. A Sudden cardiac arrest associated with early repolarization. *N Engl J Med* A 2008;358:2016â2023.
[Google Scholar](#) [Crossref](#) [Search ADS](#) [PubMed](#)
- 1018 Rosso R, Kogan E, Belhassen B, Rozovski U, Scheinman MM, Zeltser D, et al. A J-point elevation in survivors of primary ventricular fibrillation and matched control subjects: incidence and clinical significance. *J Am Coll Cardiol* A 2008;52:1231â1238.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1019 Macfarlane PW, Antzelevitch C, Haissaguerre M, Huikuri HV, Potse M, Rosso R, et al. A The early repolarization pattern: a consensus paper. *J Am Coll Cardiol* A 2015;66:470â477.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1020 Tikkanen JT, Anttonen O, Junttila MJ, Aro AL, Kerola T, Rissanen HA, et al. A Long-term outcome associated with early repolarization on electrocardiography. *N Engl J Med* A 2009;361:2529â2537.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1021 Sinner MF, Reinhard W, MÃ¼ller M, Beckmann B-M, Martens E, Perz S, et al. A Association of early repolarization pattern on ECG with risk of cardiac and all-cause mortality: a population-based prospective cohort study (MONICA/KORA). *PLoS Med* A 2010;7:e1000314.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1022 Nunn LM, Bhar-Amato J, Lowe MD, Macfarlane PW, Rogers P, McKenna WJ, et al. A Prevalence of J-point elevation in sudden arrhythmic death syndrome families. *J Am Coll Cardiol* A 2011;58:286â290.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1023 Watanabe H, Nogami A, Ohkubo K, Kawata H, Hayashi Y, Ishikawa T, et al. A Electrocardiographic characteristics and SCN5A mutations in idiopathic ventricular fibrillation associated with early repolarization. *Circ Arrhythm Electrophysiol* A 2011;4:874â881.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1024Chauveau [↗](#) S, Janin [↗](#) A, Till [↗](#) M, Morel [↗](#) E, Chevalier [↗](#) P, Millat [↗](#) G. Early repolarization syndrome caused by de novo duplication of KCND3 detected by next-generation sequencing. *HeartRhythm Case Rep* [↗](#) 2017;3:574a-578.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1025Takayama [↗](#) K, Ohno [↗](#) S, Ding [↗](#) W-G, Ashihara [↗](#) T, Fukumoto [↗](#) D, Wada [↗](#) Y, et al. [↗](#) A de novo gain-of-function KCND3 mutation in early repolarization syndrome. *Heart Rhythm* [↗](#) 2019;16:1698a-1706.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1026Rosso [↗](#) R, Glikson [↗](#) E, Belhassen [↗](#) B, Katz [↗](#) A, Halkin [↗](#) A, Steinvil [↗](#) A, et al. [↗](#) Distinguishing a benign from a malignant early repolarization: the value of the ST-segment morphology. *Heart Rhythm* [↗](#) 2012;9:225a-229.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1027Tikkanen [↗](#) JT, Junttila [↗](#) MJ, Anttonen [↗](#) O, Aro [↗](#) AL, Luttinen [↗](#) S, Kerola [↗](#) T, et al. [↗](#) Early repolarization: electrocardiographic phenotypes associated with favorable long-term outcome. *Circulation* [↗](#) 2011;123:2666a-2673.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1028Nademanee [↗](#) K, Veerakul [↗](#) G, Mower [↗](#) M, Likittanasombat [↗](#) K, Krittayapong [↗](#) R, Bhuripanyo [↗](#) K, et al. [↗](#) Defibrillator versus beta-blockers for unexplained Death in Thailand (DEBUT): a randomized clinical trial. *Circulation* [↗](#) 2003;107:2221a-2226.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1029Brugada [↗](#) J, Brugada [↗](#) R, Brugada [↗](#) P. Pharmacological and device approach to therapy of inherited cardiac diseases associated with cardiac arrhythmias and sudden death. *J Electrocardiol* [↗](#) 2000;33:41a-47.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1030Haas [↗](#) M, Sacher [↗](#) F, Nogami [↗](#) A, Komiya [↗](#) N, Bernard [↗](#) A, Probst [↗](#) V, et al. [↗](#) Characteristics of recurrent ventricular fibrillation associated with inferolateral early repolarization role of drug therapy. *J Am Coll Cardiol* [↗](#) 2009;53:612a-619.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1031Aizawa [↗](#) Y, Chinushi [↗](#) M, Hasegawa [↗](#) K, Naiki [↗](#) N, Horie [↗](#) M, Kaneko [↗](#) Y, et al. [↗](#) Electrical storm in idiopathic ventricular fibrillation is associated with early repolarization. *J Am Coll Cardiol* [↗](#) 2013;62:1015a-1019.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1032Patocskaï [↗](#) B, Barajas-Martinez [↗](#) H, Hu [↗](#) D, Gurabi [↗](#) Z, Koncz [↗](#) I, Antzelevitch [↗](#) C. Cellular and ionic mechanisms underlying the effects of cilostazol, milrinone, and isoproterenol to suppress arrhythmogenesis in an experimental model of early repolarization syndrome. *Heart Rhythm* [↗](#) 2016;13:1326a-1334.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1033Nam [↗](#) G-B, Kim [↗](#) Y-H, Antzelevitch [↗](#) C. Augmentation of J waves and electrical storms in patients with early repolarization. *N Engl J Med* [↗](#) 2008;358:2078a-2079.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1034Rodríguez-Capitán [↗](#) J, Fernández-Meseguer [↗](#) A, García-Pinilla [↗](#) JM, Calvo-Bonacho [↗](#) E, Jiménez-Navarro [↗](#) M, García-Margallo [↗](#) T, et al. [↗](#) Frequency of different electrocardiographic abnormalities in a large cohort of Spanish workers. *Europace* [↗](#) 2017;19:1855a-1863.

[Google Scholar](#) [OpenURL Placeholder Text](#) [↗](#)

1035Sun [↗](#) G-Z, Ye [↗](#) N, Chen [↗](#) Y-T, Zhou [↗](#) Y, Li [↗](#) Z, Sun [↗](#) Y-X. Early repolarization pattern in the general population: prevalence and associated factors. *Int J Cardiol* [↗](#) 2017;230:614a-618.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1036Wu [↗](#) S-H, Lin [↗](#) X-X, Cheng [↗](#) Y-J, Qiang [↗](#) C-C, Zhang [↗](#) J. Early repolarization pattern and risk for arrhythmia death: a meta-analysis. *J Am Coll Cardiol* [↗](#) 2013;61:645a-650.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

1037Malhi [↗](#) N, So [↗](#) PP, Cheung [↗](#) CC, Laksman [↗](#) ZWM, Healey [↗](#) JS, Chauhan [↗](#) VS, et al. [↗](#) Early repolarization pattern inheritance in the cardiac arrest survivors with preserved ejection fraction registry (CASPER). *JACC Clin Electrophysiol* [↗](#) 2018;4:1473a-1479.

[Google Scholar](#) [Crossref](#) [Search ADS](#) [↗](#)

- 1038Sinner Å MF, Porthan Å K, Noseworthy Å PA, Havulinna Å AS, Tikkanen Å JT, MÅ¼ller-Nurasyid Å M, et al. Å A meta-analysis of genome-wide association studies of the electrocardiographic early repolarization pattern. *Heart Rhythm* Å 2012;9:1627Å1634.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1039Adhikarla Å C, Boga Å M, Wood Å AD, Froelicher Å VF. Natural history of the electrocardiographic pattern of early repolarization in ambulatory patients. *Am J Cardiol* Å 2011;108:1831Å1835.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1040Mahida Å S, Derval Å N, Sacher Å F, Leenhardt Å A, Deisenhofer Å I, Babuty Å D, et al. Å Role of electrophysiological studies in predicting risk of ventricular arrhythmia in early repolarization syndrome. *J Am Coll Cardiol* Å 2015;65:151Å159.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1041Aizawa Å Y, Sato Å A, Watanabe Å H, Chinushi Å M, Furushima Å H, Horie Å M, et al. Å Dynamicity of the J-wave in idiopathic ventricular fibrillation with a special reference to pause-dependent augmentation of the J-wave. *J Am Coll Cardiol* Å 2012;59:1948Å1953.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1042Priori Å SG, Mazzanti Å A, Santiago Å DJ, Kukavica Å D, Trancuccio Å A, Kovacic Å JC. Precision medicine in catecholaminergic polymorphic ventricular tachycardia: JACC focus seminar 5/5. *J Am Coll Cardiol* Å 2021;77:2592Å2612.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1043Bezzina Å CR, Lahrouchi Å N, Priori Å SG. Genetics of sudden cardiac death. *Circ Res* Å 2015;116:1919Å1936.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1044Kimura Å H, Zhou Å J, Kawamura Å M, Itoh Å H, Mizusawa Å Y, Ding Å W-G, et al. Å Phenotype variability in patients carrying KCNJ2 mutations. *Circ Cardiovasc Genet* Å 2012;5:344Å353.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1045Priori Å SG, Napolitano Å C, Memmi Å M, Colombi Å B, Drago Å F, Gasparini Å M, et al. Å Clinical and molecular characterization of patients with catecholaminergic polymorphic ventricular tachycardia. *Circulation* Å 2002;106:69Å74.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1046Krahn Å AD, Gollob Å M, Yee Å R, Gula Å LJ, Skanes Å AC, Walker Å BD, et al. Å Diagnosis of unexplained cardiac arrest: role of adrenaline and procainamide infusion. *Circulation* Å 2005;112:2228Å2234.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1047Hayashi Å M, Denjoy Å I, Extramiana Å F, Maltret Å A, Buisson Å NR, Lupoglazoff Å J-M, et al. Å Incidence and risk factors of arrhythmic events in catecholaminergic polymorphic ventricular tachycardia. *Circulation* Å 2009;119:2426Å2434.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1048Leren Å IS, Saberniak Å J, Majid Å E, Haland Å TF, Edvardsen Å T, Haugaa Å KH. Nadolol decreases the incidence and severity of ventricular arrhythmias during exercise stress testing compared with Î²1-selective Î²-blockers in patients with catecholaminergic polymorphic ventricular tachycardia. *Heart Rhythm* Å 2016;13:433Å440.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1049Peltenburg Å PJ, Kallas Å D, Bos Å JM, Lieve Å KVV, Franciosi Å S, Roston Å TM, et al. Å An international multicenter cohort study on Î²-blockers for the treatment of symptomatic children with catecholaminergic polymorphic ventricular tachycardia. *Circulation* Å 2022;145:333Å344.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1050van der Werf Å C, Nederend Å I, Hofman Å N, van Geloven Å N, Ebink Å C, Frohn-Mulder Å IME, et al. Å Familial evaluation in catecholaminergic polymorphic ventricular tachycardia: disease penetrance and expression in cardiac ryanodine receptor mutation-carrying relatives. *Circ Arrhythm Electrophysiol* Å 2012;5:748Å756.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Å
- 1051Watanabe Å H, Chopra Å N, Laver Å D, Hwang Å HS, Davies Å SS, Roach Å DE, et al. Å Flecainide prevents catecholaminergic polymorphic ventricular tachycardia in mice and humans. *Nat Med*

- 1052Wang G, Zhao N, Zhong S, Wang Y, Li J. Safety and efficacy of flecainide for patients with catecholaminergic polymorphic ventricular tachycardia: a systematic review and meta-analysis. *Medicine (Baltimore)* 2019;98:e16961.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1053van der Werf C, Kannankeril PJ, Sacher F, Krahn AD, Viskin S, Leenhardt A, et al. Flecainide therapy reduces exercise-induced ventricular arrhythmias in patients with catecholaminergic polymorphic ventricular tachycardia. *J Am Coll Cardiol* 2011;57:2244a-2254.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1054Padfield GJ, AlAhmari L, Lieve KVV, AlAhmari T, Roston TM, Wilde AA, et al. Flecainide monotherapy is an option for selected patients with catecholaminergic polymorphic ventricular tachycardia intolerant of β -blockade. *Heart Rhythm* 2016;13:609a-613.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1055Roses-Noguer F, Jarman JWE, Clague JR, Till J. Outcomes of defibrillator therapy in catecholaminergic polymorphic ventricular tachycardia. *Heart Rhythm* 2014;11:58a-66.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1056De Ferrari GM, Dusi V, Spazzolini C, Bos JM, Abrams DJ, Berul CI, et al. Clinical management of catecholaminergic polymorphic ventricular tachycardia: the role of left cardiac sympathetic denervation. *Circulation* 2015;131:2185a-2193.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1057Roston TM, Jones K, Hawkins NM, Bos JM, Schwartz PJ, Perry F, et al. Implantable cardioverter-defibrillator use in catecholaminergic polymorphic ventricular tachycardia: a systematic review. *Heart Rhythm* 2018;15:1791a-1799.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1058van der Werf C, Lieve KV, Bos JM, Lane CM, Denjoy I, Roses-Noguer F, et al. Implantable cardioverter-defibrillators in previously undiagnosed patients with catecholaminergic polymorphic ventricular tachycardia resuscitated from sudden cardiac arrest. *Eur Heart J* 2019;40:2953a-2961.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1059Leenhardt A, Lucet V, Denjoy I, Grau F, Ngoc DD, Coumel P. Catecholaminergic polymorphic ventricular tachycardia in children. A 7-year follow-up of 21 patients. *Circulation* 1995;91:1512a-1519.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1060Kannankeril PJ, Moore JP, Cerrone M, Priori SG, Kertesz NJ, Ro PS, et al. Efficacy of flecainide in the treatment of catecholaminergic polymorphic ventricular tachycardia: a randomized clinical trial. *JAMA Cardiol* 2017;2:759a-766.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1061Gollob MH, Redpath CJ, Roberts JD. The short QT syndrome: proposed diagnostic criteria. *J Am Coll Cardiol* 2011;57:802a-812.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1062Thorsen K, Dam VS, Kjaer-Sorensen K, Pedersen LN, Skeberdis VA, Jurevicius J, et al. Loss-of-activity-mutation in the cardiac chloride-bicarbonate exchanger AE3 causes short QT syndrome. *Nat Commun* 2017;8:1696.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1063Mazzanti A, Kanthan A, Monteforte N, Memmi M, Bloise R, Novelli V, et al. Novel insight into the natural history of short QT syndrome. *J Am Coll Cardiol* 2014;63:1300a-1308.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1064Dhuthia H, Malhotra A, Parpia S, Gabus V, Finocchiaro G, Mellor G, et al. The prevalence and significance of a short QT interval in 18 825 low-risk individuals including athletes. *Br J Sports Med* 2016;50:124a-129.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1065Gallagher MM, Magliano G, Yap YG, Padula M, Morgia V, Postorino C, et al. Distribution and prognostic significance of QT intervals in the lowest half centile in 12,012 apparently healthy persons. *Am J*

- Cardiol 2006;98:933-935.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1066Anttonen O, Junttila MJ, Rissanen H, Reunanen A, Viitasalo M, Huikuri HV. Prevalence and prognostic significance of short QT interval in a middle-aged Finnish population. *Circulation* 2007;116:714-720.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1067Kobza R, Roos M, Niggli B, Abächerli R, Lupi GA, Frey F, et al. Prevalence of long and short QT in a young population of 41,767 predominantly male Swiss conscripts. *Heart Rhythm* 2009;6:652-657.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1068Giustetto C, Schimpf R, Mazzanti A, Scrocco C, Maury P, Anttonen O, et al. Long-term follow-up of patients with short QT syndrome. *J Am Coll Cardiol* 2011;58:587-595.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1069El-Battrawy I, Besler J, Liebe V, Schimpf R, Tschömen E, Rudic B, et al. Long-term follow-up of patients with short QT syndrome: clinical profile and outcome. *J Am Heart Assoc* 2018;7:e010073.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1070Mazzanti A, Maragna R, Vacanti G, Kostopoulou A, Marino M, Monteforte N, et al. Hydroquinidine prevents life-threatening arrhythmic events in patients with short QT syndrome. *J Am Coll Cardiol* 2017;70:3010-3015.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1071El-Battrawy I, Besler J, Li X, Lan H, Zhao Z, Liebe V, et al. Impact of antiarrhythmic drugs on the outcome of short QT syndrome. *Front Pharmacol* 2019;10:771.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1072Malik M. Drug-induced QT/QTc interval shortening: lessons from drug-induced QT/QTc prolongation. *Drug Saf* 2016;39:647-659.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1073Giustetto C, Scrocco C, Schimpf R, Maury P, Mazzanti A, Levetto M, et al. Usefulness of exercise test in the diagnosis of short QT syndrome. *Europace* 2015;17:628-634.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1074Mason JW, Ramseth DJ, Chanter DO, Moon TE, Goodman DB, Mendzelevski B. Electrocardiographic reference ranges derived from 79,743 ambulatory subjects. *J Electrocardiol* 2007;40:228-234.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1075Bun S-S, Maury P, Giustetto C, Deharo J-C. Electrical storm in short-QT syndrome successfully treated with Isoproterenol. *J Cardiovasc Electrophysiol* 2012;23:1028-1030.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1076Siu SC, Sermer M, Colman JM, Alvarez AN, Mercier LA, Morton BC, et al. Prospective multicenter study of pregnancy outcomes in women with heart disease. *Circulation* 2001;104:515-521.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1077Drenthen W, Pieper PG, Roos-Hesselink JW, van Lottum WA, Voors AA, Mulder BJM, et al. Outcome of pregnancy in women with congenital heart disease: a literature review. *J Am Coll Cardiol* 2007;49:2303-2311.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1078Roos-Hesselink JW, Ruys TPE, Stein JI, Thilén U, Webb GD, Niwa K, et al. Outcome of pregnancy in patients with structural or ischaemic heart disease: results of a registry of the European Society of Cardiology. *Eur Heart J* 2013;34:657-665.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1079Linde C, Bongiorni MG, Birgersdotter-Green U, Curtis AB, Deisenhofer I, Furokawa T, et al. Sex differences in cardiac arrhythmia: a consensus document of the European Heart Rhythm Association, endorsed by the Heart Rhythm Society and Asia Pacific Heart Rhythm Society. *Europace* 2018;20:1565-1565ao.
[Google Scholar](#) [Crossref](#) [Search ADS](#)

1080Regitz-Zagrosek Å V, Roos-Hesselink Å JW, Bauersachs Å J, Blomström-Lundqvist Å C, CÅfkovÅ; Å R, De Bonis Å M, et al. Å 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy. Eur Heart J Å 2018;39:3165åPÅDSTS3241.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1081Silversides Å CK, Harris Å L, Haberer Å K, Sermer Å M, Colman Å JM, Siu Å SC. Recurrence rates of arrhythmias during pregnancy in women with previous tachyarrhythmia and impact on fetal and neonatal outcomes. Am J Cardiol Å 2006;97:1206åPÅDSTS1212.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1082Sliwa Å K, Mebazaa Å A, Hilfiker-Kleiner Å D, Petrie Å MC, Maggioni Å AP, Laroche Å C, et al. Å Clinical characteristics of patients from the worldwide registry on peripartum cardiomyopathy (PPCM): EURObservational Research Programme in conjunction with the Heart Failure Association of the European Society of Cardiology Study Group on PPCM. Eur J Heart Fail Å 2017;19:1131åPÅDSTS1141.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1083Ware Å JS, Li Å J, Mazaika Å E, Yasso Å CM, DeSouza Å T, Cappola Å TP, et al. Å Shared genetic predisposition in peripartum and dilated cardiomyopathies. N Engl J Med Å 2016;374:233åPÅDSTS241.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1084Wang Å Y-C, Chen Å C-H, Su Å H-Y, Yu Å M-H. The impact of maternal cardioversion on fetal haemodynamics. Eur J Obstet Gynecol Reprod Biol Å 2006;126:268åPÅDSTS269.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1085Barnes Å EJ, Eben Å F, Patterson Å D. Direct current cardioversion during pregnancy should be performed with facilities available for fetal monitoring and emergency caesarean section. BJOG Å 2002;109:1406åPÅDSTS1407.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1086Miyoshi Å T, Kamiya Å CA, Katsuragi Å S, Ueda Å H, Kobayashi Å Y, Horiuchi Å C, et al. Å Safety and efficacy of implantable cardioverter-defibrillator during pregnancy and after delivery. Circ J Å 2013;77:1166åPÅDSTS1170.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1087Abello Å M, Peinado Å R, Merino Å JL, Gnoatto Å M, Mateos Å M, Silvestre Å J, et al. Å Cardioverter defibrillator implantation in a pregnant woman guided with transesophageal echocardiography. Pacing Clin Electrophysiol Å 2003;26:1913åPÅDSTS1914.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1088Hartz Å J, Clark Å BC, Ito Å S, Sherwin Å ED, Berul Å CI. Transvenous nonfluoroscopic pacemaker implantation during pregnancy guided by 3-dimensional electroanatomic mapping. HeartRhythm Case Rep Å 2017;3:490åPÅDSTS492.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1089Bauersachs Å J, KÅnig Å T, van der Meer Å P, Petrie Å MC, Hilfiker-Kleiner Å D, Mbakwem Å A, et al. Å Pathophysiology, diagnosis and management of peripartum cardiomyopathy: a position statement from the Heart Failure Association of the European Society of Cardiology Study Group on peripartum cardiomyopathy. Eur J Heart Fail Å 2019;21:827åPÅDSTS843.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1090Duncker Å D, KÅnig Å T, Hohmann Å S, Bauersachs Å J, Veltmann Å C. Avoiding untimely implantable cardioverter/defibrillator implantation by intensified heart failure therapy optimization supported by the wearable cardioverter/defibrillator-the PROLONG study. J Am Heart Assoc Å 2017;6:e004512.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1091Duncker Å D, Westenfeld Å R, Konrad Å T, Pfeffer Å T, Correia de Freitas Å CA, Pfister Å R, et al. Å Risk for life-threatening arrhythmia in newly diagnosed peripartum cardiomyopathy with low ejection fraction: a German multi-centre analysis. Clin Res Cardiol Å 2017;106:582åPÅDSTS589.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1092Jeejeebhoy Å FM, Zelop Å CM, Lipman Å S, Carvalho Å B, Joglar Å J, Mhyre Å JM, et al. Å Cardiac arrest in pregnancy: a scientific statement from the American heart association. Circulation Å 2015;132:1747åPÅDSTS1773.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Å

1093European Society of Gynecology (ESG), Association for European Paediatric Cardiology (AEPC), German Society for Gender Medicine (DGesGM), Regitz-Zagrosek Å V, Blomstrom Lundqvist Å C, Borghi Å C, et al.

- 1093 ESC Guidelines on the management of cardiovascular diseases during pregnancy: the task force on the Management of Cardiovascular Diseases during Pregnancy of the European Society of Cardiology (ESC). *Eur Heart J* 2011;32:3147-3197.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1094 Rashba EJ, Zareba W, Moss AJ, Hall WJ, Robinson J, Locati EH, et al. Influence of pregnancy on the risk for cardiac events in patients with hereditary long QT syndrome. *LQTS investigators. Circulation* 1998;97:451-456.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1095 Ishibashi K, Aiba T, Kamiya C, Miyazaki A, Sakaguchi H, Wada M, et al. Arrhythmia risk and β -blocker therapy in pregnant women with long QT syndrome. *Heart* 2017;103:1374-1379.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1096 Cheung CC, Lieve KV, Roston TM, van der Ree MH, Deyell MW, Andrade JG, et al. Pregnancy in catecholaminergic polymorphic ventricular tachycardia. *JACC Clin Electrophysiol* 2019;5:387-394.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1097 Wu L, Liang E, Fan S, Zheng L, Hu F, Liu S, et al. Effect of pregnancy in arrhythmogenic right ventricular cardiomyopathy. *Am J Cardiol* 2020;125:613-617.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1098 Bauce B, Daliento L, Frigo G, Russo G, Nava A. Pregnancy in women with arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Eur J Obstet Gynecol Reprod Biol* 2006;127:186-189.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1099 Gandjbakhch E, Varlet E, Duthoit G, Fressart V, Charron P, Himbert C, et al. Pregnancy and newborn outcomes in arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Int J Cardiol* 2018;258:172-178.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1100 Rodríguez-Mañero M, Casado-Arroyo R, Sarkozy A, Leysen E, Sieira JA, Namdar M, et al. The clinical significance of pregnancy in Brugada syndrome. *Rev Esp Cardiol (Engl Ed)* 2014;67:176-180.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1101 Benito B, Berrueto A. Brugada syndrome and pregnancy: delving into the role of sex hormones in ion channelopathies. *Rev Esp Cardiol (Engl Ed)* 2014;67:165-167.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1102 Ostrowska B, Kommata B, Malmberg H, Lundqvist CB. Arrhythmias in pregnancy. *ESC CardioMed*. 3rd ed. Oxford University Press; 2022, p2868-2870.
- 1103 Sliwa K, Blauwet L, Tibazarwa K, Libhaber E, Smedema J-P, Becker A, et al. Evaluation of bromocriptine in the treatment of acute severe peripartum cardiomyopathy: a proof-of-concept pilot study. *Circulation* 2010;121:1465-1473.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1104 Hilfiger-Kleiner D, Haghikia A, Berliner D, Vogel-Claussen J, Schwab J, Franke A, et al. Bromocriptine for the treatment of peripartum cardiomyopathy: a multicentre randomized study. *Eur Heart J* 2017;38:2671-2679.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1105 Driver K, Chisholm CA, Darby AE, Malhotra R, Dimarco JP, Ferguson JD. Catheter ablation of arrhythmia during pregnancy. *J Cardiovasc Electrophysiol* 2015;26:698-702.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1106 Wang Z, Chen YW, Jiang YH, Sun LP, Chen XJ, Tao HL, et al. Electrophysiological characteristics and therapeutic efficacy of radiofrequency ablation of focal atrial tachycardia originating from the right atrial appendage during pregnancy. *Zhonghua Xin Xue Guan Bing Za Zhi* 2020;48:859-865.
[Google Scholar](#) [OpenURL Placeholder Text](#)
- 1107 Tuzcu V, Kilinc OU. Implantable cardioverter defibrillator implantation without using fluoroscopy in a pregnant patient. *Pacing Clin Electrophysiol* 2012;35:e265-266.
[Google Scholar](#) [OpenURL Placeholder Text](#)

- 1108Frühlich Â GM, Holzmeister Â J, Häbeler Â M, Häbeler Â S, Wolfrum Â M, Enseleit Â F, et al. Â Prophylactic implantable cardioverter defibrillator treatment in patients with end-stage heart failure awaiting heart transplantation. *Heart* Â 2013;99:1158â1165.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1109Sandner Â SE, Wieselthaler Â G, Zuckermann Â A, Taghavi Â S, Schmidinger Â H, Pacher Â R, et al. Â Survival benefit of the implantable cardioverter-defibrillator in patients on the waiting list for cardiac transplantation. *Circulation* Â 2001;104:1171â1176.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1110Algarrondo Â V, Perault Â R, Bories Â M-C, Narayanan Â K, Garcia Â R, Combes Â N, et al. Â Prophylactic implantable cardioverter defibrillators for primary prevention: From implantation to heart transplantation. *Arch Cardiovasc Dis* Â 2018;111:758â765.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1111Cantero-Pérez Â EM, Sobrino-Márquez Â JM, Grande-Trillo Â A, Lage-Gallardo Â E, Rangel-Sousa Â D, Esteve-Ruiz Â IM, et al. Â Implantable cardioverter defibrillator for primary prevention in patients with severe ventricular dysfunction awaiting heart transplantation. *Transplant Proc* Â 2013;45:3659â3661.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1112Pezawas Â T, Grimm Â M, Ristl Â R, Kivaranovic Â D, Moser Â FT, Laufer Â G, et al. Â Primary preventive cardioverter-defibrillator implantation (Pro-ICD) in patients awaiting heart transplantation. A prospective, randomized, controlled 12-year follow-up study. *Transpl Int* Â 2015;28:34â41.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1113Kao Â AC, Krause Â SW, Handa Â R, Karia Â D, Reyes Â G, Bianco Â NR, et al. Â Wearable defibrillator use in heart failure (WIF): results of a prospective registry. *BMC Cardiovasc Disord* Â 2012;12:123.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1114Alba Â AC, Foroutan Â F, Hing Â NKVNF, Fan Â C-PS, Manlhiot Â C, Ross Â HJ. Incidence and predictors of sudden cardiac death after heart transplantation: a systematic review and meta-analysis. *Clin Transplant* Â 2018;32:e13206.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1115Vakil Â K, Taimeh Â Z, Sharma Â A, Abidi Â KS, Colvin Â M, Luepker Â R, et al. Â Incidence, predictors, and temporal trends of sudden cardiac death after heart transplantation. *Heart Rhythm* Â 2014;11:1684â1690.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1116Alba Â AC, Fan Â C-PS, Manlhiot Â C, Dipchand Â AI, Stehlik Â J, Ross Â HJ. The evolving risk of sudden cardiac death after heart transplant. An analysis of the ISHLT thoracic transplant registry. *Clin Transplant* Â 2019;33:e13490.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1117Eskander Â MA, Adler Â E, Hoffmayer Â KS. Arrhythmias and sudden cardiac death in post-cardiac transplant patients. *Curr Opin Cardiol* Â 2020;35:308â311.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1118Tsai Â VW, Cooper Â J, Garan Â H, Natale Â A, Ptaszek Â LM, Ellinor Â PT, et al. Â The efficacy of implantable cardioverter-defibrillators in heart transplant recipients: results from a multicenter registry. *Circ Heart Fail* Â 2009;2:197â201.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1119Klein Â HU, Meltendorf Â U. Bridging a temporary high risk of sudden arrhythmic death. Experience with the wearable cardioverter defibrillator. *PACE* Â 2010;33:353â367.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1120Corrado Â D, Zorzi Â A. Sudden death in athletes. *Int J Cardiol* Â 2017;237:67â70.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1121Rajan Â D, Garcia Â R, Svane Â J, Tfelt-Hansen Â J. Risk of sports-related sudden cardiac death in women. *Eur Heart J* Â 2021;43:1198â1206.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1122Corrado Â D, Basso Â C, Schiavon Â M, Thiene Â G. Screening for hypertrophic cardiomyopathy in young athletes. *N Engl J Med* Â 1998;339:364â369.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

- 1123Malhotra A, Dhutia H, Finocchiaro G, Gati S, Beasley I, Clift P, et al. Outcomes of cardiac screening in adolescent soccer players. *N Engl J Med* 2018;379:524-534.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1124Corrado D, Pelliccia A, Björnstad HH, Vanhees L, Biffi A, Borjesson M, et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J* 2005;26:516-524.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1125Corrado D, Schmied C, Basso C, Borjesson M, Schiavon M, Pelliccia A, et al. Risk of sports: do we need a pre-participation screening for competitive and leisure athletes? *Eur Heart J* 2011;32:934-944.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1126Baggish AL, Hutter AM, Wang F, Yared K, Weiner RB, Kupperman E, et al. Cardiovascular screening in college athletes with and without electrocardiography: a cross-sectional study. *Ann Intern Med* 2010;152:269-275.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1127Corrado D, Basso C, Pavei A, Michieli P, Schiavon M, Thiene G. Trends in sudden cardiovascular death in young competitive athletes after implementation of a preparticipation screening program. *JAMA* 2006;296:1593-1601.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1128Sharma S, Estes NAM, Vetter VL, Corrado D. Clinical decisions. Cardiac screening before participation in sports. *N Engl J Med* 2013;369:2049-2053.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1129Sarto P, Zorzi A, Merlo L, Vessella T, Pegoraro C, Giorgiano F, et al. Serial versus single cardiovascular screening of adolescent athletes. *Circulation* 2021;143:1729-1731.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1130Steinvil A, Chundadze T, Zeltser D, Rogowski O, Halkin A, Galily Y, et al. Mandatory electrocardiographic screening of athletes to reduce their risk for sudden death proven fact or wishful thinking? *J Am Coll Cardiol* 2011;57:1291-1296.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1131Corrado D, Pelliccia A, Heidbuchel H, Sharma S, Link M, Basso C, et al. Recommendations for interpretation of 12-lead electrocardiogram in the athlete. *Eur Heart J* 2010;31:243-259.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1132Sharma S, Drezner JA, Baggish A, Papadakis M, Wilson MG, Prutkin JM, et al. International recommendations for electrocardiographic interpretation in athletes. *Eur Heart J* 2018;39:1466-1480.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1133Crescenzi C, Zorzi A, Vessella T, Martino A, Panattoni G, Cipriani A, et al. Predictors of left ventricular scar using cardiac magnetic resonance in athletes with apparently idiopathic ventricular arrhythmias. *J Am Heart Assoc* 2021;10:e018206.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1134Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint task force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J* 2016;37:2315-2381.
[Google Scholar](#) [Crossref](#) [Search ADS](#)
- 1135Rizzo M, Spataro A, Cecchetelli C, Quaranta F, Livrieri S, Sperandii F, et al. Structural cardiac disease diagnosed by echocardiography in asymptomatic young male soccer players: implications for pre-participation screening. *Br J Sports Med* 2012;46:371-373.
[Google Scholar](#) [Crossref](#) [Search ADS](#)

- 1136Heidbuchel Â H, Arbelo Â E, Dâ Ascenzi Â F, Borjesson Â M, Boveda Â S, Castelletti Â S, et al.
Â Recommendations for participation in leisure-time physical activity and competitive sports of patients with arrhythmias and potentially arrhythmogenic conditions. Part 2: ventricular arrhythmias, channelopathies, and implantable defibrillators. *Europace* Â 2021;23:147â148.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1137Drezner Â JA, Peterson Â DF, Siebert Â DM, Thomas Â LC, Lopez-Anderson Â M, Suchsland Â MZ, et al.
Â Survival after exercise-related sudden cardiac arrest in young athletes: can we do better? Â *Sports Health*
Â 2019;11:91â98.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1138Drezner Â JA, Rogers Â KJ, Horneff Â JG. Automated external defibrillator use at NCAA Division II and III universities. *Br J Sports Med* Â 2011;45:1174â1178.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1139Borjesson Â M, Serratos Â L, Carre Â F, Corrado Â D, Drezner Â J, Dugmore Â DL, et al. Â Consensus document regarding cardiovascular safety at sports arenas: position stand from the European Association of Cardiovascular Prevention and Rehabilitation (EACPR), section of Sports Cardiology. *Eur Heart J*
Â 2011;32:2119â2124.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1140Drezner Â JA, Owens Â DS, Prutkin Â JM, Salerno Â JC, Harmon Â KG, Prosise Â S, et al.
Â Electrocardiographic Screening in National Collegiate Athletic Association Athletes. *Am J Cardiol*
Â 2016;118:754â759.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1141Erbel Â R, Möhlenkamp Â S, Moebus Â S, Schmermund Â A, Lehmann Â N, Stang Â A, et al. Â Coronary risk stratification, discrimination, and reclassification improvement based on quantification of subclinical coronary atherosclerosis: the Heinz Nixdorf Recall study. *J Am Coll Cardiol* Â 2010;56:1397â1406.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1142Yeboah Â J, McClelland Â RL, Polonsky Â TS, Burke Â GL, Sibley Â CT, O'Leary Â D, et al.
Â Comparison of novel risk markers for improvement in cardiovascular risk assessment in intermediate-risk individuals. *JAMA* Â 2012;308:788â795.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1143Pappone Â C, Vicedomini Â G, Manguso Â F, Saviano Â M, Baldi Â M, Pappone Â A, et al. Â Wolff-Parkinson-White syndrome in the era of catheter ablation: insights from a registry study of 2169 patients. *Circulation* Â 2014;130:811â819.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1144Obeyesekere Â MN, Leong-Sit Â P, Massel Â D, Manlucu Â J, Modi Â S, Krahn Â AD, et al. Â Risk of arrhythmia and sudden death in patients with asymptomatic preexcitation: a meta-analysis. *Circulation*
Â 2012;125:2308â2315.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1145Heidbuchel Â H, Adami Â PE, Antz Â M, Braunschweig Â F, Delise Â P, Scherr Â D, et al.
Â Recommendations for participation in leisure-time physical activity and competitive sports in patients with arrhythmias and potentially arrhythmogenic conditions: part 1: supraventricular arrhythmias. A position statement of the Section of Sports Cardiology and Exercise from the European Association of Preventive Cardiology (EAPC) and the European Heart Rhythm Association (EHRA), both associations of the European Society of Cardiology. *Eur J Prev Cardiol* Â 2020;28:1539â1551.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1146Pappone Â C, Santinelli Â V, Manguso Â F, Augello Â G, Santinelli Â O, Vicedomini Â G, et al. Â A randomized study of prophylactic catheter ablation in asymptomatic patients with the Wolff-Parkinson-White syndrome. *N Engl J Med* Â 2003;349:1803â1811.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1147Perry Â JC, Garson Â A. Supraventricular tachycardia due to Wolff-Parkinson-White syndrome in children: early disappearance and late recurrence. *J Am Coll Cardiol* Â 1990;16:1215â1220.
[Google Scholar](#) [Crossref](#) [Search ADS](#) Â
- 1148Pediatric and Congenital Electrophysiology Society (PACES), Heart Rhythm Society (HRS), American College of Cardiology Foundation (ACCF), American Heart Association (AHA), American Academy of Pediatrics (AAP), Canadian Heart Rhythm Society (CHRS), et al. Â PACES/HRS expert consensus statement on the

management of the asymptomatic young patient with a Wolff-Parkinson-White (WPW, ventricular preexcitation) electrocardiographic pattern: developed in partnership between the Pediatric and Congenital Electrophysiology Society (PACES) and the Heart Rhythm Society (HRS). Endorsed by the governing bodies of PACES, HRS, the American College of Cardiology Foundation (ACCF), the American Heart Association (AHA), the American Academy of Pediatrics (AAP), and the Canadian Heart Rhythm Society (CHRS). Heart Rhythm 2012;9:1006â1024.

[Crossref](#) [Search ADS](#) Â

1149Barsheshet Â A, Moss Â AJ, Huang Â DT, McNitt Â S, Zareba Â W, Goldenberg Â I. Applicability of a risk score for prediction of the long-term (8-year) benefit of the implantable cardioverter-defibrillator. J Am Coll Cardiol 2012;59:2075â2079.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

1150ExpÃ³sito Â V, RodrÃ­guez-MaÃ±ero Â M, GonzÃ¡lez-EnrÃ­quez Â S, Arias Â MA, SÃ¡nchez-GÃ³mez Â JM, AndrÃ©s La Huerta Â A, et al. Â Primary prevention implantable cardioverter-defibrillator and cardiac resynchronization therapy-defibrillator in elderly patients: results of a Spanish multicentre study. Europace 2016;18:1203â1210.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

1151Alhakak Â A, SÃ¸stergaard Â L, Butt Â JH, Vinther Â M, Philbert Â BT, Jacobsen Â PK, et al. Â Cause-specific death and risk factors of one-year mortality after implantable cardioverter-defibrillator implantation: a nationwide study. Eur Heart J Qual Care Clin Outcomes 2022;8:39â49.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

1152Poupin Â P, Bouleti Â C, Degand Â B, Paccalin Â M, Le Gal Â F, Bureau Â M-L, et al. Â Prognostic value of Charlson Comorbidity Index in the elderly with a cardioverter defibrillator implantation. Int J Cardiol 2020;314:64â69.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

1153Aktaâ S, Batra Â G, Wallentin Â L, Baigent Â C, Erlinge Â D, James Â S, et al. Â European society of cardiology methodology for the development of quality indicators for the quantification of cardiovascular care and outcomes. Eur Heart J Qual Care Clin Outcomes 2022;8:4â13.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

1154Minchin Â M, Roland Â M, Richardson Â J, Rowark Â S, Guthrie Â B. Quality of care in the United Kingdom after removal of financial incentives. N Engl J Med 2018;379:948â957.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

1155Song Â Z, Ji Â Y, Safran Â DG, Chernew Â ME. Health care spending, utilization, and quality 8 years into global payment. N Engl J Med 2019;381:252â263.

[Google Scholar](#) [Crossref](#) [Search ADS](#) Â

Author notes

Katja Zeppenfeld and Jacob Tfelt-Hansen chairpersons contributed equally to the document and are joint corresponding authors.

Marta de Riva and Bo Gregers Winkel task force Coordinators contributed equally to the document.

Author/task force Member affiliations are listed in Author information.

Representing the Association for European Paediatric and Congenital Cardiology (AEPC).

ESC Clinical Practice Guidelines (CPG) Committee: listed in the Appendix.

ESC subspecialty communities having participated in the development of this document:

Associations: Association for Acute CardioVascular Care (ACVC), Association of Cardiovascular Nursing & Allied Professions (ACNAP), European Association of Cardiovascular Imaging (EACVI), European Association of Percutaneous Cardiovascular Interventions (EAPCI), European Heart Rhythm Association (EHRA), Heart Failure Association (HFA).

Working Groups: Cardiac Cellular Electrophysiology, Myocardial and Pericardial Diseases.

Patient Forum

The content of these European Society of Cardiology (ESC) Guidelines has been published for personal and educational use only. No commercial use is authorized. No part of the ESC Guidelines may be translated or reproduced in any form without written permission from the ESC. Permission can be obtained upon submission of a written request to Oxford University Press, the publisher of the *European Heart Journal* and the party authorized to handle such permissions on behalf of the ESC (Journals.Permissions@oup.com).

Disclaimer: The ESC Guidelines represent the views of the ESC and were produced after careful consideration of the scientific and medical knowledge and the evidence available at the time of their publication. The ESC is not responsible in the event of any contradiction, discrepancy, and/or ambiguity between the ESC Guidelines and any other official recommendations or guidelines issued by the relevant public health authorities, in particular in relation to good use of healthcare or therapeutic strategies. Health professionals are encouraged to take the ESC Guidelines fully into account when exercising their clinical judgment, as well as in the determination and the implementation of preventive, diagnostic, or therapeutic medical strategies; however, the ESC Guidelines do not override, in any way whatsoever, the individual responsibility of health professionals to make appropriate and accurate decisions in consideration of each patient's health condition and in consultation with that patient and, where appropriate and/or necessary, the patient's caregiver. Nor do the ESC Guidelines exempt health professionals from taking into full and careful consideration the relevant official updated recommendations or guidelines issued by the competent public health authorities, in order to manage each patient's case in light of the scientifically accepted data pursuant to their respective ethical and professional obligations. It is also the health professional's responsibility to verify the applicable rules and regulations relating to drugs and medical devices at the time of prescription.

© The European Society of Cardiology 2022. All rights reserved. For permissions please e-mail: Journals.Permissions@oup.com

All experts involved in the development of these guidelines have submitted declarations of interest. These have been compiled in a report and published in a supplementary document simultaneously to the guidelines. The report is also available on the ESC website www.escardio.org/Guidelines

Published by Oxford University Press on behalf of European Society of Cardiology 2022.

This article is published and distributed under the terms of the Oxford University Press, Standard Journals Publication Model (https://academic.oup.com/journals/pages/open_access/funder_policies/chorus/standard_publication_model)