



Oregon State
University

Design Under Uncertainty

ME 615 Spring 2020


Dr. Chris Hoyle

MIME

Different Aspects of Design under Uncertainty



Oregon State University
College of Engineering




Challenges for Design under Uncertainty



Oregon State University
College of Engineering

- ❑ *Uncertainty Quantification*
- ❑ *Efficient Uncertainty Propagation (Utility & Reliability Assessments)*
- ❑ *Framework of Decision Making with Multiple Quality Attributes*



For 95% confidence, 10% error bound, 3.8×10^4 Monte Carlo samples are needed to catch 0.01 failure rate.

What is uncertainty?

- “There are known knowns. These are things we know that we know. There are known unknowns. That is to say, there are things that we know we don’t know. But there are also unknown unknowns. There are things we don’t know we don’t know.”

– **Donald Rumsfeld 2002**



Ways to Classify Uncertainties:



Oregon State University
College of Engineering

- **Classification 1**
 - **Model** (lack of knowledge)
 - **Parametric** (lack of knowledge, variability): variability in load, material property, dimensions
 - **Numerical**
 - **Testing data** (measurement)
- **Classification 2**
 - **Aleatory** (random, objective) uncertainty: inherent, irreducible uncertainty. Can also be classified as **risk**
 - **Epistemic** (subjective) uncertainty: reducible with better knowledge
 - **Ontological** uncertainty: the possibility of events occurring that we have no knowledge of
- **Classification 3**
 - **Noise factors**: Come from the environment, processes out of our control (i.e. parameters)
 - **Control factors**: Come from the factors we control (i.e. design variables)

Classifying Uncertainties



Oregon State University
College of Engineering

- “There are **known knowns**. These are things we know that we know. There are **known unknowns**. That is to say, there are things that we know we don’t know. But there are also **unknown unknowns**. There are things we don’t know we don’t know.”

– **Donald Rumsfeld 2002**

- **Aleatory** uncertainty: *known knowns*
- **Epistemic** uncertainty: *known unknowns*
- **Ontological** uncertainty: *unknown unknowns*



Our classification



Oregon State University
College of Engineering


- **Aleatory** (random, objective) uncertainty: inherent, irreducible uncertainty. Can also be classified as **risk**
 - Manufacturing tolerances
 - Material property variations
 - Assembly time variations
- **Epistemic** (subjective) uncertainty: reducible with better knowledge
 - Budget
 - Technology
 - Mission
- **Ontological** uncertainty: the possibility of events occurring that we have no knowledge of
 - Explosions
 - Fires

Aleatory Uncertainty




Oregon State University
College of Engineering


- Aleatory uncertainties
 - Probability Distribution Functions




Normal (Gaussian)



Log Normal



Beta



Uniform


Distributions




Oregon State University
College of Engineering

Imagine we created a histogram of lengths of fish caught in a river (in mm) over 1 month

We can turn that into a probability distribution function (PDF)



Frequency of fish size caught



Frequency converted to a probability density


The y axis scale is set to ensure the "area under the curve" is 1

Distributions




Oregon State University
College of Engineering

We can now just use integration compute the probability of different events:




What is the probability of catching a fish less than 185 mm?

- It is 0.16



What is the probability of catching a fish greater than 220 mm?

- It is 0.095



What is the probability of catching a fish greater than 205 and less than 215?

- It is 0.075

Epistemic Uncertainty



Oregon State University
College of Engineering


- Epistemic Uncertainties
 - Distributions with ranges of Parameters, i.e. a range of mean and variances
 - Ranges only, no distributional assumption
 - Subjective prior distribution updated as information becomes available (Bayesian approach)

Epistemic Uncertainty



Oregon State University
College of Engineering

- Subjective prior distribution updated as information becomes available (Bayesian approach)
- **We can use probability theory**



Ontological Uncertainty





- “Unknown Unknowns”
 - Not much we can do with these since they are unknown.

Representation of Uncertainty



Oregon State University
College of Engineering

- Use of Probability Theory
 - Use of Probability Density Function (**pdf**) for continuous random variables
 - Use of Probability Mass Function (**pmf**) for discrete random variables
- We will utilize **pdfs** for the course since it will enable use of simplifying mathematics.
- Examples of common distributions
 - Normal distribution
 - Lognormal distribution
 - Beta distribution
 - Uniform distribution



Representation of Uncertainty



Oregon State University
College of Engineering

- **Normal distribution**
 - Material property (Young's modulus,...), tolerance, ...
 - *Sum* of random effects, measurement errors
 - Infinite interval $[-\infty, \infty]$
- **Lognormal distribution**
 - Loading, failure by fatigue...
 - *Product* of random effects, uncertainty expressed as a factor n
 - Example: $P\left(\frac{y_0}{n} < y < y_0 n\right) = 90\%$
 - Semi-Infinite interval $[0, \infty]$
- **Beta distribution**
 - Useful for subjective estimates of uncertainty such as costs, schedules.
 - Used to fit a parametric distribution to collected data
 - Bounded Interval $[a, b]$
- **Uniform distribution**
 - Useful for subjective estimates of uncertainty such as costs, schedules.
 - Bounded Interval $[a, b]$

Representation of Uncertainty



Oregon State University
College of Engineering

- Extreme value distributions
 - Represent **maximum** or **minimum** of a number of samples of various distributions
 - Depends on the tail shape of original distribution
 - Gumbel distribution (Type I extreme value distribution for largest value)
 - From distributions with exponential tail (e.g. normal)
 - Wind load, flood levels, ...
 - Infinite interval $[-\infty, \infty]$
 - Frechet distribution (Type II extreme value distribution for largest value)
 - From distributions with polynomial tail (e.g. lognormal)
 - Earthquake load, ...
 - Semi-Infinite interval $[0, \infty]$
 - Weibull distribution (Type III extreme value distribution for smallest value)
 - From distributions with bounded ends (e.g. gamma)
 - Breaking strength of material, failure of series of identical components,...
 - Semi-Infinite interval $[0, \infty]$

Probability Space



Oregon State University
College of Engineering

- What is a probability space? It is a triple (Ω, \mathcal{F}, P) , with the components defined as:
 - A sample space, Ω , which is the set of all possible outcomes.
 - A set of events, \mathcal{F} where each event is a set containing zero or more outcomes.
 - The assignment of probabilities to the events; that is, a function P from events to probabilities.
- Let's look at an example

A coin flip



- We are tossing a “fair coin”, the sample space is:
 - $\Omega = \{H, T\}$
- The set of events is $\mathcal{F} = 2^\Omega$


Or $\mathcal{F} = \{\{\}, \{H\}, \{T\}, \{H, T\}\}$

 - $\{H\}$ (heads)
 - $\{T\}$ (tails)
 - $\{\}$ (neither heads nor tails)
 - $\{H, T\}$ (either heads or tails)
- The probability measure is using the probability function P :
 - $P(\{\}) = 0$
 - $P(\{T\}) = 0.5$
 - $P(\{H\}) = 0.5$
 - $P(\{H, T\}) = 1$

Basics of Probability



Oregon State University
College of Engineering



- Notation:
 - **U Union:** indicates the probability of either event A or event B occurring (OR)
 - **\cap Intersection:** indicates the probability that events A and B both occur (AND)
- With this notation we can define some axioms...

Axioms of Probability Theory



Oregon State University
College of Engineering

- $P(A) \geq 0$ for all events A
- $P(\Omega) = 1$
- $P(A \cap B) = P(A) P(B)$ if A and B are independent
- $P(A \cap B) = P(A/B) P(B)$ if A and B are dependent
- $P(A \cup B) = P(A) + P(B)$ for disjoint events A and B
- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ (not disjoint)
- $P(A \cup B) \leq P(A) + P(B)$

Basics of Probability



Oregon State University
College of Engineering

- Random variable

- A variable whose value is subject to variations due to chance, defined on a sample space Ω (continuous, discrete)
- A random variable $X: \Omega \rightarrow E$ is a measurable function from a set of possible outcomes Ω to a measurable space E , typically \mathbb{R}

$$X: \Omega \rightarrow \mathbb{R}$$

- A random variable does *not* return a probability
- Why is it a Function?
 - ω is the random event, e.g. landing on a head or a tail.
 - $X(\omega)$ is the function that defines a numerical quantity to the random event, e.g., the number of heads in a series of coin flips.
- Types of Random Variables:
 - Discrete Values: takes on value from countable set.
 - Continuous Values: infinite number of possible values.

Basics of Probability



Oregon State University
College of Engineering

- **Realization of X.**
 - The value $x = X(\omega)$ of a random variable X for outcome $\omega \in \Omega$ is called a realization of X .

- **Cumulative distribution function (CDF)**



$$F_X(x) = \Pr[X \leq x]$$

$$\Pr\{\omega \in \Omega : X(\omega) \leq x\}$$

- **Probability density function (PDF)**

$$\Pr[x \leq X \leq x + dx] = f_X(x)dx$$

$$f_X(x) = dF_X(x)/dx, \quad \int_{-\infty}^{\infty} f_X(x)dx = 1$$



Basics of Probability



Oregon State University
College of Engineering

- Descriptive measures of distribution
 - Expected value

$$E(x) = \int_{-\infty}^{\infty} x f_X(x) dx$$

Expectation operator is a linear operator!

- Statistical moments

Raw moments: $\mu'_k = E(x^k) = \int_{-\infty}^{\infty} x^k f_X(x) dx$

Central moments: $\mu_k = E\left[(x - \mu'_1)^k\right] = \int_{-\infty}^{\infty} (x - \mu'_1)^k f_X(x) dx$

Basics of Probability



Oregon State University
College of Engineering

Moments of Interest to us:



- **1st Moment: Mean** (expected value)
 - Measure of central location

$$E(x) = \int_{-\infty}^{\infty} xf_X(x)dx$$

- **2nd Central Moment: Variance**

- Measure of dispersion

- $Var(x) = \int_{-\infty}^{\infty} (x - \mu)^2 f_X(x)dx$



Basics of Probability




Oregon State University
College of Engineering

- 3rd Central Moment:
Skewness

- Measure for asymmetry


$$Skw(x) = \int_{-\infty}^{\infty} (x - \mu)^3 f_X(x) dx$$



- 4th Central Moment:
Kurtosis

- Measure for peakedness

$$Kurt(x) = \int_{-\infty}^{\infty} (x - \mu)^4 f_X(x) dx$$



Basics of Probability



Oregon State University
College of Engineering

Multiple random variables, i.e. a RANDOM VECTOR: $\vec{X} = [X_1, \dots, X_N]$:

- Joint probability density function $f_{XY}(x, y)$

$$P[a < X < b, c < Y < d] = \int_c^d \int_a^b f_{XY}(x, y) dx dy$$

- Marginal density function of x

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

- Conditional probability density of y given x

$$f_{Y|X}(x) = \frac{f_{XY}(x, y)}{f_X(x)}$$


- Independence

$$f_{XY}(x, y) = f_X(x) f_Y(y)$$

Two Random Variables (X,Y)



Oregon State University
College of Engineering



Basics of Probability


- Covariance

$$COV(X, Y) = E[(X - \mu_X)(Y - \mu_Y)] = E(XY) - E(X)E(Y)$$

- Correlation coefficient


$$\rho_{XY} = \frac{Cov(X, Y)}{\sigma_X \cdot \sigma_Y}$$

→ Degree of linear dependence
between two random variables



Example of correlation


- If we were to measure heights and weights in a certain population, we might see that they are correlated




Correlation




Oregon State University
College of Engineering



Positive Correlation



Negative Correlation



Uncorrelated