

# Algorithmic Methods for Singularities

## Talk 2: Invariants from the computational perspective

Anne Frühbis-Krüger and Matthias Zach

Institut für Mathematik  
Universität Oldenburg

Sao Carlos, July 12th 2022

# Dimension from definition to computation

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Dimension from definition to computation

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

- maximal length  $d$  of a chain of prime ideals

$$P_0 \subsetneq P_1 \subsetneq \dots \subsetneq P_d$$

in  $\mathbb{C}[\underline{x}]/I$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Dimension from definition to computation

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

- maximal length  $d$  of a chain of prime ideals

$$P_0 \subsetneq P_1 \subsetneq \dots \subsetneq P_d$$

in  $\mathbb{C}[\underline{x}]/I$

- minimal number of generators of an  $\mathfrak{m}$ -primary ideal

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Dimension from definition to computation

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

- ▶ maximal length  $d$  of a chain of prime ideals

$$P_0 \subsetneq P_1 \subsetneq \dots \subsetneq P_d$$

in  $\mathbb{C}[\underline{x}]/I$

- ▶ minimal number of generators of an  $\mathfrak{m}$ -primary ideal
- ▶  $\text{trdeg}_{\mathbb{C}}(\text{Quot}(\mathbb{C}[\underline{x}]/I))$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Dimension from definition to computation

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

- ▶ maximal length  $d$  of a chain of prime ideals

$$P_0 \subsetneq P_1 \subsetneq \dots \subsetneq P_d$$

in  $\mathbb{C}[\underline{x}]/I$

- ▶ minimal number of generators of an  $\mathfrak{m}$ -primary ideal
- ▶  $\text{trdeg}_{\mathbb{C}}(\text{Quot}(\mathbb{C}[\underline{x}]/I))$
- ▶ Noether normalization  $\mathbb{C}[y_1, \dots, y_d] \subset \mathbb{C}[\underline{x}]/I$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Dimension from definition to computation

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

- ▶ maximal length  $d$  of a chain of prime ideals

$$P_0 \subsetneq P_1 \subsetneq \dots \subsetneq P_d$$

in  $\mathbb{C}[\underline{x}]/I$

- ▶ minimal number of generators of an  $\mathfrak{m}$ -primary ideal
- ▶  $\text{trdeg}_{\mathbb{C}}(\text{Quot}(\mathbb{C}[\underline{x}]/I))$
- ▶ Noether normalization  $\mathbb{C}[y_1, \dots, y_d] \subset \mathbb{C}[\underline{x}]/I$
- ▶  $\deg_t(\text{HSP}(\mathbb{C}[\underline{x}]_{\mathfrak{m}}/I))$

# Dimension from definition to computation

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $X = V(I) \subseteq \mathbb{A}_{\mathbb{C}}^n$  be irreducible.

What is  $\dim(X)$  or  $\dim(\mathbb{C}[\underline{x}]/I)$  at a point  $x = V(\mathfrak{m}) \in X$ ?

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

- ▶ maximal length  $d$  of a chain of prime ideals

$$P_0 \subsetneq P_1 \subsetneq \dots \subsetneq P_d$$

in  $\mathbb{C}[\underline{x}]/I$

- ▶ minimal number of generators of an  $\mathfrak{m}$ -primary ideal
- ▶  $\text{trdeg}_{\mathbb{C}}(\text{Quot}(\mathbb{C}[\underline{x}]/I))$
- ▶ Noether normalization  $\mathbb{C}[y_1, \dots, y_d] \subset \mathbb{C}[\underline{x}]/I$
- ▶  $\deg_t(\text{HSP}(\mathbb{C}[\underline{x}]_{\mathfrak{m}}/I))$

Efficiently computable: the last two items!



# Hilbert-Samuel Polynomial

Let  $(A, \mathfrak{m})$  be a localization of an affine  $\mathbb{C}$ -algebra.

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Hilbert-Samuel Polynomial

Let  $(A, \mathfrak{m})$  be a localization of an affine  $\mathbb{C}$ -algebra.

**Hilbert-Samuel Function:**

$$HS_A(k) := \dim_{\mathbb{C}} A/\mathfrak{m}^k$$

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Hilbert-Samuel Polynomial

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Let  $(A, \mathfrak{m})$  be a localization of an affine  $\mathbb{C}$ -algebra.

## Hilbert-Samuel Function:

$$HS_A(k) := \dim_{\mathbb{C}} A/\mathfrak{m}^k$$

## Hilbert-Samuel Polynomial:

$$\exists P \in \mathbb{Q}[t] : HS_A(k) = P(k) \quad \forall k \gg 0$$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Hilbert-Samuel Polynomial

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

Let  $(A, \mathfrak{m})$  be a localization of an affine  $\mathbb{C}$ -algebra.

## Hilbert-Samuel Function:

$$HS_A(k) := \dim_{\mathbb{C}} A/\mathfrak{m}^k$$

## Hilbert-Samuel Polynomial:

$$\exists P \in \mathbb{Q}[t] : HS_A(k) = P(k) \quad \forall k \gg 0$$

Let  $P(t) = \sum_{i=0}^d a_i t^i$ , then

- ▶  $\dim(A) = d$
- ▶  $\text{mult}(A) = d! \cdot a_d$

# A monomial example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider the 3 coordinate axes in  $(\mathbb{C}^3, 0)$ :

$$I = \langle xy, xz, yz \rangle \subseteq \mathbb{C}[x, y, z]_{\langle x, y, z \rangle}$$

$$A = \mathbb{C}[x, y, z]_{\langle x, y, z \rangle} / I$$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

$k$	new monomials	total number
1	1	1
2	$x, y, z$	4
3	$x^2, y^2, z^2$	7
4	$x^3, y^3, z^3$	10
$\vdots$		

# A monomial example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider the 3 coordinate axes in  $(\mathbb{C}^3, 0)$ :

$$I = \langle xy, xz, yz \rangle \subseteq \mathbb{C}[x, y, z]_{\langle x, y, z \rangle}$$

$$A = \mathbb{C}[x, y, z]_{\langle x, y, z \rangle} / I$$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

$k$	new monomials	total number
1	1	1
2	$x, y, z$	4
3	$x^2, y^2, z^2$	7
4	$x^3, y^3, z^3$	10
$\vdots$		

Obviously,  $P(t) = 3 \cdot t - 2$

Hence:  $\dim(X) = 1$  and  $\text{mult}(X) = 3$  as expected

# From ideal to monomial ideal

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Key to computing previous example: Monomial ideal

And in the non-monomial case?

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# From ideal to monomial ideal

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Key to computing previous example: Monomial ideal

Computing  
Dimension

And in the non-monomial case? **standard bases**

Standard Bases

- ▶  $K$  a (exact computable) field
- ▶  $>$  suitable total ordering on  $\text{Mon}(\underline{x})$
- ▶  $LT(f)$  largest monomial in  $f \in K[\underline{x}]$  w.r.t.  $>$
- ▶  $L(I) := \langle LT(f) \mid f \in I \rangle$

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

$G = \{g_1, \dots, g_s\}$  standard basis of  $I$

$\Longleftrightarrow$

$L(G) = L(I)$  and  $\langle G \rangle = I$



# Standard Bases

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

$G = \{g_1, \dots, g_s\}$  standard basis of  $I$

$\Longleftrightarrow$

$L(G) = L(I)$  and  $\langle G \rangle = I$

Facts:

- ▶  $\dim(K[\underline{x}]_{\langle \underline{x} \rangle} / I) = \dim(K[\underline{x}]_{\langle \underline{x} \rangle} / L(I))$
- ▶  $\text{mult}(K[\underline{x}]_{\langle \underline{x} \rangle} / I) = \text{mult}(K[\underline{x}]_{\langle \underline{x} \rangle} / L(I))$

# Example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider

- ▶  $I = \langle x^2 + y^2, xy \rangle \subseteq \mathbb{Q}[x, y]_{\langle x, y \rangle}$
- ▶  $>$  negative degree reverse lexicographical ordering
- ▶  $L(x^2 + y^2) = x^2$ ,  $L(xy) = xy$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider

- ▶  $I = \langle x^2 + y^2, xy \rangle \subseteq \mathbb{Q}[x, y]_{\langle x, y \rangle}$
- ▶  $>$  negative degree reverse lexicographical ordering
- ▶  $L(x^2 + y^2) = x^2$ ,  $L(xy) = xy$

Obviously:

$$y^3 = y \cdot (x^2 + y^2) - x \cdot xy \in I \text{ and hence in } L(I)$$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider

- ▶  $I = \langle x^2 + y^2, xy \rangle \subseteq \mathbb{Q}[x, y]_{\langle x, y \rangle}$
- ▶  $>$  negative degree reverse lexicographical ordering
- ▶  $L(x^2 + y^2) = x^2$ ,  $L(xy) = xy$

Obviously:

$$y^3 = y \cdot (x^2 + y^2) - x \cdot xy \in I \text{ and hence in } L(I)$$

Actually, standard basis of  $I$ :

$$G = \{x^2 + y^2, xy, y^3\}$$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# A multiplicity 4 space curve example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider

- ▶  $I = \langle xy, x^2 + y^2 + z^2 \rangle \subseteq \mathbb{Q}[x, y, z]_{\langle x, y, z \rangle}$
- ▶  $>$  suitable negative degree ordering  
(e.g.  $\text{negdeglex}(z > y > x)$ )
- ▶  $L(xy) = xy$ ,  $L(x^2 + y^2 + z^2) = z^2$

This is already a standard basis.

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# A multiplicity 4 space curve example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider

- ▶  $I = \langle xy, x^2 + y^2 + z^2 \rangle \subseteq \mathbb{Q}[x, y, z]_{\langle x, y, z \rangle}$
- ▶  $>$  suitable negative degree ordering  
(e.g.  $\text{negdeglex}(z > y > x)$ )
- ▶  $L(xy) = xy$ ,  $L(x^2 + y^2 + z^2) = z^2$

This is already a standard basis.

$k$	<i>newmonomials</i>	<i>totalnumber</i>
1	1	1
2	$x, y, z$	4
3	$x^2, y^2, xz, yz$	8
4	$x^3, y^3, x^2z, y^2z$	12
$\vdots$		

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# A multiplicity 4 space curve example

First Steps

Anne  
Frühbis-Krüger  
and Matthias Zach

Consider

- ▶  $I = \langle xy, x^2 + y^2 + z^2 \rangle \subseteq \mathbb{Q}[x, y, z]_{\langle x, y, z \rangle}$
- ▶  $>$  suitable negative degree ordering  
(e.g.  $\text{negdeglex}(z > y > x)$ )
- ▶  $L(xy) = xy, L(x^2 + y^2 + z^2) = z^2$

This is already a standard basis.

$k$	$\text{newmonomials}$	$\text{totalnumber}$
1	1	1
2	$x, y, z$	4
3	$x^2, y^2, xz, yz$	8
4	$x^3, y^3, x^2z, y^2z$	12
$\vdots$		

$$P(t) = 4 \cdot t - 4, \dim = 1, \text{mult} = 4$$

Computing  
Dimension

Standard Bases

Milnor and Tjurina  
number I

Milnor and Tjurina  
number II

# Jupyter Notebook 2, Block 1