

Mathematical Proofs
A Transition to Advanced Mathematics
Chapter 14
Proofs in Calculus

Gary Chartrand Albert D. Polimeni Ping Zhang

Proofs in Calculus

The proofs that occur in calculus are considerably different than any of those we have seen thus far. The functions encountered in calculus are real-valued functions defined on sets of real numbers. That is, each function that we study in calculus is of the type

$$f : X \rightarrow \mathbf{R}, \text{ where } X \subseteq \mathbf{R}.$$

In the study of limits, we are often interested in such functions having the property that either

- (1) $X = \mathbf{N}$ and increasing values in the domain \mathbf{N} result in functional values approaching some real number L or
- (2) the function is defined for all real numbers near some specified real number a and domain values approaching a result in functional values approaching some real number L .

We begin with (1), where $X = \mathbf{N}$.

Definition

A **sequence** (of real numbers) is a real-valued function defined on the set of natural numbers; that is, a **sequence** is a function

$$f : \mathbf{N} \rightarrow \mathbf{R}.$$

If $f(n) = a_n$ for each $n \in \mathbf{N}$, then

$$f = \{(1, a_1), (2, a_2), (3, a_3), \dots\}.$$

Since only the numbers a_1, a_2, a_3, \dots are relevant in f , this sequence is often denoted only by a_1, a_2, a_3, \dots or by $\{a_n\}$.

Limits of Sequences

Definition

The numbers a_1, a_2, a_3 , etc. are called the **terms** of the sequence $\{a_n\}$, with a_1 being the first term, a_2 the second term, etc. Thus, a_n is the n th term of the sequence.

$\left\{\frac{1}{n}\right\}$ is the sequence $1, 1/2, 1/3, \dots$;

$\left\{\frac{n}{2n+1}\right\}$ is the sequence $1/3, 2/5, 3/7, \dots$.

In these two examples, the n th term of a sequence is given and, from this, we can easily find the first few terms and, in fact, any particular term. On the other hand, finding the n th term of a sequence whose first few terms are given can be challenging.

Limits of Sequences

For example, the n th term of the sequence

$$\frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \dots$$

is $1/2n$; the n th term of the sequence

$$1 + \frac{1}{2}, 1 + \frac{1}{4}, 1 + \frac{1}{8}, \dots$$

is $1 + 1/2^n$; the n th term of the sequence

$$1, \frac{3}{5}, \frac{1}{2}, \frac{5}{11}, \frac{3}{7}, \frac{7}{17}, \dots$$

is $(n + 1)/(3n - 1)$; the n th term of the sequence

$$1, -1, 1, -1, 1, -1, \dots$$

is $(-1)^{n+1}$; while the n th term of the sequence $1, 4, 9, 16, \dots$ is n^2 .

Limits of Sequences

For the sequence $\left\{\frac{1}{n}\right\}$, the larger the integer n , the closer $1/n$ is to 0; and for the sequence $\left\{\frac{n}{2n+1}\right\}$, the larger the integer n , the closer $n/(2n+1)$ is to $1/2$. On the other hand, for the sequence $\{n^2\}$, as the integer n become larger, n^2 becomes increasingly large and does not approach any real number.

Limits of Sequences

For some sequences $\{a_n\}$, there is a real number L (or at least there appears to be a real number L) such that the larger the integer n becomes, the closer a_n is to L .

Definition

A sequence $\{a_n\}$ of real numbers is said to **converge** to the real number L if for every real number $\epsilon > 0$, there exists a positive integer N such that if n is an integer with $n > N$, then $|a_n - L| < \epsilon$.

The number ϵ is a measure of how close the terms a_n are required to be to the number L and N indicates a position in the sequence beyond which the required condition is satisfied.

Definition

If a sequence $\{a_n\}$ converges to L , then $\{a_n\}$ is a **convergent sequence** and L is referred to as the **limit** of $\{a_n\}$ and we write

$$\lim_{n \rightarrow \infty} a_n = L.$$

If a sequence does not converge, it is said to **diverge**.
Consequently, if a sequence $\{a_n\}$ diverges, then there is *no* real number L such that $\lim_{n \rightarrow \infty} a_n = L$.

Limits of Sequences

For a real number x , the *ceiling* $\lceil x \rceil$ of x is the smallest integer greater than or equal to x .

$$\lceil 8/3 \rceil = 3, \lceil \sqrt{2} \rceil = 2, \lceil -1.6 \rceil = -1 \text{ and } \lceil 5 \rceil = 5.$$

Example 1

Result The sequence $\left\{ \frac{1}{n} \right\}$ converges to 0.

Proof We want to show $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$.

Let $\varepsilon > 0$.

Choose $N = \lceil \frac{1}{\varepsilon} \rceil \geq \frac{1}{\varepsilon}$. For $n > N$, we have

$$\left| \frac{1}{n} - 0 \right| = \frac{1}{n} < \frac{1}{N} \leq \varepsilon.$$



Example 2

Result The sequence $\left\{3 + \frac{2}{n^2}\right\}$ converges to 3.

Proof We want to show $\lim_{n \rightarrow \infty} 3 + \frac{2}{n^2} = 3$.

Let $\varepsilon > 0$. Choose $N = \lceil \sqrt{2/\varepsilon} \rceil \geq \sqrt{2/\varepsilon}$.

For $n > N$, we have

$$\left| 3 + \frac{2}{n^2} - 3 \right| = \frac{2}{n^2} < \frac{2}{N^2} < \frac{2}{(2/\varepsilon)} = \varepsilon.$$

Therefore, $\lim_{n \rightarrow \infty} 3 + \frac{2}{n^2} = 3$.



Limits of Sequences

Example 3

Result The sequence $\left\{ \frac{n}{2n+1} \right\}$ converges to $\frac{1}{2}$.

Proof Suppose $\varepsilon > 0$. Choose $N = \left\lceil \frac{1}{4\varepsilon} - \frac{1}{2} \right\rceil$ (so $N > \frac{1}{4\varepsilon} - \frac{1}{2}$)

For $n > N$, we have

$$\begin{aligned} \left| \frac{n}{2n+1} - \frac{1}{2} \right| &= \left| \frac{2n - 2n - 1}{2(2n+1)} \right| = \left| \frac{-1}{4n+2} \right| = \frac{1}{4n+2} \\ &< \frac{1}{4N+2} \leq \frac{1}{4\left(\frac{1}{4\varepsilon} - \frac{1}{2}\right) + 2} = \frac{1}{\frac{1}{\varepsilon} - 2 + 2} = \varepsilon. \end{aligned}$$

$$\text{Thus, } \lim_{n \rightarrow \infty} \frac{n}{2n+1} = \frac{1}{2}.$$



Example 4

Result The sequence $\{(-1)^{n+1}\}$ is divergent.

Proof By contradiction, suppose it is.

Thus, $\forall \varepsilon > 0$, $\exists N > 0$ such that $|a_n - L| < \varepsilon$
for any $n > N$.

Pick $\varepsilon = 1$. Let $k > N$ such that k is odd.

This means

$$\begin{aligned} |(-1)^{k+1} - L| < 1 &\Rightarrow |1 - L| < 1 \Rightarrow -1 < 1 - L < 1 \\ &\Rightarrow -2 < -L < 0 \Rightarrow 0 < L < 2. \end{aligned}$$

In a similar fashion, let $k > N$ be an even integer.

This means

$$\begin{aligned} |(-1)^{k+1} - L| < 1 &\Rightarrow |-1 - L| < 1 \Rightarrow |1 + L| < 1 \\ &\Rightarrow -1 < 1 + L < 1 \Rightarrow -2 < L < 0. \end{aligned}$$

Both of these imply $0 < L < 0$. This is a contradiction.

Thus $\{(-1)^{n+1}\}_{n=1}^{\infty}$ must diverge.

Limits of Sequences

A sequence $\{a_n\}$ may diverge because as n becomes larger, a_n becomes larger and eventually exceeds any given real number. If a sequence has this property, then $\{a_n\}$ is said to diverge to infinity.

Definition

More formally, a sequence $\{a_n\}$ **diverges to infinity**, written $\lim_{n \rightarrow \infty} a_n = \infty$, if for every positive number M , there exists a positive integer N such that if n is an integer such that $n > N$, then $a_n > M$.

Limits of Sequences

The sequence $\{(-1)^{n+1}\}$, although divergent, does not diverge to infinity. However, the sequence $\{n^2 + \frac{1}{n}\}$ does diverge to infinity.

Example 5

Result $\lim_{n \rightarrow \infty} \left(n^2 + \frac{1}{n} \right) = \infty.$

Proof Suppose $M > 0$. Choose $N = \lceil \sqrt{M} \rceil \geq \sqrt{M}$.

Then for $n > N$,

$$n^2 + \frac{1}{n} > n^2 > N^2 \geq (\sqrt{M})^2 = M.$$

Thus $\{n^2 + \frac{1}{n}\}_{n=1}^{\infty}$ diverges.



Definition

Let f be a real-valued function defined on a set X of real numbers and let $a \in \mathbf{R}$ such that f is defined in some deleted neighborhood of a .

A number L is the **limit** of a function $f(x)$ as x approaches a , written $\lim_{x \rightarrow a} f(x) = L$, if for every real number $\epsilon > 0$, there exists a real number $\delta > 0$ such that for every real number x with $0 < |x - a| < \delta$, it follows that $|f(x) - L| < \epsilon$.

This implies that if $0 < |x - a| < \delta$, then certainly $f(x)$ is defined.

Limits of Functions

If there exists a number L such that $\lim_{x \rightarrow a} f(x) = L$, then we say that the limit $\lim_{x \rightarrow a} f(x)$ exists and is equal to L ; otherwise, this limit does not exist.

Thus, to show that $\lim_{x \rightarrow a} f(x) = L$, it is necessary to specify $\epsilon > 0$ first and then show the existence of a real number $\delta > 0$.

Ordinarily, the smaller the value of ϵ , the smaller the value of δ . However, we must be certain that the number δ selected satisfies the requirement regardless of how small (or large) ϵ may be.

Even though our choice of δ depends on ϵ , it should not depend on which real number x with $0 < |x - a| < \delta$ is being considered.

Example 6

Result $\lim_{x \rightarrow 4} (3x - 7) = 5.$

Proof. Let $\varepsilon > 0$. Choose $\delta = \frac{\varepsilon}{3}$. Then
for $0 < |x - 4| < \delta$, we have

$$|3x - 7 - 5| = |3(x - 4)| = 3|x - 4| < 3\delta = 3 \frac{\varepsilon}{3} = \varepsilon.$$

Thus $\lim_{x \rightarrow 4} (3x - 7) = 5.$



Example 7

Result $\lim_{x \rightarrow \frac{3}{2}} \frac{4x^2 - 9}{2x - 3} = 6.$

Proof. Let $\varepsilon > 0$. Choose $\delta = \frac{\varepsilon}{2}$. Then for $0 < |x - \frac{3}{2}| < \delta$, we have,

$$\begin{aligned} \left| \frac{4x^2 - 9}{2x - 3} - 6 \right| &= \left| \frac{(2x - 3)(2x + 3)}{2x - 3} - 6 \right| = |2x + 3 - 6| \\ &= 2|x - \frac{3}{2}| < 2\delta = 2 \cdot \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Thus $\lim_{x \rightarrow \frac{3}{2}} \frac{4x^2 - 9}{2x - 3} = 6.$



Example 8

Result $\lim_{x \rightarrow 3} x^2 = 9$.

Proof. Let $\varepsilon > 0$. Choose $\delta < -3 + \sqrt{9 + \varepsilon}$. Then for $0 < |x - 3| < \delta$ we have

$$-\delta < x - 3 < \delta \Rightarrow -\delta + 3 < x < \delta + 3 \Rightarrow -\delta + 6 < x + 3 < \delta + 6,$$

so $|x + 3| < \delta + 6$. Thus

$$\begin{aligned} |x^2 - 9| &= |x + 3| \cdot |x - 3| < (\delta + 6) \delta < (3 + \sqrt{9 + \varepsilon})(-3 + \sqrt{9 + \varepsilon}) \\ &= -9 + 9 + \varepsilon \\ &= \varepsilon. \end{aligned}$$

Thus $\lim_{x \rightarrow 3} x^2 = 9$.

