

ENN595-1 Project Students Semester Two 2026

Investigation via the Engineering Literature

Assessment 1A

Student Name: Joshua Hecke

Student No# : n11585382

Research Topic Title: Investigation of Primitive-Space Scene Change Detection in Underwater 3D Gaussian Splatting Models


Project Supervisor: Niko Sünderhauf

Assessment Tasks

1. Draw and label a concept map of your research topic;
2. Translate those key concepts into a search statement;
3. Search for relevant engineering literature in a library database;
4. Identify authoritative, reputable and reliable information sources relevant to your research topic;
5. Submit a summary analysis and evaluation of three (3) key references;
6. Locate three (3) published authorities (acknowledged experts) in your field of research;
7. Identity, categorize and synthesise trends and directions found in the literature;
8. Explain what impact the literature findings have had on the design of your project;
9. Compile a reference list of the most useful literature on your topic.

This document has been prepared using the provided template on Canvas. It contains identical sections and headings to the template.

Concept Map : Draw and label a graphical outline view of your research question.



Overall Project Objective: To research and evaluate the application of primitive-space scene change detection (specifically GS-DIFF) on 3D Gaussian Splatting models reconstructed from underwater environments.

Questions to be investigated:

- 1. Can the primitive-space comparison method (GS-DIFF) handle the specific geometric and photometric drift inherent in underwater 3DGS reconstructions?
- 2. Can the integration of Surface-Aligned Gaussian Splatting (SuGaR) into the GS-DIFF pipeline improve scene change detection accuracy on complex underwater geometry by enforcing structural regularization and mitigating reconstruction noise?

Scope: Focus on newly proposed 3D Gaussian Splatting methods designed for underwater environments (such as WaterSplatting, SeaSplat, and Aqua-Splat) and the application of scene change detection within these complex marine models.

Now take the terms and phrases from your concept map to develop and refine a search strategy.

1st Concept	AND	2nd Concept	AND	3rd Concept
3D Gaussian Splatting		Change Detection		Applications & Environments

OR	Synonyms	AND	Synonyms	AND	Synonyms	OR
	3DGS, Explicit Geometry, SuGaR		GS-DIFF, Render-then-compare		Underwater, Street View, Crop Analysis	

splat*, geometr*, detect*, compar*, underwat*, crop*, street*

("3D Gaussian Splatting" OR "3DGS" OR "Explicit Geometry" OR "SuGaR")
AND ("change detection" OR "GS-DIFF" OR "render-then-compare")
AND (underwater OR "street view" OR "crop analysis")

Search for relevant engineering literature in a library database.

Database Name: IEEE Xplore

Database Platform: IEEE

Comment on the scope: IEEE Xplore provides extensive coverage of high-impact engineering and computer science proceedings, capturing the most recent advancements in robotics (ICRA, IROS, RA-L) and 3D computer vision (CVPR, ICCV) where state-of-the-art underwater 3DGS and change detection research is primarily published.

Identify relevant and authoritative information sources.

The table below summarizes the 22 information sources identified for this project so far, categorized by their types to ensure wide coverage across methodologies, applications, and procedural standards.

Category Breakdown:

- a. **Peer-reviewed journal titles / Conference publications:** The majority of the sources fall under this category, ensuring cutting-edge, peer-reviewed methodologies. Key venues include IEEE Robotics and Automation Letters (RA-L), IEEE/CVF CVPR, ICRA, and MDPI Sensors.
- b. **Texts / Handbooks / Encyclopaedias:** Comprehensive literature reviews, such as Wu et al.’s “Recent advances in 3D Gaussian splatting” [1] and Zhong et al.’s review on scene editing [2], serve as encyclopedic handbooks covering the foundation and evolution of the technology.
- c. **Standards (or regulations) / Patents:** Gordon et al.’s “Field photogrammetry in 4D” [3] provides a Standard Operational Procedure (SOP) published by the Australian Institute of Marine Science, guiding the practical capture and reconstruction of marine environments.

Ref	Author (Year)	Title	Source Type
[4]	Xu et al. (2026)	NAS-GS: Noise-Aware Sonar Gaussian Splatting	Preprint (arXiv)
[5]	Yang et al. (2026)	SonarReg-GS SLAM: Sparse Sonar-Guided Depth Regularization	Journal (Sensors)
[6]	Sethuraman et al. (2025)	SonarSplat: Novel View Synthesis of Imaging Sonar...	Journal (IEEE RA-L)
[7]	Ling et al. (2025)	Aqua-Splat: Physically-Informed Sonar-Camera Gaussian...	Journal (IEEE RA-L)
[8]	Li et al. (2025)	WaterSplatting: Fast Underwater 3D Scene Reconstruction	Preprint (arXiv)
[9]	Huang et al. (2025)	From Restoration to Reconstruction: Rethinking 3DGS...	Preprint (arXiv)
[10]	Zhang et al. (2024)	RecGS: Removing Water Caustic with Recurrent...	Preprint (arXiv)
[11]	Liu et al. (2025)	Spatiotemporal Degradation-Aware 3DGS...	Conf. (ACM MM)
[12]	Levy et al. (2023)	SeaThru-NeRF: Neural Radiance Fields in Scattering Media	Conf. (CVPR)
[13]	Fan et al. (2025)	Water-Adapted 3D Gaussian Splatting for precise...	Journal (Frontiers)
[14]	Yang et al. (2025)	SeaSplat: Representing Underwater Scenes with 3DGS...	Conf. (ICRA)
[15]	Zhang et al. (2026)	Underwater Image Restoration Integrating Monocular Depth...	Journal (JMSE)
[16]	Ngo et al. (2020)	Single Image Haze Removal from Image Enhancement...	Journal (Sensors)
[17]	Blasinski et al. (2016)	A Three Parameter Underwater Image Formation Model	Journal (Elec. Imag.)
[18]	Xu et al. (2022)	Point-NeRF: Point-based Neural Radiance Fields	Conf. (CVPR)
[1]	Wu et al. (2024)	Recent advances in 3D Gaussian splatting	Journal (CVM)
[19]	Zhou et al. (2025)	3D Scene Change Modeling With Consistent Multi-View...	Preprint (arXiv)
[20]	Galappaththige et al. (2026)	From Pixels to Primitives: Scene Change Detection...	Preprint (arXiv)
[2]	Zhong et al. (2026)	3D Gaussian Splatting for Scene Editing and Completion	Conf. (ICAIDS)
[21]	Lu et al. (2025)	3DGS-CD: 3D Gaussian Splatting-Based Change Detection...	Journal (IEEE RA-L)
[3]	Gordon et al. (2023)	Field photogrammetry in 4D. Reef Restoration (SOP)	Tech. Report (Standard)
[22]	Guédon et al. (2024)	SuGaR: Surface-Aligned Gaussian Splatting for Efficient...	Conf. (CVPR)

Evaluate and analyse three (3) key references.

Reference # 1

Citation: A. Guédon and V. Lepetit, "SuGaR: Surface-Aligned Gaussian Splatting for Efficient 3D Mesh Reconstruction and High-Quality Mesh Rendering," in *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, 2024, pp. 5354-5363 [22].

- **How does this paper relate to my topic?** This paper introduces SuGaR, a method that forces 3D Gaussians to align closely with the underlying surfaces of a scene. In the context of the project, SuGaR acts as a critical structural regularizer to clean up the chaotic and noisy geometry inherent in underwater 3DGS reconstructions before applying primitive-space change detection.
- **What key findings are reported?** The authors demonstrate that by applying a regularization term that encourages Gaussians to become flat and perfectly aligned with the surface normal, they can extract highly accurate 3D meshes directly from the 3DGS representation. This significantly reduces the typical "floaters" and disordered geometry found in standard 3DGS.
- **How does this paper relate to other work in the field?** While standard 3DGS produces excellent novel view synthesis, its underlying geometry is often highly chaotic. SuGaR bridges the gap between the rendering speed of 3DGS and the geometric consistency of traditional mesh-based or SDF-based methods like NeuS.
- **Interpret important figures/tables/illustrations (cite the page number).** Figure 2 (pg. 3) illustrates the regularization process where Gaussians are compressed along their shortest axis to align with the estimated surface normals. Figure 5 (pg. 7) shows how SuGaR completely eliminates floaters and artifacts compared to vanilla 3DGS.
- **List other references to investigate further.** GS-DIFF (Galappaththige et al., 2026) to see how these regularized Gaussians are compared, and 2DGS (Huang et al., 2024) for alternative geometric regularization.
- **General notes/observations on this paper's attributes.** This method will need to be modified for underwater 3DGS, as the surface normals may be less reliable due to scattering and attenuation. However, the underlying principle of enforcing surface alignment is likely to improve the robustness of primitive-space change detection in marine environments.

Reference # 2

Citation: D. Yang, J. J. Leonard, and Y. Girdhar, "SeaSplat: Representing Underwater Scenes with 3D Gaussian Splatting and a Physically Grounded Image Formation Model," in *2025 IEEE International Conference on Robotics and Automation (ICRA)*, 2025 [14].

- **How does this paper relate to my topic?** SeaSplat adapts 3DGS specifically for underwater scenes, providing a framework for how marine 3DGS models are reconstructed and color-restored, which acts as the foundational representation on which we intend to perform scene change detection.
- **What key findings are reported?** SeaSplat successfully combines 3D Gaussian Splatting with a physically grounded underwater image formation model. By simultaneously learning the parameters of the medium (attenuation and backscatter) alongside the 3DGS representation, it restores the scene’s true color and accurately estimates geometry without the "floaters" typical of standard 3DGS in scattering media.
- **How does this paper relate to other work in the field?** Unlike standard 3DGS or NeRF methods, SeaSplat explicitly models the underwater medium. It achieves real-time rendering and better visual quality compared to SeaThru-NeRF, while remaining computationally efficient.
- **Interpret important figures / tables / illustrations (cite the page number).** Figure 2 (pg. 3) illustrates the splatting pipeline with the medium formation model. Figure 5 (pg. 6) showcases qualitative novel view synthesis, demonstrating how SeaSplat successfully restores underlying colors compared to baselines.
- **List other references to investigate further.** SeaThru-NeRF (Levy et al., 2023) for precursor performance and "A Three Parameter Underwater Image Formation Model" (Blasinski et al., 2016) for more modeling approaches.
- **General notes/observations on this paper’s attributes.** The explicit separation of the object color and the scattering medium is highly beneficial. Change detection methods like GS-DIFF will likely perform much better if applied only to the restored "object" Gaussians rather than the medium-entangled raw representations.

Reference # 3

Citation: H. Li, W. Song, T. Xu, A. Elsig, and J. Kulhanek, "WaterSplatting: Fast Underwater 3D Scene Reconstruction Using Gaussian Splatting," *arXiv preprint arXiv:2408.08206*, 2025 [8].

- **How does this paper relate to my topic?** It presents an alternative state-of-the-art methodology for handling scattering media with 3DGS. Evaluating how WaterSplatting handles volumetric geometry vs. surface geometry is crucial for understanding how primitive-space change detection can be applied.
- **What key findings are reported?** WaterSplatting employs 3DGS for explicit geometry representation alongside a separate volumetric field (queried once per pixel) for capturing the scattering medium. This dual representation allows for rapid scene restoration and novel view synthesis, outperforming NeRF-based methods in rendering quality on underwater datasets.

- **How does this paper relate to other work in the field?** Unlike SeaSplat which embeds the medium model directly into the splatting mathematics, WaterSplatting uses a separate volumetric field for the medium. This keeps the 3DGS geometry explicit and clean, which may be more suitable for primitive-level change detection algorithms.
- **Interpret important figures / tables / illustrations (cite the page number).** Figure 1 (pg. 1) compares WaterSplatting’s real-time restored rendering with SeaThru-NeRF. Figure 2 (pg. 4) illustrates the dual representation pipeline separating medium and object components.
- **List other references to investigate further.** SeaSplat (Yang et al., 2025) to compare medium modeling approaches, the use of sonar or other sensors for depth estimation, and the original 3DGS paper (Kerbl et al., 2023) for foundational understanding.
- **General notes/observations on this paper’s attributes.** The separation of the scattering medium into a distinct volumetric field means the underlying 3DGS primitives purely represent scene geometry. This is an ideal input format for GS-DIFF’s geometric and appearance kernels.

Identify and locate published authorities (experts) on your research question.

1. **John J. Leonard (MIT):** A highly cited expert in SLAM and marine robotics. He is a co-author of several highly relevant recent papers applying 3DGS to robotics and underwater scenes, including SeaSplat and 3DGS-CD. [Google Scholar Profile Link](#).
2. **Bernhard Kerbl (University of Copenhagen):** The original co-lead author of the seminal 3D Gaussian Splatting paper. He is a recognized authority in real-time rendering and neural radiance fields, forming the foundation of the 3DGS representations used in this project. He is currently an Assistant Professor at the University of Copenhagen. [Google Scholar Profile Link](#).
3. **Lin Gao (Chinese Academy of Sciences):** An expert in computer graphics and 3D vision, and co-author of a comprehensive recent review on 3D Gaussian splatting advancements [1]. He has extensive experience in 3D modeling and neural rendering techniques. [Google Scholar Profile Link](#).

Identify, categorize and synthesize findings from your examination of the literature.

- **Adaptation of 3DGS to Marine Environments:** There is a rapid emergence of 3DGS methods tailored for underwater conditions (e.g., WaterSplatting, SeaSplat,

Aqua-Splat, SonarReg-GS, RecGS). These models specifically address marine challenges like light scattering, wavelength-dependent attenuation, and caustics, often by explicitly separating the scene geometry from the scattering medium.

- **Primitive-Space Scene Change Detection:** Concurrently, change detection in 3DGS is shifting towards direct "primitive-space" comparison (e.g., GS-DIFF) which eliminates the reliance on 2D feature extractors by comparing the Gaussians' geometric and photometric parameters directly.
- **Synthesis & Gap:** While new underwater 3DGS methods effectively restore scene geometry and color, and primitive-space change detection algorithms like GS-DIFF provide robust 3D change localization in air, the combination of these two advancements remains unexplored. There is a significant gap in understanding how primitive-space change detection performs on the unique primitive structures generated by underwater 3DGS reconstructions. This gap motivates the research project to explore the feasibility and performance of combining methods like WaterSplatting/SeaSplat with GS-DIFF for robust change detection in marine environments.

Explain the impact of your literature evaluation on the design of your research project.

The findings from the literature evaluation directly shape the risk management and methodology of the project design.

- Because underwater 3DGS reconstructions are inherently noisier suffering from severe geometric and photometric drift due to scattering, attenuation, and poor visibility—applying primitive-space change detection algorithms like GS-DIFF directly to raw underwater reconstructions is highly susceptible to false positives. To address this, the project methodology will incorporate Surface-Aligned Gaussian Splatting (SuGaR) into the pipeline. By forcing the 3D Gaussians to align with the underlying scene surfaces, SuGaR produces cleaner representations of complex underwater geometry and suppresses noisy artifacts like "floaters."
- By feeding this structurally regularized primitive data into the GS-DIFF algorithm, the project aims to significantly enhance the accuracy and robustness of primitive-space scene change detection in marine environments. The core methodology will evaluate a novel, end-to-end pipeline: **Underwater Gaussian Splatting (e.g., SeaSplat) + SuGaR + GS-DIFF**. This pipeline first leverages physically grounded underwater image formation to restore color and initial geometry, applies SuGaR to enforce surface-aligned structural regularization on the resulting splats, and finally utilizes GS-DIFF to perform direct primitive-space change detection on the complex underwater geometry, completely eliminating the need for 2D render-then-compare methodologies.

Reference List.

References

- [1] T. Wu, Y.-J. Yuan, L.-X. Zhang, J. Yang, Y.-P. Cao, L.-Q. Yan, and L. Gao, “Recent advances in 3d gaussian splatting,” *Computational Visual Media*, vol. 10, no. 4, pp. 613–642, 2024.
- [2] Q. Zhong, W. Liu, H. Zhou, P. Lu, and W. Shi, “3d gaussian splatting for scene editing and completion,” in *Proceedings of the 2nd International Conference on Artificial Intelligence, Digital Media Technology and Social Computing*, ICAIDS ’26, (New York, NY, USA), p. 14–25, Association for Computing Machinery, 2026.
- [3] S. Gordon, E. Aston, M. Lechene, J. Harianto, P. Bray, W. Figueira, M. Gonzalez Rivero, and R. Ferrari Legorreta, “Field photogrammetry in 4D. Reef Restoration and Adaption Program (EcoRRAP). Standard Operational Procedure Number 14 (1 of series): Overview and in-field workflow,” tech. rep., Australian Institute of Marine Science, Townsville, 2023.
- [4] S. Xu, J. Jiang, J. S. Willners, and S. Wang, “Nas-gs: Noise-aware sonar gaussian splatting,” 2026.
- [5] W. Yang, X. Qian, X. Liu, and J. Leng, “Sonarreg-gs slam: Sparse sonar-guided depth regularization for underwater gaussian splatting slam,” *Sensors*, vol. 26, no. 15, 2026.
- [6] A. V. Sethuraman, M. Rucker, O. Bagoren, P.-C. Kung, N. N. Amutha, and K. A. Skinner, “Sonarsplat: Novel view synthesis of imaging sonar via gaussian splatting,” *IEEE Robotics and Automation Letters*, vol. 10, no. 12, pp. 13312–13319, 2025.
- [7] Z. Ling, Y. Feng, A. Meng, R. Xiao, S. Pan, W. Lu, and L. Hu, “Aqua-splat: Physically-informed sonar-camera gaussian splatting for underwater 3d reconstruction,” *IEEE Robotics and Automation Letters*, vol. 10, no. 11, pp. 11896–11903, 2025.
- [8] H. Li, W. Song, T. Xu, A. Elsig, and J. Kulhanek, “Watersplatting: Fast underwater 3d scene reconstruction using gaussian splatting,” 2025.
- [9] G. Huang, H. Wang, Z. Qi, W. Lu, D. Bull, and N. Anantrasirichai, “From restoration to reconstruction: Rethinking 3d gaussian splatting for underwater scenes,” 2025.
- [10] T. Zhang, W. Zhi, K. Huang, J. Mangelson, C. Barbalata, and M. Johnson-Roberson, “Recgs: Removing water caustic with recurrent gaussian splatting,” 2024.
- [11] S. Liu, N. Gao, Z. Gu, H. Dou, Y. Deng, and H. Li, “Spatiotemporal degradation-aware 3d gaussian splatting for realistic underwater scene reconstruction,” in *Proceedings of the 33rd ACM International Conference on Multimedia*, MM ’25, pp. 141–150, ACM, Oct. 2025.
- [12] D. Levy, A. Peleg, N. Pearl, D. Rosenbaum, D. Akkaynak, S. Korman, and T. Treibitz, “Seathru-nerf: Neural radiance fields in scattering media,” in *2023 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pp. 56–65, 2023.

- [13] X. Fan, X. Wang, H. Ni, Y. Xin, and P. Shi, “Water-adapted 3d gaussian splatting for precise underwater scene reconstruction,” *Frontiers in Marine Science*, vol. Volume 12 - 2025, 2025.
- [14] D. Yang, J. J. Leonard, and Y. Girdhar, “Seasplat: Representing underwater scenes with 3d gaussian splatting and a physically grounded image formation model,” in *2025 IEEE International Conference on Robotics and Automation (ICRA)*, pp. 7632–7638, 2025.
- [15] T. Zhang, H. Qin, Q. Liu, and X. Liu, “Underwater image restoration integrating monocular depth estimation with a physical imaging model,” *Journal of Marine Science and Engineering*, vol. 14, no. 6, 2026.
- [16] D. Ngo, S. Lee, Q.-H. Nguyen, T. M. Ngo, G.-D. Lee, and B. Kang, “Single image haze removal from image enhancement perspective for real-time vision-based systems,” *Sensors*, vol. 20, no. 18, 2020.
- [17] H. Blasinski and J. Farrell, “A three parameter underwater image formation model,” *Electronic Imaging*, vol. 28, no. 18, pp. 1–1, 2016.
- [18] Q. Xu, Z. Xu, J. Philip, S. Bi, Z. Shu, K. Sunkavalli, and U. Neumann, “Point-nerf: Point-based neural radiance fields,” in *2022 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pp. 5428–5438, 2022.
- [19] Z. Zhou, J. Ni, S. Zhang, Y. Chen, and S. Huang, “3d scene change modeling with consistent multi-view aggregation,” 2025.
- [20] C. J. Galappaththige, J. Lai, T. Patten, D. Dansereau, N. Suenderhauf, and D. Miller, “From pixels to primitives: Scene change detection in 3d gaussian splatting,” 2026.
- [21] Z. Lu, J. Ye, and J. Leonard, “3dgs-cd: 3d gaussian splatting-based change detection for physical object rearrangement,” *IEEE Robotics and Automation Letters*, vol. 10, no. 3, pp. 2662–2669, 2025.
- [22] A. Guédon and V. Lepetit, “Sugar: Surface-aligned gaussian splatting for efficient 3d mesh reconstruction and high-quality mesh rendering,” in *2024 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pp. 5354–5363, 2024.