


# Animal Movement Course at Ekenäs

Indoor animal movement data  
from the CSI:DT project

# Lars Rönnegård




- PhD thesis on reindeer in 2003
- linear mixed models, generalized linear models, ....
- hglm package on CRAN since 2010
- Co-authored a book on hierarchical generalized linear models
- Started CSI:DT in 2020



[www.larsronnegard.se](http://www.larsronnegard.se)



# CSI:DT - Cow Social Interaction and Disease Transmission







# GEA-CowView

A real-time positioning system

tag



anchor






 Anchor

Swedish farm

Dutch farm

10 m



-  Anchor
-  Feeding Area

# Data



Illustration: Fredrik Saarkoppel

## Swedish Farm

~200 cows (2 groups)

Milking system: Milking Parlour

Breeds: Holstein, RDC, Cross

Cow traits: Parity, DIM, etc

Positions: November 2019-

## Dutch Farm

~200 cows (1 group)

Milking system: Milking Robots

Breeds: Holstein

Cow traits: Parity, DIM, etc

Positions: February 2020-



# Data



Illustration: Fredrik Saarkoppel

## Swedish Farm

~200 cows (2 groups)

Milking system: Milking Parlour

Breeds: Holstein, RDC, Cross

Cow traits: Parity, DIM, etc

Positions: November 2019-

Parity = no. of calves a cow has given birth to

## Dutch Farm

~200 cows (1 group)

Milking system: Milking Robots

Breeds: Holstein

Cow traits: Parity, DIM, etc

Positions: February 2020-

DIM = how many days has the cow been milked since its latest calf was born

... and transmission of bacteria associated with mastitis




*mastitis = udder infection*

### Swedish Farm

Bacterial samples taken every second week for 5 months (June-October 2020)

- Milk samples from all cows
- Environmental samples from different parts of the barn
- Strain typing of bacteria

### Dutch Farm



**RI  
SE**

**Anna Skarin** professor, SLU

**Keni Ren** postdoc, SLU

**Hector Marina** postdoc, SLU

**Ida Hansson** PhD student, SLU

**Freddy Fikse** Växa Sverige



**Svenja Woudstra** PhD student, University of Copenhagen

**Per Peetz Nielsen** senior researcher, RISE

**Carsten Kirkeby** senior researcher, University of Copenhagen

**Volker Krömker** professor, University of Copenhagen

**Maya Gussmann** assistant professor, University of Copenhagen

**Moudud Alam** associate professor, Dalarna University

**Lisa Beste** communication officer, SLU

## What do we want to achieve?

Focus on the social environment of cows in free-stall dairy farms

- to improve animal welfare and production
- to predict transmission of diseases (mastitis) using monitored movement and interactions between cows

## Deliverables

### To develop

- tools for summarizing animal movement and social interactions in dairy farms.
- decision-support tools for minimizing disease transmission within dairy farms.
- methodology for breeding on indirect genetic effects.

# Published papers on real-time location data and social interactions

Marina, Ren, Hansson, Fikse, Nielsen, Rönnegård (2023) **New insight into social relationships in dairy cows, and how time of birth, parity and relatedness affect spatial interactions later in life.** Journal of Dairy Science (accepted)

Hansson, I., Silvera, A., Ren, K., Woudstra, S., Skarin, A., Fikse, W. F., Nielsen, P. P., & Rönnegård, L. (2023). **Cow characteristics associated with the variation in number of contacts between dairy cows.** Journal of Dairy Science.

Ren, K., Alam, M., Nielsen, P. P., Gussmann, M. K., & Rönnegård, L. (2022). **Interpolation methods to improve data quality of indoor positioning data for dairy cattle.** Frontiers in Animal Science

Ren, K., Nielsen, P. P., Alam, M., & Rönnegård, L. (2021). **Where do we find missing data in a commercial real-time location system? Evidence from 2 dairy farms.** JDS Communications.

Churakov, M., Silvera, A. M., Gussmann, M., & Nielsen, P. P. (2021). **Parity and days in milk affect cubicle occupancy in dairy cows.** Applied Animal Behaviour Science



Contents lists available at ScienceDirect

## Applied Animal Behaviour Science

journal homepage: [www.elsevier.com/locate/applanim](http://www.elsevier.com/locate/applanim)



# Parity and days in milk affect cubicle occupancy in dairy cows

Mikhail Churakov<sup>a,1</sup>, Anna Maria Silvera<sup>b,1</sup>, Maya Gussmann<sup>c</sup>, Per Peetz Nielsen<sup>d,\*</sup>

<sup>a</sup> Department of Animal Nutrition and Management, Swedish University of Agricultural Sciences, Box 7024, SE-750 07 Uppsala, Sweden

<sup>b</sup> Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Box 7023, SE-750 07 Uppsala, Sweden

<sup>c</sup> Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, 1870 Frederiksberg, Denmark

<sup>d</sup> RISE Research Institute of Sweden, Division of Bioeconomy and health, Department of Agriculture and Food, RISE Ideon, SE-223 70 Lund, Sweden

## ARTICLE INFO

### Keywords:

Real-time location system

Lying time


Dairy cattle

## ABSTRACT


Modern dairy cattle farms are usually equipped with cubicle systems to provide cows with comfortable conditions for lying down and resting. Cows are free to choose any cubicle they want, but in reality, they do not distribute themselves uniformly throughout the barn. There are many factors that affect where a cow lies down, such as hierarchy of a cow, access to resources, cow traffic nearby, etc. In this study, we used real-time location system data from two commercial farms to examine patterns of cubicle occupancy in relation to parity and lactation stage. We summarized cubicle occupancy over several days and compared different areas of the barn. Our findings suggest that, in general, there was a higher occupancy of cubicles close to the feeding areas. High parity cows lay down more frequently in cubicles close to the milking area as opposed to first lactation cows that tend to occupy less busy areas of the barn. The overall conclusion is that cubicle occupancy is not uniform throughout the barn, and patterns related to parity and DIM are seen. This information can be important for future studies on spread of diseases and for management purposes.



## Swedish farm



## Dutch farm






## Where do we find missing data in a commercial real-time location system? Evidence from 2 dairy farms

Keni Ren,<sup>1\*</sup>  Per Peetz Nielsen,<sup>2</sup>  Moudud Alam,<sup>3</sup>  and Lars Rönnegård<sup>1,3</sup> 

**Abstract:** Real-time indoor positioning using ultra-wideband devices provides an opportunity for modern dairy farms to monitor the behavior of individual cows; however, missing data from these devices hinders reliable continuous monitoring and analysis of animal movement and social behavior. The objective of this study was to examine the data quality, in terms of missing data, in one commercially available ultra-wideband-based real-time location system for dairy cows. The focus was on detecting major obstacles, or sections, inside open freestall barns that resulted in increased levels of missing data. The study was conducted on 2 dairy farms with an existing commercial real-time location system. Position data were recorded for 6 full days from 69 cows on farm 1 and from 59 cows on farm 2. These data were used in subsequent analyses to determine the locations within the dairy barns where position data were missing for individual cows. The proportions of missing data were found to be evenly distributed within the 2 barns after fitting a linear mixed model with spatial smoothing to logit-transformed proportions (mean = 18% vs. 4% missing data for farm 1 and farm 2, respectively), with the exception of larger proportions of missing data along one of the walls on both farms. On farm 1, the variation between individual tags was large (range: 9–49%) compared with farm 2 (range: 12–38%). This greater individual variation of proportions of missing data indicates a potential problem with the individual tag, such as a battery malfunction or tag placement issue. Further research is needed to guide researchers in identifying problems relating to data capture problems in real-time monitoring systems on dairy farms. This is especially important when undertaking detailed analyses of animal movement and social interactions between animals.




**Figure 2.** Fitted proportion of lost positions (>5 s) from the ultra-wideband system within grid squares (approximate size 4 m × 1.6 m), on (a) farm 1 and (b) farm 2. Grid squares with too few observations (gray) were excluded from the analysis. The insemination area is a closed bed area for inseminating the cows.



# Interpolation Methods to Improve Data Quality of Indoor Positioning Data for Dairy Cattle

**Keni Ren<sup>1\*</sup>, Moudud Alam<sup>2</sup>, Per Peetz Nielsen<sup>3</sup>, Maya Gussmann<sup>4</sup> and Lars Rönnegård<sup>1,2</sup>**

<sup>1</sup> Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Uppsala, Sweden, <sup>2</sup> School of Information and Engineering, Statistics, Dalarna University, Falun, Sweden, <sup>3</sup> Department of Agriculture and Food, RISE Research Institutes of Sweden (RISE), Lund, Sweden, <sup>4</sup> Department of Veterinary and Animal Sciences, University of Copenhagen, Frederiksberg, Denmark



Pre = Previous last observed position

Lin = Linear interpolation

Spl = Spline interpolation

Mak = Modified Akima interpolation



**J. Dairy Sci. 106:2685–2699**

**<https://doi.org/10.3168/jds.2022-21915>**

© 2023, The Authors. Published by Elsevier Inc. and Fass Inc. on behalf of the American Dairy Science Association®.  
This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## **Cow characteristics associated with the variation in number of contacts between dairy cows**

**I. Hansson,<sup>1</sup>  A. Silvera,<sup>1</sup> K. Ren,<sup>1</sup>  S. Woudstra,<sup>2</sup>  A. Skarin,<sup>3</sup>  W. F. Fikse,<sup>4</sup>  P. P. Nielsen,<sup>5</sup>   
and L. Rönnegård<sup>1,6,7\*</sup> **

<sup>1</sup>Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Box 7023, SE-750 07 Uppsala, Sweden

<sup>2</sup>Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, 1870 Frederiksberg, Denmark

<sup>3</sup>Department of Animal Nutrition and Management, Swedish University of Agricultural Sciences, Box 7024, SE-750 07 Uppsala, Sweden

<sup>4</sup>Växa, Swedish University of Agricultural Sciences, Ulls väg 26, SE-756 51 Uppsala, Sweden

<sup>5</sup>RISE Research Institute of Sweden, Division of Bioeconomy and Health, Department of Agriculture and Food, RISE Ideon, SE-223 70 Lund, Sweden

<sup>6</sup>School of Technology and Business Studies, Dalarna University, SE-791 88 Falun, Sweden

<sup>7</sup>The Beijer Laboratory for Animal Science, Swedish University of Agricultural Sciences, Box 7024, SE-750 07 Uppsala, Sweden

### Definition of contact used in the paper

- Less than 2.5 m (one body length) apart
- Duration >10 minutes in total during a day, otherwise removed to reduce noise.

### Response variable studied

The average number of contacts for each cow was calculated as the **instantaneous number of individuals within proximity of a cow** at any time during the day.

## Traits affecting the number of contacts a cow has during a day

Trait	Feeding Area			Resting Area		
	Farm A		Farm B (n = 201)	Farm A		Farm B (n = 201)
	G1 (n= 83)	G2 (n = 80)		G1 (n = 83)	G2 (n = 80)	
Date	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001
Parity	< 0.001	0.685	0.009	0.773	0.999	< 0.001
Lactation stage	0.013	0.029	< 0.001	0.089	0.646	0.366
Breed	0.468	0.008		0.513	0.815	
Pregnancy status	0.319	0.266	0.688	< 0.001	0.583	0.725
Estrus	0.002		0.069	0.002		0.240
Udder Health	0.967	0.264		0.017	0.936	
Claw Health	0.109	0.327		0.454	0.008	



## Traits affecting the number of contacts a cow has during a day

Trait	Feeding Area			Resting Area		
	Farm A		Farm B (n = 201)	Farm A		Farm B (n = 201)
	G1 (n= 83)	G2 (n = 80)		G1 (n = 83)	G2 (n = 80)	
Date	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001
Parity	< 0.001	0.685	0.009	0.773	0.999	< 0.001
Lactation stage	0.013	0.029	< 0.001	0.089	0.646	0.366
Breed	0.468	0.008		0.513	0.815	
Pregnancy status	0.319	0.266	0.688	< 0.001	0.583	0.725
Estrus	0.002		0.069	0.002		0.240
Udder Health	0.967	0.264		0.017	0.936	
Claw Health	0.109	0.327		0.454	0.008	

SOCIAL NETWORK ANALYSIS OF DAIRY COW INTERACTIONS

**New insight into social relationships in dairy cows, and how time of birth, parity and relatedness affect spatial interactions later in life.**

**H. Marina,<sup>1\*</sup> K. Ren,<sup>1</sup> I. Hansson,<sup>1</sup> F. Fikse,<sup>2</sup> P. P. Nielsen,<sup>3</sup> and L. Rönnegård<sup>1,4,5</sup>**

<sup>1</sup> Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Box 7023, SE-750 07 Uppsala, Sweden.

<sup>2</sup> Växa, Swedish University of Agricultural Sciences, Ulls väg 26, SE-756 51 Uppsala, Sweden.

<sup>3</sup> RISE Research Institute of Sweden, Division of Bioeconomy and Health, Department of Agriculture and Food, RISE Ideon, SE-223 70 Lund, Sweden.

<sup>4</sup> School of Technology and Business Studies, Dalarna University, SE-791 88 Falun, Sweden.

<sup>5</sup> The Beijer Laboratory for Animal Science, Swedish University of Agricultural Sciences, Box 7024, SE-750 07 Uppsala, Sweden.

### Interpretive summary

Understanding social interactions between cows is crucial to improving animal welfare and production. Real-time location systems provide an opportunity to study dyadic social contacts inside dairy free-stall barns. We investigated the impact of different cow characteristics on the likelihood of the formation and persistence of social contacts in dairy cattle. **Our main finding was that cows born within seven days of each other had more consistent interactions, a kind of “kindergarten effect”, as well as cows of the same parity or related by pedigree.**

## Published papers on disease transmission

Woudstra, S., Wente, N., Zhang, Y., Leimbach, S., Gussmann, M. K., Kirkeby, C., & Krömker, V. (2023). **Strain diversity and infection durations of Staphylococcus spp. and Streptococcus spp. causing intramammary infections in dairy cows.** Journal of Dairy Science.

Woudstra, S., Lücken, A., Wente, N., Zhang, Y., Leimbach, S., Gussmann, M. K., Kirkeby, C., & Krömker, V. (2023). **Reservoirs of Corynebacterium spp. in the environment of dairy Cows.** Pathogens

Lücken, A., Woudstra, S., Wente, N., Zhang, Y., & Krömker, V. (2022). **Intramammary infections with Corynebacterium spp. in bovine lactating udder quarters.** Plos One.

... and transmission of bacteria associated with mastitis



*mastitis = udder infection*

### Swedish Farm

Bacterial samples taken every second week for 5 months (June-October 2020)

- Milk samples from all cows
- Environmental samples from different parts of the barn
- Strain typing of bacteria

### Dutch Farm

QR to our project web page



We also put our code on GitHub  
**<https://github.com/CSI-DT>**