

ClimDatDownloadR: Accessing Climate Data Repositories for Modelling

Helge Jentsch^{1,2*}, Johannes Weidinger^{2*}, Melanie Werner^{2*}, and
Maria Bobrowski^{2*}

¹ Climate Geography (CliG), Albert-Ludwigs-Universität Freiburg, Freiburg i. Breisgau, Germany ² Centrum for Earth System Research and Sustainability (CEN), University of Hamburg, Hamburg, Germany ^{*} Corresponding author ^{*} These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- Review
- Repository
- Archive

Editor: Open Journals

Reviewers:

- @openjournals

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

In partnership with



AMERICAN
ASTRONOMICAL
SOCIETY

This article and software are linked with research article DOI [10.3847/xxxxx](https://doi.org/10.3847/xxxxx) <- update this with the DOI from AAS once you know it., published in the Astrophysical Journal <- The name of the AAS journal..

Summary

Systematical accessing, downloading, and pre-processing climatological data from CHELSA (Karger et al., 2017, 2021; Karger et al., 2018) and WorldClim (Fick & Hijmans, 2017; Hijmans et al., 2005) remains a challenge in different environmental disciplines like Species Distribution Modelling (SDM) and climate studies. This package provides a set of functions that allow easy access and customized selection of climate data sets. Besides downloading the raw data, also functionalities to complete pre-processing steps like clipping, rescaling, and file management are available. The applications of the package range from one-time-use to implementing the functions in automatic processing of scientific workflows.

Statement of need

The climatology datasets CHELSA and WorldClim contribute as crucial data bases for studies in various scientific fields. Primarily used in studies with focus on ecology (~4,200 publications¹), environmental sciences (>2,200 publications), and biodiversity conservation (>1,600 publications), usages extend to a wide variety of scientific disciplines. The main usage of the datasets, however, lies in Species Distribution Modelling (SDM) and Ecological Niche Modelling (ENM). Their free availability and frequent citation in widely referenced papers on SDM and ENM strategies (e.g., Randin et al., 2020; Zurell et al., 2020) have contributed to their widespread adoption, facilitating comparability between modelling studies at different spatial and temporal scales.

The high resolution global climatological datasets (30 arc-sec. ~ 1km) include downscaled and bias-corrected data from 30-year time-periods, providing always monthly mean, minimum, and maximum values of temperature and monthly precipitation sums for analysis². Additionally, 19 bioclimatic parameters are accessible, which enable conclusions about seasonality.

Since their initial releases in 2018 (CHELSA V1.2), the CHELSA (Karger et al., 2017; Karger et al., 2018) datasets were cited in more than 2,800 peer reviewed papers, indexed on the Web of Science (source, Aug. 2025). The latest release of WorldClim 2 in 2017 (Fick & Hijmans, 2017) was cited more than 10,600 times (source, Aug. 2025).

¹Following the Web of Science Categories, citations of Karger et al. (2018) (Data from CHELSA 2.1) had 1,155 citations in the field of Ecology. The WorldClim 2 data (Fick & Hijmans, 2017) has 3,044 citations in the same Web of Science category. Both numbers are of the date 17.05.2025. The "Web of Science Categories are assigned at the journal level", meaning the publishing journal defines the category (source).

²Function Chelsa.timeseries.download supports also the download of potential evapotranspiration (PET) from CHELSA 2.1 (Karger et al., 2018)

CHELSA and WorldClim datasets are commonly utilized in models predicting the potential past, current, and future distribution of species, particularly in studies on monitoring distribution shifts under climate change (e.g., [Bobrowski et al., 2017](#); [Twala et al., 2023](#); [Werner et al., 2025](#)), tracking endangered species and planning conservation strategies (e.g., [Franklin, 2013](#); [Muscatello et al., 2021](#)), assessing the spread of invasive species ([Srivastava et al., 2019](#)), and management strategies in forestry and agriculture (e.g., [Agbezuge & Balakrishnan, 2024](#); [Pecchi et al., 2019](#)).

Recent studies have also assessed the performance of these datasets in SDM/ENM approaches, highlighting their respective strengths and limitations (e.g., [Bobrowski, Weidinger, & Schickhoff, 2021](#); [Bobrowski, Weidinger, Schwab, et al., 2021](#); [Bobrowski & Schickhoff, 2017](#); [Datta et al., 2020](#); [Rodríguez-Rey & Jiménez-Valverde, 2024](#)). Given that dataset performance may vary depending on the research scope, it is recommended to test multiple datasets to ensure their suitability for the research target and region.

For these applications, `ClimDatDownloadR` offers key advantages by enabling efficient retrieval from both dataset providers and pre-processing steps such as partial selection of parameters, months, and bioclimatic parameters, temporal subsets of timeseries, customized extent, and included file management as well as an output of the provider's respective citation file. In addition to time-saving aspects, the storage usage and management played a key role in the development of the `ClimDatDownloadR`.

The implemented data management creates a hierarchical, clear, and reproducible data structure for analyses during the processing. Downloaded data can be kept as is, deleted, or packed in a zip-archive file. All of raised *ease-of-use* add-ons contribute to the primary goal of `ClimDatDownloadR` to enable more scientists and other users or organisations to download and pre-process CHELSA and WorldClim data to gain more experience in geodata handling and applications.

Since the official release in 2023, the use of `ClimDatDownloadR` steadily increased ([Bobrowski, Weidinger, & Schickhoff, 2021](#); [Chen et al., 2025](#); [Costa-Saura et al., 2025](#); [Maitner et al., 2023](#); [Santi et al., 2024](#); [Twala et al., 2023](#); [Werner et al., 2025](#)). Further, the need of having software for downloading and pre-processing of freely available data is shown by the steady stream of interested visitors on [ResearchGate \(3,399 unique visits, 04.08.2025\)](#), [Zenodo \(>1000 views, > 150 downloads\)](#) ([Jentsch et al., 2023](#)), and citations in peer-reviewed papers.

The package implements the datasets CHELSA V1.2, V2.1, WorldClim V1.4, and V2.1. More specifically the CHELSA Climatologies, Timeseries, CRU Timeseries (CHELSAcruts), and WorldClim Histclim datasets for present data. For past data, the CHELSA PIMP3 data from CHELSA V1.2 is also available. For future data, both CHELSA and WorldClim provide datasets incorporating various CMIP 5 and 6 global circulation models with various emission scenarios and reference periods. An overview is provided below in [table 1](#) and [2](#).

Usage

The `ClimDatDownloadR` enables (1) selection of dataset and version (e.g., CHELSA 2.1), (2) selection of available variable-parameters (e.g., bioclimatic variables and precipitation sums), (3) pre-processing steps like clipping, and (4) file management options. After describing the complete extent of the workflow a working example is shown. While this introduces an overview on how to approach the usage of the download functions of the `ClimDatDownloadR`, we employ users to consult documentations on the specific functions to gain further insights on e.g., variable or model availability or additional function parameters. An up-to-date version of the manual can be accessed on the [GitHub repository](#).

(1) Start selecting the dataset with `provider.dataset.download()`. The *provider* are `Chelsa` or `WorldClim`. The following *dataset* completes the function name and specifies

the accessed dataset. Function names and accessed datasets are referenced in the leading columns of [table 1 and 2](#). The version can be specified with `version.var`.

(2) The availability of parameters differs between datasets, despite every dataset providing data on monthly temperature means, minima, and maxima, as well as precipitation sums. Bioclimatic variables are also available for all climatologies. Here, the function parameters `month.var` and `bio.var` can be used to specify a subset from the dataset. For past (last glacial maximum) and future (CMIP 5/6) model climatologies a set of function parameters to specify models (`model.var`), scenarios (`emission.scenario.var`), and time interval (`time.interval.var`) can be used. The WorldClim datasets come in different spatial resolutions (10/5/2.5 arc-min., 30 arc-sec.), while CHELSA data is always in 30 arc-sec. resolution. Therefore, `WorldClim.*.download()`-functions have an additional resolution function parameter.

(3) The pre-processing steps consist of two levels. While the processing of the provided data to interpretable values cannot be addressed by the user, clipping needs to be enabled with the eponymous function parameter and specified further with parameters like `clip.shapefile` or `clip.extent` with `buffer` as additional parameter.

(4) File-management addresses the conversion to different file formats, the decluttering of the working directory, and a citation file that includes the providers' relevant publications. Currently, the functions parameter `convert.files.to.asc` and `stacking.data` convert the data to ASCII raster and netCDF format, respectively. For a storage efficient way, the user can decide to delete the downloaded datasets (`delete.raw.data`), save them in a zip-archive file (`combine.raw.zip` (CHELSA), `keep.raw.zip` (WorldClim)), or keep them in the directory as-is. The function parameter `save.bib.file` adds a citation file of the publications from the data provider to the working directory for convenience.

In the following, the basic usage of the download functions are demonstrated by applying an example call of the `Chelsa.Clim.download()` function, aiming at downloading CHELSA V2.1 bioclimatic and precipitation data, with clipping to the European continent's extent, and file management tools:

```
ClimDatDownloadR::Chelsa.Clim.Download(
  save.location = "./",
  version.var = "2.1",
  parameter = c("bio", "prec"),
  month.var = c(4:9),
  clipping = TRUE,
  clip.extent = c(-5,25,40,62),
  buffer = 0,
  clip.shapefile = NULL,
  convert.files.to.asc = FALSE,
  stacking.data = FALSE,
  combine.raw.zip = TRUE,
  delete.raw.data = TRUE,
  save.download.table = TRUE,
  save.bib.file = TRUE
)
```

111 Tables

112 Table 1: Overview of the CHELSA function names, the respective accessible dataset, versions,
113 spatial and temporal resolutions, as well as the reference periods, and if applicable emission
114 scenarios and models. Additionally the citations and links to the documentations of the
115 datasets are listed.

Table with 11 columns: Function Name, Accessed Dataset, Version, Reference Period, Emission Scenario, Models, Spatial Resolution, Temporal Resolution, Parameters, Citation, Documentation. It lists five rows of data for different CHELSA datasets including climatology, timeseries, CRU, CMIP6, and PMIP3.

117 **Table 2:** Overview of the WorldClim functions. Similar to [table 1](#).

Function Name	Accessed Dataset	Version	Reference Period	Emission Scenario	Models	Spatial Resolution	Temporal Resolution	Parameters	Citation	Documentation
"WorldClim.HistClim.download"	WorldClim historical climate data	1.4, 2.1	1.4: 1960-1990; 2.1: 1970-2000	N/A	N/A	10 minutes to 30 arc-seconds	Monthly	Bioclim, Param. (https://www.worldclim.org/data/bioclim.html), Precip. (sum), Temp. (mean, min, max), Solar Radiation, Wind Speed, Water Vapor Pressure	Hijmans et al. (2005), Fick & Hijmans (2017)	Version 1.4: Methods (https://www.worldclim.org/data/v1.4/methods.html), Data Formats (https://www.worldclim.org/data/v1.4/formats.html); Version 2.1: Download Page (https://www.worldclim.org/data/worldclim21.html)
"WorldClim.CMIP_6.download"	WorldClim CMIP6 dataset	2.1	2021-2040, 2041-2060, 2061-2080, and 2081-2100	SSP scenarios 1 (SSP126), 2 (SSP245), 3 (SSP370), and 5 (SSP585)	Multiple CMIP5 Models, ref. 10- (https://www.worldclim.org/data/cmip5/cmip6_clim10m.html), 5- (https://www.worldclim.org/data/cmip6/cmip6_clim5m.html), 2.5- (https://www.worldclim.org/data/cmip6/cmip6_clim2.5m.html)minute s, and 30- (https://www.worldclim.org/data/cmip6/cmip6_clim30s.html)second download sites	10 minutes to 30 arc-seconds	Monthly	Bioclim, Param. (https://www.worldclim.org/data/bioclim.html), Precip. (sum), Temp. (mean, min, max)	Fick & Hijmans (2017)	Downscaling methods (https://www.worldclim.org/data/downscaling.html)
"WorldClim.CMIP_5.download"	WorldClim CMIP5 dataset	1.4	2041-2060, 2061-2080	rcp26, rcp45, rcp60, and rcp85	Multiple CMIP5 Models, ref. 10- (https://www.worldclim.org/data/v1.4/cmip5_10m.html), 5- (https://www.worldclim.org/data/v1.4/cmip5_5m.html), 2.5- (https://www.worldclim.org/data/v1.4/cmip5_2.5m.html)minutes, and 30- (https://www.worldclim.org/data/v1.4/cmip5_30s.html)second download sites	10 minutes to 30 arc-seconds	Monthly	Bioclim, Param. (https://www.worldclim.org/data/bioclim.html), Precip. (sum), Temp. (mean, min, max)	Hijmans et al. (2005), Fick & Hijmans (2017)	Downscaling methods (https://www.worldclim.org/data/downscaling.html)

118

119 Acknowledgements

120 We acknowledge the thorough testing efforts by the co-authors and Nadine Kaul, and the
121 open access efforts by the Eidg. Forschungsanstalt für Wald, Schnee und Landschaft WSL
122 [Chelsa; Karger et al. (2017)] and University of California, Davis [WorldClim; Fick & Hijmans
123 (2017)] working groups. Also we want to acknowledge the wide user group that implemented
124 the ClimDatDownloadR in their unpublished scientific work or motivated us to implement new
125 datasets or update broken links. In addition we want to thank the developers of the various
126 R-packages for their contributions that made this package possible. In alphabetical order these
127 were: curl (Ooms, 2025), httr (Wickham, 2023a), ncdf4 (Pierce, 2024), RCurl (Temple
128 Lang, 2025), RefManageR (McLean, 2014, 2017), sf (E. Pebesma, 2018; E. Pebesma & Bivand,
129 2023), stringr (Wickham, 2023b), sp (Bivand et al., 2013; E. J. Pebesma & Bivand, 2005),
130 terra (Hijmans, 2025), and the development team of R (R Core Team, 2025).

131

References

132 Agbezuge, E. Y., & Balakrishnan, P. (2024). Application of species distribution modelling in
133 agriculture: A review. In A. Swaroop, Z. Polkowski, S. D. Correia, & B. Virdee (Eds.),
134 *Proceedings of data analytics and management* (pp. 173–188). Springer Nature Singapore.
135 ISBN: 978-981-99-6547-2

136 Bivand, R. S., Pebesma, E., & Gomez-Rubio, V. (2013). *Applied spatial data analysis with R*,
137 *Second edition*. Springer, NY. <https://asdar-book.org/>

138 Bobrowski, M., Gerlitz, L., & Schickhoff, U. (2017). Modelling the potential distribution
139 of *Betula utilis* in the Himalaya. *Global Ecology and Conservation*, 11, 69–83. <https://doi.org/10.1016/j.gecco.2017.05.005>

- 140 [//doi.org/10.1016/j.gecco.2017.04.003](https://doi.org/10.1016/j.gecco.2017.04.003)
- 141 Bobrowski, M., & Schickhoff, U. (2017). Why input matters: Selection of climate data sets for
142 modelling the potential distribution of a treeline species in the Himalayan region. *Ecological*
143 *Modelling*, 359, 92–102. <https://doi.org/10.1016/j.ecolmodel.2017.05.021>
- 144 Bobrowski, M., Weidinger, J., & Schickhoff, U. (2021). Is new always better? Frontiers in
145 global climate datasets for modeling treeline species in the himalayas. *Atmosphere*, 12(543).
146 <https://doi.org/10.3390/atmos12050543>
- 147 Bobrowski, M., Weidinger, J., Schwab, N., & Schickhoff, U. (2021). Searching for ecology
148 in species distribution models in the Himalayas. *Ecological Modelling*, 458, 109693.
149 <https://doi.org/10.1016/j.ecolmodel.2021.109693>
- 150 Chen, S. H., Stevens, L., Gooden, B., Rafter, M. A., Knerr, N., Thrall, P. H., Ord, L., &
151 Schmidt-Lebuhn, A. N. (2025). PhyloControl: A phylogeny visualisation platform for risk
152 analysis in weed biological control. *Biological Control*, 105859. <https://doi.org/10.1016/j.biocontrol.2025.105859>
- 153
- 154 Costa-Saura, J. M., Midolo, G., Ricotta, C., Baudena, M., Calfapietra, C., Elia, M., Fiorucci,
155 P., Mereu, S., Sirca, C., Spano, D., Vivaldo, G., & Ottaviani, G. (2025). Are trait
156 responses of tree species across pyroregions indicative of fire-modulated plant functional
157 strategies? *Perspectives in Plant Ecology, Evolution and Systematics*, 67, 125867. <https://doi.org/10.1016/j.ppees.2025.125867>
- 158
- 159 Datta, A., Schweiger, O., & Kühn, I. (2020). Origin of climatic data can determine the
160 transferability of species distribution models. *NeoBiota : Advancing Research on Alien*
161 *Species and Biological Invasions*, 59, 61–76. <https://doi.org/10.3897/neobiota.59.36299>
- 162 Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate
163 surfaces for global land areas. *International Journal of Climatology*, 37(12), 4302–4315.
164 <https://doi.org/10.1002/joc.5086>
- 165 Franklin, J. (2013). Species distribution models in conservation biogeography: Developments
166 and challenges. *Diversity and Distributions*, 19(10), 1217–1223. <https://doi.org/10.1111/ddi.12125>
- 167
- 168 Hijmans, R. J. (2025). *Terra: Spatial data analysis* [Manual]. <https://CRAN.R-project.org/package=terra>
- 169
- 170 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high
171 resolution interpolated climate surfaces for global land areas. *International Journal of*
172 *Climatology*, 25(15), 1965–1978. <https://doi.org/10.1002/joc.1276>
- 173 Jentsch, H., Weidinger, J., & Bobrowski, M. (2023). *ClimDatDownloadR: Downloads Climate*
174 *Data from Chelsa and WorldClim*. Zenodo. <https://doi.org/10.5281/ZENODO.7924342>
- 175 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann,
176 N. E., Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's
177 land surface areas. *Scientific Data*, 4(1), 170122. <https://doi.org/10.1038/sdata.2017.122>
- 178 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann,
179 N. E., Linder, H. P., & Kessler, M. (2018). *Data from: Climatologies at high resolution for*
180 *the earth's land surface areas*. Dryad. <https://doi.org/10.5061/DRYAD.KD1D4>
- 181 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann,
182 N. E., Linder, H. P., & Kessler, M. (2021). *Climatologies at high resolution for the earth's*
183 *land surface areas*. EnviDat. <https://doi.org/10.16904/envi.dat.228>
- 184 Maitner, B., Gallagher, R., Svenning, J.-C., Tietje, M., Wenk, E. H., & Eiserhardt, W. L.
185 (2023). A global assessment of the Raunkiaeran shortfall in plants: Geographic biases in
186 our knowledge of plant traits. *New Phytologist*, 240(4), 1345–1354. <https://doi.org/10.1111/nph.15888>

- 1111/nph.18999
- McLean, M. W. (2014). *Straightforward bibliography management in R using the RefManager package* [Manual]. <https://arxiv.org/abs/1403.2036>
- McLean, M. W. (2017). RefManageR: Import and manage BibTeX and BibLaTeX references in R. *The Journal of Open Source Software*. <https://doi.org/10.21105/joss.00338>
- Muscatello, A., Elith, J., & Kujala, H. (2021). How decisions about fitting species distribution models affect conservation outcomes. *Conservation Biology : The Journal of the Society for Conservation Biology*, 35(4), 1309–1320. <https://doi.org/10.1111/cobi.13669>
- Ooms, J. (2025). *Curl: A modern and flexible web client for R* [Manual]. <https://CRAN.R-project.org/package=curl>
- Pebesma, E. (2018). Simple features for R: Standardized support for spatial vector data. *The R Journal*, 10(1), 439–446. <https://doi.org/10.32614/RJ-2018-009>
- Pebesma, E. J., & Bivand, R. S. (2005). Classes and methods for spatial data in R. *R News*, 5(2), 9–13. <https://CRAN.R-project.org/doc/Rnews/>
- Pebesma, E., & Bivand, R. (2023). *Spatial data science: With applications in R*. Chapman and Hall/CRC. <https://doi.org/10.1201/9780429459016>
- Pecchi, M., Marchi, M., Burton, V., Giannetti, F., Moriondo, M., Bernetti, I., Bindi, M., & Chirici, G. (2019). Species distribution modelling to support forest management. A literature review. *Ecological Modelling*, 411, 108817. <https://doi.org/10.1016/j.ecolmodel.2019.108817>
- Pierce, D. (2024). *Ncdf4: Interface to unidata netCDF (version 4 or earlier) format data files* [Manual]. <https://doi.org/10.32614/cran.package.ncdf4>
- R Core Team. (2025). *R: A language and environment for statistical computing* [Manual]. R Foundation for Statistical Computing. <https://doi.org/10.32614/r.manuals>
- Randin, C. F., Ashcroft, M. B., Bolliger, J., Cavender-Bares, J., Coops, N. C., Dullinger, S., Dirnböck, T., Eckert, S., Ellis, E., Fernández, N., Giuliani, G., Guisan, A., Jetz, W., Joost, S., Karger, D. N., Lembrechts, J., Lenoir, J., Luoto, M., Morin, X., ... Payne, D. (2020). Monitoring biodiversity in the Anthropocene using remote sensing in species distribution models. *Remote Sensing of Environment*, 239, 111626. <https://doi.org/10.1016/j.rse.2019.111626>
- Rodríguez-Rey, M., & Jiménez-Valverde, A. (2024). Differing sensitivity of species distribution modelling algorithms to climate data source. *Ecological Informatics*, 79, 102387. <https://doi.org/10.1016/j.ecoinf.2023.102387>
- Santi, F., Testolin, R., Zannini, P., Di Musciano, M., Micci, V., Ricci, L., Guarino, R., Bacchetta, G., Fernández-Palacios, J. M., Fois, M., Kougiumoutzis, K., Kunt, K. B., Lucchi, F., Médail, F., Nikolić, T., Otto, R., Pasta, S., Panitsa, M., Proios, K., ... Chiarucci, A. (2024). MEDIS—A comprehensive spatial database on Mediterranean islands for biogeographical and evolutionary research. *Global Ecology and Biogeography*, 33(8), e13855. <https://doi.org/10.1111/geb.13855>
- Srivastava, V., Lafond, V., & Griess, V. C. (2019). Species distribution models (SDM): Applications, benefits and challenges in invasive species management. *CABI Reviews*, 1–13. <https://doi.org/10.1079/PAVSNNR201914020>
- Temple Lang, D. (2025). *RCurl: General network (HTTP/FTP/...) client interface for R* [Manual]. <https://doi.org/10.32614/CRAN.package.RCurl>
- Twala, T. C., Fisher, J. T., & Glennon, K. L. (2023). Projecting Podocarpaceae response to climate change: We are not out of the woods yet. *AoB PLANTS*, 15(4), plad034. <https://doi.org/10.1093/aobpla/plad034>

- 234 Werner, M., Böhner, J., Oldeland, J., Schickhoff, U., Weidinger, J., & Bobrowski, M. (2025).
235 Treeline Species Distribution Under Climate Change: Modelling the Current and Future
236 Range of *Nothofagus pumilio* in the Southern Andes. *Forests*, 16(8), 1211. <https://doi.org/10.3390/f16081211>
237
- 238 Wickham, H. (2023a). *Httr: Tools for working with urls and HTTP* [Manual]. <https://doi.org/10.32614/cran.package.httr>
239
- 240 Wickham, H. (2023b). *Stringr: Simple, consistent wrappers for common string operations*
241 [Manual]. <https://doi.org/10.32614/cran.package.stringr>
- 242 Zurell, D., Franklin, J., König, C., Bouchet, P. J., Dormann, C. F., Elith, J., Fandos, G.,
243 Feng, X., Guillera-Arroita, G., Guisan, A., Lahoz-Monfort, J. J., Leitão, P. J., Park, D.
244 S., Peterson, A. T., Rapacciuolo, G., Schmatz, D. R., Schröder, B., Serra-Díaz, J. M.,
245 Thuiller, W., ... Merow, C. (2020). A standard protocol for reporting species distribution
246 models. *Ecography*, 43(9), 1261–1277. <https://doi.org/10.1111/ecog.04960>

DRAFT