

3. Prevention or Delay of Diabetes

Diabetes Care Volume 49, Supplement 1, January 2026

American Diabetes Association

Professional Practice Committee for Diabetes*

Comorbidities: Standards of Care in Diabetes—2026

Diabetes Care 2026;49(Suppl. 1):S50–S60 | <https://doi.org/10.2337/dc26-S003>

The American Diabetes Association (ADA) “Standards of Care in Diabetes” includes the ADA’s current clinical practice recommendations and is intended to provide the components of diabetes care, general treatment goals and guidelines, and tools to evaluate quality of care. Members of the ADA Professional Practice Committee for Diabetes, an interprofessional expert committee, are responsible for updating the Standards of Care annually, or more frequently as warranted. For a detailed description of ADA standards, statements, and reports, as well as the evidence-grading system for ADA’s clinical practice recommendations and a full list of Professional Practice Committee members, please refer to Introduction and Methodology. Readers who wish to comment on the Standards of Care are invited to do so at professional.diabetes.org/SOC.

For guidelines related to screening for increased risk for type 1 diabetes, prediabetes and type 2 diabetes, and other forms of diabetes, please refer to section 2, “Diagnosis and Classification of Diabetes.” For guidelines related to screening, diagnosis, and management of diabetes in children and adolescents, please refer to section 14, “Children and Adolescents.”

Recommendations

1. In people with prediabetes, monitor for the development of diabetes at least annually; modify frequency of testing based on individual risk assessment. E 2. In people with presymptomatic type 1 diabetes, monitor for disease progression using A1C approximately every 6 months and 75-g oral glucose tolerance test (i.e., fasting and 2-h plasma glucose) annually; modify frequency of monitoring and consider augmenting with other glycemic assessment tools such as continuous glucose monitoring metrics based on individual risk assessment incorporating age, number and type of autoantibodies, and glycemic metrics. E

Screening for prediabetes (which refers to hyperglycemia preceding the diagnosis of type 2 diabetes only) and type 2 diabetes risk through an assessment of risk factors (Table 2.5) or with an assessment tool, such as the American Diabetes Association risk test, which can be used by either a layperson or a health care professional (diabetes.org/diabetes-risk-test), is recommended to guide whether to perform a diagnostic test for prediabetes (Table 2.2) and type 2 diabetes (Table 2.1) (see section 2, “Diagnosis and Classification of Diabetes”). Testing high-risk individuals for prediabetes is warranted because the laboratory assessment is safe and reasonable.

*A complete list of members of the American Diabetes Association Professional Practice Committee for Diabetes can be found at <https://doi.org/10.2337/dc26-SINT>.

Duality of interest information for each contributor is available at <https://doi.org/10.2337/dc26-SDIS>.

Suggested citation: American Diabetes Association Professional Practice Committee for Diabetes. 3. Prevention or delay of diabetes and associated comorbidities: Standards of Care in Diabetes—2026. Diabetes Care 2026;49(Suppl. 1):S50–S60 © 2025 by the American Diabetes Association.

Readers may use this work for educational, noncommercial purposes if properly cited and unaltered. The version of record may be linked at <https://diabetesjournals.org/care>, but ADA permission is required to post this work on any third-party site or platform. This publication and its contents may not be reproduced, distributed, or used for text or data mining, machine learning, or similar technologies without prior written permission. Requests to reuse, adapt, or distribute this work may be sent to permissions@diabetes.org. More information is available at <https://diabetesjournals.org/journals/pages/license>.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf Downloaded

diabetesjournals.org/care # Prevention or Delay of Diabetes

In cost, and early detection of hyperglycemia affords substantial time to intervene and delay or prevent the onset of type 2 diabetes and/or its long-term complications. Indeed, once identified, several effective therapeutic approaches exist that can delay type 2 diabetes in those with prediabetes with an A1C 5.7–6.4% (39–47 mmol/mol), impaired glucose tolerance (IGT) on 75-g oral glucose tolerance test (OGTT), or impaired fasting glucose (IFG). The utility of screening with A1C for prediabetes and diabetes may be limited in the presence of certain hemoglobinopathies and conditions that affect red blood cell turnover (Table 2.3). See section 2, “Diagnosis and Classification of Diabetes,” and section 6, “Glycemic Goals, Hypoglycemia, and Hyperglycemic Crises,” for additional details on the appropriate use and limitations of A1C testing.

3.4

Prescribe an evidence-based eating pattern (e.g., Mediterranean, low carbohydrate) to individuals with prediabetes to prevent type 2 diabetes. B

3.5

Offer diabetes prevention programs to adults at high risk for type 2 diabetes. A Diabetes prevention programs should be covered by third-party payors, and inconsistencies in access should be addressed. E

3.6

Based on individual preference, certified technology-assisted diabetes prevention programs through smartphones, web-based applications, and telehealth can be effective in preventing type 2 diabetes and should be considered. B

Three stages of type 1 diabetes have been defined, with symptomatic type 1 diabetes classified as stage 3 (Table 2.4). In individuals at risk for developing clinical type 1 diabetes, younger age of seroconversion (particularly under age 3 years), the total number of diabetes-related autoantibodies (1), and the development of autoantibodies against islet antigen 2 (IA-2) have all been associated with a more rapid progression to stage 3 type 1 diabetes. While continuous glucose monitoring can predict progression to stage 3 type 1 diabetes in children and adolescents with autoantibodies (2), OGTT-based metrics are more predictive of progression compared with continuous glucose monitoring (3). The decision to perform an OGTT may depend on such factors as eligibility and interest for stage-specific treatments, interest in participation in clinical research, OGTT availability, and the burden of testing. Consensus guidance provides expert recommendations on what should be monitored and how often in people with presymptomatic type 1 diabetes (4). Cost-effectiveness of population-based screening programs for type 1 diabetes has not been established (5,6), but such screening is being implemented outside the U.S. (7).

The Diabetes Prevention Program

Several major randomized controlled trials, including the Diabetes Prevention Program (DPP) (8), the Finnish Diabetes Prevention Study (DPS) (9), and the Da Qing Diabetes Prevention Study (Da Qing study) (10), demonstrated that lifestyle/behavioral intervention with an individualized reduced-calorie meal plan is highly effective in preventing or delaying type 2 diabetes and improving other cardiometabolic risk factors such as blood pressure, lipids, and markers of inflammation (11). The strongest evidence for diabetes prevention in the U.S. comes from the DPP trial (8). The DPP demonstrated that intensive lifestyle intervention could reduce the risk of incident type 2 diabetes by 58% over 3 years. Follow-up of three large trials of lifestyle intervention for diabetes prevention showed sustained reduction in the risk of progression to type 2 diabetes: 39% reduction at 30 years in the Da Qing study (12), 43% reduction at 7 years in the Finnish DPS (9), and 34% reduction at 10 years (13), 27% reduction at 15 years (14), and 24% reduction at 21 years in the U.S. Diabetes Prevention Program Outcomes Study (DPPOS) (15,16).

The DPP lifestyle intervention was a goal-based intervention. A complete description of the DPP and how to access it is available online at coveragetoolkit.org/about-national-dpp/ndpp-overview/. Briefly, the DPP administered a 16-session structured core curriculum, completed within the first 6 months of the program, followed by an additional 6 months of a flexible maintenance program. The core curriculum included sessions on reducing calories, increasing physical activity, self-monitoring, maintaining healthy lifestyle behaviors (such as how to make healthy choices when eating out), and guidance on managing psychological, social, and motivational challenges (17).

Two major goals of the DPP were to achieve and maintain at least 7% weight loss and to obtain 150 min of moderate-intensity physical activity per week, such as brisk walking. Weight loss was the most important factor in reducing incident diabetes. Obtaining at least 150 min of physical activity per week, even without achieving the weight loss goal, reduced incidence of type 2 diabetes by 44% (18). Participants were encouraged to distribute physical activity throughout the week with a minimum frequency of three times per week and at least 10 min per session. A maximum of 75 min of strength training could be applied toward the total 150 min/week physical activity goal. The initial focus of the nutrition intervention was on reducing total fat intake rather than caloric restriction. However, the concepts of calorie balance and the need to restrict total calories and fat were also introduced. Overall, the DPP's lifestyle intervention was designed to produce a 700 kcal/week energy deficit (17).

The 7% weight loss goal was selected because it was feasible to achieve and maintain, hypothesized to reduce incident diabetes risk, and improved other cardiometabolic risk factors. Further analysis suggested greater benefit for diabetes prevention with at least 7–10% weight loss with lifestyle interventions (18). The recommended pace of weight loss was 1–2 lb/week.

While the DPP interventions were successful in preventing or delaying the onset of type 2 diabetes, major cardiovascular events were not significantly different 21 years after the DPP concluded, which may be due to the widespread use of statin, antihypertensive, and metformin therapy among all participants well after the conclusion of the study (19). However, the incidence of diabetes continued to be significantly reduced.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf Downloaded

Prevention or Delay of Diabetes # Diabetes Care Volume 49, Supplement 1, January 2026

Lower at 21 years of follow-up in both intensive lifestyle therapy and metformin groups, when compared with placebo, corresponding to increases in median diabetes-free survival of 3.5 years and 2.5 years, respectively (15). Importantly, the overall treatment effect was driven almost entirely by benefit seen during the initial DPP phase (15), underscoring the importance of early intervention and continued implementation of diabetes prevention efforts. Still, individuals can derive benefit from lifestyle interventions to prevent progression of prediabetes to diabetes irrespective of the duration of prediabetes prior to initiation of lifestyle intervention (20). Thus, there is clear potential benefit in diabetes prevention and minimal to no risk of harm with these interventions.

Nutrition

Nutrition counseling for weight loss in the DPP lifestyle intervention arm focused on reduction of total fat and calories (8,17,18). However, current evidence suggests there is no ideal percentage of calories from carbohydrate, protein, and fat to prevent diabetes; therefore, macronutrient distribution should be based on an individualized assessment of current eating patterns, preferences, and metabolic goals. A variety of eating patterns (21) can be appropriate for individuals with prediabetes or diabetes, including Mediterranean-style, plant-based (which may or may not be omnivorous), Dietary Approaches to Stop Hypertension (DASH), and low-carbohydrate (22–25). The overall quality of food consumed (as measured by the Healthy Eating Index, Alternative Healthy Eating Index, and DASH score), with an emphasis on whole grains, legumes, nuts, fruits, and vegetables and minimal refined and processed foods, is also associated with a lower risk of type 2 diabetes (26–28).

For people at risk for or living with diabetes, individualized medical nutrition therapy (see section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes,” for more detailed information) is effective in lowering A1C (29,30).

Physical Activity

Moderate-intensity physical activity, such as brisk walking for 150 min/week, has shown beneficial effects in those with prediabetes (8). Similarly, moderate-intensity physical activity has been shown to improve insulin sensitivity and reduce abdominal fat in children, adolescents, and young adults (31,32). Health care professionals are encouraged to promote a DPP-style program to all individuals who have been identified to be at an increased risk of type 2 diabetes. In addition to aerobic activity, a physical activity plan designed to

prevent diabetes can include resistance training (17,33). Breaking up prolonged sedentary time may also be encouraged, as it lowers postprandial glucose levels (34). The effects of regular physical activity participation appear to extend to the prevention of gestational diabetes mellitus (GDM) as well (35,36).

To qualify for Medicare coverage, individuals must have BMI $>25 \text{ kg/m}^2$ (or BMI $>23 \text{ kg/m}^2$ if self-identified as Asian) and glycemic testing consistent with prediabetes in the last year. Medicaid DPP coverage is also expanding on a state-by-state basis.

Delivery and Dissemination of Lifestyle Behavior Change for Diabetes Prevention

Because the intensive lifestyle intervention in the DPP was effective in preventing type 2 diabetes among those at high risk, and both DPP and adapted lifestyle behavior change programs for diabetes prevention were shown to be cost-effective across multiple applications, broader efforts to disseminate scalable DPP-like lifestyle behavior change programs for diabetes prevention with coverage by third-party payors ensued (37–42). Group delivery of DPP content in community or primary care settings has demonstrated the potential to reduce overall program costs while still producing weight loss and diabetes risk reduction (43,44). Interventions to prevent type 2 diabetes can also be delivered through workplace programming, with interventions that most closely adhere to DPP content shown to be most effective (45).

The Centers for Disease Control and Prevention (CDC) developed the National Diabetes Prevention Program (National DPP), a resource designed to bring such evidence-based lifestyle change programs for preventing type 2 diabetes to communities (cdc.gov/diabetes-prevention). To be eligible for this program, individuals must have a BMI in the overweight range and be at risk for diabetes based on laboratory testing, a previous diagnosis of GDM, or a positive risk test (cdc.gov/prediabetes/risktest/). The CDC has also developed the Diabetes Prevention Impact Tool Kit (nccd.cdc.gov/toolkit/diabetesimpact) to help organizations assess the economics of providing or assessing the effectiveness of lifestyle interventions.

Furthermore, trials involving medical nutrition therapy for adults with prediabetes found significant reductions in weight, waist circumference, and glycemia (54,55). Individuals with prediabetes can benefit from counseling by a registered dietitian nutritionist, which has been shown to help improve eating habits, increase physical activity, and achieve 7–10% weight loss (50–53).

diabetesjournals.org/care # Prevention or Delay of Diabetes

from referral to a registered dietitian constructs: quantity, quality, and timing. 3.10 Consider periodic assessment of vitamin B12 levels in individuals receiving long-term metformin therapy, especially in those with anemia or peripheral neuropathy. B

Because weight loss and maintenance through lifestyle (e.g., nutrition and physical activity) behavior changes might not be sufficient on their own (13), some people at high risk for type 2 diabetes may benefit from additional pharmacotherapy support. Various pharmacologic agents used to treat diabetes have also been evaluated for diabetes prevention. Metformin, α -glucosidase inhibitors, incretin receptor agonists (e.g., liraglutide, semaglutide, and tirzepatide), thiazolidinediones, and insulin have all been shown to lower the incidence of diabetes in specific populations (75–83).

The CDC Diabetes Prevention Recognition Program (DPRP) (<https://www.cdc.gov/diabetes-prevention/php/program-provider/program-requirements.html>) certifies technology-assisted modalities as effective vehicles for DPP-based interventions; such programs must use an approved curriculum, include interaction with a coach, and attain the DPP outcomes of participation, physical activity reporting, and weight loss. Health care professionals should consider referring adults with prediabetes to certified technology-assisted programs.

The CDC's Division of Diabetes Translation National DPP toolkit (coveragetoolkit.org/recruitment-referral-for-the-national-dpp-lifestyle-change-program) provides diabetes care team members resources for how to increase referral and uptake of the DPP. Some ways to increase DPP utilization include direct mailings, phone scripts, and text-based outreach. Using multiple methods of outreach to increase awareness among those at risk, and among health care professionals, in addition offering the DPP virtually can improve participation and uptake (65,66).

PHARMACOLOGIC INTERVENTIONS TO DELAY TYPE 2 DIABETES

Recommendations

- 3.7 Metformin for the prevention of type 2 diabetes should be considered in adults at high risk of type 2 diabetes, as typified by the Diabetes Prevention Program, especially those aged 25–59 years with BMI ≥ 35 kg/m², higher fasting plasma glucose (e.g., ≥ 110 mg/dL [≥ 6 mmol/L]), and higher A1C (e.g., $\geq 6.0\%$ [≥ 42 mmol/mol]). Importantly, in the DPP, weight loss was key to reducing the risk of progression to diabetes, with every kilogram of weight loss conferring a 16% reduction in risk of progression over 3.2 years (18). - 3.8 Consider using metformin to prevent hyperglycemia in high-risk individuals treated with a phosphatidylinositol 3-kinase α (PI3K α) inhibitor (e.g., alpelisib and inavolisib). B - 3.9 Consider using metformin to prevent hyperglycemia in high-risk individuals treated with high-dose glucocorticoids. B

Sleep Characteristics Associated With Increased Risk of Type 2 Diabetes

Sleep occupies approximately one-third of the day for most people and modulates important metabolic, endocrine, and cardiovascular processes (69). Sleep can be characterized using three key constructs: quantity, quality, and timing (i.e., chronotype). There is now established evidence for a U-shaped association between sleep duration and type 2 diabetes incidence, with the nadir typically occurring at 7 h per day, with short (typically defined as ≤ 6 h) and long (typically defined as > 9 h) sleep duration having up to a 50% increase in the risk of type 2 diabetes, including progression from prediabetes (70).

Sleep quality is defined as “an individual’s self-satisfaction with all aspects of the sleep experience” (71,72). A meta-analysis reported that poor overall sleep quality was associated with a 40% increased risk of developing type 2 diabetes (73). Chronotype preference has been linked with many chronic diseases, including type 2 diabetes. For example, for those with a preference for evenings (i.e., going to bed late and getting up late), there was a 2.5-fold higher odds ratio for type 2 diabetes than for those with a preference for mornings (i.e., going to bed early and getting up early), independent of sleep duration and sleep sufficiency (74).

More detailed information concerning sleep and the impact of sleep on prediabetes and diabetes risk and management can be found in section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes.”

There are currently no long-term data to support the use of pharmacologic treatments other than metformin for the sole purpose of preventing type 2 diabetes. However, use of glucagon-like peptide 1–based therapies for weight management in individuals with overweight or obesity—an essential component of type 2 diabetes prevention—is highly beneficial and should be considered, as discussed in detail in section 8, “Obesity and Weight Management for the Prevention and Treatment of Diabetes.”

Prevention or Delay of Diabetes # Diabetes Care Volume 49, Supplement 1, January 2026

Other medications have also been assessed for their efficacy in preventing type 2 diabetes. The antihypertensive valsartan was found to modestly reduce the incidence of diabetes among individuals with prediabetes and cardiovascular disease or risk factors, but with no difference in cardiovascular outcomes (88). Thus, we do not recommend using valsartan for the prevention of diabetes.

Testosterone therapy also reduced the risk of incident type 2 diabetes by 41% in men aged 50–74 years with central obesity and prediabetes (89). However, the benefit of testosterone was observed only when added to a structured lifestyle program. Indeed, the TRAVERSE Diabetes Study (a prespecified efficacy trial nested within the large multicenter TRAVERSE trial whose primary objective was to assess the cardiovascular outcomes of testosterone therapy in men aged 45–80 years with hypogonadism) found no benefit of testosterone as compared with placebo with respect to progression from prediabetes to diabetes or remission of diabetes among those who had prediabetes or diabetes at baseline, respectively (90). Testosterone therapy also did not alter glucose or A1C levels among participants with prediabetes or diabetes, nor were there any differences in outcomes by baseline testosterone level, preexisting cardiovascular disease, age, or race. Thus, we do not recommend testosterone therapy for the prevention of type 2 diabetes in men with hypogonadism; the decision to initiate testosterone replacement needs to be informed by individualized treatment considerations that weigh the expected benefits and harms of treatment.

There has been significant interest in understanding the potential of vitamin D therapy to prevent progression of high-risk prediabetes to type 2 diabetes in adults (91). Three randomized controlled trials tested whether

vitamin D therapy in combination with lifestyle modification reduces the risk of developing diabetes in adults with high-risk prediabetes (i.e., IGT or meeting two or three criteria for the diagnosis of prediabetes [fasting glucose, A1C, 2-h glucose after a 75-g OGTT]): the Tromsø study in Norway, with 511 participants; the Vitamin D and Type 2 Diabetes (D2d) study in the U.S., with 2,423 participants; and the Diabetes Prevention with Active Vitamin D (DPVD) study.

Although vitamin D therapy consistently appeared to modestly reduce the risk of developing diabetes in all three trials, none of the results were statistically significant, which was attributed to insufficient statistical power. Subsequently, several meta-analyses pooling the three randomized clinical trials along with other smaller studies suggested a modest (i.e., 10–15% risk reduction) potential benefit in specific populations, such as older adults, individuals with lower baseline vitamin D levels, and individuals without obesity (95,96). However, there are several concerns and uncertainties regarding recommending widespread vitamin D therapy for adults with high-risk prediabetes. First, these trials used varying dosages of vitamin D that were higher than the recommended daily allowance (i.e., 600 IU/day for those aged 18–70 years and 800 IU/day for those older than 70 years). Second, the benefit-to-risk ratio of vitamin D therapy for people with high-risk prediabetes remains uncertain. Although there were no safety concerns identified in these trials, the anticipated treatment of many millions of adults with prediabetes in the U.S. and globally could increase the risk of adverse events (such as hypercalcemia, hypercalciuria, nephrolithiasis, and kidney failure), especially treatment with unspecified doses of vitamin D without monitoring blood 25-hydroxy vitamin D levels.

Thus, the expected benefit, harm, treatment burden, and cost of using any medication to prevent type 2 diabetes must be considered for each individual as part of informed shared decision-making. This is particularly important because any pharmacologic intervention is likely to be needed to be used long-term because of waning effects after stopping the medication. At the present time, there are no pharmacologic agents approved by the U.S. Food and Drug Administration specifically for the prevention of type 2 diabetes.

Metformin

Metformin has the most robust efficacy and safety data as a pharmacologic therapy for diabetes prevention among people with prediabetes (82). Importantly, metformin was overall less effective than lifestyle modification in the DPP, although it has been demonstrated in small randomized clinical trials both in healthy individuals (100) and in individuals receiving high-dose glucocorticoids for the treatment of cancer (101) and rheumatologic/autoimmune disorders (102,103). In one trial with 40 participants receiving high-dose and/or long-duration glucocorticoid therapy, metformin can be considered to prevent the development of glucocorticoid-induced hyperglycemia in high-risk individuals, notably those receiving higher doses of glucocorticoids, with longer treatment duration, and in the presence of other risk factors for diabetes.

The efficacy of metformin to prevent hyperglycemia—particularly insulin resistance and postprandial hyperglycemia—has been demonstrated in small randomized clinical trials.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf from Downloaded

diabetesjournals.org/care # Prevention or Delay of Diabetes

doses prednisolone for the treatment of inflammatory conditions, metformin also reduced rates of infection and all-cause hospital readmissions (103).

Proactive use of metformin may also be beneficial in preventing hyperglycemia in individuals receiving phosphatidylinositol 3-kinase α (PI3K α) inhibitors (e.g., alpelisib and inavolisib) for the treatment of cancer (see section 2, “Diagnosis and Classification of Diabetes,” for additional details on diabetes induced by systemic anticancer therapy). This is particularly important because hyperglycemia is the most frequent adverse event of PI3K α inhibitor therapy that leads to treatment discontinuation (104).

In a phase 2 trial of metformin to prevent hyperglycemia in individuals receiving alpelisib for advanced breast cancer (metformin was started 1 week prior to starting alpelisib and continued for the duration of treatment), metformin prevented the development of grade 3–4 hyperglycemia in all but four participants, and none had to discontinue alpelisib therapy (105). Similar benefits were seen with implementation of a metformin hyperglycemia-prevention protocol in routine practice (106).

Individuals at highest risk for developing hyperglycemia with PI3K α inhibitor therapy include those who are older (≥ 70 years old), with obesity (BMI ≥ 30 kg/m²), concurrently treated with glucocorticoids, and with hyperglycemia at baseline (e.g., A1C $\geq 5.7\%$ [39 mmol/mol] or fasting plasma glucose ≥ 100 mg/dL [5.6 mmol/L]) (107, 108). However, it is important to weigh the benefits of preventing hyperglycemia with the potential adverse effects of metformin therapy, particularly diarrhea, which is also a frequent adverse effect of PI3K α therapy (though there is no evidence that concurrent use of these two medications further exacerbates diarrhea risk) (108).

Decreased vitamin B12 levels are a known consequence of long-term treatment with metformin (109). Periodic assessment of vitamin B12 level in those taking metformin chronically should be considered to check for possible deficiency, especially in those receiving a higher dose (e.g., $\geq 1,500$ mg/day) (110), with longer treatment duration (4–5 years) (111), with other risk factors for vitamin B12 deficiency, and if a deficiency is suspected, such as in people with anemia.

Type 2 diabetes has shown a reduction in cardiovascular risk factors and the need for medications used to treat these cardiovascular risk factors (121, 122). The lifestyle intervention in the Da Qing study was associated with lowering cardiovascular disease and mortality at 23 and 30 years of observational follow-up (10, 12).

Treatment goals and therapies for hypertension and dyslipidemia in the primary and secondary prevention of cardiovascular disease for people with prediabetes should be based on their level of cardiovascular risk. Increased vigilance is warranted to identify and treat these and other cardiovascular disease risk factors (123).

Statin use increases risk of diabetes (124, 125). In the DPP, statin use was associated with greater diabetes risk irrespective of the treatment group (pooled hazard ratio [HR] [95% CI] for incident diabetes 1.36 [1.17–1.58]) (124). In trials of primary and secondary prevention of cardiovascular disease, cardiovascular and mortality benefits of statin therapy exceed the risk of diabetes (126, 127), suggesting a highly favorable benefit-to-harm balance with statin therapy. Hence, discontinuation of statins due to concerns of diabetes risk is not recommended in this population.

Results of cardiovascular outcome trials in people without diabetes also provide important evidence for cardiovascular risk reduction in people with increased cardiometabolic risk, though not specifically prediabetes (see section 10, “Cardiovascular Disease and Risk Management,” for more details).

The IRIS (Insulin Resistance Intervention after Stroke) trial of people with a recent (<6 months) stroke or transient ischemic attack, without diabetes but with insulin resistance (as defined by a HOMA of insulin resistance index of ≥ 3.0), evaluated pioglitazone (goal dose of 45 mg daily) compared with placebo. At 4.8 years, the risk of stroke or myocardial infarction, as well as the risk of diabetes, was lower in the pioglitazone group than in the placebo group; weight gain, edema, and fractures were higher in the pioglitazone treatment group (128–130). Lower doses may mitigate the adverse effects but may also be less effective (131).

PERSON-CENTERED CARE GOALS

Recommendations

3.14 In adults with overweight or obesity at high risk of type 2 diabetes, care goals should include weight

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf from Downloaded

Prevention or Delay of Diabetes # Diabetes Care Volume 49, Supplement 1, January 2026

Loss and maintenance, minimizing the progression of hyperglycemia, and attention to cardiovascular risk.

3.15 Pharmacotherapy

(e.g., for weight management, minimizing the progression of hyperglycemia, and cardiovascular risk reduction) should be considered to support person-centered care goals.

3.16 More intensive preventive approaches

should be considered in individuals who are at particularly high risk of progression to diabetes, including individuals with BMI ≥ 35 kg/m², those with higher glucose levels (e.g., fasting plasma glucose 110–125 mg/dL

[6.1–6.9 mmol/L], 2-h post-challenge glucose 173–199 mg/dL [9.6–11.0 mmol/L], and A1C $\geq$ 6.0% [$\geq$ 42 mmol/mol]), and individuals with a history of gestational diabetes mellitus.

Individualized risk-to-benefit ratio should be considered in screening, intervention, and monitoring to lower the risk of type 2 diabetes and associated comorbidities. Multiple factors, including age, BMI, and other comorbidities, may influence the risk of progression to diabetes and lifetime risk of complications (132,133). Prediabetes is associated with increased cardiovascular disease and mortality (118), which emphasizes the importance of attending to cardiovascular risk in this population.

However, the new diagnosis of prediabetes in older adults (aged >70 years) is less relevant for progression to diabetes; indeed, regression to normoglycemia or death was more frequent than progression to diabetes in the Atherosclerosis Risk in Communities (ARIC) study (132).

Moreover, diabetes-related morbidity and mortality is lower and the benefits of treatment are not established in those who develop type 2 diabetes after age 70 years (134).

In the DPP, which enrolled high-risk individuals with IGT, elevated fasting glucose, and elevated BMI, the crude incidence of diabetes within the placebo group was 11 cases per 100 person-years, with a cumulative 3-year incidence of diabetes of 29% (8). Characteristics of individuals in the DPP and DPPOS who were at particularly high risk of progression to diabetes (crude incidence of diabetes 14–22 cases per 100 person-years) were considered in individuals at high risk of progression.

PREVENTION OR DELAY OF SYMPTOMATIC TYPE 1 DIABETES

Lifestyle and Type 1 Diabetes Progression

Observational studies have identified several factors that increase β -cell demand and risk of progression to clinical type 1 diabetes among individuals with islet autoantibodies, including lower levels of physical activity (138), higher glycemic index eating patterns (139), and total sugar intake (140). Similar associations have not been observed for the development of autoantibodies.

In The Environmental Determinants of Diabetes in the Young (TEDDY) longitudinal study, daily minutes spent in moderate to vigorous physical activity were associated with a reduced risk of progression to type 1 diabetes in children 5–15 years of age with multiple islet autoantibodies (HR 0.92 [95% CI 0.86–0.99] per 10-min increase; $P = 0.02$) (138). In the Diabetes Autoimmunity Study in the Young (DAISY), also in children with islet autoantibodies, consumption of higher glycemic index foods (HR 2.20 [95% CI 1.17–4.15]) and total sugar intake (HR 1.75 [95% CI 1.07–2.85]) (139,140) were both associated with progression to type 1 diabetes.

In nonobese diabetic mice, an animal model for the development of type 1 diabetes, sustained high-glucose drinking significantly aggravated islet inflammation and accelerated the onset of type 1 diabetes (141). However, efficacy of lifestyle interventions that modify these factors in individuals with stage 1 or stage 2 type 1 diabetes has not yet been reported.

Pharmacologic Interventions to Delay Symptomatic Type 1 Diabetes

Recommendation

3.17 Teplizumab-mzwv infusion to delay the onset of symptomatic type 1 diabetes (stage 3) should be discussed with selected individuals aged $\geq$ 8 years with stage 2 type 1 diabetes. Treatment should be in a setting with appropriately trained personnel.

Teplizumab, a CD3-directed humanized monoclonal antibody engineered to have decreased Fc receptor binding, has been...

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf from Downloaded

diabetesjournals.org/care # Prevention or Delay of Diabetes

approved to delay the onset of stage 3 type 1 diabetes in people 8 years and older with stage 2 type 1 diabetes based in part on the results of a single trial in relatives of people with type 1 diabetes (142). In this study, 44 individuals were randomized to a 14-day course of teplizumab and 32 to placebo. The median time to stage 3

type 1 diabetes diagnosis was 48.4 months in the teplizumab group and 24.4 months in the placebo group. Type 1 diabetes was diagnosed in 19 (43%) participants who received teplizumab and 23 (72%) participants who received placebo (HR 0.41 [95% CI 0.22–0.78]). In prespecified analyses, the presence of HLA-DR4, absence of HLA-DR3, and absence of antizinc transporter 8 antibody predicted response to teplizumab (HR 0.20 [95% CI 0.09–0.45], 0.18 [0.07–0.45], and 0.07 [0.02–0.26], respectively). The most common adverse reactions were transient lymphopenia (73%) followed by rash (36%).

Numerous clinical studies are being conducted to test methods for preventing or delaying the onset of stage 3 type 1 diabetes in those with evidence of autoimmunity without symptoms or for delaying loss of insulin secretory capacity after onset of stage 3, some with promising results (see ClinicalTrials.gov and TrialNet.org).

References

1. Ziegler AG, Rewers M, Simell O, et al. Seroconversion to multiple islet autoantibodies and risk of progression to diabetes in children. *JAMA* 2013;309:2473–2479
2. Steck AK, Dong F, Taki I, et al. Continuous glucose monitoring predicts progression to diabetes in autoantibody positive children. *J Clin Endocrinol Metab* 2019;104:3337–3344
3. Ylescupidez A, Speake C, Pietropaolo SL, et al. OGTT metrics surpass continuous glucose monitoring data for T1D prediction in multiple-autoantibody-positive individuals. *J Clin Endocrinol Metab* 2023;109:57–67
4. Phillip M, Achenbach P, Addala A, et al. Consensus guidance for monitoring individuals with islet autoantibody-positive pre-stage 3 type 1 diabetes. *Diabetes Care* 2024;47:1276–1298
5. Sims EK, Besser REJ, Dayan C, et al.; NIDDK Type 1 Diabetes TrialNet Study Group. Screening for type 1 diabetes in the general population: a status report and perspective. *Diabetes* 2022;71:610–623
6. Beran D, Bandini A, Bosi E, et al. Type 1 diabetes screening: need for ethical, equity, and health systems perspective. *Lancet Diabetes Endocrinol* 2025;13:175–176
7. Wentworth JM, Sing ABE, Naselli G, et al. Islet autoantibody screening throughout Australia using in-home blood spot sampling: 2-year outcome of lifestyle intervention in relation to duration of pre-diabetes: the Pathobiology and Reversibility of Prediabetes in a Biracial Cohort (PROP-ABC) study. *BMJ Open Diabetes Res Care* 2022;10:e002748
8. Dietary Guidelines Advisory Committee. Scientific Report of the 2025 Dietary Guidelines.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf from Downloaded

Prevention or Delay of Diabetes *Diabetes Care* Volume 49, Supplement 1, January 2026

overweight or obesity. *BMC Pregnancy Childbirth* 2025;25:5

1. Evert AB, Dennison M, Gardner CD, et al.; Nutrition therapy for adults with diabetes or prediabetes: a consensus report. *Diabetes Care* 2019;42:731–754
2. Raynor HA, Davidson PG, Burns H, et al. Medical nutrition therapy and weight loss questions for the evidence analysis library prevention of type 2 diabetes project: systematic reviews. *J Acad Nutr Diet* 2017;117:1578–1611
3. Khk L. Nutrition therapy for adults with diabetes or prediabetes. *ADCES Pract* 2022;10:34–38
4. Briggs Early K, Stanley K. Position of the Academy of Nutrition and Dietetics: the role of medical nutrition therapy and registered dietitian nutritionists in the prevention and treatment of prediabetes and type 2 diabetes. *J Acad Nutr Diet* 2018;118:343–353
5. Parker AR, Byham-Gray L, Denmark R, Winkle PJ. The effect of medical nutrition therapy by a registered dietitian nutritionist in patients with prediabetes participating in a randomized controlled clinical research trial. *J Acad Nutr Diet* 2014;114:1739–1748
6. Esposito K, Chiodini P, Maiorino MI, Bellastella G, Panagiotakos D, Giugliano D. Which diet for prevention of type 2 diabetes? A meta-analysis of prospective studies. *Endocrine* 2014;47:107–116
7. Powers MA, Bardsley JK, Cypress M, et al. Diabetes self-management education and support in adults with type 2 diabetes: a consensus report of the American Diabetes Association, the Association of Diabetes Care & Education Specialists, the Academy of Nutrition and Dietetics, the American Academy of Family Physicians, the American Academy of PAs, the American Association of Nurse Practitioners, and the American Pharmacists Association. *Diabetes Care* 2022;45:2753–2786
8. Mostafa SA, Mena SC, Antza C, Balanos G, Nirantharakumar K, Tahrani AA. Sleep behaviours and associated habits and the progression of prediabetes to type 2 diabetes mellitus in adults: a systematic review and meta-analysis. *Diab Vasc Dis Res* 2022;19:14791641221088824
9. Nelson KL, Davis JE, Corbett CF. Sleep quality: an evolutionary concept analysis. *Nurs Forum* 2022;57:144–151
10. Grock S, Ku J-h, Kim J, Moin T. A review of technology-assisted interventions for diabetes prevention. *Curr Diab Rep* 2017;17:107
11. Bian RR, Piatt GA, Sen A, et al. The effect of technology-mediated diabetes prevention interventions on weight: a meta-analysis. *J*

Med Internet Res 2017;19:e76 12. Moin T, Damschroder LJ, AuYoung M, et al. Results from a trial of an online diabetes prevention program intervention. *Am J Prev Med* 2018;55:583–591 13. Michaelides A, Major J, Pienkosz E, Wood M, Kim Y, Toro-Ramos T. Usefulness of a novel mobile diabetes prevention program delivery platform with human coaching: 65-week observational follow-up. *JMIR Mhealth Uhealth* 2018;6:e93 14. Kim SE, Castro Sweet CM, Cho E, Tsai J, Cousineau MR. Evaluation of a digital diabetes prevention program adapted for low-income patients, 2016-2018. *Prev Chronic Dis* 2019;16:E155 15. Vadheim LM, Patch K, Brokaw SM, et al. Telehealth delivery of the diabetes prevention program to rural communities. *Transl Behav Med* 2017;7:286–291 16. Fischer HH, Durfee MJ, Raghunath SG, Ritchie ND. Short message service text message support for weight loss in patients with prediabetes: pragmatic trial. *JMIR Diabetes* 2019;4:e12985

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf from Downloaded

diabetesjournals.org/care # Prevention or Delay of Diabetes

1. Chiasson J-L, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M, STOP-NIDDM Trial Research Group. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet* 2002;359:2072–2077 2. Wilding JPH, Batterham RL, Calanna S, et al.; STEP 1 Study Group. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med* 2021;384:989–1002 3. Jastreboff AM, Aronne LJ, Ahmad NN, et al.; SURMOUNT-1 Investigators. Tirzepatide once weekly for the treatment of obesity. *N Engl J Med* 2022;387:205–216 4. Diabetes Prevention Program Research Group. Long-term safety, tolerability, and weight loss associated with metformin in the Diabetes Prevention Program Outcomes Study. *Diabetes Care* 2012;35:731–737 5. Diabetes Prevention Program Research Group. Long-term effects of metformin on diabetes prevention: identification of subgroups that benefited most in the Diabetes Prevention Program and Diabetes Prevention Program Outcomes Study. *Diabetes Care* 2019;42:601–608 6. Dennison RA, Chen ES, Green ME, et al. The absolute and relative risk of type 2 diabetes after gestational diabetes: a systematic review and meta-analysis of 129 studies. *Diabetes Res Clin Pract* 2021;171:108625 7. Torgerson JS, Hauptman J, Boldrin MN, Sjostrom L. XENical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care* 2004;27:155–161 8. Garvey WT, Ryan DH, Henry R, et al. Prevention of type 2 diabetes in subjects with prediabetes and metabolic syndrome treated with phentermine and topiramate extended release. *Diabetes Care* 2014;37:912–921 9. Jastreboff AM, Le Roux CW, Stefanski A, et al.; SURMOUNT-1 Investigators. Tirzepatide for obesity treatment and diabetes prevention. *N Engl J Med* 2025;392:958–971 10. McMurray JJ, Holman RR, Haffner SM, et al.; NAVIGATOR Study Group. Effect of valsartan on the incidence of diabetes and cardiovascular events. *N Engl J Med* 2010;362:1477–1490 11. Wittert G, Bracken K, Robledo KP, et al. Testosterone treatment to prevent or revert type 2 diabetes in men enrolled in a lifestyle programme (T4DM): a randomised, double-blind, placebo-controlled, 2-year, phase 3b trial. *Lancet Diabetes Endocrinol* 2021;9:32–45 12. Bhasin S, Lincoff AM, Nissen SE, et al. Effect of testosterone on progression from prediabetes to diabetes in men with hypogonadism: a substudy of the TRAVERSE randomized clinical trial. *JAMA Intern Med* 2024;184:353–362 13. Demay MB, Pittas AG, Bikle DD, et al. Vitamin D for the prevention of disease: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2024;109:1907–1947 14. Jorde R, Sollid ST, Svartberg J, et al. Vitamin D 20,000 IU per week for five years does not prevent progression from prediabetes to diabetes. *J Clin Endocrinol Metab* 2016;101:1647–1655 15. Pittas AG, Dawson-Hughes B, Sheehan P, et al.; D2d Research Group. Vitamin D supplementation and prevention of type 2 diabetes. *N Engl J Med* 2019;381:520–530 16. Kawahara T, Suzuki G, Mizuno S, et al. Effect of active vitamin D treatment on development of type 2 diabetes: DPVD randomised controlled trial in Japanese population. *BMJ* 2022;377:e066222 17. Pittas AG, Kawahara T, Jorde R, et al. Vitamin D and risk for type 2 diabetes in people with prediabetes: a systematic review and meta-analysis of individual participant data from 3 randomized clinical trials. *Ann Intern Med* 2023;176:355–363 18. Shah VP, Nayfeh T, Alsawaf Y, et al. A systematic review supporting the Endocrine Society clinical practice guidelines on vitamin D. *J Clin Endocrinol Metab* 2024;109:1961–1974 19. Ratner RE, Christophi CA, Metzger BE, et al.; Diabetes Prevention Program Research Group. Prevention of diabetes in women with a history of gestational diabetes: effects of metformin and lifestyle interventions. *J Clin Endocrinol Metab* 2008;93:4774–4779 20. Aroda VR, Christophi CA, Edelstein SL, et al.; Diabetes Prevention Program Research Group. The effect of lifestyle intervention and metformin on preventing or delaying diabetes among women with and without gestational diabetes: the Diabetes Prevention Program Outcomes Study 10-year follow-up. *J Clin Endocrinol Metab* 2015;100:1646–1653 21. Kim J, Ahn CW, Fang S, Lee HS, Park JS. Association between metformin

dose and vitamin B12 deficiency in patients with type 2 diabetes. *Medicine (Baltimore)* 2019;98:e17918 22. de Jager J, Kooy A, Leher P, et al. Long term treatment with metformin in patients with type 2 diabetes and risk of vitamin B-12 deficiency: randomised placebo controlled trial. *BMJ* 2010;340:c2181 23. Griffin SJ, Bethel MA, Holman RR, et al. Metformin in non-diabetic hyperglycaemia: the GLINT feasibility RCT. *Health Technol Assess* 2018;22:1–64 24. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int* 2024;105:S117–S314 25. Jung C, Park S, Kim H. Concomitant use of metformin and proton pump inhibitors increases vitamin B12 deficiency risk in type 2 diabetes. *J Diabetes Investig* 2025;16:1020–1027 26. Chapman LE, Darling AL, Brown JE. Association between metformin and vitamin B12 deficiency in patients with type 2 diabetes: a systematic review and meta-analysis. *Diabetes Metab* 2016;42:316–327 27. Ali MK, Bullard KM, Saydah S, Imperatore G, Gregg EW. Cardiovascular and renal burdens of prediabetes in the USA: analysis of data from serial cross-sectional surveys, 1988–2014. *Lancet Diabetes Endocrinol* 2018;6:392–403 28. Pan Y, Chen W, Wang Y. Prediabetes and outcome of ischemic stroke or transient ischemic attack: a systematic review and meta-analysis. *J Stroke Cerebrovasc Dis* 2019;28:683–692 29. Huang Y, Cai X, Mai W, Li M, Hu Y. Association between prediabetes and risk of cardiovascular disease and all cause mortality: systematic review and meta-analysis. *BMJ* 2016;355:i5953 30. Yeh H-C, Duncan BB, Schmidt MI, Wang N-Y, Brancati FL. Smoking, smoking cessation, and risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med* 2010;152:10–17 31. Hu Y, Zong G, Liu G, et al. Smoking cessation, weight change, type 2 diabetes, and mortality. *N Engl J Med* 2018;379:623–632 32. Orchard TJ, Temprosa M, Barrett-Connor E, et al. Long-term effects of the Diabetes Prevention Program interventions on cardiovascular risk factors:

February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf from Downloaded

Prevention or Delay of Diabetes

Diabetes Care Volume 49, Supplement 1, January 2026

a report from the DPP Outcomes Study. *Diabet Med*

stroke or transient ischemic attack. *N Engl J Med*

disease in the Diabetes Prevention Program Outcomes Study (DPPOS). *Diabetes Care* 2019;

| 122. | Salas-Salvadó J, Díaz-López A, Ruiz-Canela M, et al.; PREDIMED-Plus investigators. Effect of a lifestyle intervention program with energy-restricted Mediterranean diet and exercise on weight loss and cardiovascular risk factors: one-year results of the PREDIMED-Plus trial. *Diabetes Care* 2019;42:777–788 | |

----- | 123. | Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019;140:e596–e646 | | 124. | Crandall JP, Mather K, Rajpathak SN, et al. Statin use and risk of developing diabetes: results from the Diabetes Prevention Program. *BMJ Open Diabetes Res Care* 2017;5:e000438 | | 125. | Cholesterol Treatment Trialists' (CTT) Collaboration. Effects of statin therapy on diagnoses of new-onset diabetes and worsening glycaemia in large-scale randomised blinded statin trials: an individual participant data meta-analysis. *Lancet Diabetes Endocrinol* 2024;12:306–319 | | 126. | Ridker PM, Pradhan A, MacFadyen JG, Libby P, Glynn RJ. Cardiovascular benefits and diabetes risks of statin therapy in primary prevention: an analysis from the JUPITER trial. *Lancet* 2012;380:565–571 | | 127. | Cai T, Abel L, Langford O, et al. Associations between statins and adverse events in primary prevention of cardiovascular disease: systematic review with pairwise, network, and dose-response meta-analyses. *BMJ* 2021;374:n1537 | | 128. | Kernan WN, Viscoli CM, Furie KL, et al.; IRIS Trial Investigators. Pioglitazone after ischemic stroke or transient ischemic attack in patients with insulin resistance. *Diabetes Obes Metab* 2022;24:1150–1158 | | 129. | Inzucchi SE, Viscoli CM, Young LH, et al.; IRIS Trial Investigators. Pioglitazone prevents diabetes in patients with insulin resistance and cerebrovascular disease. *Diabetes Care* 2016;39:1684–1692 | | 130. | Spence JD, Viscoli CM, Inzucchi SE, et al.; IRIS Investigators. Pioglitazone therapy in patients with stroke and prediabetes: a post hoc analysis of the IRIS randomized clinical trial. *JAMA Neurol* 2019;76:526–535 | | 131. | Spence JD, Viscoli C, Kernan WN, et al. Efficacy of lower doses of pioglitazone after stroke or transient ischaemic attack in patients

with insulin resistance. *Diabetes Obes Metab* 2022;24:1150–1158 || 132. | Nadeau KJ, Anderson BJ, Berg EG, et al. Youth-onset type 2 diabetes consensus report: current status, challenges, and priorities. *Diabetes Care* 2016;39:1635–1642 || 133. | Rooney MR, Rawlings AM, Pankow JS, et al. Risk of progression to diabetes among older adults with prediabetes. *JAMA Intern Med* 2021;181:511–519 || 134. | Cigolle CT, Blaum CS, Lyu C, Ha J, Kabeto M, Zhong J. Associations of age at diagnosis and duration of diabetes with morbidity and mortality among older adults. *JAMA Netw Open* 2022;5:e2232766 || 135. | Lachin JM, Christophi CA, Edelstein SL, et al.; DDK Research Group. Factors associated with diabetes onset during metformin versus placebo therapy in the diabetes prevention program. *Diabetes* 2007;56:1153–1159 || 136. | Perreault L, Pan Q, Schroeder EB, et al.; Diabetes Prevention Program Research Group. Regression from prediabetes to normal glucose regulation and prevalence of microvascular disease among people with impaired glucose tolerance: a secondary analysis of the 30 year Da Qing Diabetes Prevention Outcome Study. *Diabetologia* 2021;64:1279–1287 || 137. | Chen Y, Zhang P, Wang J, et al. Associations of progression to diabetes and regression to normal glucose tolerance with development of cardiovascular and microvascular disease among people with impaired glucose tolerance: a secondary analysis of the 30 year Da Qing Diabetes Prevention Outcome Study. *Diabetologia* 2021;64:1279–1287 || 138. | Liu X, Johnson SB, Lynch KF, et al.; TEDDY Study Group. Physical activity and the development of islet autoimmunity and type 1 diabetes in 5- to 15-year-old children followed in the TEDDY study. *Diabetes Care* 2023;46:1409–1416 || 139. | Lamb MM, Yin X, Barriga K, et al. Dietary glycemic index, development of islet autoimmunity, and subsequent progression to type 1 diabetes in young children. *J Clin Endocrinol Metab* 2008;93:3936–3942 || 140. | Lamb MM, Frederiksen B, Seifert JA, Kroehl M, Rewers M, Norris JM. Sugar intake is associated with progression from islet autoimmunity to type 1 diabetes: the Diabetes Autoimmunity Study in the Young. *Diabetologia* 2015;58:2027–2034 || 141. | Li X, Wang L, Meng G, et al. Sustained high glucose intake accelerates type 1 diabetes in NOD mice. *Front Endocrinol (Lausanne)* 2022;13:1037822 || 142. | Herold KC, Bundy BN, Long SA, et al.; Type 1 Diabetes TrialNet Study Group. An anti-CD3 antibody, teplizumab, in relatives at risk for type 1 diabetes. *N Engl J Med* 2019;381:603–613 |

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S50/848822/dc26s003.pdf Downloaded