

1. Improving Care and Promoting Health in Populations: Standards of Care in Diabetes—2026

Diabetes Care 2026;49(Suppl. 1):S13–S26 | <https://doi.org/10.2337/dc26-S001>

The American Diabetes Association (ADA) “Standards of Care in Diabetes” includes the ADA’s current clinical practice recommendations and is intended to provide the components of diabetes care, general treatment goals and guidelines, and tools to evaluate quality of care. Members of the ADA Professional Practice Committee for Diabetes, an interprofessional expert committee, are responsible for updating the Standards of Care annually, or more frequently as warranted. For a detailed description of ADA standards, statements, and reports, as well as the evidence-grading system for ADA’s clinical practice recommendations and a full list of Professional Practice Committee members, please refer to Introduction and Methodology. Readers who wish to comment on the Standards of Care are invited to do so at professional.diabetes.org/SOC.

SYSTEMS TO SUPPORT DIABETES POPULATION HEALTH

Recommendations

1. Ensure treatment decisions are timely, rely on evidence-based guidelines, address social determinants of health, and incorporate shared decision-making based on individual values, preferences, prognoses, comorbidities, and informed financial considerations. B 2. Align approaches to diabetes management with evidence-based care models. These models emphasize person-centered team care, integrated long-term treatment approaches to diabetes and comorbidities, and ongoing collaborative communication and goal setting among all team members and with people with diabetes. A 3. Care systems should facilitate in-person and virtual team-based care, include those knowledgeable and experienced in diabetes management as part of the team, and utilize patient registries, decision support tools, proactive care planning, and community involvement to meet needs of individuals with diabetes. B 4. Assess diabetes management, risk factors, and complications using reliable and relevant data metrics to improve processes of care and health outcomes, with attention to individual values, preferences, goals for care, and treatment burden, including costs of care. B 5. Health systems should adopt a culture of continuous quality improvement, implement benchmarking programs, and engage interprofessional teams to support sustainable and scalable process changes to improve quality of care and health outcomes. A

*A complete list of members of the American Diabetes Association Professional Practice Committee for Diabetes can be found at <https://doi.org/10.2337/dc26-SINT>.

Duality of interest information for each contributor is available at <https://doi.org/10.2337/dc26-SDIS>.

Suggested citation: American Diabetes Association Professional Practice Committee for Diabetes. 1. Improving care and promoting health in populations: Standards of Care in Diabetes—2026. Diabetes Care 2026;49(Suppl. 1):S13–S26 © 2025 by the American Diabetes Association. Readers may use this work for educational, noncommercial purposes if properly cited and unaltered. The version of record may be linked at <https://diabetesjournals.org/care>, but ADA permission is required to post this work on any third-party site or platform. This publication and its contents may not be reproduced, distributed, or used for text or data mining, machine learning, or similar technologies without prior written permission. Requests to reuse, adapt, or distribute this work may be sent to permissions@diabetes.org. More information is available at <https://diabetesjournals.org/journals/pages/license>.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf Downloaded

Improving Care and Promoting Health in Populations

Diabetes Care Volume 49, Supplement 1, January 2026

Population health is a multidimensional concept that refers to the health outcomes of a group of individuals and the potentially varied distribution of those outcomes within the group (1). These outcomes can be measured in terms of health indicators (mortality, morbidity, quality of life, and functional status), disease epidemiology (incidence and prevalence), and behavioral and metabolic factors (physical activity, nutrition, A1C, time in range, etc.) (1). In clinical practice, population health commonly refers to the measurement and improvement of

health outcomes for a defined group of individuals, often those receiving care within a particular health system or practice, and is central to operational “population health management,” or the proactive, team-based, and data-driven approaches to improving health care experiences and team well-being while minimizing costs and addressing inequities, in alignment with the Quadruple Aim (2).

More broadly, population health refers to the health of all people living in a geographic area or community, and it centers on upstream drivers of health that must be addressed, often through cross-sector collaboration and policy, to improve the broader conditions that influence health. Clinical practice recommendations are tools for health care professionals who seek to improve health across populations; however, for optimal outcomes, diabetes care must also be individualized for each person with diabetes and for each person's context, as well as across the life span. Thus, efforts to improve population health will require a combination of policy-level, system-level, and person-level approaches. With such an integrated approach in mind, the American Diabetes Association (ADA) highlights the importance of person-centered care, defined as care that considers an individual's comorbidities and prognoses; is respectful of and responsive to individual preferences, needs, and values; and ensures that the individual's values guide all clinical decisions (3).

Social determinants of health (SDOH)—factors often beyond an individual's direct control and potentially representing lifelong risks—play a significant role in both clinical and psychosocial outcomes. SDOH are defined as the economic, environmental, political, and social conditions in which people live, and their uneven distribution is responsible for a major part of health inequality worldwide (4). To improve health, support overall well-being, and eliminate disparities, it is crucial to address these determinants, particularly for individuals from racial and ethnic minoritized communities, underserved geographic areas (rural or urban), and those facing socioeconomic barriers to care and health (5).

State of Diabetes and Diabetes Care

In 2021, an estimated 38.4 million people were living with diabetes, representing 11.6% of the U.S. population. This includes 38.1 million adults aged 18 years and older (14.7% of U.S. adults), of whom 8.7 million were unaware of their diagnosis. An additional 97.6 million adults aged 18 years and older (38.0% of the U.S. population) were living with prediabetes (6), and over 100 million adults 20 years and older (41.9% of U.S. adults) were living with obesity (7), placing them at high risk for developing type 2 diabetes. The prevalence of diabetes is increasing among children and adolescents as well. In 2017, the year for which the most recent data are available, the prevalence of type 1 diabetes was 2.15 per 10,000 individuals aged 19 years and under, and the prevalence of type 2 diabetes was 0.67 per 1,000 individuals 10–19 years of age (prevalence among those under 10 years old was too low to be reported) (8).

Many segments of the population face additional and unique challenges to diabetes management, resulting in higher rates of diabetes complications, morbidity, and mortality. These include children, adolescents, and young adults; individuals with complex health needs, financial or other social hardships, and/or limited English proficiency; and individuals in groups and communities that have been historically marginalized (6,10,12,13). The prevalence of diabetes among adults is higher among non-Hispanic Black individuals (17.4%), non-Hispanic Asian individuals (16.7%), and Hispanic individuals (15.5%) than among White individuals (13.6%). Diabetes prevalence is also higher among adults with less than a high school education (13.1%) than among those with more than a high school education (6.9%).

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded

diabetesjournals.org/care # Improving Care and Promoting Health in Populations

among those from families living below the federal poverty level (13.1%) than among those at 500% of federal poverty level or higher (5.1%), and higher among those living in nonmetropolitan areas (9.5%) than among those living in metropolitan areas (8.1%) (6). There is also substantial county-level variation in diabetes complications, mortality, and disparities in complications and mortality experienced by minoritized racial and ethnic groups across the U.S. (13), underscoring the impacts of broader structural factors on health among historically marginalized populations.

Gaps and disparities in diabetes management and outcomes are also prevalent among children, adolescents, and young adults with diabetes in the U.S. Improving diabetes management early in life is essential for

preventing adverse health outcomes and disability and improving quality of life both in the short- and long-term. There are important and nuanced considerations in treating diabetes and supporting children, adolescents, and young adults with diabetes that are distinct from those of older adults (14). Data from SEARCH for Diabetes in Youth (SEARCH), a population-based registry network of five centers across five U.S. states, showed that in 2014–2019, mean A1C was 9.1% (76 mmol/mol) among youth and young adults with type 1 diabetes and 8.9% (74 mmol/mol) in youth and young adults with type 2 diabetes; these values increased from 8.5% (70 mmol/mol) and 8.4% (68 mmol/mol), respectively, in 2002–2007 (15).

Youth and young adults with type 1 diabetes who are non-Hispanic Black or Native American (compared with those who are non-Hispanic White), younger, not treated with insulin pump therapy, and from households with low annual income have significantly higher A1C levels (15). Data from a network of academic pediatric and adult centers across the U.S. revealed that between 2016 and 2018, mean A1C was 8.1% (65 mmol/mol) among children with type 1 diabetes at 5 years of age and 9.3% (78 mmol/mol) among children 15–18 years of age. Only 17% of youth under 18 years of age with type 1 diabetes achieved the recommended A1C goal of $\leq 7.5\%$ (≤ 58 mmol/mol), and A1C levels for non-Hispanic Black youth were higher than those for non-Hispanic or Hispanic White youth, a disparity that persisted after adjustment for socioeconomic status (16).

Diabetes and its associated health complications pose a significant financial hardship to individuals and society. It is estimated that the annual cost of diagnosed diabetes in the U.S. in 2022 was $\geq \$413$ billion, including $\geq \$307$ billion in direct health care costs and $\geq \$106$ billion in reduced productivity (17). After adjusting for inflation, the economic costs of diabetes increased by 7% between 2017 and 2022 and by 35% between 2012 and 2022 (17). This is attributed to both the increased prevalence of diabetes and the higher cost per person with diabetes. People living with diabetes also face individual financial hardship, resulting in rationing or not taking prescribed medications, delaying care, and foregoing or rationing other expenditures, among other behaviors, and ultimately correlating with higher A1C, greater diabetes distress, and depressive symptoms (18).

The growing gaps in diabetes care quality and outcomes, the high and rising costs of diabetes care across the U.S., and the disparities experienced by individuals from racial and ethnic minoritized backgrounds and those facing socioeconomic barriers to care call for urgent, substantial, and multisectoral system-level improvements to care delivery (19).

Evidence-Based Care Delivery and Payment Models to Improve Population Health

A major barrier to comprehensive high-quality diabetes care is a delivery system that is often fragmented, lacks clinical information capabilities, is not appropriately incentivized and funded, does not adequately engage people with diabetes and the communities where they live, and is poorly designed for the coordinated and longitudinal delivery of chronic care (20). Several models have been demonstrated to improve aspects of diabetes care delivery and health outcomes.

The Chronic Care Model (CCM) is a commonly used framework for describing diabetes care programs (21). It includes six core elements to optimize the care of people with chronic disease:

1. Delivery system design (moving from a reactive to a proactive care delivery system where planned visits are coordinated through a team-based approach)
2. Self-management support
3. Decision support, particularly at the point of care during a clinical encounter (basing care on evidence-based, effective care guidelines)
4. Clinical information systems (using registries that can provide person-specific and population-based support to the care team)
5. Community resources and policies (identifying or developing resources to support healthy lifestyles)
6. Health systems (to create a quality-oriented culture)

Randomized controlled trials of CCM interventions have shown that while interventions vary, programs that include core components of the CCM decrease A1C (mean difference [MD] -0.21% [95% CI -0.30 to -0.13], $P < 0.001$ compared with usual care), with greater improvements seen among adults with higher baseline A1C and with interventions that include four or more CCM elements (22). CCM-aligned programs also improved blood pressure levels and processes of diabetes care (e.g., screening for complications of diabetes), though there was no impact on cholesterol levels, tobacco use, or weight (23).

Multiple studies have examined individual components of the CCM with respect to diabetes management and have found inconsistent levels of benefit with case management, team-based care, use of electronic patient registries, clinician education, clinician and patient reminders, and patient education and promotion of individual

self-management (24). The inconsistencies in findings may be driven by heterogeneity of interventions, settings, and evaluation strategies as well as structural barriers to program utilization and needs of communities served.

Collaborative, interprofessional teams, which can bring together multiple disciplines within the health care system, payors, and community partners, are best suited to provide care for people with chronic conditions such as diabetes and to facilitate individuals' self-management (Table 1.1) (25–27). The care team, which centers around the person with diabetes, should avoid therapeutic inertia and prioritize timely and appropriate modification of behavior change interventions (nutrition and physical activity), pharmacologic therapy, technology use, and/or social and financial support systems for individuals who have not achieved recommended individualized metabolic goals or are experiencing high burden of treatment. Engagement of an endocrinology team with expertise in diabetes management is important in situations when the complexity of diabetes.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded

Improving Care and Promoting Health in Populations

Diabetes Care Volume 49, Supplement 1, January 2026

Table 1.1—Considerations for engaging interprofessional members of a comprehensive, person-centered diabetes care team to identify and meet the needs of people with diabetes across the life span

Subpopulation of a person with diabetes	Team members to engage in care	Unique care considerations
1. Type 1 diabetes 2. Type 2 diabetes 3. Gestational diabetes 4. Prediabetes 5. Diabetes in children 6. Diabetes in older adults 7. Diabetes in people with other chronic conditions 8. Diabetes in people with mental health conditions 9. Diabetes in people with limited health literacy 10. Diabetes in people with limited access to care	1. Endocrinologist 2. Primary care provider 3. Nurse practitioner 4. Registered dietitian 5. Diabetes educator 6. Pharmacist 7. Social worker 8. Behavioral health specialist 9. Community health worker 10. Care coordinator	1. Individualized care plan 2. Patient education and self-management support 3. Regular monitoring and assessment 4. Cultural and linguistic appropriateness 5. Addressing social determinants of health 6. Patient and family engagement 7. Interdisciplinary collaboration 8. Telehealth and remote monitoring 9. Patient and family education 10. Patient and family support

All adults with diabetes | Primary care clinician, CDCES, RDN, and other specialists as available and appropriate to treat comorbidities (Table 4.1) | Assess for and address social determinants of health. | | Adults treated with intensive insulin therapy, including multiple daily injections of insulin and insulin pump therapy | Clinicians and other health care team members experienced in advanced diabetes management, including technology use | - | | All children and adolescents with diabetes | Primary care clinician, pediatric endocrinologist, CDCES, RDN, other specialists as available and appropriate to treat comorbidities (Table 14.1), daycare or school nurse or other professional, behavioral health professional (as needed), and parent(s) or caregiver(s) | Assess for and address social determinants of health and barriers to safety, well-being, and academic performance in school. Engage professionals within the school and extracurricular/after-school activities to ensure safe diabetes management. An individualized diabetes medical management plan should be developed in collaboration with school professionals and parent(s) or caregiver(s). Support gradual developmentally appropriate transfer of self-management from caregivers to the children and adolescents with diabetes. | | Individuals with diabetes and diabetes-related complications or comorbidities | Specialist referrals as appropriate and available (e.g., behavioral health professional, cardiologist, endocrinologist, eye specialist, gastroenterologist or hepatologist, neurologist, nephrologist, obesity medicine specialist, or podiatrist), care coordinator/navigator or case manager, and clinical pharmacist (for those with polypharmacy or complex medication plans) | Screen for functional, cognitive, financial, and logistical barriers to self-management and evidence that self-care demands exceed capacity and available resources and support systems. | | Individuals with social and/or structural barriers to care | Care coordinator/navigator, social services professional, insurance specialist/navigator, peer-to-peer support (as available), community health worker and/or community paramedic (as available), public health professional, and interpreter (as applicable) | Consider each person's psychosocial needs, available resources, and support systems. | | Older adults | Geriatric medicine specialist, clinical pharmacist (for those with polypharmacy or complex medication plans), CDCES and/or RDN with expertise in the care for older adults, social services professional, case manager, community services provider, and physical and/or occupational therapist as available and appropriate based on functional status and independence | Consider the older adult's nutritional status, including ability to afford (financial barriers), acquire

(accessibility), prepare (cooking), and consume (oral health) nutritious food. Assess for and address needs related to vision, hearing, dexterity, cognition, mobility, and other challenges. | Individuals in long-term care settings | Long-term care facility clinicians, nurses, other health care professionals, physical and occupational therapists, and RDN | Engage professionals within the long-term care facility to ensure safe and appropriate diabetes management. | Pregnant individuals with diabetes | Maternal-fetal medicine specialist or obstetrician experienced in the care of pregnant individuals with diabetes (particularly for individuals with type 1 diabetes or requiring intensive insulin therapy), CDCES, RDN, eye specialist (particularly for individuals with preexisting type 1 or type 2 diabetes), other specialists as appropriate, and lactation consultant as appropriate | Ensure appropriate postpartum follow-up and care, including transition from obstetric care to established primary care. | Individuals with behavioral health conditions | Behavioral health professional, care coordinator/navigator, and social services professional as age and situation appropriate | Use age- and situation-appropriate screening protocols for general and diabetes-related psychosocial concerns. |

CDCES, certified diabetes care and education specialist; RDN, registered dietitian nutritionist.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf Downloaded

diabetesjournals.org/care # Improving Care and Promoting Health in Populations

treatment exceeds the capacity of the pri- and DSMES services for individuals with telehealth has demonstrated effective- ness in reducing disparities and improv- ing glycemic outcomes in racially and ethnically diverse adults with type 2 dia- betes. A systematic review and meta- analysis of nine telehealth randomized controlled trials aimed at improving A1C among Black and Hispanic adults with type 2 diabetes (>50% sample) found a -0.47% (95% CI -0.65% to -0.28%) change in A1C (60). While this systematic review and meta-analysis reported im- proved glycemic outcomes in marginalized populations, these successes typically de- pend on substantial investments in infra- structure (laptops, mobile phone access, internet access) and targeted support (digi- tal literacy education, community health workers [CHWs]) that are rarely part of routine care (61). Without this infrastruc- ture and support, attrition is high. Re- search on technology and inequities highlights that the digital divide is driven by financial (cost of data plans, high- speed internet), informational (numeracy, literacy, digital skills), and technical (inter- operability gap, complex user interfaces) barriers (62). Therefore, effective tele- health design must consider accessibility, affordability, connectivity, trust, and con- textual tailoring (62). Failure to do so risks exacerbating existing inequities.

Successful diabetes care also requires a systematic approach to supporting the be- havior change efforts of people with diabe- tes. High-quality DSMES has been shown to improve a person's self-management, satisfaction, and glycemic outcomes (see section 5, "Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes," for a detailed review of the evidence supporting DSMES). Na- tional DSMES standards call for an inte- grated approach that includes clinical content and skills, behavioral strategies (goal setting, problem-solving, etc.), and engagement with psychosocial concerns (63). Increasingly, such support is available through online or mobile platforms that can support user access and effective- ness. Findings from a 2025 systematic re- view and meta-analysis of 43 randomized controlled trials comparing DSME, diabe- tes self-management support, or DSMES delivered via mHealth interventions ver- sus usual care or attention placebo con- trol showed modest A1C benefits: DSME MD -0.3% (95% CI -0.6% to -0.1% ; $P = 0.002$), diabetes self-management sup- port MD -0.4% (95% CI -0.6% to -0.2% ;

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded

Improving Care and Promoting Health in Populations

Diabetes Care Volume 49, Supplement 1, January 2026

$P \leq 0.001$), and DSMES MD -0.2% (95% CI -0.3% to -0.0% ; $P \leq 0.001$) (64). These curricula should be tailored to the needs of their intended populations, including demographic characteristics, diabetes knowledge, emotional support, and cultural beliefs, and address the digital divide, i.e., access to the technology required for implementation (54,65,66).

Strategies for System-Level Improvement

Optimal diabetes management requires a systematic approach and coordinated team of health care professionals working in an environment where person-centered, high-quality care is a priority (23,67–69). While many diabetes care processes and access to technologies have improved nationally in the past decade, the overall quality of care for people with diabetes remains suboptimal (13). Efforts to increase the quality of diabetes care include providing care that is concordant with evidence-based guidelines (70), expanding the role of teams to implement more intensive disease management strategies (71), tracking medication-taking behavior (70), redesigning care processes (71), implementing electronic health record (EHR) population health tools (72), empowering and educating people with diabetes (73), reducing financial barriers (74), leveraging telehealth to improve access to care (51), assessing and addressing psychosocial issues (63,75), and engaging community resources and public policies that support healthy lifestyles (76). The National Diabetes Education Program maintains an online resource (cdc.gov/diabetes/php/toolkits/index.html) to help health care professionals design and implement more effective health care delivery systems for people with diabetes.

Given the pluralistic needs of people with diabetes and the challenges they experience (complex insulin treatment plans, new technologies, changes in capacity for self-management, etc.) vary over the course of disease management and life span, engagement of an interprofessional team with complementary expertise is essential (25). Findings from a systematic review and meta-analysis of 35 team-based care interventions with adults with type 1 or type 2 diabetes showed a -0.5% (95% CI -0.7 to -0.3) change in A1C, -5.5 mmHg (95% CI -8.1 to -3.0) change in systolic blood pressure, -3.2 mmHg (95% CI -4.8 to -1.5) change in diastolic blood pressure, -8.0 mg/dL (95% CI -11.8 to -4.3) change in LDL cholesterol, and -7.4 mg/dL (95% CI -13.9 to -0.9) change in total cholesterol (25). Team-based care has also been shown to reduce atherosclerotic cardiovascular disease (ASCVD) risks. Results from the Center for Integrated and Novel Approaches in Vascular-Metabolic Disease (CINEMA) Program demonstrated improvements in ASCVD risk factors and evidence-based therapies with team-based care for adults with type 2 diabetes or prediabetes via reductions in body weight (-5.5 lb [-2.5 kg]), BMI (-0.9 kg/m²), systolic (-3.6 mmHg) and diastolic (-1.2 mmHg) blood pressure, A1C (-0.5%), LDL (-9.0 mg/dL) and total cholesterol (-10.7 mg/dL), triglycerides (-13.5 mg/dL), and absolute 10-year ASCVD risk (-2.4%) (77).

Access to Care

The Affordable Care Act and Medicaid expansion have increased access to care for many individuals with diabetes, emphasizing the protection of people with preexisting conditions, health promotion, and disease prevention (86). In 2024, population-level data for the U.S. civilian noninstitutionalized population included in the National Health Interview Survey (NHIS) showed that just 4.9% of children and adolescents (aged 17 years and younger), 11.1% of adults aged 18–64 years, and 0.6% of adults aged 65 years and older were uninsured (87). People with diabetes who have consistent private or public insurance coverage (i.e., are not uninsured) are more likely to meet quality indicators for diabetes care (86). However, even individuals with insurance coverage can experience financial barriers to care, particularly if enrolled in high-deductible health plans. In 2021, 28% of individuals with employer-sponsored health plans were enrolled in high-deductible health plans (88). Such plans are increasing in popularity; by 2023, 51% of private industry employees had the option to enroll in a high-deductible health plan, although only 36% had access to health savings accounts, which can offset some of the out-of-pocket costs incurred with high-deductible plans (89). Switching to a high-deductible health plan has been shown to increase financial hardship among people with diabetes (90), decrease and delay screening for retinopathy (91), decrease monitoring of blood pressure and A1C (91), and increase the risks of experiencing both acute (severe hypoglycemia, hyperglycemic crises) (92) and chronic (myocardial infarction, stroke, hospitalization for heart disease) complications.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded

diabetesjournals.org/care # Improving Care and Promoting Health in Populations

failure, kidney failure, lower-extremity com- complications, proliferative retinopathy, and blindness) (93) diabetes complications. However, health plan adoption of preventive drug lists that reduce or eliminate out-of-pocket cost sharing for a wide range of medications used to treat chronic conditions like diabetes reduces out-of-pocket expenditures for people with diabetes and narrows gaps in diabetes care (94). Insurance coverage and formulary design influence treatment decisions; it is essential that payors cover evidence-based diabetes care with minimal cost sharing by people with diabetes. Health care teams should also discuss

insurance coverage and financial barriers to care with all individuals with diabetes and pursue therapeutic strategies that minimize financial hardship (74).

Access to primary and specialty care is essential for people with diabetes. While most adults with diabetes have access to a primary care clinician—a 2016 nationally representative population-based study found that 88% of adults with diabetes saw a primary care clinician in the prior year (95)—fewer have access to specialty endocrinology/diabetes care (96). A study of Medicare beneficiaries found that just 33% of older adults with type 1 diabetes, 14% of adults with type 2 diabetes and history of severe hypoglycemia, and 9% of other adults with type 2 diabetes saw an endocrinologist in 2019 (96). Racial and ethnic minoritized individuals, those with low income, those living in rural areas, and those residing in a long-term care facility were less likely to receive endocrinology care. Improving health outcomes for people with diabetes will therefore require improving availability of and access to primary and specialty services, and interdisciplinary care from the full diabetes care team, to meet the full range of their health care needs (Table 1.1).

COST CONSIDERATIONS

The cost of diabetes medications and devices is an ongoing barrier to achieving glycemic goals and preventing diabetes complications. Medication underuse due to cost has been termed “cost-related medication nonadherence” (here referred to as cost-related barriers to medication use). Based on a national survey conducted in 2021, 18.6% of U.S. adults with type 1 diabetes and 15.8% of adults with insulin-treated type 2 diabetes reported rationing (i.e., skipping, taking less, and/or delaying) their insulin to save money (97). The ADA Insulin Access and Affordability Working Group has recommended system-level approaches to address this issue, including concepts such as cost-sharing for insured people with diabetes based on the lowest price available, a list price for insulins that closely reflects the net price, and health plans that ensure people with diabetes can access insulin without undue administrative burden or excessive cost (98).

In 2021, the Centers for Medicare & Medicaid Services (CMS) launched the Part D Senior Savings Model (99), which required capped monthly out-of-pocket payments for covered insulins at \$35 per insulin per month. In 2022, 43% of stand-alone Part D plan enrollees and 60% of Medicare Advantage Part D plan enrollees participated in the Senior Savings Model (99). Most recently, the Inflation Reduction Act of 2022 capped out-of-pocket payments for insulin at \$35 per insulin per month for all Medicare beneficiaries. A patchwork of solutions has also been introduced for individuals with commercial insurance and those without health insurance. Over the past 5 years, 25 states and the District of Columbia have capped out-of-pocket expenditures for insulin in select state-regulated commercial health plans (100), which has resulted in modest decreases in insulin out-of-pocket spending for eligible enrollees (101).

Between 2023 and 2024, three major insulin manufacturers similarly lowered the price of insulin to \$35 per month in select circumstances (102). These programs may help reduce the financial hardship of diabetes management, though many are challenging to navigate, not all people with diabetes can benefit, and costs for insulin delivery and glucose monitoring remain high. Thus, all people with diabetes should be screened for financial hardship of treatment, cost-related barriers to medication use, and rationing of other essential services due to medical costs (103).

Health inequities related to diabetes and its complications are well documented, are heavily influenced by SDOH, and have been associated with greater risk for developing diabetes, higher disease prevalence, and worse diabetes-related outcomes (4). Greater exposure to adverse SDOH over the life course results in poor health (113). Interventions to address SDOH can improve diabetes-related outcomes (114,115). Using clinical quality data to identify inequities.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf Downloaded

S20 Improving Care and Promoting Health in Populations Diabetes Care Volume 49, Supplement 1, January 2026

and opportunities for improvement is disparities in populations is yet to be for- quality or health outcomes in people with valuable for health care professionals, mally studied, general recommendations diabetes (116,128). health systems, payors, policymakers, and from other chronic disease management A population in which such issues must people with diabetes (116). The Joint Com- and prevention models can be drawn be particularly

considered is older adults, mission requires that all accredited organizations in its ambulatory health care, diabetes (62). For example, the National Academy of Medicine has published a risk of functional dependency (129) (see critical access hospital, and hospital accreditation programs collect race and ethnicity professionals on the importance of discussion of social considerations in information and implement specific steps SDOH (123). Furthermore, there are re-older adults). to reduce health care disparities. The Joint sources available for the inclusion of standardized sociodemographic variables in screen for SDOH may help overcome organizations designate an individual or in-EHRs to facilitate the assessment of health structural barriers and communication individuals to lead efforts to reduce health disparities and the impact of intervention gaps between people with diabetes and care disparities, assess health-related social needs and provide information on community resources to meet these needs, SDOH are not consistently recognized effectiveness of identifying SDOH by using validated screening tools (131). In addition, quality and safety data using sociodemographic characteristics, develop an action plan to address health care disparities, care and health, which directly affect the around factors that significantly impact work to actively reduce health care disparities, and inform key stakeholders plans, are rarely discussed. Among people with chronic illnesses, two-thirds of Food Insecurity parities (117). The CMS Framework for those who reported not taking medication as prescribed due to cost-related economic and social condition of limited or uncertain access to adequate food (132). individual-level demographic (including race, with their physician (123). A study using In 2022, almost 13% of Americans were ethnicity, language, gender identity, sex, sexual orientation, and disability status) and adults with diabetes reported financial associated with increased risk of type 2 SDOH data as well as assessing for and addressing stress and about 20% reported food insecurity (126). Studies of both type 1 diabetes and type 2 diabetes have noted an association with higher rates of depression and diabetes outcomes among individuals with diabetes (118). Quality measures assessing SDOH screening and SDOH with health care utilization and income households, and households headed by single mothers. Additionally, those with diabetes (123,127). It is there- ing food insecurity have lower engagement (focused on food, housing, and transportation insecurity) (113) and CMS (focused on be screened for SDOH during clinical encounters and be referred to appropriate services distress, and have worse glycemic management, utility difficulties, and interpersonal clinical and community resources to address these needs (Table 1.1). Furthermore, are food secure (133,134). Older adults Other than the SDOH, there are many health systems may benefit from compiling with food insecurity are more likely to have contributors to inequities, including implicit and explicit bias, discriminatory policies and procedures, institutional practices, as necessary existing tools such as find-help.org, Unite Us, 211 (United Way), and risk for food insecurity can be assessed factors that affect health and well-being for individuals and communities (120–122). others. Policies and payment models that with a validated two-item screening tool The ADA recognizes the association between interpersonal, social, economic, and and outside the health care setting, are 1) “Within the past 12 months, we worried environmental factors and the prevention needed to ensure that these efforts are whether our food would run out before we and treatment of diabetes and has issued both feasible and sustainable. One example got money to buy more” and 2) “Within a call for research that seeks to better understand how social determinants influence incentivizes value-based care, addressing just didn’t last, and we didn’t have

money behaviors and how the relationships between SDOH and funding community-based health to get more” (137). Interventions such as between these variables can be modified for care professionals, is the Maryland Total food prescription programs are considered enhancing the prevention and management Cost of Care Model, although it is currently considered promising to address food insecurity of diabetes (114). While a comprehensive limited by a narrow focus on preventing diabetes by integrating community resources into strategy to reduce diabetes-related health outcomes and does not consider diabetes care primary care settings and directly dealing

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded

diabetesjournals.org/care # Improving Care and Promoting Health in Populations

with food deserts in underserved communities (135).

In those with diabetes and food insecurity, the priority is mitigating the increased risk for severe hyperglycemia and hypoglycemia (138,139). The reasons for the increased risk of hyperglycemia can include the consumption of inexpensive carbohydrate-rich processed foods, binge eating, financial constraints to filling diabetes medication prescriptions, anxiety and depression, and poor sleep, all contributing to hyperglycemia and poor diabetes self-care behaviors. Hypoglycemia can occur as a result of inadequate or inconsistent carbohydrate consumption following the administration of a sulfonylurea or insulin. Health care professionals should consider these factors when making treatment decisions for people with food insecurity and seek local resources to help people with diabetes and their family members obtain appropriate food more regularly (140).

Housing Insecurity

Housing insecurity has been shown to be directly associated with a person's ability to maintain appropriate diabetes self-management (141). Housing insecurity often coexists with other barriers that challenge diabetes self-management. Food insecurity, lack of health insurance, cognitive impairment, behavioral health concerns, and low literacy and numeracy skills are also factors (140). The prevalence of diabetes among people experiencing housing insecurity is estimated to be around 8% in the U.S. (142). Additionally, people with diabetes and housing insecurity need secure places to keep their diabetes medications and supplies as well as refrigerator access to safely store insulin. The risk for housing insecurity can be ascertained using a brief risk assessment tool developed and validated for use among veterans (143). Given the potential challenges, health care professionals who care for housing-insecure individuals should be familiar with resources to support these individuals or have access to social workers who can facilitate stable housing as a way to improve diabetes care (144).

Refugee Populations and Migrant and Seasonal Agricultural Workers

Refugee status, like having a diabetes diagnosis, is an independent risk factor for cardiovascular disease (145). In areas experiencing humanitarian crises, refugees are at greater risk for obstacles to optimal chronic disease management, but unfortunately, there are few quality investigations into diabetes care and its outcomes among refugees. There have been efforts to develop models of care specifically aimed at improving the health of refugee populations, but more work is needed to demonstrate effectiveness of those care models and approaches (146). Migrant and seasonal agricultural workers likely have a higher risk of type 2 diabetes than the general population. While migrant farmworker-specific data are lacking, most agricultural workers in the U.S. are Latino, a population with a high rate of type 2 diabetes. In addition, living in severe poverty brings with it food insecurity, high chronic stress, and an increased risk of diabetes; there is also an association between the exposure to certain pesticides and the incidence of diabetes (147).

Data from the Department of Labor indicate that there are approximately 2.18 million agricultural workers in the U.S. (148). These agricultural workers often travel throughout the country seasonally (149). In 2022, 175 health centers across the U.S. reported providing care to 843,071 adult migrant farmworkers, and 91,839 had encounters for diabetes (147). In a 2023 report on the National Agricultural Workers Survey, age-adjusted self-reported diabetes prevalence was 13.51% (95% CI 10.0–17.1) among migrant farmworkers and 10.8% (95% CI 9.0–12.6) among nonmigrant farmworkers (147). Migrant farmworkers and other agricultural workers encounter numerous and overlapping barriers to receiving care. Migration, which might occur as frequently as every few weeks for some, disrupts care. Common barriers to adequate diabetes care include those related to cost, culture, language, literacy, transportation, geographic distance, food access, long work hours, unfamiliarity

with new communities, the complexity of the U.S. health care system, and limited access to various other resources like medications and DSMES (149). Without regular care, farmworkers with diabetes can experience severe and often expensive complications that incur morbidity and mortality and affect quality of life. Nontraditional care delivery models, including mobile integrated health and telehealth, should be leveraged to improve access to care.

Language Barriers

Health systems and health care professionals caring for those with limited English proficiency should develop or offer educational programs and materials in culturally appropriate languages. Professional language assistance (i.e., interpreters) should be provided to individuals with limited English proficiency and/or other communication needs at no cost to them (150). Use of untrained interpreters, including family members, should be avoided when possible, as this can result in confusing or inaccurate conveyance of information. Accompanying written materials should be in the language appropriate for the individual being supported and at a reading level that is not overly complicated—typically this is defined as a sixth-grade reading level. Numerous medical translation tools are available and show promise in addressing language barriers, although further research is needed to establish their efficacy in diabetes care (151).

The National Standards for Culturally and Linguistically Appropriate Services in Health and Health Care (National CLAS Standards) provide guidance on how health care professionals can reduce language barriers by improving their cultural competency, addressing health literacy, and ensuring communication with professional language assistance (150). In addition, the National CLAS Standards website offers several resources and materials that can be used to improve the quality of care delivery to individuals with limited English proficiency.

Health Literacy and Numeracy

Health literacy is the degree to which individuals can obtain, process, and understand basic health information and services needed to make appropriate decisions (152,153). Health literacy is strongly associated with individuals engaging in complex disease management and self-care (154). Approximately 9 out of 10 American adults are estimated to have limited or low health literacy (152,155). Health care professionals and diabetes care and education specialists should provide easy-to-understand information.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf Downloaded

Improving Care and Promoting Health in Populations

Diabetes Care Volume 49, Supplement 1, January 2026

Information and reduce unnecessary complexity when developing care plans. Interventions addressing low health literacy in populations with diabetes demonstrate improvements in A1C, diabetes knowledge, and self-management behaviors. However, more research is needed to establish the most effective strategies for enhancing retention and application of diabetes knowledge among various populations of people with diabetes. Health numeracy is also essential in diabetes prevention and management. Health numeracy requires primary numeric skills, applied health numeracy, and interpretive health numeracy, which is especially important for people using diabetes technologies like insulin pumps. People with prediabetes or diabetes often need to perform numeric tasks, such as interpreting glucose readings, decoding food labels, and understanding medication doses, timings, and refills, to make treatment decisions. Thus, both health literacy and numeracy are necessary for enabling effective communication between people with diabetes and health care professionals, arriving at a treatment plan, and making diabetes self-management task decisions. If people with diabetes appear not to understand concepts associated with treatment decisions, both can be assessed using standardized screening measures. Adjunctive education and support may be indicated if limited health literacy and numeracy are barriers to optimal care decisions.

Social and Community Support

Social support, which comprises community and personal network instrumental support, promotes better health, whereas lack of social support is associated with poorer health outcomes in individuals with diabetes. Of particular concern are the social determinants of health (SDOH), including, among others, racism and

discrimination. These factors are rarely addressed in routine clinical practice but may be drivers of adverse health outcomes and lower engagement in beneficial self-care behaviors and medication use. Identifying and leveraging community resources are core components of chronic care management.

Various organizations, including the American Medical Association and the Agency for Healthcare Research and Quality, among others, promote the translation of clinical recommendations for nutrition-based diabetes care recommendations.

References

1. Silberberg M, Martinez-Bianchi V, Lyn MJ. What is population health? *Prim Care* 2019;46:475–484
2. Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med* 2014;12:573–576
3. Institute of Medicine (US) Committee on Quality of Health Care in America. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington, DC, National Academies Press, 2001
4. Hill-Briggs F, Adler NE, Berkowitz SA, et al. Social determinants of health and diabetes: a scientific review. *Diabetes Care* 2020;44:258–279
5. Haire-Joshu D, Hill-Briggs F. The next generation of diabetes translation: a path to health equity. *Annu Rev Public Health* 2019;40:391–410
6. Centers for Disease Control and Prevention. *National Diabetes Statistics Report*. Accessed 4 August 2025. Available from <https://www.cdc.gov/diabetes/php/data-research/index.html>
7. Centers for Disease Control and Prevention. *Adult Obesity Facts*. Accessed 4 August 2025. Available from <https://www.cdc.gov/obesity/adult-obesity-facts/index.html>
8. Lawrence JM, Divers J, Isom S, et al.; SEARCH for Diabetes in Youth Study Group. Trends in prevalence of type 1 and type 2 diabetes in children and adolescents in the US, 2001–2017. *JAMA* 2021;326:717–727
9. Centers for Disease Control and Prevention. *Childhood Obesity Facts*. Accessed 4 August 2025. Available from <https://www.cdc.gov/obesity/childhood-obesity-facts/childhood-obesity-facts.html>
10. Saelee R, Bullard KM, Hora IA, et al. Trends and inequalities in diabetes-related complications among U.S. Adults, 2000–2020. *Diabetes Care* 2025;48:18–28
11. Ebekozi O, Mungmode A, Sanchez J, et al. Longitudinal trends in glycemic outcomes and technology use for over 48,000 people with type 1 diabetes (2016–2022) from the T1D Exchange Quality Improvement Collaborative. *Diabetes Technol Ther* 2023;25:765–773
12. Fang M, Wang D, Coresh J, Selvin E. Trends in diabetes treatment and control in U.S. adults, 1999–2018. *N Engl J Med* 2021;384:2219–2228
13. Nassereldine H, Li Z, Compton K, et al. The burden of diabetes mortality by county, race, and ethnicity in the U.S., 2000–2019. *Diabetes Care* 2025;48:546–555
14. Chiang JL, Maahs DM, Garvey KC, et al. Type 1 diabetes in children and adolescents: a position statement by the American Diabetes Association. *Diabetes Care* 2018;41:2026–2044
15. Malik FS, Sauder KA, Isom S, et al.; SEARCH for Diabetes in Youth Study. Trends in glycemic control among youth and young adults with diabetes: the SEARCH for Diabetes in Youth Study. *Diabetes Care* 2022;45:285–294
16. Foster NC, Beck RW, Miller KM, et al. State of type 1 diabetes management and outcomes from the T1D Exchange in 2016–2018. *Diabetes Technol Ther* 2019;21:66–72

2026 January 29 on guest by Download

diabetesjournals.org/care Improving Care and Promoting Health in Populations S23

1. Parker ED, Lin J, Mahoney T, et al. Economic costs of diabetes in the U.S. in 2022. *Diabetes Care* 2024;47:26–43
2. Patel MR, Zhang G, Heisler M, et al. Measurement and validation of the Comprehensive Score for Financial Toxicity (COST) in a population with diabetes. *Diabetes Care* 2022;45:2535–2543
3. Prahalad P, Hardison H, Odugbesan O, et al.; T1D Exchange Quality Improvement Collaborative. Benchmarking diabetes technology use among 21 U.S. pediatric diabetes centers. *Clin Diabetes* 2024;42:27–33
4. Chehal PK, Selvin E, DeVoe JE, Mangione CM, Ali MK. Diabetes and the fragmented state of US health care and policy. *Health Aff (Millwood)* 2022;41:939–946
5. Stellefson M, Dipnarine K, Stopka C. The chronic care model and diabetes management in US primary care settings: a systematic review. *Prev Chronic Dis* 2013;10:E26
6. Goh LH, Siah CJR, Tam WWS, Tai ES, Young DY. Effectiveness of the chronic care model for adults with type 2 diabetes in primary care: a systematic review and meta-analysis. *Syst Rev* 2022;11:273
7. Tricco AC, Ivers NM, Grimshaw JM, et al. Effectiveness of quality improvement strategies on the management of diabetes: a systematic review and meta-analysis. *Lancet* 2012;379:2252–2261
8. Konnyu KJ, Yogasingam S, Lépine J, et al. Quality improvement strategies for diabetes care: effects on outcomes for adults living with diabetes. *Cochrane Database Syst Rev* 2023;5:CD014513
9. Levensgood TW, Peng Y, Xiong KZ, et al.; Community Preventive Services Task Force. Team-based care to improve diabetes management: a

community guide meta-analysis. *Am J Prev Med* 2019;57:e17–e26 10. Lee JK, McCutcheon LRM, Fazel MT, Cooley JH, Slack MK. Assessment of interprofessional collaborative practices and outcomes in adults with diabetes and hypertension in primary care: a systematic review and meta-analysis. *JAMA Netw Open* 2021;4:e2036725 11. Oshman L, Bhomia N, Diez HL, et al. The Michigan Collaborative for Type 2 Diabetes (MCT2D): development and implementation of a statewide collaborative quality initiative. *BMC Health Serv Res* 2024;24:1254 12. Avnat E, Chodick G, Shalev V. Identifying profiles of patients with uncontrolled type 2 diabetes who would benefit from referral to an endocrinologist. *Endocr Pract* 2023;29:855–861 13. Setji TL, Page C, Pagidipati N, Goldstein BA. Differences in achieving Hba1c goals among patients seen by endocrinologists and primary care providers. *Endocr Pract* 2019;25:461–469 14. McCoy RG, Vandergrift JL, Gray B. Patient and physician factors driving the gaps in use of drugs with cardiovascular and kidney benefits by Medicare beneficiaries with type 2 diabetes treated by endocrinologists, nephrologists, and cardiologists: population-based cohort study. *Diabetes Res Clin Pract* 2025;221:112039 15. Phan T-L, Hossain J, Lawless S, Werk LN. Quarterly visits with glycated hemoglobin monitoring: the sweet spot for glycemic control in youth with type 1 diabetes. *Diabetes Care* 2014;37:341–345 16. Parker ED, Lin J, Mahoney T, et al. Economic costs of diabetes in the U.S. in 2022. *Diabetes Care* 2024;47:26–43 17. Booth GL, Shah BR, Austin PC, Hux JE, Luo J, Lok CE. Early specialist care for diabetes: who benefits most? A propensity score-matched cohort study. *Diabet Med* 2016;33:111–118 18. Adler-Milstein J, Linden A, Hollingsworth JM, Ryan AM. Association of primary care engagement in value-based reform programs with health services outcomes: participation and synergies. *JAMA Health Forum* 2022;3:e220005 19. Bojadzievski T, Gabbay RA. Patient-centered medical home and diabetes. *Diabetes Care* 2011;34:1047–1053 20. McManus LS, Dominguez-Cancino KA, Stanek MK, et al. The patient-centered medical home as an intervention strategy for diabetes mellitus: a systematic review of the literature. *Curr Diabetes Rev* 2021;17:317–331 21. Frazee TK, Lewis VA, Tierney E, Colla CH. Quality of care improves for patients with diabetes in Medicare shared savings accountable care organizations: organizational characteristics associated with performance. *Popul Health Manag* 2018;21:401–408 22. Centers for Medicare & Medicaid Services. Accountable Health Communities Model. Accessed 8 August 2025. Available from <https://www.cms.gov/priorities/innovation/innovation-models/ahcm> 23. Centers for Medicare & Medicaid Services. Accountable Health Communities (AHC) Model Evaluation. Second Evaluation Report Accessed 8 August 2025. Available from <https://www.cms.gov/priorities/innovation/data-and-reports/2023/ahc-second-eval-rpt> 24. Wang S, Weyer G, Duru OK, Gabbay RA, Huang ES. Can alternative payment models and value-based insurance design alter the course of diabetes in the United States? *Health Aff (Millwood)* 2022;41:980–984 25. Gunter KE, Peek ME, Tanumihardjo JP, et al. Population health innovations and payment to address social needs among patients and communities with diabetes. *Milbank Q* 2021;99:928–973 26. Katula JA, Dressler EV, Kittel CA, et al. Effects of a digital diabetes prevention program: an RCT. *Am J Prev Med* 2022;62:567–577 27. Eberle C, Stichling S. Clinical improvements by telemedicine interventions managing type 1 and type 2 diabetes: systematic meta-review. *J Med Internet Res* 2021;23:e23244 28. American Telemedicine Association. Telehealth: defining 21st Century Care. Accessed 8 August 2025. Available from <https://www.americantelemed.org/resource/why-telemedicine/> 29. Health IT. Telehealth StartUp and Resource Guide Version 1.1. Accessed 8 August 2025. Available from https://www.healthit.gov/sites/default/files/telehealthguide_final_0.pdf 30. American Medical Association. AMA Telehealth Quick Guide. Accessed 8 August 2025. Available from <https://www.ama-assn.org/practice-management/digital/ama-telehealth-quick-guide> 31. Zupa MF, Vimalananda VG, Rothenberger SD, et al. Patterns of telemedicine use and glycemic outcomes of endocrinology care for patients with type 2 diabetes. *JAMA Netw Open* 2023;6:e2346305 32. Mullur RS, Hsiao JS, Mueller K. Telemedicine in diabetes care. *Am Fam Physician* 2022;105:281–288 33. Ravi S, Meyerowitz-Katz G, Yung C, et al. Effect of virtual care in type 2 diabetes management—a systematic review and meta-analysis. *J Racial Ethn Health Disparities* 2022;9:2375–2386 34. Mayberry LS, Lyles CR, Oldenburg B, Osborn CY, Parks M, Peek ME. mHealth interventions for disadvantaged and vulnerable people with type 2 diabetes. *Curr Diab Rep* 2019;19:148 35. Ebekozien O, Fantasia K, Farrokhi F, Sabharwal A, Kerr D. Technology and health inequities in diabetes management.

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf Downloaded

Improving Care and Promoting Health in Populations

Diabetes Care Volume 49, Supplement 1, January 2026

diabetes care: how do we widen access to patient-centered, team-based intervention for underserved populations and utilize technology to improve outcomes for all? *Diabetes Obes Metab* 2024;26(Suppl. 1):3–13

| Available from <<https://www.bls.gov/ebs/factsheets/patients-with-type-2-diabetes-or-prediabetes-at-high-cardiovascular-risk.htm>> | |

----- | -----
----- | 63. | Davis J, Fischl AH, Beck J, et al. 2022 National Standards for Diabetes Self-Management Education and Support. *Diabetes Care* 2022;45:484–494 | | 64. | Versluis A, Boels AM, Huijden MCG, Mijnsbergen MD, Rutten GEHM, Vos RC. Diabetes self-management education and support delivered by mobile health (mHealth) interventions for adults with type 2 diabetes—a systematic review and meta-analysis. *Diabet Med* 2025;42:e70002 | | 65. | Tanhapour M, Peimani M, Rostam Niakan Kalhori S, et al. The effect of personalized intelligent digital systems for self-care training on type II diabetes: a systematic review and meta-analysis of clinical trials. *Acta Diabetol* 2023;60:1599–1631 | | 66. | Abu S, Llahana S. Factors influencing the uptake of culturally tailored diabetes self-management education and support programmes among ethnic minority patients with type 2 diabetes: a systematic review. *Prim Care Diabetes* 2025;19:103–110 | | 67. | TRIAD Study Group. Health systems, patients factors, and quality of care for diabetes: a synthesis of findings from the TRIAD study. *Diabetes Care* 2010;33:940–947 | | 68. | Schmittiel JA, Gopalan A, Lin MW, Banerjee S, Chau CV, Adams AS. Population health management for diabetes: health care system-level approaches for improving quality and addressing disparities. *Curr Diab Rep* 2017;17:31 | | 69. | Peterson KA, Carlin CS, Solberg LI, Normington J, Lock EF. Care management processes important for high-quality diabetes care. *Diabetes Care* 2023;46:1762–1769 | | 70. | Raebel MA, Schmittiel J, Karter AJ, Konieczny JL, Steiner JF. Standardizing terminology and definitions of medication adherence and persistence in research employing electronic databases. *Med Care* 2013;51:S11–S21 | | 71. | Feifer C, Nemeth L, Nietert PJ, et al. Different paths to high-quality care: three archetypes of top-performing practice sites. *Ann Fam Med* 2007;5:233–241 | | 72. | Mungmode A, Noor N, Weinstock RS, et al. Making diabetes electronic medical record data actionable: promoting benchmarking and population health improvement using the T1D Exchange Quality Improvement Portal. *Clin Diabetes* 2022;41:45–55 | | 73. | Powell RE, Zaccardi F, Beebe C, et al. Strategies for overcoming therapeutic inertia in type 2 diabetes: a systematic review and meta-analysis. *Diabetes Obes Metab* 2021;23:2137–2154 | | 74. | Herges JR, Neumiller JJ, McCoy RG. Easing the financial burden of diabetes management: a guide for patients and primary care clinicians. *Clin Diabetes* 2021;39:427–436 | | 75. | Young-Hyman D, de Groot M, Hill-Briggs F, Gonzalez JS, Hood K, Peyrot M. Psychosocial care for people with diabetes: a position statement of the American Diabetes Association. *Diabetes Care* 2016;39:2126–2140 | | 76. | Pullen-Smith B, Carter-Edwards L, Leathers KH. Community health ambassadors: a model for engaging community leaders to promote better health in North Carolina. *J Public Health Manag Pract* 2008;14(Suppl.):S73–S81 | | 77. | Neeland IJ, Arafah A, Bourges-Sevenier B, et al. Second-year results from CINEMA: a novel, patient-centered diabetes management program. *Diabetes Care* 2023;46:1762–1769 |

2026 January 29 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded

diabetesjournals.org/care Improving Care and Promoting Health in Populations S25

therapy for type 2 diabetes management: the Set (HEDIS-MY) 2024. Accessed 6 August 2025.

135. Levi R, Bleich SN, Seligman HK. Food insecurity and diabetes: overview of intersections and potential dual solutions. *Diabetes Care* 2023; 46:1599–1608

106. McCoy RG, Van Houten HK, Deng Y, et al. Comparison of diabetes medications used by adults with commercial insurance vs Medicare Advantage, 2016 to 2019. *JAMA Netw Open* 2021; 4:e2035792

107. Benning TJ, Heien HC, Herges JR, Creo AL, Al Nofal A, McCoy RG. Glucagon fill rates and cost among children and adolescents with type 1 diabetes in the United States, 2011–2021. *Diabetes Res Clin Pract* 2023; 206:111026

108. Herges JR, Haag JD, Kosloski-Tarpenning KA, Mara KC, McCoy RG. Gaps in glucagon fills among commercially insured patients receiving a glucagon prescription. *Diabetes Res Clin Pract* 2023; 200:110720

109. Patel PM, Thomas D, Liu Z, Aldrich-Renner S, Clemons M, Patel BV. Systematic review of disparities in continuous glucose monitoring and insulin pump utilization in the United States: key themes and evidentiary gaps. *Diabetes Obes Metab* 2024; 26:4293–4301
110. Eberly LA, Yang L, Essien UR, et al. Racial, ethnic, and socioeconomic inequities in glucagon-like peptide-1 receptor agonist use among patients with diabetes in the US. *JAMA Health Forum* 2021; 2:e214182
111. Essien UR, Singh B, Swabe G, et al. Association of prescription co-payment with adherence to glucagon-like peptide-1 receptor agonist and sodium-glucose cotransporter-2 inhibitor therapies in patients with heart failure and diabetes. *JAMA Netw Open* 2023; 6:e2316290
112. Taha MB, Valero-Elizondo J, Yahya T, et al. Cost-related medication nonadherence in adults with diabetes in the United States: the National Health Interview Survey 2013–2018. *Diabetes Care* 2022; 45:594–603
113. Centers for Medicare & Medicaid Services. CMS Framework for Health Equity 2022–2032. Accessed 8 August 2025. Available from <https://www.cms.gov/files/document/cms-framework-health-equity.pdf>
114. Washington AE, Lipstein SH. The Patient-Centered Outcomes Research Institute—promoting better information, decisions, and health. *N Engl J Med* 2011; 365:e31
115. Dixon B, Pena M-M, Taveras EM. Lifecourse approach to racial/ethnic disparities in childhood obesity. *Adv Nutr* 2012; 3:73–82
116. Egede LE, Walker RJ, Linde S, et al. Nonmedical interventions for type 2 diabetes: evidence, actionable strategies, and policy opportunities. *Health Aff (Millwood)* 2022; 41:963–970
117. Jiang DH, O'Connor PJ, Huguet N, Golden SH, McCoy RG. Modernizing diabetes care quality measures. *Health Aff (Millwood)* 2022; 41:955–962
118. Joint Commission. R3 Report. Requirement, Rationale, Reference. Accessed 11 September 2025. Available from https://digitalassets.jointcommission.org/api/public/content/e57b92c0e6854a03b27302663b5c1084?v=cec3044d&_gl=1*1xqrmfh*_ga*Lg*_ga_K31T0BHP4T*cze3NTQ0OTIwNDYkbzEkZzEkdDE3NTQ0OTIxNTUkajU5JGwwJGgw
119. National Committee for Quality Assurance. Healthcare Effectiveness Data and Information Set (HEDIS-MY) 2024. Accessed 6 August 2025. Available from <https://www.ncqa.org/wp-content/uploads/HEDIS-MY-2024-Measure-Description.pdf>
120. Centers for Medicare & Medicaid Services. Quality Payment Program. Explore Measures & Activities. Accessed 6 August 2025. Available from <https://qpp.cms.gov/mips/explore-measures>
121. Ebekozien O, Mungmode A, Buckingham D, et al. Achieving equity in diabetes research: borrowing from the field of quality improvement using a practical framework and improvement tools. *Diabetes Spectr* 2022; 35:304–312
122. Odugbesan O, Addala A, Nelson G, et al. Implicit racial-ethnic and insurance-mediated bias to recommending diabetes technology: insights from T1D Exchange Multicenter Pediatric and Adult Diabetes Provider Cohort. *Diabetes Technol Ther* 2022; 24:619–627
123. Centers for Medicare & Medicaid Services. Ensuring Proper Use of Electronic Health Record Features and Capabilities. 2016. Accessed 6 August 2025. Available from <https://www.cms.gov/files/document/ehrddecisiontable062816pdf>
124. U.S. Department of Health and Human Services. Secretary's Advisory Committee on National Health Promotion and Disease Prevention Objectives for 2030. Accessed 6 August 2025. Available from <https://health.gov/healthypeople>
125. National Academy of Sciences. A Framework for Educating Health Professionals to Address the Social Determinants of Health. 2016. Accessed 6 August 2025. Available from <https://www.ncbi.nlm.nih.gov/pubmed/27854400>

126. Zakaria NI, Tehranifar P, LaFerrere B, Albrecht SS. Racial and ethnic disparities in glycemic control among insured US adults. *JAMA Netw Open* 2023; 6:e2336307
 127. Hershey JA, Morone J, Lipman TH, Hawkes CP. Social determinants of health, goals and outcomes in high-risk children with type 1 diabetes. *Can J Diabetes* 2021; 45:444–450.e1
 128. Walker RJ, Strom Williams J, Egede LE. Influence of race, ethnicity and social determinants of health on diabetes outcomes. *Am J Med Sci* 2016; 351:366–373
 129. Hashemi R, Rabizadeh S, Yadegar A, et al. High prevalence of comorbidities in older adult patients with type 2 diabetes: a cross-sectional survey. *BMC Geriatr* 2024; 24:873
 130. Huang ES. Management of diabetes mellitus in older people with comorbidities. *BMJ* 2016; 353:i2200
 131. O'Gurek DT, Henke C. A practical approach to screening for social determinants of health. *Fam Pract Manag* 2018; 25:7–12
 132. Yan AF, Chen Z, Wang Y, et al. Effectiveness of social needs screening and interventions in clinical settings on utilization, cost, and clinical outcomes: a systematic review. *Health Equity* 2022; 6:454–475
 133. Rabbitt MP, Hales LJ, Burke MP, Coleman-Jensen A. Household Food Security in the United States in 2022 (Report No. ERR-325). U.S. Department of Agriculture, Economic Research Service, 2023. Accessed 11 September 2025. Available from https://search.nal.usda.gov/discovery/fulldisplay?context=L&vid=01NAL_INST:MAIN&docid=alma-9916411232407426
 134. Kirby JB, Bernard D, Liang L. The prevalence of food insecurity is highest among Americans for whom diet is most critical to health. *Diabetes Care* 2021; 44:e131–e132
- 2026 January 29 on guest by
http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf from Downloaded
- # Improving Care and Promoting Health in Populations
- # Diabetes Care Volume 49, Supplement 1, January 2026
150. Soto S, Yoder AM, Aceves B, Nuño T, Sepulveda R, Rosales CB. Determining regional differences in barriers to accessing health care among farmworkers using the National Agricultural Workers Survey. *J Immigr Minor Health* 2023;25:324–330
 151. Thoñón F, Perrot S, Yergolkar AV, et al. Electronic tools to bridge the language gap in health care for people who have migrated: systematic review. *J Med Internet Res* 2021;23:e25131
 152. U.S. Department of Health & Human Services. National Standards for Culturally and Linguistically Appropriate Services (CLAS) in Health and Health Care. Accessed 6 August 2025. Available from https://thinkculturalhealth.hhs.gov/assets/pdfs/EnhancedNationalCLASStandards_Revised_June2025.pdf
 153. Centers for Disease Control and Prevention. Health Literacy. Talking Points About Health Literacy. Accessed 06 August 2025. Available from <https://www.cdc.gov/health-literacy/php/about/tell-others.html>
 154. Nutbeam D, Lloyd JE. Understanding and responding to health literacy as a social determinant of health. *Annu Rev Public Health* 2021;42:159–173
 155. Marciaño L, Camerini A-L, Schulz PJ. The role of health literacy in diabetes knowledge, self-care, and glycemic control: a meta-analysis. *J Gen Intern Med* 2019;34:1007–1017
 156. Schaffler J, Leung K, Tremblay S, et al. The effectiveness of self-management interventions for individuals with low health literacy and/or low income: a descriptive systematic review. *J Gen Intern Med* 2018;33:510–523
 157. Butayeva J, Ratan ZA, Downie S, Hosseinzadeh H. The impact of health literacy interventions on glycemic control in patients with type 2 diabetes: a systematic review and meta-analysis. *Lancet* 2023;402(Suppl 1):S40
 158. Baumeister A, Aldin A, Chakraverty D, et al. Interventions for improving health literacy in migrants. *Cochrane Database Syst Rev* 2023;11:CD013303

159. Schapira MM, Fletcher KE, Gilligan MA, et al. A framework for health numeracy: how patients use quantitative skills in health care. *J Health Commun* 2008;13:501–517
160. Agency for Healthcare Research and Quality. Clinical-community linkages. Accessed 6 August 2025. Available from <https://archive.ahrq.gov/ncepcr/tools/community/index.html>
161. Herges JR, Matulis JC, Kessler ME, Ruehmann LL, Mara KC, McCoy RG. Evaluation of an enhanced primary care team model to improve diabetes care. *Ann Fam Med* 2022;20:505–511
162. Kasper AL, Myers LA, Carlson PN, et al. Diabetes management for community paramedics: development and implementation of a novel curriculum. *Diabetes Spectr* 2022;35:367–376
163. Evans J, White P, Ha H. Evaluating the effectiveness of community health worker interventions on glycaemic control in type 2 diabetes: a systematic review and meta-analysis. *Lancet* 2023;402(Suppl 1):S40
164. Ducharme-Smith A, Junttunen M, Espinoza A, et al. Pilot pragmatic trial of a community paramedic diabetes self-management education program for adults with diabetes. *Diabetes Care* 2025;48:2089–2094
165. Azmiardi A, Murti B, Febriñasari RP, Tamtomo DG. The effect of peer support in diabetes self-management education on glycemic control in patients with type 2 diabetes: a systematic review and meta-analysis. *Lancet* 2023;402(Suppl 1):S40
166. Fisher EB, Boothroyd RI, Elstad EA, et al. Peer support of complex health behaviors in prevention and disease management with special reference to diabetes: systematic reviews. *Clin Diabetes Endocrinol* 2017;3:4
167. Foster G, Taylor SJC, Eldridge SE, Ramsay J, Griffiths CJ. Self-management education programmes by lay leaders for people with chronic conditions. *Cochrane Database Syst Rev* 2007: CD005108
168. Piatt GA, Rodgers EA, Xue L, Zgibor JC. Integration and utilization of peer leaders for diabetes self-management support: results from Project SEED (Support, Education, and Evaluation in Diabetes). *Diabetes Educ* 2018;44:373–382
169. Rosenthal EL, Rush CH, Allen CG. Understanding scope and competencies: a contemporary look at the United States community health worker field: progress report of the Community Health Worker (CHW) Core Consensus (C3) Project: building national consensus on CHW core roles, skills, and qualities. CHW Central, 2016. Accessed 6 August 2025. Available from <https://files.ctctcdn.com/a907c850501/1c1289f0-88cc-49c3-a238-66def942c147.pdf>
170. Guide to Community Preventive Services. Community health workers help patients manage diabetes. Updated 25 October 2022. Accessed 06 August 2025. Available from <https://www.thecommunityguide.org/content/community-health-workers-help-patients-manage-diabetes>
171. Cuellar AE, Calonge BN. The Community Preventive Services Task Force: 25 years of effectiveness, economics, and equity. *Am J Prev Med* 2022;62:e371–e373
- 2026 January 29 on guest by
http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S13/848832/dc26s001.pdf